

UCLA

UCLA Previously Published Works

Title

Ventricular Tachycardia Storm in a Young Adult Post-ALCAPA Repair.

Permalink

<https://escholarship.org/uc/item/2tx405j1>

Journal

JACC: Case Reports, 29(16)

Authors

Jeong, Sun

Hayase, Justin

Moore, Jeremy

et al.

Publication Date

2024-08-21

DOI

10.1016/j.jaccas.2024.102418

Peer reviewed

MINI-FOCUS ISSUE ON CONGENITAL HEART DISEASE

CASE REPORT: CLINICAL CASE

Ventricular Tachycardia Storm in a Young Adult Post-ALCAPA Repair



Sun Young Jeong, MD, MPH,^a Justin Hayase, MD,^b Jeremy Moore, MD,^b Aron Bender, MD,^b Duc Do, MD,^b Daniel R. Sanchez, MD,^a Jonathan M. Doris, MD^c

ABSTRACT

Anomalous left coronary artery to pulmonary artery (ALCAPA) is a leading cause of pediatric myocardial ischemia. This paper presents a case of a young man presenting with ventricular tachycardia storm 18 years after ALCAPA repair. Clinicians should recognize the risk of ventricular tachycardia in this patient population. (JACC Case Rep 2024;29:102418)
© 2024 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

HISTORY OF PRESENTATION

An 18-year-old man with a remote history of anomalous left coronary artery to pulmonary artery (ALCAPA) repair in infancy presented to the emergency department (ED) for recurrent palpitations. His heart rate was 200 beats/min with blood pressure of 73/64 mm Hg. He was diaphoretic and tachycardic on

examination. Laboratory data were notable for high-sensitivity cardiac troponin I level of 228 pg/mL (normal ≤ 20 pg/mL). Electrocardiogram (ECG) showed wide complex tachycardia (WCT) with an atypical right bundle branch block (RBBB) and inferior axis at a rate of 200 beats/min, consistent with monomorphic ventricular tachycardia (MMVT) (Figure 1A). The patient was electrically converted to sinus rhythm and admitted to the cardiac care unit (CCU). An implantable cardioverter-defibrillator (ICD) was placed, and he was discharged on metoprolol succinate. Less than 48 hours later, he returned to the ED with ventricular tachycardia (VT) storm with multiple ICD shocks and was readmitted to the CCU.

LEARNING OBJECTIVES

- To recognize the risk of VT in adults despite early and successful ALCAPA repair in infancy.
- To distinguish VT from other forms of wide complex tachycardia.
- To formulate an appropriate surveillance protocol to mitigate the risk of sudden cardiac death in this high-risk patient population.

PAST MEDICAL HISTORY

As a neonate, the patient developed recurrent respiratory distress, ultimately leading to the diagnosis of

From the ^aDepartment of Cardiology, Kaiser Permanente Los Angeles Medical Center, Los Angeles, California, USA; ^bCardiac Arrhythmia Center, University of California-Los Angeles, Los Angeles, California, USA; and the ^cDepartment of Cardiac Electrophysiology, Kaiser Permanente Los Angeles Medical Center, Los Angeles, California, USA.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

Manuscript received March 28, 2024; revised manuscript received May 9, 2024, accepted May 15, 2024.

**ABBREVIATIONS
AND ACRONYMS**

AAD	= antiarrhythmic drug
ALCAPA	= anomalous left coronary artery to pulmonary artery
CCU	= cardiac care unit
ECG	= electrocardiogram
ED	= emergency department
ICD	= implantable cardioverter-defibrillator
LV	= left ventricular
MMVT	= monomorphic ventricular tachycardia
RBBB	= right bundle branch block
SVT	= supraventricular tachycardia
VT	= ventricular tachycardia
WCT	= wide complex tachycardia

dilated cardiomyopathy and ALCAPA at 5 weeks of age. He underwent repair via coronary reimplantation and was noted to have a large anterolateral myocardial infarction intraoperatively. The patient was followed by pediatric cardiology growing up, with regular echocardiographic surveillance and Holter monitoring. Left ventricular (LV) systolic function was normal throughout this time.

Nine months prior to the current episode, he presented to the ED with chest pain, and was found to have a non-ST elevation myocardial infarction with a peak high-sensitivity cardiac troponin I level of 2,084 pg/mL (normal ≤ 20 pg/mL). ECG showed normal sinus rhythm with right axis deviation and incomplete RBBB (Figure 1B). Cardiac catheterization showed no anastomotic strictures, kinking, or obstruction of the coronary artery but mildly reduced LV systolic function. He was discharged on guideline-directed medical therapy, including carvedilol.

This event was followed by multiple instances of palpitations over the subsequent months. ECGs were interpreted as sinus tachycardia with RBBB or as supraventricular tachycardia (SVT) with aberrancy in the 190s (tracings not available), requiring cardioversions in the field by paramedics. He was discharged each time after an unremarkable evaluation from the ED.

DIFFERENTIAL DIAGNOSIS

The differential diagnosis for a young patient presenting with WCT includes VT, SVT with aberrancy, SVT with antegrade conduction via an accessory pathway, and antidromic atrioventricular reentrant tachycardia.

INVESTIGATIONS

Cardiac magnetic resonance performed prior to ICD placement revealed LV ejection fraction of 42% and late gadolinium enhancement in the anterolateral LV wall (Figure 2). After presenting with VT storm, his ICD was interrogated and showed 26 appropriate shocks for VT.

MANAGEMENT


After readmission to the hospital for VT storm, the patient required amiodarone and lidocaine infusions, intubation, and sedation. He underwent endocardial VT ablation. No VT could be induced; however, electroanatomic maps showed scar along the anterolateral left ventricle with local abnormal ventricular activities and decrement-evoked potentials along the basal aspect of the scar (Figures 3 and 4). These areas were targeted for ablation with extensive substrate modification. Postprocedure, the patient was extubated, and intravenous antiarrhythmic drugs (AADs) were discontinued. He was discharged on mexiletine and carvedilol.

DISCUSSION

ALCAPA is a congenital heart defect wherein the left coronary artery originates from the pulmonary trunk instead of the aorta.¹ Without treatment, 90% of infants die in the first year of life. There are a scarcity of data on long-term survival outcomes after pediatric ALCAPA repair. However, various cohort studies show that ALCAPA has a favorable prognosis for mortality with early repair, with survival ranging from 86% to 100% at 10 years.² Late sequelae postrepair includes mitral regurgitation, ischemic cardiomyopathy, scar-mediated arrhythmias, and sudden cardiac death. Life-long monitoring is recommended due to the risk of late complications, which remains largely unquantified by large-scale studies.³

This case highlights the importance of the past medical history in risk stratification. He had a late ALCAPA repair as demonstrated by the presence of a large anterolateral myocardial infarction during the original surgery (and persistent scar in the same territory by cardiac magnetic resonance 18 years later). Because myocardial scar is a well-recognized substrate for arrhythmogenic sequelae and sudden cardiac death, lifelong surveillance was warranted. Palpitations in this patient should have been aggressively evaluated, particularly after a non-ST-segment elevation myocardial infarction. His multiple presentations of WCT, with associated hemodynamic instability, were unfortunately dismissed as benign due to his young age, a form of anchoring

FIGURE 1 Electrocardiograms




(A) Monomorphic ventricular tachycardia with atypical right bundle branch block, right inferior axis, and atrioventricular dissociation (best seen in V₁), with initial dominant R in aVR, monophasic R waves in V₁ and aVR, and rS in V₂ to V₆. (B) Sinus rhythm and incomplete right bundle branch block.

bias, and his baseline RBBB. Had his past medical history been taken into more careful consideration, the diagnosis of VT could have been made much sooner.

His past medical history aside, the ECG demonstrating WCT is most consistent with VT. Several


features of the ECG help distinguish between VT and SVT with aberrancy in the setting of a WCT. In this case, factors favoring VT includes regular rate, wide QRS interval >140 milliseconds in an atypical RBBB, right inferior axis, and atrioventricular dissociation, with initial dominant R in

FIGURE 2 Cardiac Magnetic Resonance


Late gadolinium enhancement is seen in the anterolateral wall of the left ventricle (yellow arrow).

aVR, monophasic R waves in V_1 and aVR, and rS in V_2 to V_6 .

Per the American College of Cardiology/American Heart Association guidelines, the initial management of sustained MMVT in patients with structural heart disease includes direct current cardioversion (Class I) and intravenous AADs such as intravenous procainamide (Class IIa), amiodarone, or sotalol (Class IIb), along with sedation and anesthesia.⁴ VT occurring after coronary surgery warrants imaging to exclude any anatomic abnormalities (eg, anastomotic stricture, kinking of the vessel). In this case, this was accomplished with coronary angiography and cardiac computed tomography. Management of VT in young patients can be particularly challenging because the toxicity associated with many AADs is not ideal for this age group. VT refractory to medications and antitachycardia pacing has a Class I recommendation for catheter ablation. We favored an aggressive approach in this young patient in VT storm with substantial myocardial scar, opting for

FIGURE 3 Electroanatomic Voltage Map

Voltage maps (NavX) during sinus rhythm at standard scar settings (0.5-1.5 mV) with AP (left) and LAO cranial (right) views showing areas of low voltage along basal superior aspect and along the entire anterolateral left ventricular wall. AP = anterior-posterior; LAO = left anterior oblique.



ablation during his second admission to the CCU in lieu of trialing him on AADs.

FOLLOW-UP

At 1-month follow-up, the patient had SVT but no VT recurrence as confirmed by ICD interrogations. Cardiac computed tomography showed no stricture, kinking, or obstruction of the reimplanted left coronary artery with normalized LV ejection fraction (Figure 5).

CONCLUSIONS

We describe a case of a young adult man with a history of ALCAPA repair presenting with VT storm

mediated by LV scar 18 years after his original surgery. Ischemic insults and resultant myocardial scars sustained as an infant became the substrate for VT. Patients with a history of repaired congenital heart disease who present with WCT warrant a high index of suspicion for VT despite their age. Managing VT in young patients can be difficult due to the long-term adverse effects of many AADs. When VT is resistant to medical therapy, catheter ablation becomes an important option. There is currently a lack of consensus regarding the optimal surveillance strategy for VT in young patients post-ALCAPA repair, suggesting a need for the inclusion of ALCAPA surveillance strategies in future expert consensus guidelines.

FIGURE 5 Cardiac Computed Tomography

There was no kinking or strictures of the reimplanted left coronary artery.

FUNDING SUPPORT AND AUTHOR DISCLOSURES

The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

ADDRESS FOR CORRESPONDENCE: Dr Sun Young Jeong, Department of Cardiology, Kaiser Permanente Los Angeles Medical Center, 1526 North Edgemont Street, Los Angeles, California 90027, USA. E-mail: sunyoung.jeong@kp.org. X handle: [@SunnyJeongMD](https://twitter.com/SunnyJeongMD).

REFERENCES

1. Azakie A. Repair of anomalous coronary artery from the pulmonary artery by aortic implantation. *Oper Tech Thorac Cardiovasc Surg*. 2015;20(2):135-147.
2. Naimo PS, Fricke TA, d'Udekem Y, et al. Surgical intervention for anomalous origin of left coronary artery from the pulmonary artery in children: a long-term follow-up. *Ann Thorac Surg*. 2016;101(5):1842-1848.
3. Kanoh M, Inai K, Shinohara T, Tomimatsu H, Nakanishi T. Outcomes from anomalous origin of the left coronary artery from the pulmonary artery repair: long-term complications in relation to residual myocardial abnormalities. *J Cardiol*. 2017;70(5):498-503.
4. Al-Khatib SM, Stevenson WG, Ackerman MJ, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. 2018;72(14):e91-e220. <https://doi.org/10.1016/j.jacc.2017.10.054>

KEY WORDS congenital heart disease, myocardial ischemia, ventricular arrhythmia