

# UCLA

## UCLA Previously Published Works

### Title

Genomic signatures of host-associated divergence and adaptation in a coral-eating snail, *Coralliophila violacea* (Kiener, 1836)

### Permalink

<https://escholarship.org/uc/item/2vg5526v>

### Journal

Ecology and Evolution, 10(4)

### ISSN

2045-7758

### Authors

Simmonds, Sara E  
Fritts-Penniman, Allison L  
Cheng, Samantha H  
et al.

### Publication Date

2020-02-01

### DOI

10.1002/ece3.5977

Peer reviewed

## ORIGINAL RESEARCH

# Genomic signatures of host-associated divergence and adaptation in a coral-eating snail, *Coralliophila violacea* (Kiener, 1836)

Sara E. Simmonds<sup>1</sup>  | Allison L. Fritts-Penniman<sup>1</sup> | Samantha H. Cheng<sup>1,2</sup> | Gusti Ngurah Mahardika<sup>3</sup> | Paul H. Barber<sup>1</sup>

<sup>1</sup>Department of Ecology and Evolutionary Biology, University of California Los Angeles, Los Angeles, CA, USA

<sup>2</sup>Center for Biodiversity and Conservation, American Museum of Natural History, New York, NY, USA

<sup>3</sup>Animal Biomedical and Molecular Biology Laboratory, Faculty of Veterinary Medicine, Udayana University Bali, Denpasar, Indonesia

**Correspondence**

Department of Biological Sciences, Boise State University, Boise, ID 83725, USA.  
Email: sarasimmonds@boisestate.edu

**Funding information**

Sigma Xi; University of California, Los Angeles; Office of International Science and Engineering, Grant/Award Number: OISE-0730256 and OISE-1243541; Division of Ocean Sciences, Grant/Award Number: OCE-0349177; United States Agency for International Development, Grant/Award Number: 497-A-00-10-00008-00

**Abstract**

The fluid nature of the ocean, combined with planktonic dispersal of marine larvae, lowers physical barriers to gene flow. However, divergence can still occur despite gene flow if strong selection acts on populations occupying different ecological niches. Here, we examined the population genomics of an ectoparasitic snail, *Coralliophila violacea* (Kiener 1836), that specializes on *Porites* corals in the Indo-Pacific. Previous genetic analyses revealed two sympatric lineages associated with different coral hosts. In this study, we examined the mechanisms promoting and maintaining the snails' adaptation to their coral hosts. Genome-wide single nucleotide polymorphism (SNP) data from type II restriction site-associated DNA (2b-RAD) sequencing revealed two differentiated clusters of *C. violacea* that were largely concordant with coral host, consistent with previous genetic results. However, the presence of some admixed genotypes indicates gene flow from one lineage to the other. Combined, these results suggest that differentiation between host-associated lineages of *C. violacea* is occurring in the face of ongoing gene flow, requiring strong selection. Indeed, 2.7% of all SNP loci were outlier loci (73/2,718), indicative of divergence with gene flow, driven by adaptation of each *C. violacea* lineage to their specific coral hosts.

**KEYWORDS**

adaptation, coral reefs, ecological divergence, gastropods, population genomics, RAD-seq

## 1 | INTRODUCTION

While ecological speciation has been documented for almost three decades across a wide variety of organisms on land (Case & Willis, 2008; Feder et al., 1994; Jiggins, 2008; Martin et al., 2013; Schluter, 2009; Seehausen et al., 2008; Sorenson, Sefc, & Payne, 2003; Thorpe, Surget-Groba, & Johansson, 2010; Waser

& Campbell, 2004) and in freshwater (Hatfield & Schluter, 1999; Langerhans, Gifford, & Joseph, 2007; Puebla, 2009; Seehausen et al., 2008; Seehausen & Wagner, 2014), ecological speciation in the ocean was thought to be rare, and only recently has that viewpoint begun to change (Bird, Fernandez-Silva, Skillings, & Toonen, 2012; Bird, Holland, Bowen, & Toonen, 2011; Bowen, Rocha, Toonen, Karl, & ToBo Laboratory, 2013; Foote & Morin,

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.


2015; Hurt, Silliman, Anker, & Knowlton, 2013; Ingram, 2010; Litsios et al., 2012; Rocha, Robertson, Roman, & Bowen, 2005). There are a number of reasons for this reassessment. First, absolute physical barriers in the sea are exceedingly rare (Ludt & Rocha, 2015; Rocha & Bowen, 2008; Rocha et al., 2005). As a result, speciation must often proceed with varying levels of gene flow and aided by divergent selection in different habitats or hosts (Palumbi, 1994). Second, the strong interspecific interactions that can promote ecological speciation in terrestrial species (e.g., host–parasite, mutualisms) are also common in certain marine ecosystems (Blackall, Wilson, & van Oppen, 2015; Stella, Jones, & Pratchett, 2010). For example, reef-building corals have tight ecological associations with a wide variety of invertebrate taxa (Zann, 1987), including ~900 named species of sponges, copepods, barnacles, crabs, shrimp, worms, bivalves, nudibranchs, and snails (reviewed by Stella et al., 2010). This wide array of symbiotic relationships creates tremendous potential for host shifting and the development of host specificity that can lead to sympatric speciation.

Evidence from traditional genetic markers (i.e., microsatellites, RFLPs, allozymes, nuclear, mitochondrial, and ribosomal genes) demonstrates the potential for ecological speciation in marine taxa exhibiting symbiotic relationships (Bowen et al., 2013; Miglietta, Faucci, & Santini, 2011; Peijnenburg & Goetze, 2013; Potkamp & Fransen, 2019), including amphipods on macroalgae (Sotka, 2005), coral-dwelling barnacles (Tsang, Chan, Shih, Chu, & Allen Chen, 2009), coral-eating nudibranchs (Faucci, Toonen, & Hadfield, 2007; Fritts-Penniman, Gosliner, Mahardika, & Barber, 2020), parasitic snails (Gittenberger & Gittenberger, 2011; Reijnen, Hoeksema, & Gittenberger, 2010), anemone-associated shrimp (Hurt et al., 2013), anemone fish (Litsios et al., 2012), and coral-dwelling gobies (Duchene, Klanten, Munday, Herler, & van Herwerden, 2013; Munday, van Herwerden, & Dudgeon, 2004).

While encouraging, there are gaps in our knowledge that with the expansion of genomic technologies, we are now in a position to begin to fill. Detecting signatures of natural selection in populations where there is likely ongoing gene flow is now possible using genome-wide data, lending insight into the mechanisms of ecological speciation (Bernal, Gaither, Simison, & Rocha, 2017; Campbell, Poelstra, & Yoder, 2018; Puebla, Bermingham, & McMillan, 2014; Westram et al., 2018). To date, however, no studies examining the genomic signatures of ecological divergence in marine host–parasite systems have been conducted.

The ~6 million km<sup>2</sup> Coral Triangle region is home to over 500 species of reef-building corals (Veron et al., 2011) and thousands of unique species of fishes and invertebrates (Barber & Boyce, 2006; Briggs, 2003), making it the global center of marine biodiversity (Cowman & Bellwood, 2011; Hoeksema, 2007). Most of the literature examining the evolution of this biodiversity hotspot has focused on allopatric processes such as divergence across geological and oceanographic features such as the Sunda Shelf or Halmahera Eddy during Pleistocene low sea levels stands (for reviews, see Barber, Cheng, Erdmann, Tenggardjaja, & Ambariyanto 2011; Carpenter et al., 2011; Gaither & Rocha, 2013). Allopatric divergence is clearly an important factor in the biodiversity of the Coral Triangle. However, the extraordinary diversity in this region, combined with the prevalence of strong species–species interactions on coral reefs, makes it likely that ecological speciation also contributes to the evolution of biodiversity in this hotspot.

The marine snail, *Coralliophila violacea* (Figure 1), is an obligate ectoparasite, living, feeding, and reproducing exclusively on corals in Poritidae, a highly abundant and diverse coral family (Kitahara, Cairns, Stolarski, Blair, & Miller, 2010), which is found in shallow reefs across the tropical Indo-Pacific. The snails attach themselves to their host, form feeding aggregations, and drain energy from their host as it tries to repair damaged tissues (Oren, Brickner, &



**FIGURE 1** Violet coral snails, (a) *Coralliophila violacea* (Kiener, 1836), are obligate ectoparasites of corals in the family Poritidae. Their shells are usually fouled with crustose coralline algae because of their sedentary lifestyle, making them difficult to spot on their host corals. They are commonly found living among the branches of species such as (b) *Porites cylindrica* and can form aggregations on massive coral species like (c) *P. lobata*. (Photos by S.E. Simmonds)


**TABLE 1** *Coralliophila violacea* collection locations, latitude, longitude, coral host species, and number of samples collected

Location	Country	Province	Latitude	Longitude	Coral host species	
					<i>Porites lobata</i>	<i>Porites cylindrica</i>
1. Pemuteran	Indonesia	Bali	-8.1400	114.6540	-	7
2. Nusa Penida	Indonesia	Bali	-8.6750	115.5130	11	10
3. Pulau Mengyatan	Indonesia	East Nusa Tenggara	-8.5570	119.6850	4	3
4. Lembeh	Indonesia	North Sulawesi	1.4790	125.2510	7	1
5. Bunaken	Indonesia	North Sulawesi	1.6120	124.7830	9	6
6. Dumaguete	Philippines	Negros Oriental	9.3320	123.3120	2	7
Total N					33	34

Loya, 1998). They are sequential hermaphrodites, a common trait of parasitic mollusks (Heller, 1993), and breed with conspecifics on their host coral colony. Two genetically distinct lineages of *C. violacea* occur sympatrically on reefs of the Coral Triangle, but each lineage occupies one of two groups of *Porites* corals, suggesting ecological divergence (Simmonds et al., 2018). A lack of evidence of genetic structure within each lineage of *C. violacea* inside the Coral Triangle precludes physical isolation as an explanation for the observed divergence. Host specificity commonly results from preferential larval settlement (Ritson-Williams, Shjegstad, & Paul, 2003, 2007, 2009). This genetic evidence combined with observations of adult preference for specific coral hosts (unpubl. data S.

Simmonds) strongly suggests ecological divergence driven by host association.

To determine where diverging populations of *C. violacea* lie on the continuum of the speciation process (i.e., host-associated lineages, sibling species or good species), it is important to examine patterns of realized gene flow between the divergent coral host-associated lineages. Effective contemporary gene flow should result in linkage disequilibria between host-associated marker loci in populations utilizing different hosts. However, if lower rates of gene flow (<1% per generation) are found, then populations should be considered incipient species (Drès & Mallet, 2002; Malaua et al., 2007).

**FIGURE 2** Collection locations for *Coralliophila violacea* from coral host species *Porites lobata* and *P. cylindrica*. 1. Pemuteran, 2. Nusa Penida, 3. Pulau Mengyatan, 4. Lembeh, 5. Bunaken, 6. Dumaguete. Map made with vector and raster map data available at [naturalearthdata.com](http://naturalearthdata.com)

Genomic tests of selection are key to distinguishing between these possibilities. If divergence among *C. violacea* lineages results purely from neutral processes, genetic drift and migration should have approximately equal effects on all parts of the genome (Nielsen, 2005), and frequencies of neutral loci should show similar levels of differentiation (Via, 2009). However, if divergent selection is driving diversification of *C. violacea* lineages, there should be clear signatures of divergent selection (Feder et al., 1994; Nasil, Funk, & Ortiz-Barrientos, 2009), because natural selection affects non-neutral parts of the genome, as well as linked loci, to a greater extent (Smith & Haigh, 1974). As such, frequencies of loci under selection (outlier loci) or linked loci should either be unusually high or unusually low, in host-associated populations, depending on the type of selection occurring (Beaumont & Nichols, 1996).

In this study, we use genome-wide single nucleotide polymorphisms (SNPs) to investigate the possibility of ecological divergence with gene flow in populations of a corallivorous gastropod, *C. violacea*, from the Coral Triangle. Specifically, we (a) test for reduced gene flow between sympatric lineages of host-associated snails, (b) identify outlier loci under putative selection between hosts, and (c) annotate possible functions of linked genes that might be necessary for adaptation to hosts.

## 2 | MATERIALS AND METHODS

### 2.1 | Sample collection

We collected snails on snorkel during 2011–2013 from six sympatric populations of two lineages of *C. violacea* representing unique parasite–host groups (Table 1, Figure 2, Appendix S1). We chose snails from the most abundant *Porites* species from each group (*P. lobata*, *P. cylindrica*, Dana, 1846, Figure 1) to maximize the number of samples and reduce potentially confounding effects of differences among hosts within the same group. To further reduce confounding effects resulting from taxonomic complexity within *P. lobata* (Forsman, Barshis, Hunter, & Toonen, 2009; Prada et al., 2014), we used coral host species identifications from Simmonds et al. (2018) that were confirmed through RAD-seq data.

### 2.2 | Creation of RAD libraries

We extracted genomic DNA from 20 mg of foot tissue from 67 individual *C. violacea* (34 from *P. cylindrica* and 33 from *P. lobata*; Table 1) using a DNeasy® Blood and Tissue Kit (QIAGEN), following manufacturer's instructions, save for elution of DNA with molecular grade H<sub>2</sub>O rather than AE buffer. We estimated initial DNA concentrations using a NanoDrop™ 2000 Spectrophotometer (Thermo Scientific™) and visualized DNA quality on a 1% agarose gel stained with SYBR® Safe DNA Gel Stain (Invitrogen™). We used only high-quality DNA with a bright high molecular weight

band and minimal smearing. We dried DNA extractions using a SpeedVac™ (Thermo Scientific™) on medium heat and reconstituted using molecular grade H<sub>2</sub>O to a final uniform 250 ng/μl DNA concentration.

We created reduced representation libraries to survey SNP variation following published protocols (Wang, Meyer, McKay, & Matz, 2012) as updated by Dr. Eli Meyer (<http://people.oregonstate.edu/~meyere/docs/Preparing2bRAD.pdf>). Alfl restriction enzyme digest reduced representation (1/16th) libraries were labeled with individual barcodes and subjected to 18–20 PCR amplification cycles. The number of PCR cycles varied based on the optimal number determined in the test-scale PCR to find the minimum number of cycles to produce a visible product at 166 bp. We electrophoresed products on a 2% agarose gel in 1 × TBE buffer and ran at 150 V for 90 min, visualized target bands (165 bp) with SYBR® Safe DNA Gel Stain (Invitrogen™), and excised them from the gel. Then, we purified the excised bands using a QIAquick® Gel Extraction Kit (QIAGEN). A final cleaning step used Agencourt® AMPure® XP beads (Beckman Coulter). QB3 Genomics at the University of California, Berkeley performed quality checks (qPCR, BioAnalyzer) and sequencing, multiplexing 10–20 snails per lane in 5 lanes of a 50 bp Single-End run on an Illumina HiSeq 2000 sequencer.

### 2.3 | RAD-seq data processing

To prepare raw sequence data for SNP identification, we truncated all raw sequence reads to the insert size (36 bp), filtered for quality (PHRED scores >20), and discarded empty constructs. We then processed the resulting data using custom scripts written by Misha Matz, available on the 2bRAD GitHub site ([https://github.com/z0on/2bRAD\\_denovo](https://github.com/z0on/2bRAD_denovo)). First, we counted unique tag sequences (minimum sequencing depth 5×) and the number of sequences in reverse-complement orientation and then merged these tags into one table. Then, we clustered all sequences in CD-HIT (Fu, Niu, Zhu, Wu, & Li, 2012) using a 91% similarity threshold. Next, we defined the most abundant sequence in the cluster as a reference sequence and then filtered a locus-annotated table from the previous two steps, excluding reads below 5× depth and those exhibiting strand bias. Lastly, we flipped the orientation of the resulting clustered sequences to match the most abundant tag in a cluster.

To call genotypes (as population-wide RAD-tag haplotypes), we used GATK (McKenna et al., 2010) and applied mild allele filters (10× total depth, allele bias cutoff 10, and strand bias cutoff 10), with the additional requirement that alleles appear in at least two individuals. We then applied locus filters allowing a maximum of 50% heterozygotes at a locus, no more than two alleles, genotyped in 30% of samples and polymorphic. Finally, we removed loci with the fraction of heterozygotes >75% (potential lumped paralogs) and missing >70% of genotypes. The final set of SNPs was then thinned to one per tag (that with the highest minor allele frequency) for  $F_{ST}$  and STRUCTURE analysis to remove linked loci.

## 2.4 | Individual sample filtering steps

To maximize the quality of the final dataset, we further filtered out individuals ( $N = 11$ ) with low genotyping rates, indicating low DNA quality, by taking the  $\log_{10}$  of the number of sites genotyped per individual, and removing any individuals that were outside one standard deviation ( $SD$ ) of the mean. We used VCFtools (Danecek et al., 2011) to estimate inbreeding coefficients and removed individuals ( $N = 5$ ) with inbreeding coefficients outside the normal range ( $\pm 2 SD$  of mean  $F$ ) indicating possible low coverage sequencing or lumped paralogs ([https://github.com/z0on/2bRAD\\_de-novo](https://github.com/z0on/2bRAD_de-novo)). The remaining 51 individuals were used in analyses of population genetic structure. The final data file was in VCF format and converted to other formats using PGDSpider v2.0.8.0 (Lischer & Excoffier, 2012).


## 2.5 | Genetic structure

To test whether the patterns observed in a mitochondrial locus were present in loci genome-wide, we inferred the population genetic structure of the full RAD-seq dataset (2,718 loci), outlier loci only (73 loci), and neutral loci only (2,645 loci), from 51 individuals using two methods. First, we ran the Bayesian model-based clustering method STRUCTURE (Pritchard, Stephens, & Donnelly, 2000) using a burn-in period of 20,000 followed by 50,000 MCMC replicates for  $K = 1-12$ , and 10 runs for each  $K$ . We used the admixture model, with allele frequencies correlated among populations. The results from STRUCTURE were then analyzed in CLUMPAK v1.1 (Kopelman, Mayzel, Jakobsson, Rosenberg, & Mayrose, 2015) to select for the best  $K$  and graphically display the results.

## 2.6 | Outlier analyses

To test for evidence of natural selection in relation to coral host, we compared SNPs between lineages of snails on different hosts, pooled across six localities, with two datasets: (a) including all individuals and (b) excluding migrants and admixed individuals that we identified using STRUCTURE. First, we performed an outlier loci analysis using BayeScan v2.1 (Foll & Gaggiotti, 2008) with a prior of 10, a sample size of 5,000, and 100,000 iterations, using a burn-in of 50,000, and 20 pilot runs of 5,000 each. To explore the impact of misleading data, we employed a 10% false discovery rate.

To further explore outlier loci, we used a second method to detect loci under selection (FDIST2) as implemented in ARLEQUIN (Excoffier & Lischer, 2010). We ran 100 demes per group and 50 groups for 50,000 simulations. This model compares a simulated neutral distribution of  $F_{ST}$  to the observed distribution and identifies outliers. Loci with significant  $F_{ST}$   $p$  values ( $< 0.01$ ) were considered to be under selection (Excoffier & Lischer, 2010).



**FIGURE 3** Histogram of variation in  $F_{ST}$  between lineages of *Coralliophila violacea* on two different coral hosts (*Porites lobata* and *P. cylindrica*) across all SNPs, excluding migrants and admixed individuals.  $F_{ST}$  calculated using FDIST in ARLEQUIN. Red line indicates the mean  $F_{ST}$  value (0.075)

## 2.7 | Candidate gene identification and annotation


To annotate the putative functions of genes linked to outlier loci, we aligned sequences containing SNP outlier loci to nucleotide collections (nr/nt) available on the NCBI website, in Blast2GO 5 Basic version (October 7, 2019) using the BLASTn algorithm (Altschul et al., 1997) with a taxonomic filter for Mollusca (taxid:6447). We adjusted parameters (expected threshold 10, word size 7, no low complexity filter, no mask for look-up table only) to accommodate short read sequences. We only examined hits with a high query coverage ( $> 80\%$ ). Then, we identified and annotated any associated genes using NCBI and GeneCards®.

## 3 | RESULTS

After removing empty constructs and filtering for quality, we obtained an average of 5,710,091 unique sequence reads per individual at a minimum  $5\times$  depth. In total, we sequenced and genotyped 17,676 high-quality RAD-seq loci with  $\geq 25\times$  coverage, in 67 snails collected from two different coral host species, at six locations. After filtering for 30% maximum missing data per locus, this total decreased to 5,999 loci and then to 2,718 SNPs following thinning to one SNP per loci to remove any physically linked SNPs for STRUCTURE and  $F_{ST}$  analyses. Next, we removed 16 individuals that had either low DNA quality (missing data  $\geq +1SD$  from the mean) or potential contamination issues (inbreeding coefficient  $\geq +2SD$  from the mean), leaving 51 individuals.

### 3.1 | Genetic structure

Tests of genetic differentiation between sympatric snail lineages on different coral hosts revealed moderate but significant structure (mean  $F_{ST} = 0.047$ , weighted  $F_{ST} = 0.090$  (Weir & Cockerham, 1984)), between host-associated lineages of snails (Figure 3). CLUMPAK



**FIGURE 4** Bar plot of Bayesian assignment probability from STRUCTURE for  $K = 2$  using 2,718 loci from 51 *Coralliophila violacea*. Each vertical bar corresponds to an individual. The proportion of each bar represents an individual's assignment probability to cluster one (green) or two (gold), shown grouped by coral host and then by location as numbered in Table 1, Figure 2

analysis of the STRUCTURE results indicated  $K = 2$  as the best  $K$  value (Appendix S2). At  $K = 2$ , the majority (88%) of all snails grouped by their coral host (Figure 4). Grouping by host was stronger in snails collected from *P. lobata* (97%) than from *P. cylindrica* (79%). Neutral loci (2,645) and outlier loci only (73) showed similar patterns of population structure in STRUCTURE to the full dataset of SNPs (Appendix S3).

### 3.2 | Migration and admixture

Inferring the ancestry of individuals in STRUCTURE, using host as a prior, revealed strong differences among *C. violacea* living on different coral hosts (*P. lobata* and *P. cylindrica*, Figure 4), despite some migration and admixing between sympatric lineages. Moreover, migration rates were strongly asymmetric between snails living on these two hosts. In total, 19% (5 of 26 samples) of the snails collected from *P. cylindrica* had *P. lobata* genetic ancestry, while no snails (0 of 25 samples) with *P. cylindrica* ancestry were ever found on *P. lobata* (Appendix S4 and S5). Admixed individuals were only found at locations where migration was also observed (Dumaguete and Pulau Mengyatan; Appendix S5). After excluding migrants and admixed individuals, the mean  $F_{ST}$  across all loci increased from 0.047 to 0.075 and the weighted  $F_{ST}$  from 0.090 to 0.150.

### 3.3 | Host-specific directional selection


Because STRUCTURE identified 9/51 individuals that were either migrants from one coral host to the other, or of admixed ancestry (Appendix S5), we used two different datasets for detecting host-specific selection: (a) all individuals in the filtered dataset and (b) excluding migrants and admixed individuals. We then searched for loci under selection using two methods. The first involved a Bayesian model, BayeScan (Foll & Gaggiotti, 2008). Using the default false discovery rate (FDR) of 10%, we identified six loci as outliers (pairwise  $F_{ST} = 0.241$ –0.354, mean  $F_{ST} = 0.305$ , Figure 5a, Table 2) in the dataset with all snails. Three of these outlier loci (tag21753, tag39884, tag52997) had  $\log_{10}(PO) > 1$  giving substantial-to-strong support as candidate loci, based on criteria from (Jeffreys, 1961). After excluding all admixed and migrant

individuals, the number of outlier loci only increased to eight (pairwise  $F_{ST} = 0.419$ –0.543, mean  $F_{ST} = 0.480$ , Figure 5b, Table 2). Four of these outlier loci (tag21753, tag28478, tag39884, and tag25141) had  $\log_{10}(PO) > 1$  giving substantial-to-strong support as candidate loci, based on criteria from (Jeffreys, 1961). All outlier loci had positive alpha values, indicating they are under directional selection between snails on different coral hosts.

In the second method, FDIST2, we used the infinite island model of migration to identify 51 outlier loci (pairwise  $F_{ST} = 0.177$ –0.729, mean  $F_{ST} = 0.492$ , Figure 5c) in the dataset with all snails. After removing migrants and admixed individuals, the number of outliers increased to 65 with higher  $F_{ST}$  values (pairwise  $F_{ST} = 0.320$ –0.925, mean  $F_{ST} = 0.620$ , Figure 5d) indicating directional selection, resulting in a combined total of 73 outlier loci across the two methods and datasets. Of these 73, a total of 43 outlier loci were shared between the two datasets; 8 were unique to the all-individual dataset, and 22 were unique to the dataset that excluded migrants and admixed individuals (Table 2). Three outlier loci (tag28478, tag21753, and tag39884) were common among all datasets and methods (Table 2).

### 3.4 | Mapping and annotation of outlier loci

The majority (78%) of putative outlier loci did not align to any other mollusk sequences currently available in the NCBI database (11/2019, Table 2). Sixteen outlier loci DNA sequences aligned with a variety of mollusks including four gastropods (*Aplysia californica*, *Littorina saxatilis*, *Lottia gigantea*, and *Pomacea canaliculata*), three bivalves (*Mizuhopecten yessoensis*, *Crassostrea gigas*, and *C. virginica*), and two cephalopods (*Octopus bimaculoides* and *O. vulgaris*) (Table 2). Of these loci, 7 mapped to hypothetical or uncharacterized proteins. The remaining 9 loci mapped to gene regions with predicted functions. The annotated genes had various associated gene ontology terms including lipid metabolism, metal-ion binding, methyltransferase activity, immune response, chromatin binding, DNA binding, and serine/threonine-protein kinase. The top two hits (lowest e-values) were a neurotransmitter gene (tag15079, *SLC6A7* gene) that plays a role in gastropod feeding behavior (Miller, 2019), and a hormone receptor gene (tag28347, *HR96* gene) involved in the regulation of xenobiotic detoxification (Lindblom & Dodd, 2006; Richter & Fidler, 2014). At tag28347, there were two alleles that occurred



**FIGURE 5** (a)–(b). Results from BayeScan analysis of full RAD-seq dataset (2,718 loci) from *Coralliophila violacea*. Filled gray dots are  $F_{ST}$  outlier loci. (a) All individuals, 6 outlier loci identified FDR = 0.10, (b) excluding migrants and admixed individuals, 8 outlier loci identified FDR = 0.10. (c)–(d). Results from FDIST2 analysis implemented in ARELQUIN using the hierarchical island model of migration. Full RAD-seq dataset (2,718 loci) from *Coralliophila violacea*. Filled black dots are  $F_{ST}$  outlier loci above the 99% quantile (red line). (c) All individuals, 51 outliers, (d) excluding migrants and admixed individuals, 65 outliers

in almost equal frequency (43%, 57%) in the *P. lobata*-associated lineage of snails but were nearly fixed (97%) for one allele in the *P. cylindrica*-associated lineage of snails. Another gene of interest (tag13930, *DRPR* gene) codes for receptors involved in larval locomotory behavior (Freeman, Delrow, Kim, Johnson, & Doe, 2003).

## 4 | DISCUSSION

Genome-wide SNP data from six sympatric populations of *C. violacea* revealed two clearly differentiated clusters that were largely concordant with coral host, consistent with results from mitochondrial DNA (Simmonds et al., 2018). As with insects (Jean & Jean-Christophe, 2010; Simon et al., 2015), this genome-wide differentiation supports the conclusion of ecological divergence based on host association and adds to a small but growing literature on ecological divergence in marine environments (Fritts-Penniman et al., 2020; Potkamp & Fransen, 2019; Titus, Blischak, & Daly, 2019).

While SNP data reveal significant divergence between host-specific lineages of *C. violacea*, divergence was substantially lower in genome-wide SNPs compared to mtDNA ( $F_{ST} = 0.047$  vs.  $\Phi_{CT} = 0.561$ ).

This result may partially be a function of the smaller effective population size of the mitochondrial genome (Palumbi, Cipriano, & Hare, 2001). However, lower divergence values also suggest intermediate levels of gene flow between distinct host-associated lineages ( $Nm > 10$ ), values that are similar to other cases of sympatric host-associated divergence (e.g., Gouin et al., 2017; Peccoud, Ollivier, Plantegenest, & Simon, 2009; Smadja et al., 2012). Divergence with gene flow is further supported by the presence of admixed genotypes and unidirectional gene flow from one host lineage to the other. Moreover, considerable detection of outlier loci under directional selection (2.7% of all SNP loci; 73/2,718) strongly suggests that selection by coral host is likely contributing to the partitioning of *C. violacea* lineages.

### 4.1 | Divergence with gene flow

In parasitic species such as *C. violacea*, divergence with gene flow likely occurs through two mechanisms of premating isolation (Nosil, Vines, & Funk, 2005). The first is host preference for egg laying and/or recruitment to their host (either individual or species). Divergence occurs when mating takes place solely on that host, eventually



Table 2 Outlier loci analysis from *Coralliphila violacea* found on different coral hosts (*Porites lobata*, *P. cylindrica*), BLAST hits, and functional annotations

Outlier analysis		BLAST search results										Gene ontology		
Dataset	Method	FST	log <sub>10</sub> (PO)	Tag ID	DNA sequence	Organism	Description	Score	Coverage (%)	E-value	Identity (%)	Gene symbol	GO terms	Predicted function
all ind.	FDIST2	0.716		21753	AGGTCTCTCTGG CACTGAGCTGCCA AGCTTCCACA	<i>Mizuhopecten yessoensis</i>	Prosaposin-like	35.6	80%	0.23	86%	PSAPL1	Lipid metabolic process Adenylate cyclase-inhibiting G protein-coupled receptor signaling pathway Sphingolipid metabolic process Regulation of metabolic process	NA
no mig./ adm.	FDIST2	0.885												
no mig./ adm.	Bayescan	0.474	1.125											
all ind.	FDIST2	0.665		28478	CATCCCCTCTAT GCAACAGTATGC AAGTCCCCTCT									
all ind.	Bayescan	0.241	0.585											
no mig./ adm.	FDIST2	0.948												
no mig./ adm.	Bayescan	0.534	2.446											
all ind.	FDIST2	0.718		39884	GGGTTGGCTGTAG CAACCTGCTGCC CCCAAAACCTT									
all ind.	Bayescan	0.3511	2.2244											
no mig./ adm.	FDIST2	0.905												
no mig./ adm.	Bayescan	0.484	1.2823											
all ind.	FDIST2	0.659	1.743	52997	CCAGGATCAGC AGTCTCCTGCC ACTGTTCCACAAG	<i>Aplysia californica</i>	Hemocyanin 1	34.6	86%	0.81	84%	KLH1	Metal-ion binding Oxidoreductase activity	NA
no mig./ adm.	FDIST2	0.91												
all ind.	Bayescan	0.507												
all ind.	FDIST2	0.654	1.456	25141	GGTAAAAAGT TGCAATGCTGTG CACAAAAAGTTCA									
no mig./ adm.	FDIST2	0.869												
no mig./ adm.	Bayescan	0.488												

(Continues)

TABLE 2 (Continued)

Outlier analysis		BLAST search results						Gene ontology						
Dataset	Method	FST	log <sub>10</sub> (PO)	Tag ID	DNA sequence	Organism	Description	Score	Coverage (%)	E-value	Identity (%)	Gene symbol	GO terms	Predicted function
all ind.	FDIST2	0.633		14249	AGACAAATTGCC GCACACACATGC AGACAAAACACA	<i>Aplysia californica</i>	Histone-lysine N-methyltransferase 2D-like	38.3	80%	0.066	90%	KMT2D	Metal-ion binding Methyltransferase activity	NA
all ind.	Bayescan	0.321	1.378										Transcription coactivator activity	
no mig./ adm.	FDIST2	0.798											DNA binding	
all ind.	FDIST2	0.702		19628	GGCTATGGGTTT GCAAGGGAGTG CACTCTGCAATCA									
no mig./ adm.	FDIST2	0.893												
no mig./ adm.	Bayescan	0.403	0.603											
all ind.	FDIST2	0.54		36127	TGATCAAGCTT CGCATCGGTCTG CGCTCTCTCTTC									
no mig./ adm.	FDIST2	0.869												
no mig./ adm.	Bayescan	0.419	0.508											
all ind.	FDIST2	0.588		30631	AGCAAGAGAATT GCACAAGGATGC GACCACAGAATG									
no mig./ adm.	FDIST2	0.83												
all ind.	FDIST2	0.65		37258	GATGATCCTGCAG CAGTGTA CTGCC TCTCTCTCTCT	<i>Lotia gigantea</i>	Hypothetical protein	36.5	100%	0.23	84%	Hypothetical protein	NA	NA
no mig./ adm.	FDIST2	0.823												
all ind.	FDIST2	0.478		10161	CACCCCTCTATGC AACAAATGCAC GTCCCCCTCT									
no mig./ adm.	FDIST2	0.795												
all ind.	FDIST2	0.627		30668	AGTGTCTCTTAG CAGGTGACTGC ATGTTGTGACG									
no mig./ adm.	FDIST2	0.794												
all ind.	FDIST2	0.461		21640	AGCTGGATACTG CAGTAACCTGCTT TACAGGAGCA									
no mig./ adm.	FDIST2	0.788												

(Continues)

TABLE 2 (Continued)

Outlier analysis				BLAST search results				Gene ontology						
Dataset	Method	FST	log <sub>10</sub> (PO)	Tag ID	DNA sequence	Organism	Description	Score	Coverage (%)	E-value	Identity (%)	Gene symbol	GO terms	Predicted function
all ind.	FDIST2	0.515		24247	AGTTGCGGCAGG									
no mig./ adm.	FDIST2	0.784			GCAGACTACTGC ATTGACGATCCC									
all ind.	FDIST2	0.572		38182	CGACGGCTAGTGG	<i>Lottia gigantea</i>	Hypothetical protein	32.8	83%	2.8	83%	Hypothetical	NA	NA
no mig./ adm.	FDIST2	0.775			CAATGCTTTTGA ATCGAACATCA									
all ind.	FDIST2	0.55		17358	CAGAAATGTTTCATG	<i>Mizuhopecten yessoensis</i>	Uncharacterized	37.4	83%	0.066	87%	Uncharacterized	NA	NA
no mig./ adm.	FDIST2	0.768			CAGTCCCATGCC ATGTCTCAACT									
all ind.	FDIST2	0.541		38553	AGCACACGACATG									
no mig./ adm.	FDIST2	0.742			CATTCTGTGCC TGAGAAATGCC									
all ind.	FDIST2	0.485		33555	AGGCCTTCATCAG									
no mig./ adm.	FDIST2	0.735			CATCCACAGTGCA TCTCAGGAACA									
all ind.	FDIST2	0.518		22329	TGCTAACACAAGG	<i>Crassostrea gigas</i>	Uncharacterized	38.3	91%	0.066	85%	Uncharacterized	NA	NA
no mig./ adm.	FDIST2	0.729			CATAGTATTGCCGA CATATAACCC									
all ind.	FDIST2	0.536		21872	CGACTCGCGAATG									
no mig./ adm.	FDIST2	0.727			CATTCTTTTGCT GCCTCTTTTTC									
all ind.	FDIST2	0.456		39420	TGTTTGGCTATGG									
no mig./ adm.	FDIST2	0.721			CAGCTGTGTGC TACAACAGAATT									
all ind.	FDIST2	0.468		33550	TGAGAAACACA									
no mig./ adm.	FDIST2	0.705			GCATTAGTTTGC AAATTTAATTCT									
all ind.	FDIST2	0.415		30176	AGGCCTTTTATG									
no mig./ adm.	FDIST2	0.679			GCAAACAGCTG CAACATACTGCCA									
all ind.	FDIST2	0.526		32580	CACCGTTATCTGG									
no mig./ adm.	FDIST2	0.673			CACAACAGTGCG ACGCCTGAAC									

(Continues)

TABLE 2 (Continued)

Outlier analysis				BLAST search results				Gene ontology						
Dataset	Method	FST	log <sub>10</sub> (PO)	Tag ID	DNA sequence	Organism	Description	Score	Coverage (%)	E-value	Identity (%)	Gene symbol	GO terms	Predicted function
all ind.	FDIST2	0.525		28305	TGCTTGCAACATG									
no mig./ adm.	FDIST2	0.67			CAGCATATGCA CACACAAAAC									
all ind.	FDIST2	0.471		10755	GGTGTGAAATTGG									
no mig./ adm.	FDIST2	0.659			CAGC AAAATGCC TTACTCATCCT									
all ind.	FDIST2	0.498		24085	GGATAAAAAGCGG	<i>Pomacea canalicularata</i>	PR domain zinc finger protein 8-like	30.1	86%	9.9	81%	PRDM8	Metal-ion binding	NA
no mig./ adm.	FDIST2	0.652			CACCAAAAATGCG CATAATTTCT								Histone methyltransferase activity Chromatin binding	
all ind.	FDIST2	0.462		32708	TGTGATACTCTTGC	<i>Octopus bimaculoides</i>	AP2-associated protein kinase 1-like	35.6	91%	0.23	85%	AAK1	Kinase, serine/threonine-protein kinase, transferase	NA
no mig./ adm.	FDIST2	0.646			ACTTTACTGCAA AGGCCATGTT								DNA binding, ATP binding, endocytosis	
all ind.	FDIST2	0.57		24158	GGCCTGATCACTG									
no mig./ adm.	FDIST2	0.634			CAGGATCTTGCT GGTATTTGTCA									
all ind.	FDIST2	0.429		28347	AGAAAAGAGGC	<i>Aplysia californica</i>	Nuclear hormone receptor HR96-like	39.2	100%	0.019	83%	HR96	Metal-ion binding	Xenobiotic detoxification
no mig./ adm.	FDIST2	0.617			AGAGAAAAGATAT GGGAGAAAGACA								DNA binding Receptor	
all ind.	FDIST2	0.417		37421	AACTCAAAAATCG									
no mig./ adm.	FDIST2	0.614			CATTTGTTTGCT TTAGTTGCGCT									
all ind.	FDIST2	0.463		22275	TGCAATTGCGAAG									
no mig./ adm.	FDIST2	0.611			CAAATGCTGCT CTGGTGCGCCG									
all ind.	FDIST2	0.404		24087	TGCATATTGTGTGC									
no mig./ adm.	FDIST2	0.599			AGTGCCTTGCGAG AGTATATGCC									

(Continues)

TABLE 2 (Continued)

Outlier analysis				BLAST search results				Gene ontology						
Dataset	Method	FST	log <sub>10</sub> (PO)	Tag ID	DNA sequence	Organism	Description	Score	Coverage (%)	E-value	Identity (%)	Gene symbol	GO terms	Predicted function
all ind.	FDIST2	0.427		16452	AGTGACTGGAGAG	<i>Littorina saxatilis</i>	NA	41	88%	0.005	88%	Uncharacterized		
no mig./adm.	FDIST2	0.587			CACITGTTTGGC GCCTATGTTCC									
all ind.	FDIST2	0.432		27928	CGTGACAACGCCG									
no mig./adm.	FDIST2	0.557			CAACAGAGTGCC TTGGGGACGCC									
all ind.	FDIST2	0.458		48048	GACAGCAAACTG									
no mig./adm.	FDIST2	0.556			CAGCCAGTTGC TTCCCTTGATCG									
all ind.	FDIST2	0.414		17029	TGGTGTTACCTTG									
no mig./adm.	FDIST2	0.554			CAGTCAACTGCA TTTATTCTCT									
all ind.	FDIST2	0.374		34705	AGCAGTCTCACTG									
no mig./adm.	FDIST2	0.526			CAGTTTCTGCA CTGCATAAACT									
all ind.	FDIST2	0.34		20904	TGGCAAGACCTGG									
no mig./adm.	FDIST2	0.522			CAACAGCTGCT GAGATGGGACC									
all ind.	FDIST2	0.372		20142	AGATTCATGCCAG									
no mig./adm.	FDIST2	0.52			CACAATCCTGCA AGACACTATCC									
all ind.	FDIST2	0.388		21098	TGAGAAAAAGTTG									
no mig./adm.	FDIST2	0.516			CATGTGAGTGCG TGCAATGGCCCG									
all ind.	FDIST2	0.334		27266	TGCAATGAAAACA									
no mig./adm.	FDIST2	0.471			CATAAAAACACC TGTGTGCACTC									
all ind.	FDIST2	0.407		15079	GGCTGAGCAGAGG	<i>Pomacea canaliculata</i>	Sodium-dependent proline transporter-like	43.7	86%	0.002	90%	SLC6A7	Neurotransmitter Sodium symporter activity	Gastropod feeding behavior
no mig./adm.	FDIST2	0.451			CAGACGGCTGCG GAGCAGGAGGA									
no mig./adm.	FDIST2	0.748		42043	CGCAATCGTATTGC AAAAATGTGCAAT TGCTCCACT									

(Continues)

TABLE 2 (Continued)

Outlier analysis		BLAST search results					Gene ontology							
Dataset	Method	FST	log <sub>10</sub> (PO)	Tag ID	DNA sequence	Organism	Description	Score	Coverage (%)	E-value	Identity (%)	Gene symbol	GO terms	Predicted function
no mig./ adm.	FDIST2	0.676		31609	CGAACAGATGTGG CAAAGACTGCTG CCTTGGACCA									
no mig./ adm.	FDIST2	0.651		22586	AGACACAGATTGC ATCCCTTTGCCGC GCACTCACC	<i>Octopus vulgaris</i>	Uncharacterized	30.1	100%	9.9	78%	Uncharacterized	NA	NA
no mig./ adm.	FDIST2	0.636		22561	TGTGTGTGTGTTG CACCTACATGCACC TAAGTTACG									
no mig./ adm.	FDIST2	0.624		31557	CGGAGTTTGTAGC AGAGCCTTGCCTG CCATAGTCT	<i>Aplysia californica</i>	Neurogenic protein mastermind-like	31.9	83%	2.8	87%	MAM	Developmental protein, neurogenesis, differentiation	NA
no mig./ adm.	FDIST2	0.559		21042	AGGCTTTGAAGTGC ATGCATGTGCAGC CGTCTGTCA									
no mig./ adm.	FDIST2	0.555		33474	TGACACTAGTCAGC AGATAGATGCCAG GGATGGCCC									
no mig./ adm.	FDIST2	0.514		11613	GGTCCGTGGCTTG ACAGGGATGCAAT GCAATGTCT									
no mig./ adm.	FDIST2	0.492		15069	TGAACATGTCCAG CACCCTTTTGCG CTAAAGAACCCT									
no mig./ adm.	FDIST2	0.486		18108	CACATCCATCTCGCA TAGTTCTGCTGATC CAGAGCA	<i>Crassostrea gigas</i>	NA	39.2	86%	0.019	87%	Uncharacterized	NA	NA
no mig./ adm.	FDIST2	0.478		27744	GAAGTTACACAAGC ACTGCCATGCGTA AAAATGACT									
no mig./ adm.	FDIST2	0.476		32951	TACCTTGGGTATG CAACCCGATGCC AAGACCAAGAT									
no mig./ adm.	FDIST2	0.448		33996	CACGTCCTGCACG CACAAAACCTGCA CTGATGTCTCT									

(Continues)

TABLE 2 (Continued)

Outlier analysis				BLAST search results				Gene ontology						
Dataset	Method	FST	log <sub>10</sub> (PO)	Tag ID	DNA sequence	Organism	Description	Score	Coverage (%)	E-value	Identity (%)	Gene symbol	GO terms	Predicted function
no mig./ adm.	FDIST2	0.44		16737	TGTGTTGTGTGTC AGTTTCATGCAGCT GATTGGTG									
no mig./ adm.	FDIST2	0.431		13930	AGTGAAATAAAGCA ATGAAATGCAGGG CCGTGTCA	<i>Pomacea canaliculata</i>	Protein draper-like		91%	0.81	82%	DRPR	Transmembrane receptor, phagocytosis	Larval locomotory behavior
no mig./ adm.	FDIST2	0.428		34999	GGATCTGTCTGCA AAAGCTGCCTG CTGATCTTG									
no mig./ adm.	FDIST2	0.424		27749	TGAGACGTTAACGCA TACGGCTGCTTTT AAGTAGCC									
no mig./ adm.	FDIST2	0.424		17800	TGTGCTTCCTTGGC AGAACCCCTGCAAA AATAATCTG									
no mig./ adm.	FDIST2	0.407		13296	AGAAAATCTTGGCA CTGTGCTGCTATT GCTTATCA									
no mig./ adm.	FDIST2	0.404		17181	AGCACACAGCACGCA CGTGTTCACAC CAAGAGCA									
no mig./ adm.	FDIST2	0.373		16929	GGTAATCCAAAGCA ACTCAGTGCCTTAC CCCCCCT									
no mig./ adm.	FDIST2	-0.033		23096	CACCCCTCTATGCA AAGTCATGCAAGT CTGCCTCT									
all ind.	FDIST2	0.638		21172	GGTACTAAAAAGCA ACCGTATGCGTAAT CGTCTCA									
all ind.	Bayescan	0.255	0.655											
all ind.	FDIST2	0.593		20062	CACCATGTCTATGC ACGTGCATGCAG ACACTGGCA									
all ind.	FDIST2	0.491		38482	AGGGCACACAGGGC ACACAGATGCACA TCTTACTCA									

(Continues)

TABLE 2 (Continued)

Outlier analysis		BLAST search results						Gene ontology						
Dataset	Method	FST	log <sub>10</sub> (PO)	Tag ID	DNA sequence	Organism	Description	Score	Coverage (%)	E-value	Identity (%)	Gene symbol	GO terms	Predicted function
all ind.	FDIST2	0.417		32340	GAGTTGTCCAAGGC AAAATTCTGCAGA AAGGAACA									
all ind.	FDIST2	0.366		33003	TGAGGCTATTTGC ATGCAGCTGCTA GATCTCTTCC									
all ind.	FDIST2	0.323		9230	TGCAAGCTTTTGGCA TTCCTTTGCAAAT CGAAGGCT									
all ind.	FDIST2	0.225		19533	TGCTCATTACTCGCA TACTGTTGCTCTG TTCAGACT									
all ind.	FDIST2	0.195		11006	CGCAGAAGGAAGG CAAGCAGATGCCT AATAATCGCT									

Note: Only the results that met cutoff statistics are shown.

Abbreviations: adm., admixed; ind., individuals; mig., migrants.



leading to speciation (Funk, Filchak, & Feder, 2002; Hawthorne & Via, 2001). Second is host adaptation, where selection acts against immigrants from another host via immigrant inviability (Nosil, 2007; Nosil et al., 2005; Porter & Benkman, 2017). Our study suggests that both mechanisms may be occurring in *C. violacea*. All migrants were individuals that genetically sorted to the lineage associated with *P. lobata* but were instead living on *P. cylindrica*. Additionally, only admixed individuals were observed on *P. lobata*. This pattern suggests that gene flow and admixture between host-associated lineages are unidirectional—from *lobata* to *cylindrica*. Such unidirectional gene flow could result from two possible scenarios, either the failure of larvae to recruit, or the failure of recruited larvae to survive.

Larval recruitment processes could promote asymmetrical gene flow if the lineage associated with *P. cylindrica* strongly prefers *P. cylindrica* as a host over *P. lobata* or does not respond to chemical settlement cues from *P. lobata*, preventing the recruitment of *P. cylindrica*-associated larvae to *P. lobata*. In addition, larvae from *P. lobata* would need to be less selective in their recruitment, occasionally landing on *P. cylindrica* rather than *P. lobata*. Such a mechanism makes sense, given that there are twice as many coral species ( $N = 8$ ) in the clade of *Porites* to which *P. lobata* belongs, than in that to which *P. cylindrica* belongs.

An alternative, but not mutually exclusive explanation is that asymmetry in gene flow and admixture could result from postsettlement processes. For example, if larvae from *P. cylindrica*-associated individuals settle on *P. lobata*, but are less likely to survive and reproduce, this could lead to immigrant inviability (Ingley & Johnson, 2016; Nosil et al., 2005; Richards & Ortiz-Barrientos, 2016) and asymmetry in admixture. Under such a scenario, genes beneficial to snails living on *P. cylindrica* would likely be less helpful on *P. lobata* and we should see some indication of a selective sweep in the derived lineage with respect to the standing genetic variation of the ancestral lineage (Przeworski, Coop, & Wall, 2005). Indeed, results showed some outlier loci (e.g., *HR96*, detoxification gene) that were in equal proportions in *P. lobata* (43%, 57%) but were at near fixation in *P. cylindrica* (97%), indicating a soft sweep on standing genetic variation at that locus.

Regardless of whether the limited misalignment of snails and coral hosts results from pre- or postrecruitment processes, the fact that the vast majority of snails sort by host coral in the face of hybridization and gene flow indicates that natural selection must be relatively strong to counteract gene flow of  $Nm > 10$  (Funk, Egan, & Nosil, 2011). Moreover, the high fidelity of the snails occupying *P. cylindrica* and lower fidelity of snails occupying *P. lobata*, combined with selective sweeps in *P. cylindrica*, suggest that snails parasitizing *P. lobata* are the ancestral lineage. This conjecture is consistent with the observation that specialist species often evolve from generalist ancestors (Nosil, 2002), likely because specialization constrains further evolution by reducing genetic variation (Moran, 1988). If it is generally true that specialists evolve from generalists (Kawecki, 1996, 1998), then host specialization could be an important mechanism of divergence within the Coral Triangle (Briggs, 2005) as

increased diversity should raise niche partitioning, leading to more opportunities for host specialization (Janz, Nylin, & Wahlberg, 2006).

## 4.2 | Candidate genes involved in adaptation to host

Outlier loci can provide insights into the targets of natural selection (Storz, 2005) and are a useful starting point for determining how selection may be acting on lineages diverging on different hosts. Our analysis revealed 73 putative gene regions with  $F_{ST}$  values significantly higher than neutral expectations, suggesting that they are likely under selection and could be involved in adaptation to coral hosts, or linked to such genes via hitchhiking (Via, 2012).

There is no a priori information on the types of genes involved in the adaptation of mollusks to different hosts and, due to a lack of genomic resources for *C. violacea*, only 9 of 73 outlier loci mapped to gene regions with predicted functions. However, a useful comparison can be found in ectoparasitic phloem-feeding insects adapting to different host plants (Oren et al., 1998). Genes under selection in these insect–plant interactions include genes involved in sensing hosts, that protect insects against plant defenses and facilitate feeding, and that code for digestive and detoxifying enzymes to neutralize plant toxins (e.g., metal-ion binding, Simon et al., 2015).

Experimental evidence suggests genes with metal-ion binding functions are repeatedly under selection in stick insects adapting to different host plants (Soria-Carrasco et al., 2014). Indeed, four of the *C. violacea* candidate genes we identified in outlier tests are involved in metal-ion binding (*KTM2D*, *KLH1*, *PRDM8*, and *HR96*). Very little is known about how corals and their algal symbionts chemically defend themselves against or react to parasites and predators. *Symbiodinium* species do produce toxins—Zooxanthellatoxins—(Gordon & Leggat, 2010), but it is unknown whether these toxins are upregulated in response to parasites or predators.

Additional evidence for detoxification playing a role in host divergence comes from *HR96*, a nuclear hormone receptor involved in xenobiotic detoxification (Richter & Fidler, 2014). Interestingly, *HR96* was nearly fixed for one allele in *C. violacea* from *P. cylindrica* (97%) but was at 50% in *C. violacea* from *P. lobata*, which indicates a selective sweep at that locus. This result, combined with the four metal-ion binding gene regions, suggests that there may be important differences in host-associated detoxification processes in the different *C. violacea* lineages. If adaptation to host-specific toxins drives host specificity, mismatches between snail metabolic abilities and coral hosts could explain the strong asymmetry in snails being found on an atypical coral host.

While the above results suggest a putative detoxification role for some outlier loci, two other genes with predicted functions, a neurotransmitter (*SLC6A7*) important for gastropod feeding behavior (Miller, 2019) and a transmembrane receptor (*DRPR*) involved in larval locomotory behavior, indicate a possible role of behavior in adaptation (Freeman et al., 2003). Notably, this is only the first genomic exploration of *C. violacea* and a broader survey of genomic diversity would be needed to pin down areas of the genome that

are crucial for adaptations to coral hosts. Future work would benefit from a fully annotated genome of *C. violacea* that would allow us to examine the genomic architecture of divergence with gene flow and quantitative trait loci. In turn, this would allow us to better pinpoint regions of the genome under selection, and the specific functions of genes involved in adapting to different hosts.

### 4.3 | Ecological divergence in the sea

John Briggs originally proposed the idea of sympatric speciation as an important diversification mechanism within the Coral Triangle (i.e., “Center of Origin” hypothesis), as well as in the export of species formed under intense competition within the region (Briggs, 1999, 2005). To support his hypothesis, he pointed to multiple cases of sympatric sibling species with distributions centered on the Coral Triangle, where the older of the two species has a wide range, while the younger has a much more restricted range limited to the Coral Triangle (Briggs, 1999). Our study provides the first genomic evidence to support his assertion that ecological divergence with gene flow could be generating biodiversity in the Coral Triangle. In addition, spatial patterning of *C. violacea* sympatric host lineages also matches the pattern Briggs described, with the ancestral *P. lobata* host lineage having a broad geographic distribution, and the derived *P. cylindrica* host lineage restricted to the Coral Triangle (Simmonds et al., 2018).

As the global epicenter of marine biodiversity, there is a large and diverse literature on the processes shaping the Coral Triangle (Barber, Erdmann, & Palumbi, 2006; Bowen et al., 2013; Carpenter et al., 2011; Gaither et al., 2011; Hoeksema, 2007; Kochzius & Nuryanto, 2008; Tornabene, Valdez, Erdmann, & Pezold, 2015). While there is ongoing debate (Evans, McKenna, Simpson, Tournois, & Genner, 2016; Huang, Goldberg, Chou, & Roy, 2018; Di Martino, Jackson, Taylor, & Johnson, 2018; Matias & Riginos, 2018), there is clearly a multiplicity of processes driving diversification in this region (Barber & Meyer, 2015). Given the results of this study, it is important to expand our thinking beyond models that focus solely on allopatry to advance our understanding of marine speciation and origins of the Coral Triangle biodiversity hotspot.

### ACKNOWLEDGMENTS

We are grateful to Dr. Eli Meyer and Dr. Misha Matz for the use of their scripts to process and analyze 2b-RAD sequence data. We acknowledge the Indonesian government, including the Indonesian Ministry of Research and Technology (RISTEK), Indonesian Institute of Sciences (LIPI), Nature Conservation Agency (BKSDA), and the National Marine Park offices of Bunaken and Wakatobi for their support. We are also grateful to the Indonesian Biodiversity Research Center at Udayana University and the Institute for Environmental and Marine Sciences at Silliman University for institutional support. This work was funded by three National Science Foundation programs (OISE-0730256, OISE-1243541, and OCE-0349177) and a grant from the United States Agency for International Development (497-A-00-10-00008-00). The UCLA Department of Ecology and Evolutionary Biology, Lemelson

Foundation, Conchologists of America, and Sigma Xi gave additional funding to S. Simmonds. Sampling was covered under research permits obtained in the Philippines (Department of Agriculture—Bureau of Fisheries and Aquatic Resources) and Indonesia (RISTEK 2011, 198/SIP/FRP/SMNI/2012, 187/SIP/FRP/SM/VI/2013).

### AUTHOR CONTRIBUTIONS

SES conceived of and designed the study. SES, AFP, and SHC collected samples, prepared libraries, and analyzed genomic data. All authors worked on and approved of the manuscript.

### DATA AVAILABILITY STATEMENT

Raw single-end Illumina HiSeq 2000 reads and RAD-seq loci datasets are archived on Dryad: <https://doi.org/10.5068/D1995V>.

### ORCID

Sara E. Simmonds  <https://orcid.org/0000-0003-3438-3559>

### REFERENCES

- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Research*, 25(17), 3389–3402. <https://doi.org/10.1093/nar/25.17.3389>
- Barber, P., & Boyce, S. L. (2006). Estimating diversity of Indo-Pacific coral reef stomatopods through DNA barcoding of stomatopod larvae. *Proceedings. Biological Sciences/the Royal Society*, 273(1597), 2053–2061. <https://doi.org/10.1098/rspb.2006.3540>
- Barber, P. H., Cheng, S. H., Erdmann, M. V., & Tengardjaja, K. (2011). Evolution and conservation of marine biodiversity in the Coral Triangle: Insights from stomatopod Crustacea. In C. Held, S. Koenemann & C.D. Schubart (Eds.), *Phylogeography and population genetics in Crustacea. Crustean Issues* (pp. 264–277). Boca Raton, FL: CRC Press.
- Barber, P. H., Erdmann, M. V., & Palumbi, S. R. (2006). Comparative phylogeography of three codistributed stomatopods: Origins and timing of regional lineage diversification in the Coral Triangle. *Evolution*, 60(9), 1825–1839. <https://doi.org/10.1111/j.0014-3820.2006.tb00526.x>
- Barber, P. H., & Meyer, C. P. (2015). Pluralism explains diversity in the Coral Triangle. In C. Mora (Ed.), *Ecology of fishes on coral reefs* (pp. 258–263). Cambridge, UK: Cambridge University Press. <https://doi.org/10.1017/CBO9781316105412.032>
- Beaumont, M. A., & Nichols, R. A. (1996). Evaluating loci for use in the genetic analysis of population structure. *Proceedings. Biological Sciences/the Royal Society*, 263(1377), 1619–1626. <https://doi.org/10.1098/rspb.1996.0237>
- Bernal, M. A., Gaither, M. R., Simison, W. B., & Rocha, L. A. (2017). Introgression and selection shaped the evolutionary history of sympatric sister-species of coral reef fishes (Genus: *Haemulon*). *Molecular Ecology*, 26(2), 639–652. <https://doi.org/10.1111/mec.13937>
- Bird, C. E., Fernandez-Silva, I., Skillings, D. J., & Toonen, R. J. (2012). Sympatric speciation in the post “Modern Synthesis” era of evolutionary biology. *Evolutionary Biology*, 39(2), 158–180. <https://doi.org/10.1007/s11692-012-9183-6>
- Bird, C. E., Holland, B. S., Bowen, B. W., & Toonen, R. J. (2011). Diversification of sympatric broadcast-spawning limpets (*Cellana* spp.) within the Hawaiian archipelago. *Molecular Ecology*, 20(10), 2128–2141. <https://doi.org/10.1111/j.1365-294X.2011.05081.x>
- Blackall, L. L., Wilson, B., & van Oppen, M. J. H. (2015). Coral—the world’s most diverse symbiotic ecosystem. *Molecular Ecology*, 24(21), 5330–5347. <https://doi.org/10.1111/mec.13400>

- Bowen, B. W., Rocha, L. A., Toonen, R. J., Karl, S. A., & ToBo Laboratory. (2013). The origins of tropical marine biodiversity. *Trends in Ecology & Evolution*, 28(6), 359–366. <https://doi.org/10.1016/j.tree.2013.01.018>
- Briggs, J. C. (1999). Modes of speciation: Marine Indo-West Pacific. *Bulletin of Marine Science*, 65(3), 645–656.
- Briggs, J. C. (2003). Marine centres of origin as evolutionary engines. *Journal of Biogeography*, 30(1), 1–18. <https://doi.org/10.1046/j.1365-2699.2003.00810.x>
- Briggs, J. C. (2005). The marine East Indies: Diversity and speciation. *Journal of Biogeography*, 32(9), 1517–1522. <https://doi.org/10.1111/j.1365-2699.2005.01266.x>
- Campbell, C. R., Poelstra, J. W., & Yoder, A. D. (2018). What is speciation genomics? The roles of ecology, gene flow, and genomic architecture in the formation of species. *Biological Journal of the Linnean Society. Linnean Society of London*, 124(4), 561–583. <https://doi.org/10.1093/biolinnean/bly063>
- Carpenter, K. E., Barber, P. H., Crandall, E. D., Ablan-Lagman, M. C. A., Ambariyanto, Mahardika, G. N., ... Toha, A. H. A. (2011). Comparative phylogeography of the Coral Triangle and implications for marine management. *Journal of Marine Biology*, 2011, 1–14. <https://doi.org/10.1155/2011/396982>
- Case, A. L., & Willis, J. H. (2008). Hybrid male sterility in *Mimulus* (Phrymaceae) is associated with a geographically restricted mitochondrial rearrangement. *Evolution*, 62(5), 1026–1039. <https://doi.org/10.1111/j.1558-5646.2008.00360.x>
- Cowman, P. F., & Bellwood, D. R. (2011). Coral reefs as drivers of cladogenesis: Expanding coral reefs, cryptic extinction events, and the development of biodiversity hotspots. *Journal of Evolutionary Biology*, 24(12), 2543–2562. <https://doi.org/10.1111/j.1420-9101.2011.02391.x>
- Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., ... 1000 Genomes Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics*, 27(15), 2156–2158. <https://doi.org/10.1093/bioinformatics/btr330>
- Di Martino, E., Jackson, J. B. C., Taylor, P. D., & Johnson, K. G. (2018). Differences in extinction rates drove modern biogeographic patterns of tropical marine biodiversity. *Science Advances*, 4(4), eaaq1508. <https://doi.org/10.1126/sciadv.aaq1508>
- Drès, M., & Mallet, J. (2002). Host races in plant-feeding insects and their importance in sympatric speciation. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 357(1420), 471–492. <https://doi.org/10.1098/rstb.2002.1059>
- Duchene, D., Klanten, S. O., Munday, P. L., Herler, J., & van Herwerden, L. (2013). Phylogenetic evidence for recent diversification of obligate coral-dwelling gobies compared with their host corals. *Molecular Phylogenetics and Evolution*, 69(1), 123–132. <https://doi.org/10.1016/j.ympev.2013.04.033>
- Evans, S. M., McKenna, C., Simpson, S. D., Tournois, J., & Genner, M. J. (2016). Patterns of species range evolution in Indo-Pacific reef assemblages reveal the Coral Triangle as a net source of transoceanic diversity. *Biology Letters*, 12(6), 20160090. <https://doi.org/10.1098/rsbl.2016.0090>
- Excoffier, L., & Lischer, H. E. L. (2010). Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources*, 10(3), 564–567. <https://doi.org/10.1098/rspb.2006.3685>
- Fauci, A., Toonen, R. J., & Hadfield, M. G. (2007). Host shift and speciation in a coral-feeding nudibranch. *Proceedings Biological Sciences/the Royal Society*, 274(1606), 111–119. <https://doi.org/10.1098/rspb.2006.3685>
- Feder, J. L., Opp, S. B., Wlazlo, B., Reynolds, K., Go, W., & Spisak, S. (1994). Host fidelity is an effective pre-mating barrier between sympatric races of the apple maggot fly. *Proceedings of the National Academy of Sciences of the United States of America*, 91(17), 7990–7994. <https://doi.org/10.1073/pnas.91.17.7990>
- Foll, M., & Gaggiotti, O. (2008). A genome-scan method to identify selected loci appropriate for both dominant and codominant markers: A Bayesian perspective. *Genetics*, 180(2), 977–993. <https://doi.org/10.1534/genetics.108.092221>
- Footo, A. D., & Morin, P. A. (2015). Sympatric speciation in killer whales? *Heredity*, 114(6), 537–538. <https://doi.org/10.1038/hdy.2014.120>
- Forsman, Z. H., Barshis, D. J., Hunter, C. L., & Toonen, R. J. (2009). Shape-shifting corals: Molecular markers show morphology is evolutionarily plastic in *Porites*. *BMC Evolutionary Biology*, 9, 45. <https://doi.org/10.1186/1471-2148-9-45>
- Freeman, M. R., Delrow, J., Kim, J., Johnson, E., & Doe, C. Q. (2003). Unwrapping glial biology: Gcm target genes regulating glial development, diversification, and function. *Neuron*, 38(4), 567–580. [https://doi.org/10.1016/S0896-6273\(03\)00289-7](https://doi.org/10.1016/S0896-6273(03)00289-7)
- Fritts-Penniman, A. L., Gosliner, T. M., Mahardika, G. N., & Barber, P. H. (2020). Cryptic ecological and geographic diversification in coral-associated nudibranchs. *Molecular Phylogenetics and Evolution*, 144, 106698, ISSN 1055-7903. <https://doi.org/10.1016/j.ympev.2019.106698>
- Fu, L., Niu, B., Zhu, Z., Wu, S., & Li, W. (2012). CD-HIT: Accelerated for clustering the next-generation sequencing data. *Bioinformatics*, 28(23), 3150–3152. <https://doi.org/10.1093/bioinformatics/bts565>
- Funk, D. J., Egan, S. P., & Nosil, P. (2011). Isolation by adaptation in *Neochlamisus* leaf beetles: Host-related selection promotes neutral genomic divergence. *Molecular Ecology*, 20, 4671–4682. <https://doi.org/10.1111/j.1365-294X.2011.05311.x>
- Funk, D. J., Filchak, K. E., & Feder, J. L. (2002). Herbivorous insects: Model systems for the comparative study of speciation ecology. *Genetica*, 116(2–3), 251–267. [https://doi.org/10.1007/978-94-010-0265-3\\_10](https://doi.org/10.1007/978-94-010-0265-3_10)
- Gaither, M. R., Bowen, B. W., Bordenave, T.-R., Rocha, L. A., Newman, S. J., Gomez, J. A., ... Craig, M. T. (2011). Phylogeography of the reef fish *Cephalopholis argus* (Epinephelidae) indicates Pleistocene isolation across the Indo-Pacific barrier with contemporary overlap in the Coral Triangle. *BMC Evolutionary Biology*, 11(1), 189. <https://doi.org/10.1186/1471-2148-11-189>
- Gaither, M. R., & Rocha, L. A. (2013). Origins of species richness in the Indo-Malay-Philippine biodiversity hotspot: Evidence for the centre of overlap hypothesis. *Journal of Biogeography*, 40(9), 1638–1648. <https://doi.org/10.1111/jbi.12126>
- Gittenberger, A., & Gittenberger, E. (2011). Cryptic, adaptive radiation of endoparasitic snails: Sibling species of *Leptoconchus* (Gastropoda: Coralliophilidae) in corals. *Organisms Diversity & Evolution*, 11(1), 21–41. <https://doi.org/10.1007/s13127-011-0039-1>
- González, A. M., Prada, C. A., Ávila, V., & Medina, M. (2018). Ecological speciation in corals. In M. Oleksiak & O. Rajora (Eds.), *Population genomics* (pp. 1–22). Cham, Switzerland: Springer.
- Gordon, B. R., & Leggat, W. (2010). Symbiodinium-invertebrate symbioses and the role of metabolomics. *Marine Drugs*, 8(10), 2546–2568. <https://doi.org/10.3390/md8102546>
- Gouin, A., Bretaudeau, A., Nam, K., Gimenez, S., Aury, J.-M., Duvic, B., ... Fournier, P. (2017). Two genomes of highly polyphagous lepidopteran pests (*Spodoptera frugiperda*, Noctuidae) with different host-plant ranges. *Scientific Reports*, 7(1), 11816. <https://doi.org/10.1038/s41598-017-10461-4>
- Hatfield, T., & Schluter, D. (1999). Ecological speciation in sticklebacks: Environment-dependent hybrid fitness. *Evolution*, 53(3), 866–873. <https://doi.org/10.1111/j.1558-5646.1999.tb05380.x>
- Hawthorne, D. J., & Via, S. (2001). Genetic linkage of ecological specialization and reproductive isolation in pea aphids. *Nature*, 412(6850), 904–907.
- Heller, J. (1993). Hermaphroditism in molluscs. *Biological Journal of the Linnean Society. Linnean Society of London*, 48(1), 19–42. <https://doi.org/10.1111/j.1095-8312.1993.tb00874.x>
- Hoeksema, B. W. (2007). Delineation of the Indo-Malayan centre of maximum marine biodiversity: The Coral Triangle. In W. Renema (Ed.),

- Biogeography, time, and place: Distributions, barriers, and islands* (pp. 117–178). Dordrecht, Netherlands: Springer.
- Huang, D., Goldberg, E. E., Chou, L. M., & Roy, K. (2018). The origin and evolution of coral species richness in a marine biodiversity hotspot. *Evolution*, 72(2), 288–302. <https://doi.org/10.1111/evo.13402>
- Hurt, C., Silliman, K., Anker, A., & Knowlton, N. (2013). Ecological speciation in anemone-associated snapping shrimps (*Alpheus armatus* species complex). *Molecular Ecology*, 22(17), 4532–4548. <https://doi.org/10.1111/mec.12398>
- Ingle, S. J., & Johnson, J. B. (2016). Divergent natural selection promotes immigrant inviability at early and late stages of evolutionary divergence. *Evolution*, 70(3), 600–616. <https://doi.org/10.1111/evo.12872>
- Ingram, T. (2010). Speciation along a depth gradient in a marine adaptive radiation. *Proceedings of the Royal Society B: Biological Sciences*, 278(1705), 613–618. <https://doi.org/10.1098/rspb.2010.1127>
- Janz, N., Nylin, S., & Wahlberg, N. (2006). Diversity begets diversity: Host expansions and the diversification of plant-feeding insects. *BMC Evolutionary Biology*, 6, 4. <https://doi.org/10.1186/1471-2148-6-4>
- Jean, P., & Jean-Christophe, S. (2010). The pea aphid complex as a model of ecological speciation. *Ecological Entomology*, 35, 119–130. <https://doi.org/10.1111/j.1365-2311.2009.01147.x>
- Jeffreys, H. (1961). *Theory of Probability* (3rd). Oxford, UK: Clarendon Press.
- Jiggins, C. D. (2008). Ecological speciation in mimetic butterflies. *BioScience*, 58(6), 541–548. <https://doi.org/10.1641/B580610>
- Kawecki, T. J. (1996). Sympatric speciation driven by beneficial mutations. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 263(1376), 1515–1520. <https://doi.org/10.1098/rspb.1996.0221>
- Kawecki, T. J. (1998). Red queen meets Santa Rosalia: Arms races and the evolution of host specialization in organisms with parasitic lifestyles. *The American Naturalist*, 152(4), 635–651. <https://doi.org/10.1086/286195>
- Kitahara, M. V., Cairns, S. D., Stolarski, J., Blair, D., & Miller, D. J. (2010). A comprehensive phylogenetic analysis of the Scleractinia (Cnidaria, Anthozoa) based on mitochondrial CO1 sequence data. *PLoS ONE*, 5(7), e11490. <https://doi.org/10.1371/journal.pone.0011490>
- Kochzius, M., & Nuryanto, A. (2008). Strong genetic population structure in the boring giant clam, *Tridacna crocea*, across the Indo-Malay Archipelago: Implications related to evolutionary processes and connectivity. *Molecular Ecology*, 17(17), 3775–3787. <https://doi.org/10.1111/j.1365-294x.2008.03803.x>
- Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). Clumpak: A program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources*, 15(5), 1179–1191. <https://doi.org/10.1111/1755-0998.12387>
- Langerhans, R. B., Gifford, M. E., & Joseph, E. O. (2007). Ecological speciation in *Gambusia* fishes. *Evolution*, 61(9), 2056–2074. <https://doi.org/10.1111/j.1558-5646.2007.00171.x>
- Lindblom, T. H., & Dodd, A. K. (2006). Xenobiotic detoxification in the nematode *Caenorhabditis elegans*. *Journal of Experimental Zoology. Part A, Comparative Experimental Biology*, 305(9), 720–730. <https://doi.org/10.1002/jez.a.324>
- Lischer, H. E. L., & Excoffier, L. (2012). PGDSpider: An automated data conversion tool for connecting population genetics and genomics programs. *Bioinformatics*, 28(2), 298–299. <https://doi.org/10.1093/bioinformatics/btr642>
- Litsios, G., Sims, C. A., Wüest, R. O., Pearman, P. B., Zimmermann, N. E., & Salamin, N. (2012). Mutualism with sea anemones triggered the adaptive radiation of clownfishes. *BMC Evolutionary Biology*, 12, 212. <https://doi.org/10.1186/1471-2148-12-212>
- Ludt, W. B., & Rocha, L. A. (2015). Shifting seas: The impacts of Pleistocene sea-level fluctuations on the evolution of tropical marine taxa. *Journal of Biogeography*, 42(1), 25–38. <https://doi.org/10.1111/jbi.12416>
- Malaua, T., Dalecky, A., Ponsard, S., Audiot, P., Streiff, R., Chaval, Y., & Bourguet, D. (2007). Genetic structure and gene flow in French populations of two *Ostrinia* taxa: Host races or sibling species? *Molecular Ecology*, 16(20), 4210–4222. <https://doi.org/10.1111/j.1365-294x.2007.03457.x>
- Martin, S. H., Dasmahapatra, K. K., Nadeau, N. J., Salazar, C., Walters, J. R., Simpson, F., ... Jiggins, C. D. (2013). Genome-wide evidence for speciation with gene flow in *Heliconius* butterflies. *Genome Research*, 23(11), 1817–1828.
- Matias, A. M. A., & Riginos, C. (2018). Revisiting the “Centre Hypotheses” of the Indo-West Pacific: Idiosyncratic genetic diversity of nine reef species offers weak support for the Coral Triangle as a centre of genetic biodiversity. *Journal of Biogeography*, 45(8), 1806–1817. <https://doi.org/10.1111/jbi.13376>
- McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernysky, A., ... DePristo, M. A. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, 20(9), 1297–1303. <https://doi.org/10.1101/gr.107524.110>
- Miglietta, M. P., Faucci, A., & Santini, F. (2011). Speciation in the sea: Overview of the symposium and discussion of future directions. *Integrative and Comparative Biology*, 51(3), 449–455. <https://doi.org/10.1093/icb/acr024>
- Miller, M. W. (2019). GABA as a neurotransmitter in gastropod molluscs. *The Biological Bulletin*, 236(2), 144–156. <https://doi.org/10.1086/701377>
- Moran, N. A. (1988). The evolution of host-plant alternation in aphids: Evidence for specialization as a dead end. *The American Naturalist*, 132(5), 681–706. <https://doi.org/10.1086/284882>
- Munday, P. L., van Herwerden, L., & Dudgeon, C. L. (2004). Evidence for sympatric speciation by host shift in the sea. *Current Biology*, 14(16), 1498–1504. <https://doi.org/10.1016/j.cub.2004.08.029>
- Nielsen, R. (2005). Molecular signatures of natural selection. *Annual Review of Genetics*, 39, 197–218. <https://doi.org/10.1146/annurev.ev.genet.39.073003.112420>
- Nosil, P. (2002). Transition rates between specialization and generalization in phytophagous insects. *Evolution*, 56(8), 1701–1706. <https://doi.org/10.1111/j.0014-3820.2002.tb01482.x>
- Nosil, P. (2007). Divergent host plant adaptation and reproductive isolation between ecotypes of *Timema cristinae* walking sticks. *The American Naturalist*, 169(2), 151–162.
- Nosil, P., Funk, D. J., & Ortiz-Barrientos, D. (2009). Divergent selection and heterogeneous genomic divergence. *Molecular Ecology*, 18(3), 375–402. <https://doi.org/10.1111/j.1365-294x.2008.03946.x>
- Nosil, P., Vines, T. H., & Funk, D. J. (2005). Perspective: Reproductive isolation caused by natural selection against immigrants from divergent habitats. *Evolution*, 59(4), 705–719. <https://doi.org/10.1554/04-428>
- Oren, U., Brickner, I., & Loya, Y. (1998). Prudent sessile feeding by the corallivore snail, *Coralliophila violacea* on coral energy sinks. *Proceedings of the Royal Society B: Biological Sciences*, 265(1410), 2043–2050. <https://doi.org/10.1098/rspb.1998.0538>
- Palumbi, S. R. (1994). Genetic divergence, reproductive isolation, and marine speciation. *Annual Review of Ecology and Systematics*, 25(1), 547–572. <https://doi.org/10.1146/annurev.es.25.110194.002555>
- Palumbi, S. R., Cipriano, F., & Hare, M. P. (2001). Predicting nuclear gene coalescence from mitochondrial data: The three-times rule. *Evolution*, 55(5), 859–868. [https://doi.org/10.1554/0014-3820\(2001\)055\[0859:PNGCFM\]2.0.CO;2](https://doi.org/10.1554/0014-3820(2001)055[0859:PNGCFM]2.0.CO;2)
- Peccoud, J., Ollivier, A., Plantegenest, M., & Simon, J.-C. (2009). A continuum of genetic divergence from sympatric host races to species in the pea aphid complex. *Proceedings of the National Academy of Sciences of the United States of America*, 106(18), 7495–7500. <https://doi.org/10.1073/pnas.0811117106>

- Peijnenburg, K. T. C. A., & Goetze, E. (2013). High evolutionary potential of marine zooplankton. *Ecology and Evolution*, 3(8), 2765–2781. <https://doi.org/10.1002/ece3.644>
- Porter, C. K., & Benkman, C. W. (2017). Assessing the potential contributions of reduced immigrant viability and fecundity to reproductive isolation. *The American Naturalist*, 189(5), 580–591. <https://doi.org/10.1086/691191>
- Potkamp, G., & Fransen, C. M. (2019). Speciation with gene flow in marine systems. *Contributions to Zoology*, 88(2), 133–172. <https://doi.org/10.1163/18759866-20191344>
- Prada, C., DeBiasse, M. B., Neigel, J. E., Yednock, B., Stake, J. L., Forsman, Z. H., ... Hellberg, M. E. (2014). Genetic species delineation among branching Caribbean *Porites* corals. *Coral Reefs*, 33(4), 1019–1030. <https://doi.org/10.1007/s00338-014-1179-5>
- Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, 155(2), 945–959.
- Przeworski, M., Coop, G., & Wall, J. D. (2005). The signature of positive selection on standing genetic variation. *Evolution*, 59(11), 2312–2323. <https://doi.org/10.1554/05-273.1>
- Puebla, O. (2009). Ecological speciation in marine v. freshwater fishes. *Journal of Fish Biology*, 75(5), 960–996. <https://doi.org/10.1111/j.1095-8649.2009.02358.x>
- Puebla, O., Bermingham, E., & McMillan, W. O. (2014). Genomic atolls of differentiation in coral reef fishes (*Hypoplectrus* spp., Serranidae). *Molecular Ecology*, 23(21), 5291–5303. <https://doi.org/10.1111/mec.12926>
- Reijnen, B. T., Hoeksema, B. W., & Gittenberger, E. (2010). Host specificity and phylogenetic relationships among Atlantic Ovulidae (Mollusca: Gastropoda). *Contributions to Zoology*, 79(2), 69–78. <https://doi.org/10.1163/18759866-07902002>
- Richards, T. J., & Ortiz-Barrientos, D. (2016). Immigrant inviability produces a strong barrier to gene flow between parapatric ecotypes of *Senecio lautus*. *Evolution*, 70(6), 1239–1248. <https://doi.org/10.1111/evo.12936>
- Richter, I., & Fidler, A. E. (2014). Marine invertebrate xenobiotic-activated nuclear receptors: Their application as sensor elements in high-throughput bioassays for marine bioactive compounds. *Marine Drugs*, 12(11), 5590–5618. <https://doi.org/10.3390/md12115590>
- Ritson-Williams, R., Shjegstad, S., & Paul, V. (2003). Host specificity of four corallivorous *Phestilla* nudibranchs (Gastropoda: Opisthobranchia). *Marine Ecology Progress Series*, 255, 207–218. <https://doi.org/10.3354/meps255207>
- Ritson-Williams, R., Shjegstad, S. M., & Paul, V. J. (2007). Larval metamorphic competence in four species of *Phestilla* (Gastropoda: Opisthobranchia). *Journal of Experimental Marine Biology and Ecology*, 351(1), 160–167. <https://doi.org/10.1016/j.jembe.2007.06.010>
- Ritson-Williams, R., Shjegstad, S. M., & Paul, V. J. (2009). Larval metamorphosis of *Phestilla* spp. in response to waterborne cues from corals. *Journal of Experimental Marine Biology and Ecology*, 375(1), 84–88. <https://doi.org/10.1016/j.jembe.2009.05.010>
- Rocha, L. A., & Bowen, B. W. (2008). Speciation in coral-reef fishes. *Journal of Fish Biology*, 72(5), 1101–1121. <https://doi.org/10.1111/j.1095-8649.2007.01770.x>
- Rocha, L. A., Robertson, D. R., Roman, J., & Bowen, B. W. (2005). Ecological speciation in tropical reef fishes. *Proceedings of the Royal Society B: Biological Sciences*, 272(1563), 573–579. <https://doi.org/10.1098/2004.3005>
- Schluter, D. (2009). Evidence for ecological speciation and its alternative. *Science*, 323(5915), 737–741. <https://doi.org/10.1126/science.1160006>
- Seehausen, O., Terai, Y., Magalhaes, I. S., Carleton, K. L., Mrosso, H. D. J., Miyagi, R., ... Okada, N. (2008). Speciation through sensory drive in cichlid fish. *Nature*, 455(7213), 620–626. <https://doi.org/10.1038/nature07285>
- Seehausen, O., & Wagner, C. E. (2014). Speciation in freshwater fishes. *Annual Review of Ecology, Evolution, and Systematics*, 45(1), 621–651. <https://doi.org/10.1146/annurev-ecolsys-120213-091818>
- Simmonds, S. E., Chou, V., Cheng, S. H., Rachmawati, R., Calumpang, H. P., Mahardika, G. N., & Barber, P. H. (2018). Evidence of host-associated divergence from coral-eating snails (genus *Coralliophila*) in the Coral Triangle. *Coral Reefs*, 37(2), 355–371. <https://doi.org/10.1007/s00338-018-1661-6>
- Simon, J.-C., d'Alençon, E., Guy, E., Jacquin-Joly, E., Jaquier, J., Nouhaud, P., ... Streiff, R. (2015). Genomics of adaptation to host-plants in herbivorous insects. *Briefings in Functional Genomics*, 14(6), 413–423. <https://doi.org/10.1093/bfpg/elv015>
- Smadja, C. M., Canbäck, B., Vitalis, R., Gautier, M., Ferrari, J., Zhou, J.-J., & Butlin, R. K. (2012). Large-scale candidate gene scan reveals the role of chemoreceptor genes in host plant specialization and speciation in the pea aphid. *Evolution*, 66(9), 2723–2738. <https://doi.org/10.1111/j.1558-5646.2012.01612.x>
- Smith, J. M., & Haigh, J. (1974). The hitch-hiking effect of a favourable gene. *Genetical Research*, 23(1), 23–35. <https://doi.org/10.1017/S0016672300014634>
- Sorenson, M. D., Sefc, K. M., & Payne, R. B. (2003). Speciation by host switch in brood parasitic indigobirds. *Nature*, 424(6951), 928–931. <https://doi.org/10.1038/nature01863>
- Soria-Carrasco, V., Gompert, Z., Comeault, A. A., Farkas, T. E., Parchman, T. L., Johnston, J. S., ... Nosil, P. (2014). Stick insect genomes reveal natural selection's role in parallel speciation. *Science*, 344(6185), 738–742. <https://doi.org/10.1126/science.1252136>
- Sotka, E. E. (2005). Local adaptation in host use among marine invertebrates. *Ecology Letters*, 8, 448–459. <https://doi.org/10.1111/j.1461-0248.2004.00719.x>
- Stella, J. S., Jones, G. P., & Pratchett, M. S. (2010). Variation in the structure of epifaunal invertebrate assemblages among coral hosts. *Coral Reefs*, 29(4), 957–973. <https://doi.org/10.1007/s00338-010-0648-8>
- Storz, J. F. (2005). Using genome scans of DNA polymorphism to infer adaptive population divergence. *Molecular Ecology*, 14(3), 671–688. <https://doi.org/10.1111/j.1365-294x.2005.02437.x>
- Titus, B. M., Blishchak, P. D., & Daly, M. (2019). Genomic signatures of sympatric speciation with historical and contemporary gene flow in a tropical anthozoan (Hexacorallia: Actiniaria). *Molecular Ecology*, 28(15), 3572–3586. <https://doi.org/10.1111/mec.15157>
- Thorpe, R. S., Surget-Groba, Y., & Johansson, H. (2010). Genetic tests for ecological and allopatric speciation in anoles on an island archipelago. *PLoS Genetics*, 6(4), e1000929. <https://doi.org/10.1371/journal.pgen.1000929>
- Tornabene, L., Valdez, S., Erdmann, M., & Pezold, F. (2015). Support for a “Center of Origin” in the Coral Triangle: Cryptic diversity, recent speciation, and local endemism in a diverse lineage of reef fishes (Gobiidae: *Eviota*). *Molecular Phylogenetics and Evolution*, 82(Pt A), 200–210. <https://doi.org/10.1016/j.ympev.2014.09.012>
- Tsang, L. M., Chan, B. K. K., Shih, F.-L., Chu, K. H., & Allen Chen, C. (2009). Host-associated speciation in the coral barnacle *Wanella milleporae* (Cirripedia: Pyrgomatidae) inhabiting the *Millepora* coral. *Molecular Ecology*, 18(7), 1463–1475. <https://doi.org/10.1111/j.1365-294x.2009.04090.x>
- Veron, J. C. E. N., DeVantier, L. M., Turak, E., Green, A. L., Kininmonth, S., Stafford-Smith, M., & Peterson, N. (2011). The Coral Triangle. In Z. Dubinsky & N. Stambler (Eds.), *Coral reefs: An ecosystem in transition* (pp. 47–55). Dordrecht, Netherlands: Springer.
- Via, S. (2009). Natural selection in action during speciation. *Proceedings of the National Academy of Sciences of the United States of America*, 106(Suppl. 1), 9939–9946. <https://doi.org/10.1073/pnas.0901397106>
- Via, S. (2012). Divergence hitchhiking and the spread of genomic isolation during ecological speciation-with-gene-flow. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 367(1587), 451–460. <https://doi.org/10.1098/rstb.2011.0260>

- Wang, S., Meyer, E., McKay, J. K., & Matz, M. V. (2012). 2b-RAD: A simple and flexible method for genome-wide genotyping. *Nature Methods*, 9(8), 808–810. <https://doi.org/10.1038/nmeth.2023>
- Waser, N. M., & Campbell, D. R. (2004). Ecological speciation in flowering plants. In U. Dieckmann, M. Doebeli, J. Metz & D. Tautz(Eds.), *Adaptive speciation (Cambridge Studies in Adaptive Dynamics)* (pp. 264–277). Cambridge, UK: Cambridge University Press. <https://doi.org/10.1017/CBO9781139342179.015>
- Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population structure. *Evolution*, 38(6), 1358–1370. <https://doi.org/10.2307/2408641>
- Westram, A. M., Rafajlović, M., Chaube, P., Faria, R., Larsson, T., Panova, M., ... Butlin, R. (2018). Clines on the seashore: The genomic architecture underlying rapid divergence in the face of gene flow. *Evolution Letters*, 2(4), 297–309. <https://doi.org/10.1002/evl3.74>
- Zann, L. P. (1987). A review of macrosymbiosis in the coral reef ecosystem. *International Journal for Parasitology*, 17(2), 399–405. [https://doi.org/10.1016/0020-7519\(87\)90115-9](https://doi.org/10.1016/0020-7519(87)90115-9)

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Simmonds SE, Fritts-Penniman AL, Cheng SH, Mahardika GN, Barber PH. Genomic signatures of host-associated divergence and adaptation in a coral-eating snail, *Coralliophila violacea* (Kiener, 1836). *Ecol Evol*. 2020;10:1817–1837. <https://doi.org/10.1002/ece3.5977>