UCLA

UCLA Previously Published Works

Title

Response to Letter to the Editor: Psychiatric Disease Among Patients with Takotsubo Syndrome.

Permalink

<https://escholarship.org/uc/item/2w30m0gh>

Journal

Psychosomatics, 59(1)

ISSN

0033-3182

Authors

Nayeri, Arash
Rafla-Yuan, Eric
Farber-Eger, Eric
et al.

Publication Date

2018

DOI

10.1016/j.psych.2017.10.006

Peer reviewed

Psychosomatics 2017 <http://dx.doi.org/10.1016/j.psych.2017.04.008>

2. Wittstein IS, Thiemann DR, Lima JA, et al: Neurohumoral features of myocardial stunning due to sudden emotional stress. *N Engl J Med* 2005; 352(6):539–548
3. Sobue Y, Watanabe E, Ichikawa T, et al: Physically triggered Takotsubo cardiomyopathy has a higher in-hospital mortality rate. *Int J Cardiol* 2017; 235:87–93
4. Konstantinos G, El-Battrawy I, Schramm K, et al: Comparison and outcome analysis of patients with takotsubo cardiomyopathy triggered by emotional stress or physical stress. *Front Psychol* 2017; 8:527
5. Ghadri JR, Cammann VL, Napp LC, et al: Differences in the clinical profile and outcomes of typical and atypical Takotsubo syndrome: data from the International Takotsubo Registry. *JAMA Cardiol* 2016; 1(3):335–340

Response to Letter to the Editor: Psychiatric Disease Among Patients with Takotsubo Syndrome

TO THE EDITOR: We read with great interest the letter to the editor on behalf of Drs. El-Battrawy et al. regarding our recent publication in *Psychosomatics* on the prognostic significance of pre-existing psychiatric illness in Takotsubo cardiomyopathy (TC).^{1,2} In response to the letter, the pattern of wall motion abnormality in TC was not included in our study's multivariable models assessing mortality or recurrence. However, in univariate analysis, there was no significant difference in the rate of "atypical" TC patterns between patients with or without pre-existing psychiatric illness.² As El-Battrawy et al. note, the pattern of TC is likely another prognostic

indicator, as evidenced by findings from the International Takotsubo Registry.³

We also reviewed the publications cited in the letter regarding the potential relationships of TC stressors (emotional vs physical) with risk of recurrent cardiomyopathy. Sobue et al. showed that physical trigger of TC was associated with in-hospital mortality; however, they did not assess for any relationship with TC recurrence.⁴ Giannakopoulos et al. assessed a cohort of 84 patients with TC and were able to show that physical stressors were associated with a composite endpoint of several outcome variables including mortality, rehospitalization, and recurrence of TC.⁵ However, they did not assess the individual relationship between physical or emotional stressors with recurrence. Our study also did not show an association between mode of TC trigger and recurrence.²

In fact, to our knowledge, there has been no study published to date that has demonstrated a link between emotional or physical triggers and risk of recurrence in TC. There is significant concern for low statistical power to detect demographic or clinical associations with recurrent TC given the relatively low rates of incidence and subsequent recurrence of the disorder. We suggest multi-institution and preferably prospective studies to better investigate psychiatric illness in TC in addition to other potential predictors of recurrent cardiomyopathy.

Arash Nayeri, M.D.
Boback Ziaeeian, M.D.
Martin Cadeiras, M.D.
University of California, Los Angeles, CA
Eric Rafla-Yuan, M.D.
University of California, San Diego, CA

Eric Farber-Eger, B.S.
Marcia Blair, M.S.
John A. McPherson, M.D.
Quinn S. Wells, M.D.
Vanderbilt University Medical Center, Nashville, TN

* Send correspondence and reprint requests to Arash Nayeri, M.D., University of California, 757 Westwood Plaza, Suite 7501, Los Angeles, CA.

References

1. El-Battrawy I, Borggrefe M, Akin I: Psychiatric disease among patients with Takotsubo syndrome. *Psychosomatics* 2017
2. Nayeri A, Rafla-Yuan E, Farber-Eger E, et al: Pre-existing psychiatric illness is associated with increased risk of recurrent Takotsubo cardiomyopathy. *Psychosomatics* 2017; 58:527–532
3. Ghadri JR, Cammann VL, Napp LC, et al: Differences in the clinical profile and outcomes of typical and atypical Takotsubo syndrome: data from the international Takotsubo registry. *JAMA Cardiol* 2016; 1(3):335–340
4. Sobue Y, Watanabe E, Ichikawa T, et al: Physically triggered Takotsubo cardiomyopathy has a higher in-hospital mortality rate. *Int J Cardiol* 2017; 235:87–93
5. Giannakopoulos K, El-Battrawy I, Schramm K, et al: Comparison and outcome analysis of patients with Takotsubo cardiomyopathy triggered by emotional stress or physical stress. *Front Psychol* 2017; 8:527

Increasing the Safety of Clozapine Management in Hospitalized Patients With or Without Infection: Still Much to Learn ... and Teach

TO THE EDITOR: Ruan et al.¹ describe more evidence via a case