UCLA

UCLA Previously Published Works

Title

Longitudinal quantitative assessment of coronary plaque progression related to body mass index using serial coronary computed tomography angiography.

Permalink

<https://escholarship.org/uc/item/2wb5g9xn>

Journal

European heart journal. Cardiovascular Imaging, 20(5)

ISSN

2047-2404

Authors

Won, Ki-Bum
Lee, Sang-Eun
Lee, Byoung Kwon
et al.

Publication Date

2019-05-01

DOI

10.1093/ehjci/je192

Peer reviewed

Longitudinal quantitative assessment of coronary plaque progression related to body mass index using serial coronary computed tomography angiography

Ki-Bum Won^{1,2†}, Sang-Eun Lee^{2,3†}, Byoung Kwon Lee^{4*}, Hyung-Bok Park^{3,5}, Ran Heo^{3,6}, Asim Rizvi⁷, Martin Hadamitzky⁸, Yong-Jin Kim⁹, Ji Min Sung^{2,3}, Edoardo Conte¹⁰, Daniele Andreini¹⁰, Gianluca Pontone¹⁰, Matthew J. Budoff¹¹, Ilan Gottlieb¹², Eun Ju Chun¹³, Filippo Cademartiri¹⁴, Erica Maffei¹⁵, Hugo Marques¹⁶, Jonathon A. Leipsic¹⁷, Sanghoon Shin¹⁸, Jung Hyun Choi¹⁹, Renu Virmani²⁰, Habib Samady²¹, Peter H. Stone²², Daniel S. Berman²³, Jagat Narula²⁴, Leslee J. Shaw²¹, Jeroen J. Bax²⁵, James K. Min⁷, and Hyuk-Jae Chang^{2,3}

¹Division of Cardiology, Ulsan University Hospital, University of Ulsan College of Medicine, Ulsan, South Korea; ²Division of Cardiology, Severance Cardiovascular Hospital, Yonsei University College of Medicine, Yonsei University Health System, Seoul, South Korea; ³Yonsei-Cedars-Sinai Integrative Cardiovascular Imaging Research Center, Yonsei University College of Medicine, Yonsei University Health System, Seoul, South Korea; ⁴Gangnam Severance Hospital, Yonsei University College of Medicine, 211 Eonju-ro, Gangnam-gu, Seoul 06237, South Korea; ⁵Division of Cardiology, Catholic Kwandong University International St. Mary's Hospital, Incheon, South Korea; ⁶Division of Cardiology, Hanyang University Seoul Hospital, Hanyang University College of Medicine, Seoul, South Korea; ⁷Dalio Institute of Cardiovascular Imaging, New York-Presbyterian Hospital and Weill Cornell Medical College, New York, NY, USA; ⁸Division of Radiology and Nuclear Medicine, German Heart Center, Munich, Germany; ⁹Seoul National University Hospital, Seoul, South Korea; ¹⁰Centro Cardiologico Monzino, IRCCS, Milan, Italy; ¹¹Department of Medicine, Harbor UCLA Medical Center, Los Angeles, CA, USA; ¹²Department of Radiology, Casa de Saude São Jose, Rio de Janeiro, Brazil; ¹³Seoul National University, Bundang Hospital, Seoul, South Korea; ¹⁴Cardiovascular Imaging Center, IRCCS SDN, Naples, Italy; ¹⁵Department of Radiology, Area Vasta 1/ASUR Marche, Urbino, Italy; ¹⁶Hospital da Luz, Lisbon, Portugal; ¹⁷Department of Radiology, St Paul's Hospital, University of British Columbia, Vancouver, Canada; ¹⁸National Health Insurance Service Ilsan Hospital, Gyeonggi-do, South Korea; ¹⁹Busan University Hospital, Busan, South Korea; ²⁰Department of Pathology, CVPPath Institute, Gaithersburg, MD, USA; ²¹Division of Cardiology, Emory University School of Medicine, Atlanta, GA, USA; ²²Division of Cardiovascular Medicine, Brigham and Women's Hospital, Boston, MA, USA; ²³Department of Imaging, Burns and Allen Research Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA; ²⁴Icahn School of Medicine at Mount Sinai, Mount Sinai Heart, Zena and Michael A. Wiener Cardiovascular Institute, Marie-Josée and Henry R. Kravis Center for Cardiovascular Health, New York, NY, USA; and ²⁵Department of Cardiology, Leiden University Medical Center, Leiden, The Netherlands

Received 17 October 2017; editorial decision 13 November 2018; accepted 26 November 2018

Aims

This study explored the coronary plaque volume change (PVC) according to the change of percent body mass index (BMI) and categorical BMI group using serial coronary computed tomography angiography (CCTA).

Methods and results

A total of 1568 subjects who underwent serial CCTA with available BMI at baseline (CCTA1) and follow-up (CCTA2) were included. Median inter-scan period was 3.3 (interquartile range: 2.6–4.6) years. Quantitative assessment of coronary plaque was performed at both scans. All participants were categorized into three BMI (kg/m²) groups: normal: <25.0; overweight: 25.0–29.9; and obesity: ≥30.0. During follow-up, there were no significant differences in annualized PVC according to the 5% change of BMI in all BMI groups. Among 1424 (90.8%) subjects in the same BMI group at CCTA1 and CCTA2, a significant difference in annualized (PVC) was observed among the three groups. In 144 (9.2%) subjects with the change in their BMI group at CCTA2 compared their results at CCTA1, annualized PVC was not different compared with subjects in the same BMI group during follow-up. The percent change of BMI was not significantly related to the annualized PVC after adjusting confounding factors.

* Corresponding author. Tel: +82 (2) 2019 3307; Fax: +82 (2) 3463 3882. E-mail: cardiobk@yuhs.ac

† The first two authors contributed equally to manuscript preparation.

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author(s) 2019. For permissions, please email: journals.permissions@oup.com.

categorical variables were compared using a χ^2 test. Comparison of continuous variables between CCTA1 and CCTA2 was performed using a paired *t*-test. Generalized linear model was used to identify the association between clinical variables and annualized PVC. Logistic regression analysis was performed to identify the independent predictors of coronary plaque progression. Independent variables included traditional cardiovascular risk factors including age, gender, hypertension, diabetes, dyslipidaemia, and current smoking, baseline plaque volume, baseline categorical BMI group, and statin use. Variables with $P < 0.05$ in the univariate analysis were considered confounding variables and entered into multivariate regression analysis. All statistical analyses were performed using the Statistical Package for the Social Sciences version 19 (SPSS, Chicago, IL, USA) and SAS (version 9.1.3; SAS Institute Inc., Cary, NC, USA). A P -value < 0.05 was considered significant for all analyses.

Results

Baseline characteristics of study population and coronary arteries

The overall mean age was 61 ± 9 years, 62.4% were male, and the median inter-scan period was 3.3 years (2.6–4.6). Hypertension, diabetes mellitus, and dyslipidaemia were present in 54.6%, 21.4%, and 43.9% of the subjects, respectively. In addition, 17.8% of the patients were current smokers. Baseline characteristics according to BMI groups are described in Table 1. The prevalence of male gender, hypertension, and dyslipidaemia were significantly different in each BMI group. However, there were significant differences in the

vessel volume, lumen volume, and plaque volume among the three BMI groups.

Clinical and laboratory variables at CCTA1 and CCTA2

Changes in clinical and laboratory variables during the follow-up periods are presented in Table 2. There were significant differences in systolic and diastolic blood pressure and the levels of triglyceride (TG) and high-density lipoprotein cholesterol (HDL-C) according to the three BMI groups at CCTA1 and CCTA2. However, the levels of total cholesterol (T-chol) and low-density lipoprotein cholesterol (LDL-C) were not different between each of the BMI groups at both CCTAs. During the inter-scan period, the use of statins increased considerably after CCTA1 (45.2% vs. 57.4%). Compared with CCTA1, systolic and diastolic blood pressure and the levels of T-chol, TG, HDL-C, and LDL-C were significantly decreased in all subjects at CCTA2. Shifts from one BMI group to another between CCTA1 and CCTA2 were observed in only 9.2% of the subjects, with 4.4% shifting into a decreased BMI group and 4.8% shifting to an increased BMI group. Subjects with change in BMI more than 5% between CCTA1 and CCTA2 were observed in 15.5%; subjects with decrease and increase more than 5% were 7.0% and 8.5%, respectively.

Change of plaque volume according to the 5% percent change of BMI

There were no significant differences in either PVC or annualized PVC between subjects with a change in BMI of less than 5%,

Table 1 Baseline characteristics

	Normal (n = 752)	Overweight (n = 675)	Obese (n = 141)	P-value
Age (years)	62 (55–67)	61 (55–67)	61 (54–67)	0.468
Male	430 (57.2)	453 (67.1)	95 (67.4)	<0.001
Initial BMI (kg/m ²)	23.3 (22.0–24.2)	26.8 (26.0–27.9)	31.3 (30.5–33.0)	<0.001
Hypertension	367 (49.1)	395 (59.5)	94 (67.1)	<0.001
Diabetes mellitus	162 (21.6)	143 (21.5)	30 (21.3)	0.994
Dyslipidaemia	299 (40.0)	316 (47.7)	74 (52.9)	0.002
Current smoking	137 (18.3)	116 (17.5)	26 (18.7)	0.894
Coronary artery				
Segment length (mm)	379.1 (270.9–464.7)	389.2 (272.7–473.0)	379.4 (274.9–456.1)	0.816
Vessel volume (mm ³)	1890.0 (1277.1–2549.5)	2098.9 (1449.4–2946.6)	2391.8 (1714.3–3127.8)	<0.001
Lumen volume (mm ³)	1783.3 (1212.1–2438.6)	1976.2 (1353.0–2826.3)	2256.4 (1533.6–2995.7)	<0.001
Plaque volume (mm ³)	45.5 (3.2–127.6)	58.7 (9.2–150.4)	80.8 (14.9–190.3)	0.004
Fibrous	18.8 (0.4–53.2)	26.0 (4.9–66.4)	37.6 (6.3–82.8)	<0.001
Fibrous fatty	1.9 (0–15.9)	5.2 (0.0–22.5)	7.8 (0.4–31.5)	<0.001
Necrotic core	0.0 (0.0–0.6)	0.0 (0.0–1.5)	0.2 (0.0–2.3)	<0.001
Dense calcium	8.3 (0.0–45.4)	10.0 (0.0–41.5)	16.7 (0.4–53.8)	0.187
Inter-scan period (years)	3.3 (2.6–4.6)	3.4 (2.6–4.6)	3.3 (2.7–4.8)	0.813

Values are expressed as median (interquartile range) or n (%).
BMI, body mass index.

those with decreases in BMI of more than 5%, and those with increases in BMI of more than 5% at CCTA2 compared with CCTA1 in all BMI groups (Figure 2). Coronary artery characteristics at CCTA2 according to BMI groups are described in [Supplementary data online, Table S3](#).

Change of plaque volume according to the categorical BMI group change

The results of changes in coronary plaque volume according to BMI group are presented in Figure 3. In subjects without any change in BMI group between CCTA1 and CCTA2, a significant difference in PVC was observed among the three groups (normal: 49.1 ± 83.0 mm³ vs. overweight: 59.0 ± 89.3 mm³ vs. obesity: 70.8 ± 87.8 mm³; $P = 0.014$). Similarly, the annualized PVC was significantly different among the three groups (normal: 14.6 ± 24.0 mm³ vs. overweight: 17.0 ± 24.7 mm³ vs. obesity: 21.7 ± 29.4 mm³; $P = 0.009$) (Figure 3A). In subjects who shifted to an increased or decreased BMI group between CCTA1 and CCTA2, both PVC and annualized PVC were not significantly different compared to subjects with no change in BMI group at both CCTA1 and CCTA2 (Figure 3B–D).

Association between clinical variables and change of plaque volume

The results of generalized linear model for analysing the association between clinical variables and annualized PVC were present in Table 3. Hypertension, diabetes mellitus, and baseline plaque volume were independently associated with annualized PVC. However, the percentage change of BMI did not have significant association with annualized PVC.

Independent predictors for coronary plaque progression

The results of univariate and multivariate regression analyses to identify the independent predictors for coronary plaque progression were present in Table 4. In the univariate regression analysis, age [odds ratio (OR): 1.03; 95% confidence interval (CI): 1.02–1.04], male gender (OR: 1.76; 95% CI: 1.37–2.25), hypertension (OR: 1.72; 95% CI: 1.34–2.20), diabetes mellitus (OR: 1.71; 95% CI: 1.22–2.38), dyslipidaemia (OR: 1.43; 95% CI: 1.11–1.84), current smoking (OR: 1.43; 95% CI: 1.01–2.02), baseline plaque volume (OR: 1.08; 95% CI: 1.06–1.10), baseline overweight or obesity (OR: 1.54; 95% CI: 1.21–1.98), and statin use (OR: 1.77; 95% CI: 1.37–2.30) were significantly associated with plaque progression (all $P < 0.05$). Multivariate regression analysis showed that male (OR: 1.38; 95% CI: 1.05–1.81), baseline coronary plaque volume (OR: 1.07; 95% CI: 1.05–1.09), and baseline overweight or obesity (OR: 1.35; 95% CI: 1.04–1.77) were independently associated with coronary plaque progression (All $P < 0.05$).

Discussion

The results of this prospective, multicentre, international study demonstrated that both 5% change of BMI and categorical BMI group change were not related to the coronary PVC during the near-term period. This study could identify that male gender, baseline plaque volume, and baseline overweight or obesity were independently associated with coronary plaque progression.

The advancements in CT technology enable evaluations of coronary atherosclerosis including plaque morphology, quantification of plaque volume, and stenosis severity.^{14–16} In addition, the semi-

Table 2 Changes in clinical and laboratory variables between CCTA1 and CCTA2

	Total (n = 1568)	Normal (n = 752)	Overweight (n = 675)	Obese (n = 141)	P-value
CCTA1					
Systolic blood pressure (mmHg)	129 ± 18	126 ± 17	131 ± 18*	138 ± 20*,**	<0.001
Diastolic blood pressure (mmHg)	78 ± 11	76 ± 10	79 ± 11*	81 ± 12*	<0.001
Fasting glucose (mg/dL)	110 ± 34	108 ± 36	111 ± 32	113 ± 32	0.235
T-chol (mg/dL)	187 ± 40	185 ± 40	189 ± 40	190 ± 42	0.213
TG (mg/dL)	143 ± 85	130 ± 80	155 ± 90*	161 ± 75*	<0.001
HDL-C (mg/dL)	51 ± 14	53 ± 15	49 ± 12*	48 ± 11*	<0.001
LDL-C (mg/dL)	114 ± 34	113 ± 34	115 ± 34	114 ± 41	0.467
Statin use	708 (45.2)	334 (46.8)	307 (49.4)	67 (53.6)	0.306
CCTA2					
Systolic blood pressure (mmHg)	127 ± 17***	125 ± 16	128 ± 17*,***	137 ± 20*,**	<0.001
Diastolic blood pressure (mmHg)	76 ± 10***	75 ± 10***	77 ± 10*,***	79 ± 12*,**	<0.001
Fasting glucose (mg/dL)	110 ± 28	108 ± 28	110 ± 29	115 ± 25	0.162
T-chol (mg/dL)	171 ± 38***	170 ± 37***	173 ± 39***	178 ± 41***	0.118
TG (mg/dL)	129 ± 77***	118 ± 71***	139 ± 82*,***	138 ± 80***	<0.001
HDL-C (mg/dL)	50 ± 13***	52 ± 14***	48 ± 12*,***	49 ± 12	<0.001
LDL-C (mg/dL)	100 ± 33***	98 ± 32***	101 ± 34***	106 ± 34	0.134
Statin use	900 (57.4)	442 (63.5)	384 (62.3)	74 (62.2)	0.896
Changed obesity group between CCTA1 and CCTA2					<0.001
No change	1424 (90.8)	701 (93.2)	606 (89.8)	117 (83.0)	
Decreased	68 (4.4)	0 (0.0)	44 (6.5)	24 (17.0)	
Increased	76 (4.8)	51 (6.8)	25 (3.7)	0 (0.0)	
Cut-offs of 5% BMI change between CCTA1 and CCTA2					<0.001
Within 5% BMI change	1325 (84.5)	655 (87.1)	570 (84.4)	100 (70.9)	
Decrease more than 5%	110 (7.0)	34 (4.5)	52 (7.7)	24 (17.0)	
Increased more than 5%	133 (8.5)	63 (8.4)	53 (7.9)	17 (12.1)	
Change of BMI (kg/m ²)	0.03 ± 1.43	0.15 ± 1.21	0.00 ± 1.08	-0.52 ± 3.02*	<0.001

Values are expressed as mean ± standard deviation or n (%).

*P < 0.05 vs. normal.

**P < 0.05 vs. overweight.

***P < 0.05 vs. index CCTA.

BMI, body mass index; CCTA, coronary computed tomography angiography; CCTA1, baseline CCTA; CCTA2, follow-up CCTA; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; T-chol, total cholesterol; TG, triglyceride.

automated plaque quantification methods allow for the faster measurement of plaque volumes with only minor modifications compared to the traditional manual assessment methods. Plaque volumes measured using the semi-automated methods are highly reproducible and well correlated with those measured by invasive intravascular ultrasound (IVUS).^{17,18} Therefore, CCTA can be considered a useful non-invasive tool for identifying the natural course of coronary atherosclerosis in subjects with low-to-intermediate cardiovascular risk.

Obesity is an important public health issue, and the prevalence of obesity is rapidly increasing. Clearly, obesity is strongly associated with adverse CAD events.^{19,20} In the MESA (multi-ethnic study of atherosclerosis) study, asymptomatic obese subjects had an increased prevalence of CAD as defined by the coronary artery calcium score.²¹ Moreover, subjects with increased BMI showed a greater prevalence, extent, and severity of CAD among subjects with suspected CAD who were referred for CCTA in the CONFIRM (COronary CT Angiography EvaluationN For Clinical Outcomes: An

InterNational Multicentre) study.⁵ However, these studies evaluated only the association between baseline CCTA findings and clinical outcome according to obese status. The PROSPECT (Providing Regional Observations to Study Predictors of Events in the Coronary Tree) study evaluated the natural history of coronary atherosclerosis using CCTA, but the study cohort was very small and all subjects were evaluated after acute coronary syndrome.¹² Thus, data on the change in patients' lifestyles after performing CCTA and the progression of coronary atherosclerosis according to BMI status in subjects with low-to-intermediate risk have been scarce in real clinical practice. The recent guidelines have strongly recommended both modest weight reduction of 5–10% of initial body weight, not absolute body weight, for the overweight or obese populations. It is well-known that the absolute body weight could be significantly influenced on the temporal lifestyle change (i.e. voiding, defecation, dehydration, and so on). To minimize these confounding effects, the present study focused on evaluating the PVC and annualized PVC according to the cut-off of 5% BMI change and categorical BMI group change together.

Figure 2 Coronary plaque progression according to the 5% of BMI change at CCTA2. (A) Normal, (B) overweight, and (C) obesity at CCTA1.

In the present study, the levels of LDL at CCTA2 were lower than those at CCTA1 without significant difference according to obese status. Likewise, the incidence of statin use increased considerably after the CCTAs in all groups. This might have been influenced by the results of previous studies demonstrating that strict LDL control had a beneficial effect on the coronary atherosclerosis.^{22–24} During the median 3.3 years of follow-up, the change in BMI group between CCTA1 and CCTA2 was observed in approximately 9% of the overall subjects. However, 17% of obese subjects had shifted to a decreased BMI group between CCTA1 and CCTA2. In addition, 17% of obese subjects had achieved more than 5% reduction of BMI at CCTA2 compared with the baseline BMI at CCTA1. Considering the negligible weight loss over 1 year in the overweight or obese survivors after acute myocardial infarction in

the PREMIER (Prospective Registry Evaluating Myocardial Infarction: Event and Recovery) study,²⁵ this proportion of obese subjects might be meaningful in the present study. Notably, baseline overweight or obesity was independently related to the coronary plaque progression. However, no significant difference in the annualized PVC according to 5% of BMI change was observed in normal, overweight, and obesity group. In addition, subgroup analysis for subjects with obesity at CCTA1 showed that there were no significant differences in either PVC or annualized PVC between subjects with decreases in BMI of more than 10%, those with a change in BMI of less than 10%, and those with increases in BMI of more than 10% at CCTA2 compared to CCTA1 (Supplementary data online, Figure S1). This finding could be said to indicate that the beneficial effect of body weight reduction on coronary

Figure 3 Comparison of coronary plaque progression based on BMI group at CCTA1 and CCTA2. (A) Same BMI group at CCTA1 and CCTA2; normal vs. overweight vs. obesity at CCTA2. (B) Normal at CCTA1; normal vs. overweight or obesity at CCTA2. (C) Overweight at CCTA1; overweight vs. normal vs. obesity at CCTA2. (D) Obesity at CCTA1; obesity vs. normal or overweight at CCTA2.

atherosclerosis was not fully reflected during the near-term period. To attenuate coronary plaque volume progression, medical therapy for traditional cardiovascular risk factors should be emphasized in overweight and obese subjects with low-to-intermediate risk in clinical practice.

The present study has some limitations. First, only baseline and final BMI data were available; therefore, longitudinal changes in BMI between CCTA could not be confirmed. Second, larger plaque volumes at baseline might lead to larger increases in plaque volume. However, when the baseline plaque volume was adjusted for, we were able to identify that baseline overweight or obesity markedly influenced plaque progression in addition to the baseline plaque volume. Third, despite the application of strict and standardized criteria for the CCTA assessment, atherosclerosis findings might be influenced by the Hounsfield unit density, which is affected by a myriad of factors. In addition, the different kVp setting according to obesity status could potentially have a confounding effect. Fourth, this study was performed in multi-ethnic population but participants were

overwhelmingly composed of Caucasian and East Asian subjects. Finally, there was minimal change in BMI because of the observational design of the present study. Thus, further prospective and randomized studies with larger sample sizes are necessary to confirm the results of the present study. Despite these limitations, this study used CCTA to estimate plaque volume in a large cohort and identified the progression of coronary atherosclerosis according to the percent BMI change and categorical BMI group change in subjects with low-to-intermediate cardiovascular risk.

Conclusions

The present study demonstrated that neither 5% change of BMI nor categorical BMI group change were associated with coronary PVC based on serial quantitative assessment by CCTA during near-term period. This study could identify that male gender, baseline plaque volume, and baseline overweight or obesity had independent impact on coronary plaque progression. Further large prospective and randomized studies with longer follow-up duration might be necessary to confirm the results of present study.

Supplementary data

Supplementary data are available at *European Heart Journal - Cardiovascular Imaging* online.

Funding

This research was supported by Leading Foreign Research Institute Recruitment Program through the National Research Foundation of Korea funded by the Ministry of Science, ICT & Future Planning (grant No. 2012027176), and GE Healthcare.

Conflict of interest: none declared.

Table 3 Generalized linear model for the association between clinical variables and annualized PVC

	Multivariate beta	P-value
Age (years)	0.59	0.442
Male	0.10	0.749
Percentage change of BMI	3.25	0.072
Hypertension	4.10	0.043
Diabetes mellitus	4.53	0.034
Current smoking	0.39	0.531
Baseline plaque volume (mm ³)	419.48	<0.001
Statin use	0.63	0.426

BMI, body mass index; PVC, plaque volume change.

Table 4 Independent predictors for coronary plaque progression

	Univariate		Multivariate	
	OR (95% CI)	P-value	OR (95% CI)	P-value
Age (years)	1.03 (1.02 - 1.04)	<0.001	1.01 (1.00 - 1.03)	0.099
Male	1.76 (1.37 - 2.25)	<0.001	1.38 (1.05 - 1.81)	0.022
Hypertension	1.72 (1.34 - 2.20)	<0.001	1.23 (0.94 - 1.63)	0.138
Diabetes mellitus	1.71 (1.22 - 2.38)	0.002	1.32 (0.93 - 1.88)	0.126
Dyslipidaemia	1.43 (1.11 - 1.84)	0.005	1.00 (0.738 - 1.35)	0.979
Current smoking	1.43 (1.01 - 2.02)	0.044	1.24 (0.85 - 1.81)	0.258
Baseline plaque volume (per 10 mm ³)	1.08 (1.06 - 1.10)	<0.001	1.07 (1.05 - 1.09)	<0.001
Baseline categorical BMI group				
Normal	1		1	
Overweight or obesity	1.54 (1.21 - 1.98)	0.001	1.35 (1.04 - 1.77)	0.027
Statin use	1.77 (1.37 - 2.30)	<0.001	1.34 (0.99 - 1.81)	0.054

BMI, body mass index; CI, confidence interval; OR, odds ratio.

References

- Poirier P, Giles TD, Bray GA, Hong Y, Stern JS, Pi-Sunyer FX et al. Obesity and cardiovascular disease: pathophysiology, evaluation, and effect of weight loss: an update of the 1997 American Heart Association Scientific Statement on Obesity and Heart Disease from the Obesity Committee of the Council on Nutrition, Physical Activity, and Metabolism. *Circulation* 2006;**113**:898–918.
- Lavie CJ, Milani RV, Ventura HO. Obesity and cardiovascular disease: risk factor, paradox, and impact of weight loss. *J Am Coll Cardiol* 2009;**53**:1925–32.
- Smith SC Jr, Jackson R, Pearson TA, Fuster V, Yusuf S, Faergeman O et al. Principles for national and regional guidelines on cardiovascular disease prevention: a scientific statement from the World Heart and Stroke Forum. *Circulation* 2004;**109**:3112–21.
- Naghavi M, Wang H, Lozano R, Davis A, Liang X, Zhou M et al. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;**385**:117–71.
- Labounty TM, Gomez MJ, Achenbach S, Al-Mallah M, Berman DS, Budoff MJ et al. Body mass index and the prevalence, severity, and risk of coronary artery disease: an international multicenter study of 13,874 patients. *Eur Heart J Cardiovasc Imaging* 2013;**14**:456–63.
- Dores H, de Araújo Gonçalves P, Carvalho MS, Sousa PJ, Ferreira A, Cardim N et al. Body mass index as a predictor of the presence but not the severity of coronary artery disease evaluated by cardiac computed tomography. *Eur J Prev Cardiol* 2014;**21**:1387–93.
- Yumuk V, Tsigos C, Fried M, Schindler K, Busetto L, Micic D et al. European guidelines for obesity management in adults. *Obes Facts* 2015;**8**:402–24.
- Lee SE, Chang HJ, Rizvi A, Hadamitzky M, Kim YJ, Conte E et al. Rationale and design of the Progression of Atherosclerotic Plaque Determined by Computed Tomographic Angiography IMaging (PARADIGM) registry: a comprehensive exploration of plaque progression and its impact on clinical outcomes from a multicenter serial coronary computed tomographic angiography study. *Am Heart J* 2016;**182**:72–9.
- Leipsic J, Abbara S, Achenbach S, Cury R, Earls JP, Mancini GJ et al. SCCT guidelines for the interpretation and reporting of coronary CT angiography: a report of the Society of Cardiovascular Computed Tomography Guidelines Committee. *J Cardiovasc Comput Tomogr* 2014;**8**:342–58.
- Abbara S, Blanke P, Maroules CD, Cheezum M, Choi AD, Han BK et al. SCCT guidelines for the performance and acquisition of coronary computed tomographic angiography: a report of the society of Cardiovascular Computed Tomography Guidelines Committee: endorsed by the North American Society for Cardiovascular Imaging (NASCI). *J Cardiovasc Comput Tomogr* 2016;**10**:435–49.
- Motoyama S, Ito H, Sarai M, Kondo T, Kawai H, Nagahara Y et al. Plaque characterization by coronary computed tomography angiography and the likelihood of acute coronary events in mid-term follow-up. *J Am Coll Cardiol* 2015;**66**:337–46.
- de Graaf MA, Broersen A, Kitslaar PH, Roos CJ, Dijkstra J, Lelieveldt BP et al. Automatic quantification and characterization of coronary atherosclerosis with computed tomography coronary angiography: cross-correlation with intravascular ultrasound virtual histology. *Int J Cardiovasc Imaging* 2013;**29**:1177–90.
- Achenbach S, Moselewski F, Ropers D, Ferencik M, Hoffmann U, MacNeill B et al. Detection of calcified and noncalcified coronary atherosclerotic plaque by contrast-enhanced, submillimeter multidetector spiral computed tomography: a segment-based comparison with intravascular ultrasound. *Circulation* 2004;**109**:14–7.
- Hoffmann U, Moselewski F, Nieman K, Jang IK, Ferencik M, Rahman AM et al. Noninvasive assessment of plaque morphology and composition in culprit and stable lesions in acute coronary syndrome and stable lesions in stable angina by multidetector computed tomography. *J Am Coll Cardiol* 2006;**47**:1655–62.
- Motoyama S, Kondo T, Sarai M, Sugiura A, Harigaya H, Sato T et al. Multislice computed tomographic characteristics of coronary lesions in acute coronary syndromes. *J Am Coll Cardiol* 2007;**50**:319–26.
- Motoyama S, Sarai M, Harigaya H, Anno H, Inoue K, Hara T et al. Computed tomographic angiography characteristics of atherosclerotic plaques subsequently resulting in acute coronary syndrome. *J Am Coll Cardiol* 2009;**54**:49–57.
- Papadopoulou SL, Neefjes LA, Schaap M, Li HL, Capuano E, van der Giessen AG et al. Detection and quantification of coronary atherosclerotic plaque by 64-slice multidetector CT: a systematic head-to-head comparison with intravascular ultrasound. *Atherosclerosis* 2011;**219**:163–70.
- Papadopoulou SL, Garcia-Garcia HM, Rossi A, Girasis C, Dharampal AS, Kitslaar PH et al. Reproducibility of computed tomography angiography data analysis using semiautomated plaque quantification software: implications for the design of longitudinal studies. *Int J Cardiovasc Imaging* 2013;**29**:1095–104.
- Hubert HB, Feinleib M, McNamara PM, Castelli WP. Obesity as an independent risk factor for cardiovascular disease: a 26-year follow-up of participants in the Framingham Heart Study. *Circulation* 1983;**67**:968–77.
- Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, Halsey J et al. Body mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet* 2009;**373**:1083–96.
- Folsom AR, Kronmal RA, Detrano RC, O'Leary DH, Bild DE, Bluemke DA et al. Coronary artery calcification compared with carotid intima-media thickness in the prediction of cardiovascular disease incidence: the multi-ethnic study of atherosclerosis (MESA). *Arch Intern Med* 2008;**168**:1333–9.
- Nissen SE, Nicholls SJ, Sipahi I, Libby P, Raichlen JS, Ballantyne CM et al. Effect of very high-intensity statin therapy on regression of coronary atherosclerosis: the ASTEROID trial. *JAMA* 2006;**295**:1556–65.
- Inoue K, Motoyama S, Sarai M, Sato T, Harigaya H, Hara T et al. Serial coronary CT angiography-verified changes in plaque characteristics as an end point: evaluation of effect of statin intervention. *JACC Cardiovasc Imaging* 2010;**3**:691–8.
- Zeb I, Li D, Nasir K, Malpeso J, Batool A, Flores F et al. Effect of statin treatment on coronary plaque progression—a serial coronary CT angiography study. *Atherosclerosis* 2013;**231**:198–204.
- Fadl YY, Krumholz HM, Kosiborod M, Masoudi FA, Peterson PN, Reid KJ et al. Predictors of weight change in overweight patients with myocardial infarction. *Am Heart J* 2007;**154**:711–7.