

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

CONTINUOUS AND DISCONTINUOUS DISAPPEARANCE OF CAPILLARY SURFACES

### Permalink

<https://escholarship.org/uc/item/2wm6r022>

### Authors

Concus, P.  
Finn, R.

### Publication Date

1985-12-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

## Physics Division

JUN 30 1986

LIBRARY AND  
DOCUMENTS SECTION

Mathematics Department


To be submitted for publication

CONTINUOUS AND DISCONTINUOUS DISAPPEARANCE  
OF CAPILLARY SURFACES

P. Concus and R. Finn

December 1985

**TWO-WEEK LOAN COPY**  
*This is a Library Circulating Copy  
which may be borrowed for two weeks.*



*LBL-21002  
c.2*

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

**Continuous and Discontinuous Disappearance  
of Capillary Surfaces<sup>1</sup>**

Paul Concus

Lawrence Berkeley Laboratory  
and  
Department of Mathematics  
University of California  
Berkeley, California 94720

Robert Finn

Department of Mathematics  
Stanford University  
Stanford, California 94305

December 1985

---

<sup>1</sup>This work was supported in part by NASA grant NAG3-146, by NSF grant MCS83-07826, and by the Applied Mathematical Sciences subprogram of the Office of Energy Research, U.S. Department of Energy under contract DE-AC03-76SF00098.

## Continuous and Discontinuous Disappearance of Capillary Surfaces

Paul Concus and Robert Finn

1. We consider the problem of finding a capillary surface  $u(x)$  in a cylinder  $Z$  with section  $\Omega$ , in the absence of gravity. The surface is to meet  $Z$  in (constant) contact angle  $\gamma$  and project simply onto  $\Omega$ . It is known that to each  $\Omega$  there exists a critical angle  $\gamma_0 \in [0, \pi/2]$  such that a surface exists if  $\gamma_0 < \gamma \leq \pi/2$ , while no surface exists if  $0 \leq \gamma < \gamma_0$ . We show here that if  $0 < \gamma_0 < \pi/2$  and if  $\Omega$  is smooth then there is no surface at  $\gamma = \gamma_0$ , while if  $\Omega$  has corners a surface can in some cases be found. In the former case, the surface disappears "continuously" and always becomes unbounded in a subdomain  $\Omega^*$  of positive measure, as  $\gamma \downarrow \gamma_0$ . In the latter case the surface can be bounded, and even analytic with the exception of the corner points. Some applications of the result are given and the exceptional case  $\gamma_0 = 0$  is discussed.

2. In the presence of a gravity field the free surface  $u(x)$  in a capillary tube of section  $\Omega$  is determined as a solution of the governing equations under very general conditions on the form of  $\Omega$ , see, e.g. [1, 2, 3]. If the gravity field is removed, there may be no surface satisfying the physical requirements, even for convex analytic  $\Omega$ , see [4, 5, 6].

In the absence of gravity, the governing equations take the form

$$\operatorname{div} Tu = \frac{\Sigma}{\Omega} \cos \gamma, \quad Tu = \frac{Du}{\sqrt{1+|Du|^2}} \quad (1)$$

in  $\Omega$ , and

$$\nu \cdot Tu = \cos \gamma, \quad (2)$$

on  $\Sigma = \partial\Omega$ . Here  $\gamma$  is the prescribed boundary angle,  $\nu$  is unit exterior normal on  $\Sigma$ . We use the symbols,  $\Sigma, \Omega, \dots$  to denote alternatively a set or its measure. We may

assume  $0 \leq \gamma \leq \pi/2$ , see, e.g. [4].

The material of this paper derives from the following condition for existence, see [6, Theorem 7.1].

*Suppose that an inequality*

$$\int_{\Sigma} |f| \, ds \leq \mu \int_{\Omega} |Df| + \Upsilon \int_{\Omega} |f| \quad (3)$$

*holds for every  $f$  of bounded variation,  $f \in BV(\Omega)$ , with  $\mu < 1/\cos \gamma$ ,  $\Upsilon = \text{const}$  depending only on  $\Omega$ . Suppose further than an isoperimetric inequality*

$$\min(\Omega^*, \Omega \setminus \Omega^*) < C(\Omega \cap \partial\Omega^*)^2 \quad (4)$$

*holds for every Cacciopoli set  $\Omega^* \subset \Omega$ . Then a solution of the capillary problem (1, 2) exists if and only if the functional*

$$\Phi[\Omega^*; \gamma] \equiv \Gamma - \Sigma^* \cos \gamma + \left( \frac{\Sigma}{\Omega} \cos \gamma \right) \Omega^* \quad (5)$$

*satisfies  $\Phi > 0$  for every Cacciopoli set  $\Omega^* \subset \Omega$  with  $\Omega^* \neq \emptyset, \Omega$ .*

Here

$$\Gamma = \int_{\Omega} |D\phi| \quad , \quad \Sigma^* = \int_{\Sigma} \phi \, ds \quad ,$$

where  $\phi$  is the characteristic function of  $\Omega^*$ ; in the second integral,  $\phi$  is taken to be the trace on  $\Sigma^*$  from within  $\Omega$ , see [7, ch. 1]. When it exists, the solution is real analytic and bounded in  $\Omega$ , satisfies (1) strictly and (2) in a generalized sense, and is unique up to an additive constant among all solutions in that class, in particular among all functions that satisfy (1, 2) strictly.

An inequality of the form (3) appears first in Emmer [1], who proved it for a Lipschitz domain with  $\mu = \sqrt{1+L^2}$ ,  $L =$  Lipschitz constant. Thus (3) holds for any smooth domain for any  $\mu > 1$ . An inequality (4) holds in considerable generality, and certainly for smooth domains, cf. [7, p. 25]. Thus, we conclude from the above theorem and from the form of (5) that to every smooth  $\Omega$  there corresponds  $\gamma_0 \in [0, \pi/2]$  such that existence holds if  $\gamma_0 < \gamma \leq \pi/2$ , whereas existence fails if  $0 \leq \gamma < \gamma_0$ . Simple examples show that the case  $0 < \gamma_0 < \pi/2$  occurs commonly; that is, it must be expected that solutions of (1, 2) will exist for some values of  $\gamma$  in the admissible physical range, but not for others.

3. We ask what happens at  $\gamma = \gamma_0$  and find a result that depends in an unusual way on the smoothness of  $\Omega$ .

**THEOREM.** *Let  $\Omega$  be such that  $0 < \gamma_0 < \pi/2$ . Suppose (3) holds with  $\mu < 1/\cos \gamma_0$  and (4) also holds. Then the zero-gravity capillary problem (1, 2) for  $\Omega$  admits no solution when  $\gamma = \gamma_0$ .*

*Proof.* From the condition for existence given in Sec. 2, we find that if  $\gamma < \gamma_0$  then there exists  $\Omega^* \subset \Omega$  with  $\Phi[\Omega^*; \gamma] \leq 0$ ,  $\Omega^* \neq \emptyset$ ,  $\Omega$ . Taking now a sequence  $\gamma_j \uparrow \gamma_0$ , we find that the  $\Omega_j^*$  form a family of sets of bounded perimeter in  $\Omega$ , hence there is a subsequence that converges to  $\Omega_0^*$ , in the sense that the characteristic functions converge in  $L^1(\Omega)$ .

Suppose  $\Omega_0^* = \emptyset$ . We choose  $f$  in (3) to be the characteristic function  $\phi_j$  of  $\Omega_j^*$ , obtaining

$$\Phi[\Omega_j^*; \gamma_j] \geq (1 - \mu \cos \gamma_j) \Gamma - C \Omega_j^*$$

for some fixed constant  $C$ . By (4) we have, since  $\Omega_j^* \rightarrow \emptyset$ , that  $\Omega_j^* < C \sqrt{\Omega_j^*} \Gamma$  for all

sufficiently large  $j$ . Since  $\mu < 1/\cos \gamma_0$  there follows  $\Phi[\Omega_j^*; \gamma_j] > 0$  for large enough  $j$ , which is a contradiction.

Suppose  $\Omega_0^* = \Omega$ . We introduce the "adjoint" functional

$$\Psi[\Omega^*; \gamma] \equiv \Gamma + \Sigma^* \cos \gamma - \left( \frac{\Sigma}{\Omega} \cos \gamma \right) \Omega^*$$

and observe that  $\Phi[\Omega^*; \gamma] = \Psi[\Omega \setminus \Omega^*; \gamma]$ . Using (4), we thus find

$$\Phi[\Omega_j^*; \gamma] \geq (1 - C \sqrt{\Omega \setminus \Omega_j^*}) \Gamma > 0$$

for large enough  $j$ , again a contradiction.

Thus  $\Omega_0^* \neq \emptyset, \Omega$ . We now use an observation that first appears in Gerhardt [8].

We set  $\beta_0 = \cos \gamma_0$ ,  $\beta_j = \cos \gamma_j$  and consider

$$\begin{aligned} \Phi_0 - \Phi_j &= \int_{\Omega} |D \phi_0| - \int_{\Omega} |D \phi_j| - \beta_0 \int_{\Sigma} (\phi_0 - \phi_j) ds \\ &\quad + (\beta_j - \beta_0) \int_{\Sigma} \phi_j ds \\ &\quad + \frac{\Sigma}{\Omega} \beta_j \int_{\Omega} (\phi_0 - \phi_j) dx - \frac{\Sigma}{\Omega} (\beta_j - \beta_0) \int_{\Omega} \phi_0 dx. \end{aligned}$$

Let  $A_\delta \subset \Omega$  denote a strip of width  $\delta$  adjacent to  $\Sigma$ . Let  $\eta(x) \in C^\infty(\bar{\Omega})$ , with  $\eta(x) \equiv 1$  on  $\Sigma$ ,  $\eta(x) \equiv 0$  in  $\Omega_\delta = \Omega \setminus A_\delta$ ,  $0 \leq \eta \leq 1$  in  $\Omega$ . Since  $\beta_j \rightarrow \beta_0$ ,  $0 < \beta_j < 1$ , and  $\phi_j \rightarrow \phi_0$  in  $L^1(\Omega)$ , we may write

$$\begin{aligned} \Phi_0 - \Phi_j &\leq \int_{\Omega_\delta} |D \phi_0| - \int_{\Omega_\delta} |D \phi_j| + \int_{A_\delta} |D \phi_0| - \int_{A_\delta} |D \phi_j| \\ &\quad + \beta_0 \int_{\Sigma} |(\phi_0 - \phi_j)| ds + \epsilon_j, \end{aligned}$$

where  $\epsilon_j \rightarrow 0$  with  $j$ . To the integral over  $\Sigma$  we apply (3), obtaining



$$\begin{aligned} \beta_0 \int_{\Sigma} |(\phi_0 - \phi_j)\eta| ds &\leq (1 - \epsilon) \int_{A_\delta} |D \phi_0| + (1 - \epsilon) \int_{A_\delta} |D \phi_j| \\ &+ C(\Omega; \delta) \int_{\Omega} |\phi_0 - \phi_j| dx \end{aligned}$$

for some  $\epsilon > 0$ . We have further

$$\limsup_j \left\{ \int_{\Omega_\delta} |D \phi_0| - \int_{\Omega_\delta} |D \phi_j| \right\} \leq 0$$

by the semicontinuity of the length integral (cf [7], Theorem 1.9). Thus

$$\limsup_j \{\Phi_0 - \Phi_j\} \leq (2 - \epsilon) \int_{A_\delta} |D \phi_0| .$$

Since  $\delta$  is arbitrary, we find

$$\Phi[\Omega_0; \gamma_0] \leq \limsup_j \Phi[\Omega_j; \gamma_j] \leq 0 ,$$

and since  $\Omega_0 \neq \emptyset, \Omega$ , the existence condition in Sec. 1 yields that no solution exists at  $\gamma_0$ .

The theorem is proved.

Since  $0 < \cos \gamma_0 < 1$ , we conclude immediately from the Theorem and from the remarks preceding it that for a smooth domain there is no surface of the type sought for the critical angle  $\gamma_0$ . It is remarkable that if  $\Omega$  is allowed to contain corners, then a solution at  $\gamma_0$  can in some cases be found. A simple example is obtained by choosing for  $\Omega$  a regular polygon with interior corner angle  $2\alpha$ . A lower hemisphere whose equatorial circle circumscribes  $\Omega$  then provides an explicit (bounded) solution of (1, 2), analytic except at the corner, and with  $\gamma = \pi/2 - \alpha$ . It is shown in [4] that in the case of an interior boundary angle  $2\alpha$  no solution can exist if  $\gamma < \pi/2 - \alpha$ . Thus in the example considered,  $\gamma_0 = \pi/2 - \alpha$  and a solution exists at  $\gamma_0$ .

In such a situation the solution is seen to disappear discontinuously as  $\gamma$  decreases through  $\gamma_0$ . In the former (nonexistence) case, e.g., for smooth boundaries, we may regard the disappearance as continuous, in the sense that each  $\gamma$  for which existence holds lies in a  $\gamma$ -interval of existence.

The Theorem is sharp, in the sense that the result can fail when  $\mu = 1/\cos \gamma_0$ ; that can be seen from the above example of the regular polygon. Nevertheless, the result can hold in much more generality than the case of Lipschitz continuity that seems to be indicated. For example, an inward cusp with opening angle  $2\pi$  can be admissible; see, e.g., the discussion in Chapter 6 of [9]. Precise geometric conditions for admissibility have not yet been established.

4. As an illustration of continuous disappearance, consider the configuration of Figure 1, bounded by line segments  $L_1, L_2$  and by tangent circular arcs  $C_1, C_2$  of radius 1,  $\rho$  respectively. It can be shown that if  $h$  is sufficiently large then there is a unique  $\gamma_0$  and circular arc  $C_0$  meeting the boundary  $\Sigma$  in angle  $\gamma_0$  (as in Figure 1), such that  $\Phi[\Omega^*; \gamma_0] = 0$ . If  $\gamma > \gamma_0$  a solution of (1, 2) exists, while if  $\gamma \leq \gamma_0$  there is no solution. As  $\gamma \downarrow \gamma_0$  the solution  $u(x; \gamma)$  can be normalized to converge to a solution  $u^0(x)$  throughout  $\Omega \setminus \Omega^*$  and to infinity throughout  $\Omega^*$ .

In the above example, for given  $\rho$  there holds  $\lim_{\alpha \rightarrow 0} \gamma_0 = \pi/2$ . Thus, the more "parallel" the segments  $L_1, L_2$  are to each other, the smaller is the range of  $\gamma$  for existence. Nevertheless, if  $\alpha = 0$  a solution exists for any  $\gamma$ , regardless of  $h$ . This singular behavior can be seen in the curves of Figure 2 relating  $\gamma_0$  to  $\rho$  for differing values of  $\alpha$ . (The computed tabular points for the curves are spaced with increment 0.025 in  $\rho$  and connected with straight line segments). As  $\alpha \rightarrow 0$  the curves tend to the




Figure 1

Critical  $\gamma$  vs.  $\rho$  for  
 $\alpha = 85, 70, 45, 20, 10, 5, 2.5, 1, .5, .1, .01$




Figure 2

upper and right hand boundary segments; however  $\alpha = 0$  yields only the indicated single point (1,0).

5. The varied behavior that can occur in the case  $\gamma = 0$  is illustrated further by the example of Figure 3, in which  $C_1$  is a semicircle of radius 1 tangent to the two parallel line segments, and  $C_2$  is a circular arc of radius  $\rho$ .

One can show that there exists a unique  $\rho_0 = 1.974\dots$  such that if  $\rho < \rho_0$  a solution surface exists for  $\gamma = 0$  (and, *a fortiori*, for all  $\gamma > 0$ ) for any value of  $h$ . If  $\rho = \rho_0$  then an "extremal"  $\Omega_0^*$ , for which  $\Phi[\Omega_0^*; 0] = 0$ , is obtained by inscribing a semicircular arc of radius 1 at any point in the strip as indicated. Thus, no solution can exist at  $\gamma_0 = 0$ . If  $\gamma > 0$ , one can show that  $\Phi[\Omega^*, \gamma] > 0$  for any possible "extremal"  $\Omega^*$ , regardless of  $h$ , so that a solution exists for this case. One can show, as for the example of Sec. 4, that as  $\gamma \downarrow 0$  the solution  $u(x; \gamma)$  can be normalized to converge to a solution  $u^0(x)$  throughout  $\Omega \setminus \Omega^*$ , and to infinity throughout the  $\Omega^*$  that is determined in the strip by the arc  $C_0$ .

6. An elaboration of the above configuration yields the example indicated in Figure 4. Here  $\delta = 60^\circ$ ,  $ABC$  is a semicircle;  $AF$ ,  $CG$  are circular arcs of unit radius.

If  $\rho = 1.974$ , calculations show that for  $h = 10$  a solution exists if  $\gamma > \gamma_0 \approx 13.76^\circ$ . At  $\gamma = \gamma_0$  an "extremal" arc  $AC$  with  $\Phi = 0$  appears, of radius  $\rho_1 \approx 1.028$ . A sequence  $u^j$  of solutions corresponding to data  $\gamma_j \downarrow \gamma_0$  can be normalized so that  $u^j$  tends to a solution to the left of  $AC$ , and to infinity to the right of  $AC$ .

If  $\rho = 1.984$ , a solution exists for  $h = 10$  if  $\gamma > \gamma_0 \approx 15.25^\circ$ . At this angle an "extremal" arc  $DE$  with  $\Phi = 0$  appears, of radius  $\rho_2 \approx 1.039$ . A sequence  $u^j$  of solutions corresponding to data  $\gamma_j \downarrow \gamma_0$  can be normalized so that  $u^j$  tends to a solution to




Figure 3




Figure 4

the left of  $DE$ , and to infinity throughout the region to the right of  $DE$ .

Thus, the region in which the solution becomes infinite can be made to shift, essentially discontinuously, with small changes in the domain and data.

We wish to thank E. Giusti for comments that have led to an improvement of the original results. This work was supported in part by NASA grant NAG3-146, by NSF grant MCS83-07826, and by the Applied Mathematical Sciences subprogram of the Office of Energy Research, U.S. Department of Energy under contract DE-AC03-76SF00098.

#### REFERENCES

- [1] Emmer, M.: *Esistenza, unicità e regolarità delle superfici di equilibrio nei capillari*, Ann. Univ. Ferrara, 18 (1983), 79-94.
- [2] Finn, R. and C. Gerhardt: *The internal sphere condition and the capillary problem*, Ann. Mat. Pura Appl., 112 (1977), 13-31.
- [3] Giusti, E.: *Boundary value problems for non parametric surfaces of prescribed mean curvature*, Ann. Scuola Norm. Sup. Pisa 3 (1976), 501-548.
- [4] Concus, P. and R. Finn: *On capillary free surfaces in the absence of gravity*, Acta Math. 132 (1974), 177-198.
- [5] Finn, R.: *Existence criteria for capillary free surfaces without gravity*, Indiana Univ. Math. J. 32 (1983), 439-460.
- [6] Finn, R.: *A subsidiary variational problem and existence criteria for capillary surfaces*, J. Reine Angew. Math. 353 (1984), 196-214.
- [7] Giusti, E.: *Minimal Surfaces and Functions of Bounded Variation*, Monographs in Mathematics, vol. 80, Birkhäuser, 1984.
- [8] Gerhardt, C.: *Global regularity of the solution to the capillary problem*. Ann. Scuola Norm. Sup. Pisa 3 (1976), 157-175.
- [9] Finn, R.: *Equilibrium Capillary Surfaces*, Springer-Verlag, to appear.



This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY  
TECHNICAL INFORMATION DEPARTMENT  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720*