

UCSF

UC San Francisco Previously Published Works

Title

REM sleep behavior disorder, as assessed by questionnaire, in G2019S LRRK2 mutation PD and carriers

Permalink

<https://escholarship.org/uc/item/2x5152hv>

Journal

Movement Disorders, 30(13)

ISSN

0885-3185

Authors

Saunders-Pullman, Rachel
Alcalay, Roy N
Mirelman, Anat
[et al.](#)

Publication Date

2015-11-01

DOI

10.1002/mds.26413

Peer reviewed



Published in final edited form as:

Mov Disord. 2015 November ; 30(13): 1834–1839. doi:10.1002/mds.26413.

REM Sleep Behavior Disorder, as assessed by the RBDSQ, in G2019S LRRK2 mutation PD and unaffected carriers

Rachel Saunders-Pullman, MD, MPH^{1,2,*}, Roy N Alcalay, MD, MSc^{3,4}, Anat Mirelman, PhD⁵, Cuiling Wang, PhD⁶, Marta San Luciano, MD, MS⁷, Roberto A. Ortega, MS¹, Amanda Glickman, BA¹, Deborah Raymond, MS¹, Helen Mejia-Santana, BS³, Nancy Doan, BA¹, Brooke Johannes, MS¹, Kira Yasinovsky, BSc⁵, Laurie Ozelius, PhD^{2,8}, Lorraine Clark,

*Correspondence: Rachel Saunders-Pullman, MD, MPH. The Alan and Barbara Mirken Department of Neurology, Mount Sinai Beth Israel Medical Center, 10 Union Square East, Suite 5J, New York, NY 10003 Phone: 212-844-8719 Fax: 212-844-8461
rsaander@bethisraelny.org.

Disclosures: The authors report no conflicts of interest relating to this study

Authors' roles: 1. Research project: A. Conception, B. Organization, C. Execution; 2. Statistical Analysis: A. Design, B. Execution, C. Review and Critique; 3. Manuscript Preparation: A. Writing of the first draft, B. Review and Critique

R.S-P. 1A, 1B, 1C, 2A, 2C, 3A, 3B

R.N.A. 1A, 1B, 3B

A.M. 1C, 3B

C.W. 2A, 2B, 2C, 3B

M.S. 2A, 2B, 2C, 3B

R.O. 1C, 2B, 2C, 3B

A.G. 1C, 2B, 3B

D.R. 1B, 1C, 3B

H.M-S. 1B, 1C

N.D. 1B, 1C

B.J. 1B, 1C

K.Y. 1B, 1C

L.O. 1B, 1C, 3B

L.C. 1B, 1C, 3B

A.O-U. 1A, 1B, 1C, 3B

K.M. 1A, 3B

N.G. 1A, 1B, 1C, 3B

S.B.B. 1A, 1B, 1C, 3B

Disclosure: The authors report no conflicts of interest. Dr. Saunders-Pullman, Dr. Alcalay, Dr. Mirelman, Dr. Wang, Dr. San Luciano, Mr. Ortega, Ms. Glickman, Ms. Raymond, Ms. Mejia-Santana, Ms. Doan, Ms. Johannes, Ms. Yasinovsky, Dr. Ozelius, Dr. Clark, Dr. Orr-Urtreger, Dr. Marder, Dr. Giladi and Dr. Bressman have nothing to disclose.

Dr. Saunders-Pullman served on the Scientific Advisory Board of the Dystonia Medical Research Foundation. She receives research support from the NIH (K02 NS073836), the Michael J Fox Foundation for Parkinson's Research, the Bachmann-Strauss Dystonia and Parkinson's Foundation, the Marclad Foundation, and the Genzyme Generations Program.

Dr. Alcalay receives research support from the NIH (K02NS080915), the Parkinson's disease Foundation, the Smart Foundation and the Michael J Fox foundation.

Dr. Ozelius receives salary support from NIH [NS037409, NS075881, DC011805]. She is a current member of the scientific advisory boards of the National Spasmodic Dysphonia Association, the Benign Essential Blepharospasm Research Foundation and Tourette Syndrome Association, Inc. Dr. Ozelius receives royalty payments from Athena Diagnostics related to patents.

Dr. Clark receives research support from the NIH [NINDS #R01 NS060113 (principal investigator), NINDS #R01 NS073872 (Co-principal investigator), NIA #5P50AG008702(Project 3, principal investigator), and NINDS #NS36630 (coinvestigator) and 2P50NS038370-11 (Co-Investigator)], and the Parkinson's Disease Foundation (principal investigator) and the Michael J Fox Foundation (co-investigator).

Dr. Bressman serves on the advisory boards of the Michael J. Fox Foundation, the Dystonia Medical Research Foundation, the Bachmann Strauss Dystonia and Parkinson's Foundation, and the Board of We Move. She has consulted for Bristol Meyer Squibb. She has received research support from the Michael J. Fox Foundation, National Institutes of Health (NIH), and Dystonia Medical Research Foundation.

Dr. Marder receives research support from the NIH [#NS036630 (PI), 1UL1 RR024156-01(Director PCIR), PO412196- G (Co-I), and PO412196-G (Co-I)]. She received compensation for participating on the steering committee for U01NS052592 and from the Parkinson Disease Foundation, Huntington's Disease Society of America, the Parkinson Study Group, CHDI, and the Michael J Fox foundation.

PhD^{4,9}, Avi Orr-Utreger, MD, PhD¹⁰, Karen Marder, MD, MS^{3,4,11,12}, Nir Giladi, MD⁵, Susan B Bressman, MD^{1,2}, and the AJ LRRK2 Consortium

¹Department of Neurology, Mount Sinai Beth Israel Medical Center, New York, NY, USA

²Department of Neurology, Icahn School of Medicine at Mount Sinai, New York, NY, USA

³Department of Neurology, College of Physicians and Surgeons, Columbia University, New York, NY, USA

⁴Taub Institute for Research on Alzheimer's Disease and the Aging Brain, College of Physicians and Surgeons, Columbia University, New York, NY, USA

⁵Movement Disorders Unit, Department of Neurology, Tel-Aviv Medical Center, Sieratzki Chair of Neurology, Department of Neurology and Neurosurgery, Sackler School of Medicine, Sagol School of Neuroscience, Tel-Aviv University, Tel-Aviv, Israel

⁶Department of Epidemiology and Social Medicine, Albert Einstein College of Medicine, Bronx, NY, USA

⁷Department of Neurology, University of California San Francisco, San Francisco, CA, USA

⁸Department of Genetics, Icahn School of Medicine at Mount Sinai, New York, NY, USA

⁹Department of Pathology and Cell Biology, College of Physicians and Surgeons, Columbia University, New York, NY, USA

¹⁰Genetic Institute, Tel-Aviv Medical Center, Sackler School of Medicine, Sagol School of Neuroscience, Tel-Aviv University, Tel-Aviv, Israel

¹¹Gertrude H. Sergievsky Center, College of Physicians and Surgeons, Columbia University, New York, NY, USA

¹²Department of Psychiatry, Columbia University Medical Center, New York, NY, USA

Abstract

Background—Rapid eye movement sleep Behavior Disorder occurs with idiopathic Parkinson disease (PD), and often precedes PD. Its frequency in *LRRK2*-PD and utility as a pre-clinical marker has not been established.

Methods—144 idiopathic PD, 142 *LRRK2* G2019S mutation PD, 117 non-manifesting carriers, 93 related non-carriers, and 40 controls completed the Rapid eye movement sleep Behavior Disorder Screening Questionnaire.

Results—30.6% idiopathic PD, 19.7% *LRRK2*-PD, 6% non-manifesting carriers, 20.4% related non-carriers, and 15% controls met cut-scores. The likelihood of abnormal scores was decreased in *LRRK2*-PD vs. idiopathic PD (OR=0.55, p=0.03), non-manifesting carriers vs. related non-carriers (OR=0.25, p<0.01), and PD <3 years duration, 1/19 *LRRK2*-PD vs. 14/41 idiopathic PD (p<0.05).

Conclusions—There is a lower frequency of abnormal Questionnaire scores in *LRRK2*-PD, especially in early *LRRK2*-PD, and in non-manifesting carriers. Therefore, the Rapid eye

movement sleep Behavior Disorder Questionnaire is unlikely to serve as a pre-clinical marker for phenoconversion to PD.

Keywords

Parkinson's disease; LRRK2; REM Behavior Disorder; RBDSQ; pre-clinical

INTRODUCTION

In the search for markers of pre-clinical parkinsonism, REM sleep behavior disorder (RBD) holds promise, as it is highly specific for phenoconversion to parkinsonism. As many as 90% of RBD cases develop a synucleinopathy, either idiopathic PD (IPD), Parkinson disease with dementia, dementia with Lewy bodies, or multiple system atrophy¹⁻⁴. The time course to motor phenoconversion may exceed 10 years^{3, 5}, suggesting a long at-risk “pre-clinical” period during which potential therapeutics might be applied to lessen pathologic burden before development of full-fledged clinical disease^{6, 7}. The gold-standard assessment for RBD is polysomnography (PSG). However, questionnaires regarding sleep behavior have good correlation with cases diagnosed in sleep laboratories,⁸ and are more practical as screening tools. Harboring the G2019S *LRRK2* mutation is also a marker of pre-clinical disease, but only a third of carriers develop PD in their lifetimes^{9, 10}. Therefore, combining a questionnaire marker for RBD with carriage of the *LRRK2* mutation might detect individuals who are at higher risk for phenoconversion. The REM Sleep Behavior Disorder Screening Questionnaire (RBDSQ)⁸ is one such questionnaire. In a smaller group of *LRRK2*-PD, there was a trend for RBDSQ abnormalities to be less frequent in *LRRK2*-PD than IPD³². Further, in a subset of the population evaluated here, continuous RBDSQ scores did not differ between carriers without PD (non-manifesting carriers (NMC)) and family members without PD (non-carrier family members, NCF)¹⁵. Herein we report the largest group of G2019S mutation *LRRK2*-PD assessed with RBDSQ, and because of the sample size, are able to report on a subgroup with early disease.

METHODS

536 participants from the *LRRK2* Ashkenazi Jewish Consortium centers in Tel Aviv (Tel Aviv Medical Center, TA n=249,) and New York (Mount Sinai Beth Israel, MSBI n=155, and Columbia University Medical Center, CU n=155), self-administered the RBDSQ, a questionnaire that evaluates frequency of dreams and nocturnal behavior, self and bed partner injurious behavior, specific movements and activities during sleep, nocturnal awakenings and overall sleep quality. Subjects included 144 PD without G2019S or any of the eight major Ashkenazi Jewish *GBA1* mutations (IPD), 142 G2019S mutation carriers with PD (*LRRK2*-PD), 117 G2019S mutation carriers without PD (NMC), 93 non-PD, non-mutation-carrying family members (NC-F), and 40 unrelated controls. Continuous scores from a subset of the RBDSQ data comparing NMC to NC-F were previously reported.¹⁵

Univariate demographic and RBDSQ comparisons were performed using Wilcoxon rank sum and Fisher's exact or Chi-squared tests both for the groups overall and evaluating factors that might be related to RBD. Generalized estimating equation (GEE) logistic

regression models evaluated likelihood of RBD by the RBDSQ cutoff-scores (defined as 6 for PD³⁷ and 5 for non-PD⁸), and adjusted for age, gender, UPDRS-III, and site (NY or TA), accounting for correlation between family members.

Further secondary analyses were performed within groups to determine whether features segregated with RBD in the LRRK2-PD and IPD groups, including duration of disease, UPDRS-III, motor subtype (postural instability gait predominant, PIGD)¹⁶, Montreal Cognitive Assessment (MoCA) and hyposmia (defined as <15th percentile for age and gender with the 40-item University of Pennsylvania Smell Identification Test (UPSIT)).

RESULTS

Demographics (Table 1)

LRRK2-PD were more likely to be male than controls and NMC but not than IPD or NC-F. Mean age of IPD (64.9 ± 11.3) was not different from LRRK2-PD (67.2 ± 9.8). Both IPD and LRRK2-PD were older than NMC (53.3 ± 15.9) and NC-F (52.8 ± 18.5). IPD were younger than controls (69.8 ± 10.6) but LRRK2-PD were not. Age at onset did not differ between IPD and LRRK2-PD (58.0 ± 10.9 and 57.4 ± 11.0), nor did UPDRS-III, but duration of PD was longer in the LRRK2-PD group (7.36 ± 7.53 vs. 9.98 ± 7.10 years) ($p < 0.01$).¹⁷

RBDSQ Scores (Table 1)

30.6% of IPD met criteria for abnormal RBDSQ score vs. 19.7% of LRRK2-PD, 6.0% of NMC, 20.4% of non-carrier family members and 15.0% of controls. In the main GEE model adjusting for age, site (NY vs. TA) and gender, the likelihood of having abnormal RBDSQ was decreased in LRRK2-PD vs. IPD (OR=0.55, $p=0.03$) and increased in LRRK2-PD vs. NMC (OR=3.24, $p=0.01$) but not different in LRRK2-PD vs. controls (OR 1.50, $p=0.42$). It was less in NMC than NC-F (OR=0.25, $p < 0.01$), and did not differ between NMC and controls (OR=2.17, $p=0.21$). Among subjects with early PD (<3 years duration), abnormal scores were present in only 1/19 cases with LRRK2-PD and in 12/41 cases with IPD ($p < 0.05$). In post hoc univariate comparisons in IPD (Table 2), abnormal RBDSQ scores were associated with worse UPSIT score, PIGD phenotype. In LRRK2-PD, duration of disease and anxiety were significant.

There was no difference between the IPD and LRRK2-PD groups in current use of medications implicated as triggers of RBD, such as MAO-B inhibitor, tricyclic antidepressant (TCA), or selective serotonin reuptake inhibitor (SSRI) (IPD=82.6% vs. LRRK2-PD=73.9%, $p=0.07$) or those used for treatment of RBD, such as clonazepam (IPD=6.9% vs. LRRK2-PD=7.8%, $p=0.80$). In a GEE model limited to patients with PD adjusting for reported current use of trigger or treatment medication, age, site, gender, and disease duration, there was no change in the direction or the overall magnitude of the odds ratio for abnormal RBDSQ between LRRK2-PD and IPD.

DISCUSSION

RBD should meet several criteria to serve as a marker to detect a highly at-risk LRRK2-PD subset: it should 1) have easily applied and relevant instruments that discern the marker and these should 2) be abnormal in at least a subset of affected individuals in the disease of interest (LRRK2-PD), 3) be abnormal early in disease, and 4) be present in at least a subset of NMC prior to phenoconversion. While PSG is the gold-standard, the RBDSQ is readily applied and inexpensive and, in our study, the proportion of IPD with RBD was similar to another study¹⁸. However, while present in LRRK2 carriers, abnormal RBDSQ scores occurred in only one of the 19 LRRK2-PD with early disease, compared with 14/41 IPD ($p=0.05$). This supports that even in a cross-sectional *LRRK2* study, abnormal RBDSQ is not a frequent feature in early LRRK2-PD. Further, the paucity of NMC who meet criteria for RBDSQ does not allow for assessment of a subcluster, which might be present in a small subgroup of NMC close to phenoconversion.¹⁹

The decreased frequency of abnormal RBDSQ scores in LRRK2-PD compared with IPD is consistent with its infrequency in certain genetic forms of PD. RBD is more common in IPD with neuropsychological deficits,^{20–24} akinetic-rigid disease,^{25–27} orthostatic changes,^{3, 28} and greater heart-rate variability²⁹. With the exception of MoCA, which did not correlate with abnormal RBDSQ scores in this cohort, we observed similar trends suggesting possible subgroups relating RBD and non-motor features in both IPD and LRRK2-PD. In the IPD group, RBDSQ was associated with worse olfaction and PIGD phenotype. In the LRRK2-PD group, RBD was associated with more prominent trait ($p<0.01$) and state ($p=0.03$) anxiety.^{30,31,19}

There are several limitations to our study. Measurement of RBD by questionnaire does not capture RBD in the same manner as the gold standard polysomnography, and the RBDSQ is just one of several validated questionnaires, each with their respective strengths. Additionally, we did not require sleep partners to participate in all questionnaires, and the information regarding whether a bed partner participated in the questionnaire was not consistently noted, so that a proportion of cases might be underestimated or mis-classified. However, in support of our methodology and findings, our results are consistent with the trends seen using other assessment methods, where the rates of RBD were lower in LRRK2-PD (range: 11.1–21.2%) compared to IPD (range: 29 to 42%)^{32–34}. Further, the RBDSQ performs well overall, particularly with normal volunteer controls, with 96% sensitivity and 92% specificity⁸. This group is more similar to NMC, NC-F and controls than a symptomatic RBD group would be. Only one study reports the gold standard for RBD, PSG, and reports six cases of LRRK2-PD with RBD¹⁴. However rates were not compared between LRRK2-PD and IPD. Further, PSG is time-intensive and expensive. We cannot exclude that rates might differ were PSG utilized, as clinical reports may underreport as many as half of the cases^{36, 37}.

In summary, RBD, as measured by RBDSQ, is not a common feature of early LRRK2 PD and *LRRK2* gene carrier status overall. Longitudinal study with PSG will facilitate a more precise assessment of the association between RBD and *LRRK2* expression over time.

Acknowledgments

Funding: This study was funded by the Michael J Fox Foundation for Parkinson's Research, the National Institute of Health (through Grant Numbers K02-NS073836, R56NS036630, K02NS080915, NS050487, NS060113, UL1 TR000040, UL1 RR024156 and NINDS 10628097) as well as by grants from the Tel Aviv Sourasky Medical Center Grant of Excellence, Khan Foundation, and the Israel Science Foundation Heritage Legacy. The funding sources had no involvement in the design, interpretation or writing of this manuscript.

We thank the participants of this project for their invaluable participation and the staff of the AJ *LRRK2* consortium for their contribution:

Mount Sinai Beth Israel Medical Center, New York, NY : Vicki Lynn Shanker, MD; Mark Groves, MD; Christine Palmese, PhD; Erin Deegan, MD; Adam Drobnis, MD; Jeffrey Ratliff, MD; Lawrence Severt, MD; Naomi Lubarr, MD; Rivka Sachdev, MD; Amanda Pomerantz, BA

Columbia University, Columbia Presbyterian Medical Center, New-York, NY: Stanley Fahn, MD; Lucien Cote, MD; Paul Greene, MD; Cheryl Waters, MD; Pietro Mazzoni, MD, PhD; Blair Ford, MD; Elan Louis, MD MSc; Oren Levy, MD PhD; Ming Xin Tang, MD; Brian Rakitin, PhD; Ernest Roos, MD; Martha Orbe Riley, MD; Lency Rosado, MD; Carol Moskowitz, RNC; Tsvyatko Dorovski; Xinmin Liu, PhD; Sergey Kisselev, MS; Itsik Peer, PhD; Vladimir Vacic, PhD

Tel Aviv Sourasky Medical Center, Tel Aviv, Israel: Yaacov Balash, MD; Shabtai Hertzal, MD; Ariella Hillel, RN; Ziv Gan Or, PhD; Hila Kobo, MSc; Noa Bregman, MD; Meir Kestenbaum, MD; Talma Hendler, MD, PhD; Hedva Lehrman, MD; Einat Even Sapir, MD, PhD; Maayan Zelis, DMD; Michal Shtagman, RN; Eti Shimony; Michaela Victor; Marina Grumberg; Ora Assais

References

1. Ahmed Z, Asi YT, Sailer A, et al. The neuropathology, pathophysiology and genetics of multiple system atrophy. *Neuropathology and applied neurobiology*. 2012; 38:4–24. [PubMed: 22074330]
2. Iranzo A, Tolosa E, Gelpi E, et al. Neurodegenerative disease status and post-mortem pathology in idiopathic rapid-eye-movement sleep behaviour disorder: an observational cohort study. *The Lancet Neurology*. 2013; 12:443–453. [PubMed: 23562390]
3. Postuma RB, Gagnon JF, Vendette M, Fantini ML, Massicotte-Marquez J, Montplaisir J. Quantifying the risk of neurodegenerative disease in idiopathic REM sleep behavior disorder. *Neurology*. 2009; 72:1296–1300. [PubMed: 19109537]
4. Schenck CH, Boeve BF, Mahowald MW. Delayed emergence of a parkinsonian disorder or dementia in 81% of older men initially diagnosed with idiopathic rapid eye movement sleep behavior disorder: a 16-year update on a previously reported series. *Sleep medicine*. 2013; 14:744–748. [PubMed: 23347909]
5. Boeve BF, Silber MH, Ferman TJ, et al. Clinicopathologic correlations in 172 cases of rapid eye movement sleep behavior disorder with or without a coexisting neurologic disorder. *Sleep medicine*. 2013; 14:754–762. [PubMed: 23474058]
6. Postuma RB, Aarsland D, Barone P, et al. Identifying prodromal Parkinson's disease: pre-motor disorders in Parkinson's disease. *Movement disorders : official journal of the Movement Disorder Society*. 2012; 27:617–626. [PubMed: 22508280]
7. Postuma RB, Gagnon JF, Montplaisir J. Rapid eye movement sleep behavior disorder as a biomarker for neurodegeneration: the past 10 years. *Sleep medicine*. 2013; 14:763–767. [PubMed: 23058689]
8. Stiasny-Kolster K, Mayer G, Schafer S, Moller JC, Heinzel-Gutenbrunner M, Oertel WH. The REM sleep behavior disorder screening questionnaire--a new diagnostic instrument. *Movement disorders : official journal of the Movement Disorder Society*. 2007; 22:2386–2393. [PubMed: 17894337]
9. Marder K, Tang M, Alcalay R, et al. Age Specific Penetrance of the LRRK2 G2019S Mutation in the Michael J Fox Ashkenazi Jewish (AJ) LRRK2 Consortium (S17.002). *Neurology*. 2014; 82:S17.002.

10. Ozelius LJ, Senthil G, Saunders-Pullman R, et al. LRRK2 G2019S as a cause of Parkinson's disease in Ashkenazi Jews. *The New England journal of medicine*. 2006; 354:424–425. [PubMed: 16436782]
11. Stiasny-Kolster K, Sixel-Doring F, Trenkwalder C, et al. Diagnostic value of the REM sleep behavior disorder screening questionnaire in Parkinson's disease. *Sleep medicine*. 2015; 16:186–189. [PubMed: 25534709]
12. The Parkinson Progression Marker Initiative (PPMI). *Progress in neurobiology*. 2011; 95:629–635. [PubMed: 21930184]
13. Mahajan A, Rosenthal LS, Gamaldo C, et al. REM Sleep Behavior and Motor Findings in Parkinson's Disease: A Cross-sectional Analysis. *Tremor and other hyperkinetic movements (New York, NY)*. 2014; 4:245.
14. Healy DG, Falchi M, O'Sullivan SS, et al. Phenotype, genotype, and worldwide genetic penetrance of LRRK2-associated Parkinson's disease: a case-control study. *The Lancet Neurology*. 2008; 7:583–590. [PubMed: 18539534]
15. Mirelman A, Alcalay RN, Saunders-Pullman R, et al. Non-motor symptoms in healthy Ashkenazi Jewish carriers of the G2019S mutation in the LRRK2 gene. *Movement disorders : official journal of the Movement Disorder Society*. 2015 In Press.
16. Jankovic J, McDermott M, Carter J, et al. Variable expression of Parkinson's disease: a base-line analysis of the DATATOP cohort. *The Parkinson Study Group. Neurology*. 1990; 40:1529–1534. [PubMed: 2215943]
17. Alcalay RN, Mejia-Santana H, Mirelman A, et al. Neuropsychological performance in LRRK2 G2019S carriers with Parkinson's disease. *Parkinsonism & related disorders*. 2015; 21:106–110. [PubMed: 25434972]
18. Hu Y, Yu SY, Zuo LJ, et al. Parkinson disease with REM sleep behavior disorder: features, alpha-synuclein, and inflammation. *Neurology*. 2015; 84:888–894. [PubMed: 25663225]
19. Saunders-Pullman R, Mirelman A, Wang C, et al. Olfactory identification in LRRK2 G2019S mutation carriers: a relevant marker? *Annals of clinical and translational neurology*. 2014; 1:670–678. [PubMed: 25493281]
20. Gagnon JF, Vendette M, Postuma RB, et al. Mild cognitive impairment in rapid eye movement sleep behavior disorder and Parkinson's disease. *Annals of neurology*. 2009; 66:39–47. [PubMed: 19670440]
21. Marion MH, Qurashi M, Marshall G, Foster O. Is REM sleep behaviour disorder (RBD) a risk factor of dementia in idiopathic Parkinson's disease? *Journal of neurology*. 2008; 255:192–196. [PubMed: 18217187]
22. Naismith SL, Terpening Z, Shine JM, Lewis SJ. Neuropsychological functioning in Parkinson's disease: differential relationships with self-reported sleep-wake disturbances. *Movement disorders : official journal of the Movement Disorder Society*. 2011; 26:1537–1541. [PubMed: 21469205]
23. Sinforiani E, Zangaglia R, Manni R, et al. REM sleep behavior disorder, hallucinations, and cognitive impairment in Parkinson's disease. *Movement disorders : official journal of the Movement Disorder Society*. 2006; 21:462–466. [PubMed: 16228998]
24. Vendette M, Gagnon JF, Decary A, et al. REM sleep behavior disorder predicts cognitive impairment in Parkinson disease without dementia. *Neurology*. 2007; 69:1843–1849. [PubMed: 17984452]
25. Kumru H, Santamaria J, Tolosa E, Iranzo A. Relation between subtype of Parkinson's disease and REM sleep behavior disorder. *Sleep medicine*. 2007; 8:779–783. [PubMed: 17904419]
26. Lee JE, Kim KS, Shin HW, Sohn YH. Factors related to clinically probable REM sleep behavior disorder in Parkinson disease. *Parkinsonism & related disorders*. 2010; 16:105–108. [PubMed: 19716333]
27. Postuma RB, Gagnon JF, Vendette M, Charland K, Montplaisir J. REM sleep behaviour disorder in Parkinson's disease is associated with specific motor features. *Journal of neurology, neurosurgery, and psychiatry*. 2008; 79:1117–1121.

28. Postuma RB, Gagnon JF, Vendette M, Charland K, Montplaisir J. Manifestations of Parkinson disease differ in association with REM sleep behavior disorder. *Movement disorders : official journal of the Movement Disorder Society*. 2008; 23:1665–1672. [PubMed: 18709686]
29. Postuma RB, Lanfranchi PA, Blais H, Gagnon JF, Montplaisir JY. Cardiac autonomic dysfunction in idiopathic REM sleep behavior disorder. *Movement disorders : official journal of the Movement Disorder Society*. 2010; 25:2304–2310. [PubMed: 20669316]
30. Pouloupoulos M, Cortes E, Vonsattel JP, et al. Clinical and pathological characteristics of LRRK2 G2019S patients with PD. *Journal of molecular neuroscience : MN*. 2012; 47:139–143. [PubMed: 22194196]
31. Kalia LV, Lang AE, Hazrati LN, et al. Clinical Correlations With Lewy Body Pathology in LRRK2-Related Parkinson Disease. *JAMA neurology*. 2015; 72:100–105. [PubMed: 25401511]
32. Gaig C, Vilas D, Infante J, et al. Nonmotor symptoms in LRRK2 G2019S associated Parkinson's disease. *PloS one*. 2014; 9:e108982. [PubMed: 25330404]
33. Ruiz-Martinez J, Gorostidi A, Goyenechea E, et al. Olfactory deficits and cardiac 123I-MIBG in Parkinson's disease related to the LRRK2 R1441G and G2019S mutations. *Movement disorders : official journal of the Movement Disorder Society*. 2011; 26:2026–2031. [PubMed: 21611983]
34. Trinh J, Amouri R, Duda JE, et al. Comparative study of Parkinson's disease and leucine-rich repeat kinase 2 p.G2019S parkinsonism. *Neurobiology of aging*. 2014; 35:1125–1131. [PubMed: 24355527]
35. Alcalay RN, Mirelman A, Saunders-Pullman R, et al. Parkinson disease phenotype in Ashkenazi Jews with and without LRRK2 G2019S mutations. *Movement disorders : official journal of the Movement Disorder Society*. 2013; 28:1966–1971. [PubMed: 24243757]
36. Eisensehr I, v Lindeiner H, Jager M, Noachtar S. REM sleep behavior disorder in sleep-disordered patients with versus without Parkinson's disease: is there a need for polysomnography? *Journal of the neurological sciences*. 2001; 186:7–11. [PubMed: 11412865]
37. Gagnon JF, Bedard MA, Fantini ML, et al. REM sleep behavior disorder and REM sleep without atonia in Parkinson's disease. *Neurology*. 2002; 59:585–589. [PubMed: 12196654]
38. Nomura T, Inoue Y, Kagimura T, Uemura Y, Nakashima K. Utility of the REM sleep behavior disorder screen questionnaire (RBDSQ) in Parkinson's disease patients. *Sleep Medicine*. 2011; 12:711–713. [PubMed: 21700495]

Table 1
Demographics and REM Sleep Behavior Disorder Screening Questionnaire (RBDSQ) Scores

n	IPD ^a	LRRK2-PD ^b	NMC ^c	NC-F ^d	Control ^e
144	142	117	93	40	
Women %	35.4% ^{cde}	45.8% ^{ce}	59.8% ^{ab}	50.5% ^{ae}	72.5% ^{abd}
Age years	64.9 ± 11.3 ^{cde}	67.2 ± 9.8 ^{cd}	53.3 ± 15.9 ^{abe}	52.8 ± 18.5 ^{abe}	69.8 ± 10.6 ^{abcd}
UPDRS-III	20.1 ± 11.8 ^{cde}	21.1 ± 13.1 ^{cde}	2.5 ± 3.5 ^{ab}	2.5 ± 3.5 ^{ab}	2.1 ± 3.4 ^{ab}
Age onset	58.0 ± 10.9	57.4 ± 11.0	-	-	-
PD Duration years	7.4 ± 7.5 ^b	10.0 ± 7.1 ^a	-	-	-
RBDSQ score median (IQR)	4.0 (2.0-7.0) ^{cde}	3.0 (2.0-5.0) ^{cde}	2.0 (1.0-3.0) ^{ab}	2.0 (1.0-4.0) ^{ab}	2.0 (1.0-4.0) ^{ab}
RBD+ %	30.6% ^{bcd}	19.7% ^{ac}	6.0% ^{abd}	20.4% ^{ac}	15.0% ^a

Odds Ratios for Abnormal RBDSQ Scores Among

All Groups**	OR	95% CI	P
n = 536			
Age	1.01	.99-1.02	0.56
Sex (Women)	0.64	.41-1.01	0.06
Site (NY)	1.57	.97-2.54	0.07
Group:			
Control (ref)	---	---	---
IPD	2.75	.03-7.35	0.04
LRRK2-PD	1.50	.56-4.04	0.42
NMC	0.46	.14-1.54	0.21
NC-F	1.89	.65-5.51	0.25
<i>Further group comparisons*</i>			
IPD vs. LRRK2-PD	1.83	.05-3.18	0.03
NMC vs. LRRK2-PD	0.31	.13-0.75	0.01
NC-F vs. LRRK2-PD	0.80	.40-1.59	0.52
NMC vs. NC-F	0.25	.10-0.61	<0.01

Odds Ratios for RBDSQ Restricted to PD Groups

n = 275	OR	95% CI	P
IPD vs. LRRK2-PD	2.38	.29–4.38	<0.01
Age	1.01	.99–1.04	0.34
Sex (Women)	0.54	.29–1.00	0.05
Site (NY)	1.37	.75–2.52	0.31
Duration	1.08	.03–1.14	<0.01
UPDRS-III	0.99	.97–1.02	0.20

* Additional group comparisons were deduced from the main model; Values provided are number (%) or mean ± SD, as appropriate, unless otherwise stated; Superscripts denote significant differences from the superscripted group (p<0.05); ORs from GEE models adjusted for age, site (NY vs. Israel), gender and family membership Definitions: LRRK2-PD=LRRK2 G2019S mutation carriers with PD; IPD=idiopathic PD; NMC=LRRK2 mutation carriers without PD; NC-F=non-carrier family members; UPDRS-III=Unified Parkinson Disease Rating Scale Motor Score; RBDSQ=REM Sleep Behavior Disorder Screening Questionnaire; RBD+=total RBDSQ score 5 (non-PD) and total RBDSQ score 6 (PD).

** In post-hoc analysis using the main GEE model to assess site differences, all differences from the main model were maintained when limited to participants from either NY or TA, except that NMC from NY were not different from LRRK2-PD (OR=0.34, p=0.07) or NC-F (OR=0.30, p=0.06) and, although the magnitude of the OR was maintained, in TA the odds of RBD was not significantly reduced in LRRK2-PD vs. IPD (OR=0.73, p=0.49).

Table 2

Features associated with RBDSQ Cut-scores in Parkinson disease

	RBDSQ+	RBDSQ-	<i>P</i>
LRRK2-PD (n=142)	28 (19.7%)	114 (80.3%)	---
Women	9 (32.1%)	56 (49.1%)	0.14
Age	68.2 ± 9.1	67.0 ± 10.0	0.72
Duration	14.6 ± 7.6	8.8 ± 6.5	<0.01
UPSIT Score (125)	20.9 ± 8.6	23.5 ± 8.6	0.19
Hyposmia	15 (60.0%)	53 (53.0%)	0.53
UPDRS-III (135)	24.2 ± 12.9	20.3 ± 13.1	0.07
PIGD phenotype (133)	22 (81.5%)	67 (63.2%)	0.11
MoCA Score (131)	24.8 ± 3.6	25.4 ± 2.8	0.67
STAI (42)			
State	43.4 ± 11.8	33.7 ± 13.4	0.03
Trait	48.8 ± 14.6	33.4 ± 11.7	<0.01
IPD (n=144)	44 (30.6%)	100 (69.4%)	---
Women	11 (25.0%)	40 (40.0%)	0.08
Age	67.5 ± 10.1	63.8 ± 11.7	0.06
Duration	7.8 ± 6.6	7.2 ± 7.9	0.48
UPSIT Score (117)	17.1 ± 7.6	19.5 ± 6.9	<0.05
Hyposmia	26 (86.7%)	65 (74.1%)	0.21
UPDRS-III (142)	21.2 ± 12.0	19.7 ± 11.8	0.51
PIGD phenotype (141)	31 (70.5%)	45 (46.4%)	<0.01
MoCA Score (134)	24.7 ± 3.2	25.1 ± 3.9	0.28
STAI (36)			
State	40.2 ± 7.1	34.6 ± 10.5	0.06
Trait	38.6 ± 9.8	36.9 ± 12.6	0.48

Values provided are number (%) or mean ± SD, as appropriate, unless otherwise stated.

RBD+=total RBDSQ score \leq 5 (non-PD) and total RBDSQ score \geq 6 (PD); LRRK2-PD = *LRRK2* G2019S mutation carriers with PD; IPD = idiopathic PD; UPSIT = University of Pennsylvania Smell Identification Test; Hyposmic = UPSIT percentile score $<$ 15; UPDRS-III = Unified Parkinson Disease Rating Scale Motor Score; PIGD = Postural Instability and Gait Disorder subtype; MoCA = Montreal Cognitive Assessment; STAI = State-Trait Anxiety Inventory