Lawrence Berkeley National Laboratory

Recent Work

Title

NOTE ON THE ION EXCHANGE SEPARATION OF EUROPIUM, GADOLINIUM, AND TERBIUM

Permalink

<https://escholarship.org/uc/item/2xd2w23g>

Authors

Higgins, G.R.

K.

Street, Jr.

Publication Date

1950-04-24

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

UNCLASSIFIED

Contract No. W-7405-eng-48

Note on the Ion Exchange Separation of

Europium, Gadolinium, and Terbium

G. H. Higgins and K. Street, Jr.

April 24, 1950

Berkeley, California

-2-

<u>INSTALLATION:</u>	<u>No. of Copies</u>
Argonne National Laboratory	8
Armed Forces Special Weapons Project	1
Atomic Energy Commission, Washington	2
Battelle Memorial Institute	1
Brush Beryllium Company	1
Brookhaven National Laboratory	8
Bureau of Medicine and Surgery	1
Bureau of Ships	1
Carbide & Carbon Chemicals Div., Union Carbide & Carbon Corp. (K-25 Plant)	4
Carbide & Carbon Chemicals Div., Union Carbide & Carbon Corp. (Y-12 Plant)	4
Chicago Operations Office	1
Cleveland Area Office, AEC	1
Columbia University (J. R. Dunning)	1
Columbia University (G. Failla)	1
Dow Chemical Company	1
H. K. Ferguson Company	1
General Electric, Richland	6
Harshaw Chemical Corporation	1
Idaho Operations Office	1
Iowa State College	2
Kansas City Operations Branch	1
Kellex Corporation	2
Knolls Atomic Power Laboratory	4
Los Alamos Scientific Laboratory	3
Mallinckrodt Chemical Works	1
Massachusetts Institute of Technology (A. Gaudin)	1
Massachusetts Institute of Technology (A. R. Kaufmann)	1
Mound Laboratory	3
National Advisory Committee for Aeronautics	2
National Bureau of Standards	2
Naval Radiological Defense Laboratory	2
New Brunswick Laboratory	1
New York Operations Office	5
North American Aviation, Inc.	1
Oak Ridge National Laboratory	8
Patent Branch (Washington)	1
Rand Corporation	1
Sandia Laboratory	1
Santa Fe Operations Office	1
Sylvania Electric Products, Inc.	1
Technical Information Division (Oak Ridge)	15
USAF, Air Surgeon (R. H. Blount)	1
USAF, Director of Armament (C. I. Browne)	1
USAF, Director of Plans and Operations (R. L. Applegate)	1
USAF, Director of Research and Development (F. W. Bruner, ... and R. J. Mason)	2
USAF, Eglin Air Force Base (K. K. Compton)	1

-2a-

<u>INSTALLATION:</u>	<u>No. of Copies</u>
USAF, Kirtland Air Force Base (H. G. Montgomery, Jr.)	1
USAF, Maxwell Air Force Base (F. N. Moyers)	1
USAF, NEPA Office	2
USAF, Office of Atomic Energy (A. A. Fickel, H. C. Donnelly)	2
USAF, Offutt Air Force Base (H. R. Sullivan, Jr.)	1
USAF, Wright-Patterson Air Force Base (Rodney Nudenberg)	1
U. S. Army, Atomic Energy Branch (A. W. Betts)	1
U. S. Army, Army Field Forces (James Kerr)	1
U. S. Army, Commanding General, Chemical Corps Technical Command (J. A. MacLaughlin thru Mrs. G. Benjamin)	1
U. S. Army, Chief of Ordnance (A. R. Del Campo)	1
U. S. Army, Commanding Officer Watertown Arsenal (Carroll H. Deitrick)	1
U. S. Army, Director of Operations Research (Ellis Johnson)	1
U. S. Army, Office of Engineers (B. D. Jones)	1
U. S. Army, Office of the Chief Signal Officer (Curtis T. Clayton thru George C. Hunt)	1
U. S. Army, Office of the Surgeon General (W. S. Stone)	1
U. S. Geological Survey (T. B. Nolan)	1
U. S. Public Health Service	1
University of California at Los Angeles	1
University of California Radiation Laboratory	5
University of Rochester	2
University of Washington	1
Western Reserve University	2
Westinghouse Electric Company	4
Univ. of Rochester, Physics Department (R. E. Marshak)	1
Total	<u>145</u>

Information Division
 Radiation Laboratory
 University of California
 Berkeley, California

Note on the Ion Exchange Separation of Europium, Gadolinium, and Terbium

G. H. Higgins and K. Street, Jr.
Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California
April 24, 1950

In connection with the general problem of comparing the behavior of the actinide elements with their rare earth homologues, it was desirable to know the relative rates of elution of the rare earth triad europium, gadolinium, and terbium on eluting with citrate solutions from Dowex-50 cation resin. Ketelle and Boyd¹

¹ B. H. Ketelle and G. E. Boyd, J. Am. Chem. Soc. 69, 2800 (1947)

in Fig. 6c show an elution (run at 100°C) of the heavy rare earth elements from Dowex-50 cation resin using 5% ammonium citrate solution of pH 3.25. In their experiment the activity peak that eluted in about 0.7 the time required to elute their europium peak is labelled Gd thus implying a large separation between europium and gadolinium. B. B. Cunningham and H. G. Hicks² ran europium and gadolinium under

² B. B. Cunningham and H. G. Hicks, unpublished work.

somewhat different conditions (room temperature and pH 3.05 citrate) and found practically no separation of europium and gadolinium. In addition, J. O. Rasmussen³

³ J. O. Rasmussen, unpublished work.

ran europium, gadolinium, and terbium under still different conditions (75°C and pH 3.2 citrate) and again found essentially no separation of europium and gadolinium. In addition, he found that terbium was eluted well ahead of europium and gadolinium. It thus appeared that there was a real discrepancy between this later work and that of Ketelle and Boyd.

In order to check this point a run was made under conditions very close to those used by Ketelle and Boyd. Fig. 1 shows the results of an elution of yttrium,

Fig. 1-- Elution of Y, Tb, Gd, and Eu with
pH 3.25 ammonium citrate at 97°C.

terbium, gadolinium, and europium made at 97°C . using 5% ammonium citrate solution of pH 3.25. Tracer Y^{91} , Gd^{154} , and $\text{Eu}^{152-154}$ were used and in addition approximately 200 ug. amounts of stable terbium, gadolinium, and europium were added to make spectrographic identification of the peaks possible. The column used was 20 cm. long and 5 mm. in diameter and a flow rate of $0.3 \text{ ml/cm}^2/\text{min}$. was maintained. The Eu and Gd activities appear as a single broad peak, however spectrographic analyses make possible the construction of the dashed curves for Eu, Gd, and Tb. It is evident on comparing this elution with Fig. 6c of Ref. 1 that the activity labelled Gd by Ketelle and Boyd is very likely a terbium activity and that any gadolinium activities they may have had are under the front of the europium peak.

It should be noted that the very small separation of gadolinium from europium and the much greater separation of terbium from gadolinium agrees very well with the expected effect of the half-filled 4f electron configuration as was first pointed out by Boyd.¹

This work was performed under the auspices of the U. S. Atomic Energy Commission.

FIG. 1

Mu 160