

Lawrence Berkeley National Laboratory

LBL Publications

Title

Taxonomic and environmental distribution of bacterial amino acid auxotrophies

Permalink

<https://escholarship.org/uc/item/2zm68073>

Journal

Nature Communications, 14(1)

ISSN

2041-1723

Authors

Ramoneda, Josep
Jensen, Thomas BN
Price, Morgan N
et al.

Publication Date

2023

DOI

10.1038/s41467-023-43435-4

Peer reviewed

Taxonomic and environmental distribution of bacterial amino acid auxotrophies

Received: 7 July 2023

Accepted: 8 November 2023

Published online: 22 November 2023

 Check for updates

Josep Ramoneda ¹✉, Thomas B. N. Jensen ^{1,2}, Morgan N. Price ³,
Emilio O. Casamayor ⁴ & Noah Fierer ^{1,5}✉

Many microorganisms are auxotrophic—unable to synthesize the compounds they require for growth. With this work, we quantify the prevalence of amino acid auxotrophies across a broad diversity of bacteria and habitats. We predicted the amino acid biosynthetic capabilities of 26,277 unique bacterial genomes spanning 12 phyla using a metabolic pathway model validated with empirical data. Amino acid auxotrophy is widespread across bacterial phyla, but we conservatively estimate that the majority of taxa (78.4%) are able to synthesize all amino acids. Our estimates indicate that amino acid auxotrophies are more prevalent among obligate intracellular parasites and in free-living taxa with genomic attributes characteristic of ‘streamlined’ life history strategies. We predicted the amino acid biosynthetic capabilities of bacterial communities found in 12 unique habitats to investigate environmental associations with auxotrophy, using data compiled from 3813 samples spanning major aquatic, terrestrial, and engineered environments. Auxotrophic taxa were more abundant in host-associated environments (including the human oral cavity and gut) and in fermented food products, with auxotrophic taxa being relatively rare in soil and aquatic systems. Overall, this work contributes to a more complete understanding of amino acid auxotrophy across the bacterial tree of life and the ecological contexts in which auxotrophy can be a successful strategy.

Microbial auxotrophy (i.e. the inability of microorganisms to synthesize the compounds they require for growth) has been identified in taxa isolated from many environments^{1–11}. The loss of biosynthetic genes can, under certain conditions, confer a selective advantage due to the corresponding reduction in metabolic and energetic costs^{12–14}. Auxotrophy can be particularly advantageous when the essential metabolites can be readily obtained from the surrounding environment, or from nearby cells, leading to the expectation that in environments with abundant nutrients or close-range interactions, auxotrophy will be an adaptive trait¹⁵. For example, under laboratory conditions, *E. coli* supplied with amino acids can evolve amino acid

auxotrophies in under 2000 generations and outcompete its ancestral prototrophic relatives (i.e. taxa that have the ability to synthesize all amino acids¹⁶). Another example are obligate intracellular parasites, which have among the smallest genomes of all bacteria and are commonly auxotrophic for vitamins and certain amino acids available from their host¹⁷.

Microorganisms can be auxotrophic for multiple types of metabolites. The most frequent auxotrophies are those for vitamins^{3,6,18–21}, amino acids^{22–27}, and diverse cofactors (e.g. heme groups²⁸). Here, we focus on amino acid auxotrophies because amino acids can be important both as energy sources and as building blocks of the

¹Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado, Boulder, CO, USA. ²Center for Microbial Communities, Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark. ³Environmental Genomics and Systems Biology, Lawrence Berkeley National Laboratory, Berkeley, CA, USA. ⁴Spanish Research Council (CSIC), Center for Advanced Studies of Blanes (CEAB), Blanes, Spain. ⁵Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO, USA. ✉e-mail: ramoneda.massague@gmail.com; noah.fierer@colorado.edu

proteome, the costs associated with synthesizing amino acids are reasonably well-constrained¹², and because the amino acid biosynthetic capabilities of many bacteria can be inferred with recent improvements in our understanding of biosynthetic pathways and the bioinformatic tools to infer amino acid auxotrophies^{29–32}. In synthetic assemblages, amino acid cross-feeding can be an ecologically stable strategy when interacting partners complement each other in their metabolic capabilities³³. Thus, it is often assumed that auxotrophic interactions and the cross-feeding of amino acids are a key factor structuring microbial communities¹⁵. While there is limited evidence for auxotrophy-mediated amino acid exchange in microbial communities found in natural systems, previous work has suggested that this phenomenon likely occurs in microbial consortia responsible for hydrocarbon degradation⁸, methanogenesis³⁴, and anammox³⁵.

Auxotrophy is expected to be more common in habitats where the essential metabolites are more readily available and diffusible. For example, protein-rich environments such as dairy products contain a high availability of amino acids³⁶, and are dominated by well-known amino acid auxotrophs such as bacteria from the genus *Lactobacillus*²⁴. The physical structure of microbial habitats can also influence the availability of essential metabolites. Auxotrophies may be particularly prevalent among bacteria living in biofilms or in well-mixed systems, where metabolites can more readily be exchanged between taxa primarily due to their spatial proximity^{37,38}. Generally, we would expect that communities from different environments should vary with respect to the prevalence of auxotrophies due to differences in the amounts and types of metabolites available. For example, we would expect bacterial amino acid auxotrophs to be more common in host-associated systems where amino acid availability is reasonably high, such as the human gut^{22,26,27}. However, the broader prevalence of auxotrophic bacteria in other types of microbial systems (including soil and aquatic systems) remains largely undetermined.

Using genomic information alone, it is possible to predict the metabolic capabilities of many bacterial taxa^{31,32,39–41}. These metabolic pathway models rely on a priori knowledge of the genes involved in the metabolic pathways of interest and allow for the prediction of auxotrophy in any taxon for which high quality genomic information is available. For example, D'Souza et al.¹³ used genomic information from 949 full genomes to estimate that 76% of bacterial taxa were auxotrophic for at least one essential metabolite. The frequent application of metabolic pathway models contrasts with the paucity of experiments that empirically validate the predictions of these models. The experimental validation of auxotrophy typically requires challenging and time consuming in vitro assays that are, by definition, difficult to conduct on the large fraction of bacterial taxa that remain uncultured⁴². Those studies that have attempted to empirically validate predictions of auxotrophy show that genome-based models largely underestimate the metabolic capabilities of bacterial taxa^{29–31,43}. For example, Price et al.²⁹ studied 10 bacterial genera that were predicted to be auxotrophic for several amino acids, but found that these taxa could grow on minimal media in the absence of externally supplied amino acids. Using genome-wide mutant fitness data, the authors identified genes for 9 of the 11 missing steps in amino acid biosynthesis. While many biosynthetic pathways remain poorly understood⁴⁴, new empirical findings and conservative bioinformatic approaches make it possible to infer bacterial auxotrophies^{31,32,43}.

Here, we predicted the prevalence of amino acid auxotrophies across a broad diversity of bacteria by analyzing 26,277 genomes representing 12 different bacterial phyla. We also compared the predicted prevalence of amino acid auxotrophies from 13,523 representative taxa found in 12 different habitats, ranging from soils, freshwater, and marine waters, to engineered systems such as activated sludge and food products, and to host-associated systems including the human gut, skin, and plant leaf surfaces. We validated the predictions of a metabolic pathway model of bacterial auxotrophy³¹ by

compiling empirical information on the metabolic capabilities of diverse bacterial taxa to minimize the overestimation of auxotrophy. Finally, we evaluated which genomic features are more frequently associated with bacterial amino acid auxotrophy to characterize the broader life history strategies that differentiate amino acid auxotrophs from prototrophs. By covering a broad range of taxa and habitats we provide a comprehensive view on the taxonomic and environmental signatures of amino acid auxotrophies in bacteria.

Results and discussion

Model validation

To test our ability to infer amino acid auxotrophy from genomic analyses, we first validated our model after predicting the amino acid biosynthesis capabilities of 171 taxa that can make all amino acids (prototrophs). Doing so allowed us to quantify how many genes need to be missing from an amino acid biosynthesis pathway in a certain organism to be considered auxotrophic for that amino acid. To minimize the overestimation of auxotrophy, we found that at least 40% of the genes needed to be missing in a given amino acid biosynthesis pathway to obtain a very low 0.4% rate of false positives (i.e. erroneously predicted auxotrophies). This means that our model predictions were correct in ~99% of the cases in which an organism was able to synthesize a given amino acid. Only for serine and cysteine (4% error) did our model incorrectly predict amino acid auxotrophies (i.e. inferring auxotrophies when the taxa were actually capable of synthesizing those amino acids, Supplementary Fig. 1). In the case of serine, 6 of the 7 genomes that were misclassified as auxotrophic belonged to taxa from the phylum Desulfobacteria, which are typically sulfate-reducers (the remaining genome belonged to a green sulfur bacterium from the Chlorobiaceae, Bacteroidetes; Supplementary Data 1). A group of sulfate-reducing bacteria, including *Desulfovibrio* and related genera, appear to produce serine from pyruvate or related compounds as in the standard pathway⁴⁵, but the genes involved are not known. The phylum Desulfobacteria was not included in the analyses presented below. Similarly, all the genomes that were misclassified as cysteine auxotrophs belonged to phyla not included in this study such as the Desulfobacteria and the Aquificae, also characterized by having sulfur-related metabolisms (Supplementary Data 1). We found that these genomes contained the cysteine synthase gene (*cysK*), which makes it unlikely that these taxa synthesize cysteine via alternative pathways. Together, these results suggest that our decision to require at least 40% of the genes to be missing to infer auxotrophies for cysteine and serine auxotrophy primarily affected less abundant phyla not included in the study.

We then quantified the rate of false negatives (i.e. inferring prototrophy for amino acids that taxa cannot synthesize) using genomes from taxa with experimentally determined auxotrophies compiled from the literature (Supplementary Table 1). Applying our threshold that a minimum of 40% of genes from a pathway had to be missing to consider a genome auxotrophic for a given amino acid led to false negative rate of 20% (i.e. the proportion of amino acids in each genome for which our model predicted taxa to be prototrophic when they were auxotrophic, Supplementary Fig. 2). On a per genome basis (i.e. predicting whether a given genome is auxotrophic for 1 or more amino acids versus prototrophic), our model correctly infers prototrophy in 93% of the cases, and infers that a taxon is auxotrophic for at least 1 amino acid correctly in 95% of the cases. This means that, although the model tends to underestimate the number of amino acids that a given taxon is unable to synthesize, we can accurately identify when a taxon is generally auxotrophic or prototrophic. We recognize that our current understanding of amino acid biosynthesis pathway derives from taxa that have been cultured, and that improved knowledge beyond those taxa is required to improve our inferences of auxotrophies in particular groups.




Fig. 1 | Amino acid auxotrophy across the predominant bacterial phyla.

A Prevalence of amino acid auxotrophy in bacterial taxa from the most common phyla ($N = 26,277$ genomes). **B** Prevalence of amino acid auxotrophy in genomes derived from environmental metagenomes (MAGs) or single cells (SAGs) (Assembled), and in genomes obtained from bacterial isolates (Isolate). The mean number of amino acid auxotrophies in **A**, **B** is indicated with white diamonds. **C** Proportion of taxa that are auxotrophic for each of the 17 amino acids and chorismate out of the total number of auxotrophic taxa ($N = 3613$ genomes). Numbers in brackets in

panel A indicate the number of genomes for which we predicted amino acid auxotrophy, numbers in brackets in panel B indicate the number of assembled and isolate genomes included, and numbers in brackets in **C** indicate the subset of taxa within each phylum that were predicted to be auxotrophs for at least one amino acid. Letters in panel A indicate statistical differences ($P < 0.05$) between phyla based on Mann-Whitney U tests with P -values Bonferroni-corrected for multiple comparisons.

Previous genome-based studies have largely overestimated amino acid auxotrophy, despite mounting evidence that most of these inaccurate predictions come from knowledge gaps or from lack of awareness of alternative biosynthetic pathways^{29,30}. A number of studies have used focused culturing efforts to identify auxotrophies in experimental isolates^{24,25,29,32,46} and high-throughput culturing techniques make it possible to screen for bacterial growth across a wide range of media types^{47,48}. We recognize that our approach likely misses a number of auxotrophies, but it does provide a more conservative perspective on the actual amino acid biosynthesis capabilities of most bacterial taxa. The fact that we only found 19 taxa with genomic data available and known amino acid auxotrophy profiles highlights the difficulties of conducting in vitro experiments to confirm amino acid auxotrophies³². Future work could benefit from advances in high-throughput cultivation-based approaches to experimentally identify auxotrophies⁴⁹ and expand the datasets needed for validation of genome-based models⁵⁰. Dedicated efforts combining extensive media testing, whole genome sequencing, and comparative genomics will further reduce uncertainty around amino acid biosynthesis in bacteria. Until then, we are confident that our approach is conservative, recognizing that we are likely underestimating the occurrence of some amino acid auxotrophies.

Prevalence of amino acid auxotrophies in bacteria

We used our genome-based approach to predict amino acid auxotrophies in 26,277 bacterial taxa from the 12 phyla with >100 non-chimeric representative genomes estimated to be >95% complete in the Genome Taxonomy Database (GTDB, release 207)⁵¹. A large majority of taxa (78.4%), each represented by a single genome, were inferred to be able to synthesize all amino acids (i.e. were completely prototrophic; Fig. 1A). This prediction contrasts with the previous comprehensive study of amino acid auxotrophy in bacteria, which was based on 949 sequenced genomes with the authors reporting that only 24% of bacterial taxa were able to synthesize all amino acids¹³. There are many reasons this discrepancy may exist, but it does suggest that the GapMind predictive framework applied here yields a more conservative estimate of amino acid auxotrophies (as explained above) and is less likely to incorrectly infer auxotrophies when specific biosynthetic genes are not detected in genomes.

Even though our model estimated that 78.4% of the 26,277 bacterial taxa were deemed to be completely prototrophic, there was a high degree of variation in the distribution of amino acid auxotrophies across bacterial taxa. We observed the lowest proportion of auxotrophs in the Cyanobacteria (0.9%) and the highest proportion in the Tenericutes (99.2%). The phyla with the largest numbers of representative genomes all contained large numbers of both auxotrophs

and prototrophs, with members of the Actinobacteria (8.6%) and Proteobacteria (10.4%) having significantly lower proportions of auxotrophs than Bacteroidetes (37.2%) and Firmicutes (37.0%) (Fig. 1A; Supplementary Table 2). Our finding that the Bacteroidetes and Firmicutes phyla contain higher proportions of auxotrophs than most other phyla is in agreement with previous work^{13,15,30,52}. Similarly, our finding that most Cyanobacteria are prototrophic for all amino acids is in line with previous work suggesting that Cyanobacteria are able to synthesize all amino acids⁵³ and our observation that only 0.8% of the Tenericutes are prototrophic is to be expected given that auxotrophies are widely observed in this group, which is mostly represented by obligate, intracellular parasites^{54,55}.

Our analysis of the prevalence of auxotrophies at the family level emphasizes the broad taxonomic distribution of auxotrophs. We predicted the prevalence and identity of amino acid auxotrophies across the predominant bacterial families (51 families from the 12 phyla with at least 100 available genomes; Supplementary Fig. 3). Less than a quarter (21.6%) of the families contained more auxotrophic than prototrophic taxa. The Mycoplasmataceae was the only family where all bacterial members were predicted to be auxotrophs, as expected for this group of intracellular parasites that obtain required nutrients from their host⁵⁶. All families where over 80% of their members were predicted to be auxotrophs contained predominantly host-associated taxa, including Coriobacteriaceae⁵⁷, Lactobacillaceae⁵⁸, and Streptococcaceae⁵⁹ (Supplementary Fig. 3). On the opposite end of the spectrum, 54.9% of the 51 families had less than 10% auxotrophic taxa (Supplementary Fig. 3). The least auxotrophic families were the Streptomycetaceae (0.1%), Paenibacillaceae (0.2%), and the Pseudomonadaceae (0.3%).

Associations between amino acid auxotrophy, genome size, and genome origin

We found that the prevalence of auxotrophs was significantly lower for genomes derived from bacterial isolates compared to those genomes assembled from environmental metagenomes (MAGs) and single cells (SAGs) (Mann-Whitney U, $p < 0.001$; Fig. 1B). Note that all MAG/SAG genomes included in the study were thoroughly filtered for completeness (>95% complete), absence of chimerism, and were required to contain an assembled 16 S rRNA gene. We also found that MAGs/SAGs, >95% of which represent uncultivated taxa, had generally smaller genomes and higher predicted minimal doubling times than genomes derived from cultured isolates (Welch two-sample t-test, $p < 0.001$; Supplementary Fig. 4A, B), in agreement with previous findings⁶⁰. Crucially, the number of amino acids that taxa were unable to synthesize was inversely proportional to their genome size ($r = -0.40$, $p < 0.001$; Supplementary Fig. 4C). This general negative association between genome size and auxotrophy across phyla suggests that the higher number of auxotrophies observed in MAGs/SAGs is likely due to evolutionary processes associated with genome size reduction, and not potential annotation or completeness biases. Isolate-derived genomes had higher completeness (99.2% average completeness) than those from MAGs/SAGs (97.6%), but this difference alone is likely insufficient to result in a sizeable difference in the number of estimated auxotrophies. We also verified that the potential impact of genome completeness on predicted amino acid auxotrophy was minor based on the weak correlation between genome completeness and the number of auxotrophies per genome (within MAGs/SAGs $r = -0.07$; within isolates $r = -0.14$). We also verified that the phyla with the highest proportions of auxotrophic taxa did not typically contain a larger proportion of MAG/SAG genomes (Supplementary Fig. 5). These results suggest that many bacterial taxa are not readily cultivated because they have life history strategies characterized by slow growth and complex external nutrient requirements that impair growth under laboratory conditions⁴². This seems unsurprising as phyla with low proportions of auxotrophs (e.g. Cyanobacteria or Actinobacteria) tend

to have larger genomes compared to phyla with higher proportions of auxotrophs⁶¹, and genome reduction by loss of biosynthetic genes has previously been associated with auxotrophy across bacterial groups⁶² (see below for further discussion of this point).

Amino acid auxotrophies associated with specific bacterial phyla

We next investigated which specific amino acid auxotrophies were most common across bacteria. Auxotrophic bacteria were most frequently auxotrophic for leucine (58.5%), valine (57.8%), and isoleucine (54.9%) (branched-chain amino acids), and were the least likely to be auxotrophic for asparagine (7.0%), glycine (7.2%), and glutamine (9.3%) (Fig. 1C). The availability of branched-chain amino acids controls the virulence gene expression in diverse host-associated bacteria, and auxotrophy for these amino acids has been suggested to be an adaptation to regulate bacterial metabolic activity with changes in external nutrient levels⁶³. Generally, the amino acid auxotrophic profiles were primarily dictated by the identity of the amino acids rather than the taxonomic affiliation of the genomes in question, meaning that most phyla were more auxotrophic for the same amino acids (Fig. 1C). There were some exceptions to this pattern. For example, in the Actinobacteria (91.4% prototrophs) 61.6% of the auxotrophic taxa could not synthesize tryptophan (Fig. 1C). Notably, 41.6% of those actinobacterial tryptophan auxotrophs belonged to the gut-associated genera *Collinsella* and *Olsenella*⁶⁴. We verified that the number of genes in a given amino acid biosynthesis pathway was not strongly correlated with the proportion of auxotrophic taxa for that amino acid ($r = -0.43$, $p = 0.100$). Note that the predicted auxotrophy for serine in the Deinococcus-Thermus phylum is likely due to a novel phosphoserine phosphatase in *Thermus thermophilus*, which has not been incorporated into GapMind⁶⁵.

In contrast to previous studies, we did not find a significant correlation between the proportion of auxotrophic taxa for each amino acid and the metabolic cost of each amino acid calculated from the number of P-bonds required to synthesize a given amino acid ($r = -0.24$, $p = 0.4$; Supplementary Fig. 6A)¹². When we explored this relationship within each of the predominant phyla, we only found a significant correlation in the phylum Spirochaetes ($r = 0.71$, $p = 0.001$; Supplementary Fig. 6B).

Prevalence of amino acid auxotrophy across habitats

We analyzed representative genomes from bacterial taxa found across 12 different habitats to assess general patterns in amino acid auxotrophies (Table 1). The habitats included in our analyses covered a broad range of habitat types, including terrestrial (bulk soil, rhizosphere soil), aquatic (freshwater lakes, marine surface waters), engineered (activated sludge and residential plumbing), host-associated habitats (phyllosphere, human gut, human skin, and human oral cavity), and fermented foods (cheese and sourdough). We identified between 148 (cheese) and 2949 (phyllosphere) representative genomes per habitat (13,523 genomes in total) (Table 1, see Methods). The proportion of taxa that were capable of synthesizing all amino acids was highly variable across habitats. More than 95% of bacteria found in rhizosphere soils, residential plumbing, and bulk soils were capable of synthesizing all amino acids (Fig. 2A; Table 1). In contrast, less than half of the bacteria in the human gut (41.6%) and oral cavity (24.7%) were prototrophic for all amino acids (Fig. 2A). The habitat-specific patterns in auxotrophy prevalence were still evident even when we restricted our analyses to the phylum Proteobacteria, the most ubiquitous phylum across habitats and a phylum with biosynthetic pathways that have been relatively well-studied³¹. These proteobacterial-specific analyses also show that the human gut and oral cavity were inferred to have the highest proportions of auxotrophic taxa (Supplementary Fig. 7).

The differences in the prevalence of amino acid auxotrophies across different habitats matched differences in the taxonomic

Table 1 | Attributes of the datasets included in the study

Habitat	#Samples	#Genomes	% ASVs with representative genomes	Proportional abundance of all ASVs with representative genomes (%)	% prototrophs	% with only 1 amino acid auxotrophy	Refs.
Rhizosphere	230	2886	48.3	70.0	97.5	23.6	106
Residential plumbing	471	733	57.5	84.5	97.4	42.1	107
Soil	255	448	18.1	35.3	96.4	62.5	108
Freshwater	299	688	45.5	57.4	95.9	17.9	109
Marine	365	772	32.9	33.9	91.5	65.2	110
Activated sludge	514	1369	32.2	54.2	89.6	30.1	111
Phyllosphere	128	2949	43.6	41.9	89.4	9.9	112
Cheese	98	148	69.6	87.6	81.8	18.5	113
Sourdough	421	218	93.8	82.7	75.2	5.6	114
Human skin	335	1778	93.9	98.1	69.7	10.8	115
Human gut	350	979	66.3	89.1	41.6	25.7	116
Human oral cavity	347	571	83.7	97.9	24.7	9.1	115

Reference genomes were obtained by matching the 16 S rRNA gene amplicon sequences to the Genome Taxonomy Database (GTDB) allowing a single nucleotide mismatch. Genomes with a completeness lower than 95% were discarded. Only amplicon sequence variants (ASV) with more than 10 reads in a given habitat occurring in at least 10% of the samples were included. The proportion of single amino acid auxotrophs was calculated from the total number of auxotrophic taxa. The table is sorted by increasing proportion of amino acid auxotrophs in each habitat.

composition of the communities found in those habitats (Fig. 2B). Habitats dominated by the phylum Proteobacteria were the least auxotrophic, and habitats dominated by the Firmicutes were the most auxotrophic (Fig. 2B). These results agreed with the patterns we observed in the analysis across phyla (Fig. 1A), with families in the Firmicutes like *Lactobacillaceae* and *Streptococcaceae* being more auxotrophic than proteobacterial families like the *Pseudomonadaceae* or *Burkholderiaceae* (Supplementary Fig. 3). These results are unlikely to be biased by knowledge gaps in the amino acid biosynthesis pathways of the Firmicutes, as the Firmicutes is a well-studied phylum (see e.g. ref. 66). Since we observed that assembled genomes had more auxotrophies than genomes from cultured isolates, we verified that the differences in the prevalence of auxotrophy across habitats were not driven by the proportion of assembled genomes and genomes derived from isolates across those habitats (Supplementary Fig. 8). Since the proportion of representative genomes recovered differed among habitats (Table 1), we also verified this proportion did not correlate with the proportion of auxotrophic taxa in those habitats ($r = -0.17$, $P = 0.6$).

As there are numerous examples of auxotrophic bacteria that have been isolated from soil^{67,68}, aquatic environments¹, food²⁴, plants³, and the human gut^{21,69,70}, it has been assumed amino acid auxotrophy is a widespread trait across habitats. Our results indicate that amino acid auxotrophies are rather uncommon in non-host associated systems, and are only relatively common in host-associated systems (skin, gut, or oral cavity) and some fermented foods (cheese and sourdough) (Fig. 2A). The mean number of amino acids that bacterial taxa were unable to synthesize ranged between nearly zero in rhizosphere soils, residential plumbing, bulk soil, freshwater lakes, and marine surface waters, to 2–3 amino acids in taxa from the oral cavity, the human gut, and sourdough starter microbiomes (Fig. 2A). Host-associated habitats and fermented foods not only contained more auxotrophic taxa but those auxotrophs were unable to synthesize a larger number of amino acids (Table 1), suggesting that these environments generally support auxotrophic taxa¹³. Host-associated habitats often share a high and temporally stable supply of amino acids both from the host and ingested food⁷¹, and fermented foods can have a high availability of peptides rich in amino acids such as milk proteins⁷². For example, in *Clostridium* species (phylum Firmicutes) amino acid auxotrophies have been associated with toxin production, which increases the availability of amino acids in the gut lumen⁷³. We detected multiple amino acid auxotrophies in *Clostridium* species, which are capable of obtaining energy via the oxidation and reduction of amino acids using the

Stickland reaction in amino acid-rich environments⁷⁴. Overall, our analyses suggest that amino acid auxotrophy might be most beneficial under conditions of temporally stable and (mostly) abundant amino acid supply, conditions which are not likely to be common in soils and aquatic environments. However, there are notable exceptions in these non-host associated environments. For example, while soils generally select for prototrophic bacteria (96.4% of soil taxa in our analyses were prototrophic, Table 1)⁷⁵, the common soil bacterium *Candidatus Udaeobacter* has a ‘streamlined’ genome with multiple amino acid auxotrophies that make it unique among soil bacterial taxa⁷⁶. *Candidatus Udaeobacter* is considered a nutrient scavenger that likely benefits from the locally abundant nutrients provided by decaying bacterial biomass^{76,77} (Supplementary Fig. 9). As another example, we found amino acid auxotrophies to be widespread among soil-dwelling Bdellovibrionaceae (Supplementary Fig. 9) and the predatory lifestyles of members of this group may allow amino acids to be obtained from ingested prey^{78,79}. *Pelagibacter ubique*, an abundant pelagic bacterium with a highly streamlined genome^{80,81,2}, is another example of an organism with a free-living lifestyle where auxotrophy (in this case glycine auxotrophy¹) is a successful strategy owing to the local abundance of glycolate (a precursor of glycine) from neighboring phytoplankton⁸².

Signatures of genome streamlining in amino acid auxotrophs

As noted above, we found that auxotrophic taxa tend to have smaller genomes than prototrophic taxa and genome size was negatively correlated with the number of amino acid auxotrophies per genome (Supplementary Fig. 4C). This pattern is, in part, a product of obligate intracellular parasites having smaller genomes as a product of genetic drift⁸³, as would be the case for Spirochaetes and Tenericutes (Fig. 1A; Supplementary Fig. 3). However, this pattern could also be driven by auxotrophic free-living bacteria being more likely to have ‘streamlined’ genomes⁸⁴. In other words, there is selection for amino acid auxotrophy in free-living taxa with smaller genomes that minimize cell complexity to more efficiently use the resources required to sustain growth. To test this ‘streamlining’ hypothesis, we focused our analyses on two phyla, Bacteroidetes and Firmicutes, with high proportions of auxotrophic taxa (37.2% and 37.0%, respectively), and we identified gene categories (COG categories⁸⁵) that were differentially abundant across auxotrophic versus prototrophic members of each phylum (Fig. 3). In this analysis, we considered auxotrophic taxa to be only those taxa that were unable to synthesize two or more amino acids. In both phyla, genome size was negatively correlated with the number of




Fig. 2 | Amino acid auxotrophy in bacteria across habitats. **A** Prevalence of amino acid auxotrophy in representative bacterial taxa from 12 different habitats ($N = 13,523$ genomes). The mean number of amino acid auxotrophies of representative bacterial taxa in each habitat is shown as red diamonds in the main panel, and as horizontal bars in the top subpanel. **B** Relative abundance of the most dominant phyla across habitats. The x-axis is sorted by increasing numbers of auxotrophic taxa in each habitat. Letters at the top of **A** indicate statistical differences ($P < 0.05$) between habitats based on Mann-Whitney U tests with Bonferroni-corrected p -values.

amino acid auxotrophies per genome (Fig. 3A, B), in agreement with the general expectation from streamlining theory⁸⁴. Likewise, as expected for streamlined taxa, genes for translation, protein turnover, and post-translational modification were all overrepresented in the genomes of auxotrophic taxa (Fig. 3C). These and other functional gene categories, such as nucleotide transport and metabolism and DNA replication, recombination and repair have all been previously linked to genome streamlining and associated life history strategies across a broad range of bacterial taxa^{75,86,87}. The genes overrepresented in the genomes of prototrophic taxa were also consistent with our expectations and previous findings: genes for the transport and metabolism of carbohydrates, amino acids, and lipids, and genes for

transcription and signal transduction were all overrepresented in the genomes of prototrophic taxa (Fig. 3C)^{86,87}. Together, these findings indicate that amino acid auxotrophy is part of the general life history strategy that characterizes bacteria with ‘streamlined’ genomes.

Conclusions

Amino acid auxotrophy is broadly distributed across the bacterial tree of life, but it is likely less common than previously assumed. We observed appreciable taxon-specific and habitat-specific differences in the prevalence of amino acid auxotrophies, whereby amino acid auxotrophy seems to be most prevalent in host-associated systems or habitats where amino acid availability is expected to be relatively high. In free-living taxa, amino acid auxotrophy likely arises as a product of the genome streamlining process, whereby taxa are adapted for efficient growth sustained on temporally stable supplies of nutrients. This strategy is likely a characteristic of the majority of bacterial taxa that remain uncultured⁴², emphasizing the need for directing culturing efforts towards bacteria with traits such as auxotrophy and small genomes. Overall, our comprehensive investigation of bacterial amino acid auxotrophies highlights that we still have insufficient experimental evidence to confirm amino acid auxotrophies across many bacterial groups. Dedicated culturing and testing of growth requirements across diverse bacterial taxa would further our understanding of the links between auxotrophy and the specific bacterial life history strategies that make amino acid auxotrophy an ecologically successful strategy.

Methods

Study design

We compiled the full sequences of the ~62,000 unique bacterial genomes (‘species clusters’) available in the Genome Taxonomy Database (GTDB) (release 207)⁵³. We restricted our analyses to only those bacterial phyla with more than 100 representative genomes available in GTDB (12 phyla in total) and only included genomes estimated to be >95% complete based on CheckM (v1.1.6)⁸⁸. We also removed all metagenome-assembled genomes (MAGs) that lacked 16S rRNA genes, as well as those with signals of chimerism based on GUNC (Genome Unclutterer)⁸⁹, yielding 26,277 genomes in total. We then ran the automated amino acid biosynthesis annotation tool GapMind on all of these genomes³¹. GapMind identifies candidates for steps in amino acid biosynthesis by using a database of 1849 proteins that have been experimentally shown to be involved in amino acid biosynthesis (taken primarily from MetaCyc⁹⁰, SwissProt⁹¹ and BRENDA⁹²), as well as 145 protein families (144 TIGRFams⁹³ and 1 Pfam⁹⁴). GapMind then searches genomes for candidates in the reference biosynthesis pathways using *ublast* (for similar proteins⁹⁵) or *HMMER* (for members of the same protein family⁹⁶), providing confidence of matches based on sequence identity and coverage³¹. At this step, GapMind uses *ublast* to check if these candidates are similar to any of 113,704 experimentally-characterized proteins that could have alternative functions to amino acid biosynthesis. Candidates are considered valid if the bit score of the alignment to proteins involved in amino acid biosynthesis is higher than the bit score of the alignment to proteins with other functions. We considered a biosynthetic step to be present if it had at least a medium-confidence candidate, which for protein candidates based on similarity to a characterized protein means either (1) at least 40% identity and 70% coverage to a characterized protein, or (2) at least 30% identity and 80% coverage and more similar to protein(s) with this function than to another characterized protein in the database of the 113,704 proteins. We predicted the biosynthesis capabilities for 17 amino acids and chorismate (a precursor of aromatic amino acids), but excluded alanine, aspartate, and glutamate because these amino acids can be produced via the transamination of intermediates from central metabolism, and annotating the substrates of transaminases is inherently challenging²⁹.




Fig. 3 | Evidence for genome streamlining and related functional features in bacterial taxa that are auxotrophic for amino acids. **A, B** Relationship between genome size and the number of amino acid auxotrophies in bacterial taxa from the phylum Bacteroidetes **A** and Firmicutes **B**, respectively. **C** Functional categories that are consistently overrepresented in auxotrophic and prototrophic genomes

from the phyla Bacteroidetes and Firmicutes. Functional categories were defined as Clusters of Orthologous Genes (COGs). We displayed those categories where the Bacteroidetes and Firmicutes had non-statistically significant trends in grey font based on Mann-Whitney U tests ($P < 0.01$). Pearson's correlation coefficients (r) are displayed on **A, B**. $N_{\text{Bacteroidetes}} = 3232$ genomes, $N_{\text{Firmicutes}} = 4674$ genomes.

In addition to predicting amino acid auxotrophy across bacterial phyla, we also investigated how the prevalence of amino acid auxotrophy varies across different bacterial habitats. To do so, we used 16 S rRNA gene sequencing data from 12 different habitats (one dataset per habitat, Table 1), to identify the predominant bacterial taxa found in each of the 12 habitats. We selected 12 publicly available 16 S rRNA gene sequence datasets that each had >100 samples, with each dataset including a broad range of sample types representative of the habitat. These datasets were analyzed using the same bioinformatic pipeline. Briefly, we used *cutadapt* (v1.18)⁹⁷ to remove primers, adapters and ambiguous bases from the 16 S rRNA gene reads. We then quality-filtered the sequences, inferred amplicon sequence variants (ASVs) using the DADA2 pipeline (v1.14.1)⁹⁸, and removed chimeric sequences. Taxonomic affiliations were determined against the SILVA SSU database (release 138)⁹⁹. We used the *phyloseq* R package (v1.38.0)¹⁰⁰ for downstream analyses. From each dataset we obtained representative genomes by matching the 16 S rRNA gene sequences of individual taxa to genomes in GTDB, allowing a single base mismatch (i.e. 99.6% sequence similarity for 250 bp fragments), following the approach used previously to investigate the genomic attributes of bacteria across environmental gradients¹⁰¹. We only included ASVs that had

more than 10 reads in a given habitat and occurred in at least 10% of the samples from each dataset as we wanted to focus on representative genomes from those taxa that are reasonably ubiquitous in each of the 12 habitats. We ran the GapMind pipeline on these representative genomes to infer the completeness of the amino acid biosynthesis profiles for those bacterial community members in each habitat.

Validation of amino acid auxotrophy predictions

Since many of the genes involved in amino acid biosynthesis are not well described²⁹, genome-based inferences can significantly overestimate the prevalence of auxotrophies. Thus, to validate our approach, we compiled genomic information from 171 taxa that are known to grow in minimal media without the external supply of amino acids (i.e. prototrophs, compiled in Price et al.³¹; Supplementary Data 1) and ran GapMind on those genomes to quantify biases in our predictions. We also estimated the accuracy of the predictions for specific auxotrophies by compiling genomic information for 19 taxa with experimentally determined auxotrophies (compiled from^{31,102}; Supplementary Table 1). This validation allowed us to determine the number of genes that need to be missing in any given amino acid biosynthesis pathway to consider that taxon auxotrophic for a given amino acid.

Associations between functional genes and amino acid auxotrophy

We investigated associations between amino acid auxotrophy and broad functional gene categories by testing the prevalence of Clusters of Orthologous Genes (COGs) in the genomes of auxotrophic and prototrophic taxa⁸⁵. We conducted these analyses on the phyla Bacteroidetes and Firmicutes as the metabolic pathways of these phyla are relatively well-studied, contain >3000 taxa with available genomes, and these phyla include sizeable proportions of auxotrophs for robust statistical analyses. We annotated genomes into COG categories using eggNOG-mapper v2¹⁰³, and calculated the genome size-corrected prevalence of each COG category per genome. In order to have a conservative classification of auxotrophy, we only classified those taxa that contained 2 or more amino acid auxotrophies as auxotrophs, and those taxa with no auxotrophies as prototrophs. We obtained minimal doubling times for all genomes based on the predictions established by Weissman et al.¹⁰⁴ (gRodon R package; <https://github.com/jlw-ecoevo/gRodon>), by matching the genome accessions of the taxa in the EGGO database (<https://github.com/jlw-ecoevo/eggo>).

Statistical analyses

We verified the non-normality of the data using the Shapiro-Wilk test and compared the number of auxotrophic taxa between phyla and habitats using Mann-Whitney U tests using the `wilcox.test()` R function with Bonferroni correction of *p*-values for multiple comparisons. We used Pearson's correlation tests to determine whether bacteria were more auxotrophic for amino acids with higher biosynthetic energy costs. The same test was used to investigate correlations of auxotrophy with genome size. We obtained information on the energy (P-bonds) required for amino acid biosynthesis from Akashi and Gojobori¹². We used multiple Mann-Whitney U tests with Bonferroni correction for multiple comparisons to investigate whether particular COG categories were overrepresented in genomes from auxotrophic versus prototrophic taxa. We represented the results as the log₂-fold ratio. Finally, we investigated associations between the estimated bacterial minimal doubling times and genome origin using Mann-Whitney U tests, and tested differences in genome size between assembled genomes and genomes from cultured isolates using Welch two-sample two-sided *t*-tests. All statistical analyses were performed in R (v4.1.3)¹⁰⁵.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

All sequence data analyzed for this study had already been deposited in open repositories and can be accessed through the specific works cited in this work. The source data to reproduce the findings of this study has been made publicly available on Figshare (<https://doi.org/10.6084/m9.figshare.24101742.v1>). The genome data included in this study can be found in the Genome Taxonomy Database (GTDB, <https://data.gtdb.ecogenomic.org/releases/release207/207.0/>). Information on predicted doubling times in bacteria can be found in the EGGO database (<https://github.com/jlw-ecoevo/eggo>). Functional gene annotations were based on the Database of Clusters of Orthologous Genes (COGs, <https://www.ncbi.nlm.nih.gov/research/cog>).

Code availability

The code to reproduce the findings of this study has been made publicly available on Figshare (<https://doi.org/10.6084/m9.figshare.24101742.v1>).

References

1. Tripp, H. J. et al. SAR11 marine bacteria require exogenous reduced sulphur for growth. *Nature* **452**, 741–744 (2008).

2. Yu, X. J., Walker, D. H., Liu, Y. & Zhang, L. Amino acid biosynthesis deficiency in bacteria associated with human and animal hosts. *Infect. Genet. Evol.* **9**, 514–517 (2009).
3. Ryback, B., Bortfeld-Miller, M. & Vorholt, J. A. Metabolic adaptation to vitamin auxotrophy by leaf-associated bacteria. *ISME J.* **16**, 2712–2724 (2022).
4. Bertrand, E. M. & Allen, A. E. Influence of vitamin B auxotrophy on nitrogen metabolism in eukaryotic phytoplankton. *Front. Microbiol.* **3**, 375 (2012).
5. Thakur, K., Tomar, S. K. & De, S. Lactic acid bacteria as a cell factory for riboflavin production. *Microb. Biotechnol.* **9**, 441–451 (2016).
6. Romine, M. F., Rodionov, D. A., Maezato, Y., Osterman, A. L. & Nelson, W. C. Underlying mechanisms for syntrophic metabolism of essential enzyme cofactors in microbial communities. *ISME J.* **11**, 1434–1446 (2017).
7. Paerl, R. W. et al. Prevalent reliance of bacterioplankton on exogenous vitamin B1 and precursor availability. *Proc. Natl Acad. Sci. USA* **115**, 10447–10456 (2018).
8. Liu, Y. F. et al. Metabolic capability and in situ activity of microorganisms in an oil reservoir. *Microbiome* **6**, 1–12 (2018).
9. Jiang, X. et al. Impact of spatial organization on a novel auxotrophic interaction among soil microbes. *ISME J.* **12**, 1443–1456 (2018).
10. Johnson, W. M. et al. Auxotrophic interactions: a stabilizing attribute of aquatic microbial communities? *FEMS Microbiol. Ecol.* **96**, 115 (2020).
11. Yu, J. S. L. et al. Microbial communities form rich extracellular metabolomes that foster metabolic interactions and promote drug tolerance. *Nat. Microbiol.* **7**, 542–555 (2022).
12. Akashi, H. & Gojobori, T. Metabolic efficiency and amino acid composition in the proteomes of *Escherichia coli* and *Bacillus subtilis*. *Proc. Natl Acad. Sci. USA* **99**, 3695–3700 (2002).
13. D'Souza, G. et al. Less is more: selective advantages can explain the prevalent loss of biosynthetic genes in bacteria. *Evolution* **68**, 2559–2570 (2014).
14. Puente-Sánchez, F., Pascual-García, A., Bastolla, U., Pedrós-Alió, C. & Tamames, J. Cross-biome microbial networks reveal functional redundancy and suggest genome reduction through functional complementarity. *bioRxiv* 2022.09.11.507163; <https://doi.org/10.1101/2022.09.11.507163> (2022).
15. Zengler, K. & Zaramela, L. S. The social network of microorganisms — how auxotrophies shape complex communities. *Nat. Rev. Microbiol.* **16**, 383–390 (2018).
16. D'Souza, G. & Kost, C. Experimental evolution of metabolic dependency in bacteria. *PLoS Genet.* **12**, e1006364 (2016).
17. Suthers, P. F. et al. A genome-scale metabolic reconstruction of *Mycoplasma genitalium* iPS189. *PLoS Comput. Biol.* **5**, e1000285 (2009).
18. Hockney, R. C. & Scott, T. A. The isolation and characterization of three types of vitamin B6 auxotrophs of *Escherichia coli* K12. *J. Gen. Microbiol.* **110**, 275–283 (1979).
19. Tang, Y. Z., Koch, F. & Gobler, C. J. Most harmful algal bloom species are vitamin B1 and B12 auxotrophs. *Proc. Natl Acad. Sci. USA* **107**, 20756–20761 (2010).
20. Rodionova, I. A. et al. Genomic distribution of B-vitamin auxotrophy and uptake transporters in environmental bacteria from the Chloroflexi phylum. *Environ. Microbiol. Rep.* **7**, 204–210 (2015).
21. Soto-Martin, E. C. et al. Vitamin biosynthesis by human gut butyrate-producing bacteria and cross-feeding in synthetic microbial communities. *MBio* **11**, 1–18 (2020).
22. Sebald, M. & Costilow, R. N. Minimal growth requirements for *Clostridium perfringens* and isolation of auxotrophic mutants. *Appl. Microbiol.* **29**, 1–6 (1975).

23. Barth, A. L. & Pitt, T. L. Auxotrophic variants of *Pseudomonas aeruginosa* are selected from prototrophic wild-type strains in respiratory infections in patients with cystic fibrosis. *J. Clin. Microbiol.* **33**, 37–40 (1995).
24. Christensen, J. E. & Steele, J. L. Impaired growth rates in milk of *Lactobacillus helveticus* peptidase mutants can be overcome by use of amino acid supplements. *J. Bacteriol.* **185**, 3297–3306 (2003).
25. Ferrario, C. et al. Exploring amino acid auxotrophy in *Bifidobacterium bifidum* PRL2010. *Front. Microbiol.* **6**, 1331 (2015).
26. Veith, N. et al. Using a genome-scale metabolic model of *Enterococcus faecalis* V583 to assess amino acid uptake and its impact on central metabolism. *Appl. Environ. Microbiol.* **81**, 1622–1633 (2015).
27. Devendran, S. et al. *Clostridium scindens* ATCC 35704: Integration of nutritional requirements, the complete genome sequence, and global transcriptional responses to bile acids. *Appl. Environ. Microbiol.* **85**, e00052 (2019).
28. Kim, S. et al. Heme auxotrophy in abundant aquatic microbial lineages. *Proc. Natl Acad. Sci. USA* **118**, e2102750118 (2021).
29. Price, M. N. et al. Filling gaps in bacterial amino acid biosynthesis pathways with high-throughput genetics. *PLoS Genet.* **14**, e1007147 (2018).
30. Tramontano, M. et al. Nutritional preferences of human gut bacteria reveal their metabolic idiosyncrasies. *Nat. Microbiol.* **3**, 514–522 (2018).
31. Price, M. N., Deutschbauer, A. M. & Arkin, A. P. GapMind: automated annotation of amino acid biosynthesis. *mSystems* **5**, e00291 (2020).
32. Seif, Y. et al. Metabolic and genetic basis for auxotrophies in Gram-negative species. *Proc. Natl Acad. Sci. USA* **117**, 6264–6273 (2020).
33. Mee, M. T., Collins, J. J., Church, G. M. & Wang, H. H. Syntrophic exchange in synthetic microbial communities. *Proc. Natl Acad. Sci. USA* **111**, 2149–2156 (2014).
34. Embree, M., Liu, J. K., Al-Bassam, M. M. & Zengler, K. Networks of energetic and metabolic interactions define dynamics in microbial communities. *Proc. Natl Acad. Sci. USA* **112**, 15450–15455 (2015).
35. Lawson, C. E. et al. Metabolic network analysis reveals microbial community interactions in anammox granules. *Nat. Commun.* **8**, 1–12 (2017).
36. Walzem, R. L., Dillard, C. J. & German, J. B. Whey components: millennia of evolution create functionalities for mammalian nutrition: what we know and what we may be overlooking. *Crit. Rev. Food Sci. Nut.* **42**, 353–375 (2002).
37. Stewart, P. S. Diffusion in biofilms. *J. Bacteriol.* **185**, 1485–1491 (2003).
38. Flemming, H. C. et al. Biofilms: an emergent form of bacterial life. *Nat. Rev. Microbiol.* **14**, 563–575 (2016).
39. Chen, I. M. A. et al. Improving microbial genome annotations in an integrated database context. *PLoS ONE* **8**, e54859 (2013).
40. Monk, J. M. et al. Genome-scale metabolic reconstructions of multiple *Escherichia coli* strains highlight strain-specific adaptations to nutritional environments. *Proc. Natl Acad. Sci. USA* **110**, 20338–20343 (2013).
41. Machado, D., Andrejev, S., Tramontano, M. & Patil, K. R. Fast automated reconstruction of genome-scale metabolic models for microbial species and communities. *Nucleic Acids Res.* **46**, 7542–7553 (2018).
42. Solden, L., Lloyd, K. & Wrighton, K. The bright side of microbial dark matter: lessons learned from the uncultivated majority. *Curr. Opin. Microbiol.* **31**, 217–226 (2016).
43. Price, M. Erroneous predictions of auxotrophies by CarveMe. *Nat. Ecol. Evol.* **7**, 194–195 (2022).
44. De Crécy-Lagard, V. Variations in metabolic pathways create challenges for automated metabolic reconstructions: Examples from the tetrahydrofolate synthesis pathway. *Comput. Struct. Biotechnol. J.* **10**, 41–50 (2014).
45. Tang, Y. et al. Pathway confirmation and flux analysis of central metabolic pathways in *Desulfovibrio vulgaris* Hildenborough using gas chromatography-mass spectrometry and Fourier transform-ion cyclotron resonance mass spectrometry. *J. Bacteriol.* **189**, 940–949 (2007).
46. Christiansen, J. K. et al. Phenotypic and genotypic analysis of amino acid auxotrophy in *Lactobacillus helveticus* CNRZ 32. *Appl. Environ. Microbiol.* **74**, 416–423 (2008).
47. Andresen, L. et al. Auxotrophy-based high throughput screening assay for the identification of *Bacillus subtilis* stringent response inhibitors. *Sci. Rep.* **6**, 1–8 (2016).
48. Watterson, W. J. et al. Droplet-based high-throughput cultivation for accurate screening of antibiotic resistant gut microbes. *Elife* **9**, e56998 (2020).
49. Huang, Y. et al. High-throughput microbial culturomics using automation and machine learning. *Nat. Biotechnol.* **41**, 1424–1433 (2023).
50. Madin, J. S. et al. A synthesis of bacterial and archaeal phenotypic trait data. *Sci. Data* **7**, 1–8 (2020).
51. Parks, D. H. et al. GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Res.* **50**, 785–794 (2022).
52. Mee, M. T. & Wang, H. H. Engineering ecosystems and synthetic ecologies. *Mol. Biosyst.* **8**, 2470–2483 (2012).
53. Norena-Caro, D. & Benton, M. G. Cyanobacteria as photoautotrophic biofactories of high-value chemicals. *J. CO₂ Util.* **28**, 335–366 (2018).
54. Razin, S., Yogeve, D. & Naot, Y. Molecular biology and pathogenicity of Mycoplasmas. *Microbiol. Mol. Biol. Rev.* **62**, 1094–1156 (1998).
55. Yus, E. et al. Impact of genome reduction on bacterial metabolism and its regulation. *Science* **326**, 1263–1268 (2009).
56. Vanyushkina, A. A., Fisunov, G. Y., Gorbachev, A. Y., Kamashev, D. E. & Govorun, V. M. Metabolomic analysis of three mollusc species. *PLoS ONE* **9**, e89312 (2014).
57. Clavel, T., Lepage, P. & Charrier, C. The family Coriobacteriaceae in: *The Prokaryotes: Actinobacteria* 201–238; https://doi.org/10.1007/978-3-642-30138-4_343 (2014).
58. Walter, J. Ecological role of lactobacilli in the gastrointestinal tract: Implications for fundamental and biomedical research. *Appl. Environ. Microbiol.* **74**, 4985–4996 (2008).
59. Bessen, D. E., Smeesters, P. R. & Beall, B. W. Molecular epidemiology, ecology, and evolution of Group A Streptococci. *Microbiol. Spectr.* **6**; <https://doi.org/10.1128/microbiolspec.CPP3-0009-2018> (2018).
60. Albright, S. & Louca, S. Trait biases in microbial reference genomes. *Sci. Data* **10**, 1–17 (2023).
61. Martinez-Gutierrez, C. A. & Aylward, F. O. Genome size distributions in bacteria and archaea are strongly linked to evolutionary history at broad phylogenetic scales. *PLoS Genet.* **18**, e1010220 (2022).
62. Morris, J. J., Lenski, R. E. & Zinser, E. R. The black queen hypothesis: Evolution of dependencies through adaptive gene loss. *MBio* **3**, e00036 (2012).
63. Kaiser, J. C. & Heinrichs, D. E. Branching out: Alterations in bacterial physiology and virulence due to branched-chain amino acid deprivation. *MBio* **9**, e01188 (2018).
64. Doden, H. L. et al. Completion of the gut microbial epi-bile acid pathway. *Gut Microbes* **13**, 1–20 (2021).
65. Chiba, Y. et al. Discovery and analysis of a novel type of the serine biosynthetic enzyme phosphoserine phosphatase in *Thermus thermophilus*. *FEBS J.* **286**, 726–736 (2019).

66. van der Kaaij, H., Desiere, F., Mollet, B. & Germond, J. E. L-alanine auxotrophy of *Lactobacillus johnsonii* as demonstrated by physiological, genomic, and gene complementation approaches. *Appl. Environ. Microbiol.* **70**, 1869–1873 (2004).
67. Iwasaki, Y., Ichino, T. & Saito, A. Transition of the bacterial community and culturability of chitinolytic bacteria in chitin-treated upland soil: from *Streptomyces* to methionine-auxotrophic *Lyso-bacter* and other genera. *Microbes Environ.* **35**, ME19070 (2020).
68. Kuykendall, L. D. & Elkan, G. H. *Rhizobium japonicum* derivatives differing in nitrogen-fixing efficiency and carbohydrate utilization. *Appl. Environ. Microbiol.* **32**, 511–519 (1976).
69. Tenover, F. C. & Patton, C. M. Naturally occurring auxotrophs of *Campylobacter jejuni* and *Campylobacter coli*. *J. Clin. Microbiol.* **25**, 1659–1661 (1987).
70. Ottman, N. et al. Genomescale model and omics analysis of metabolic capacities of *Akkermansia muciniphila* reveal a preferential mucin-degrading lifestyle. *Appl. Environ. Microbiol.* **83**, 1014–1031 (2017).
71. Neis, E. P. J. G., Dejong, C. H. C. & Rensen, S. S. The role of microbial amino acid metabolism in host metabolism. *Nutrients* **7**, 2930–2946 (2015).
72. Liepke, C. et al. Human milk provides peptides highly stimulating the growth of bifidobacteria. *Eur. J. Biochem.* **269**, 712–718 (2002).
73. Fletcher, J. R. et al. *Clostridioides difficile* exploits toxin-mediated inflammation to alter the host nutritional landscape and exclude competitors from the gut microbiota. *Nat. Commun.* **12**, 1–14 (2021).
74. Bouillaut, L., Self, W. T. & Sonenshein, A. L. Proline-dependent regulation of *Clostridium difficile* stickland metabolism. *J. Bacteriol.* **195**, 844–854 (2013).
75. Cobo-Simón, M. & Tamames, J. Relating genomic characteristics to environmental preferences and ubiquity in different microbial taxa. *BMC Genomics* **18**, 1–11 (2017).
76. Brewer, T. E., Handley, K. M., Carini, P., Gilbert, J. A. & Fierer, N. Genome reduction in an abundant and ubiquitous soil bacterium ‘*Candidatus Udaeobacter copiosus*’. *Nat. Microbiol.* **2**, 1–7 (2016).
77. Willms, I. M. et al. Globally abundant ‘*Candidatus Udaeobacter*’ benefits from release of antibiotics in soil and potentially performs trace gas scavenging. *mSphere* **5**, e00186-20 (2020).
78. Rendulic, S. et al. A predator unmasked: life cycle of *Bdellovibrio bacteriovorus* from a genomic perspective. *Science* **303**, 689–692 (2004).
79. Pasternak, Z. et al. By their genes ye shall know them: genomic signatures of predatory bacteria. *ISME J.* **7**, 756–769 (2012).
80. Giovannoni, S. J. et al. Genetics: genome streamlining in a cosmopolitan oceanic bacterium. *Science* **309**, 1242–1245 (2005).
81. Carini, P., Steindler, L., Beszteri, S. & Giovannoni, S. J. Nutrient requirements for growth of the extreme oligotroph ‘*Candidatus Pelagibacter ubique*’ HTCC1062 on a defined medium. *ISME J.* **7**, 592–602 (2012).
82. Kieft, B. et al. Phytoplankton exudates and lysates support distinct microbial consortia with specialized metabolic and ecophysiological traits. *Proc. Natl Acad. Sci. USA* **118**, e2101178118 (2021).
83. Kuo, C. H., Moran, N. A. & Ochman, H. The consequences of genetic drift for bacterial genome complexity. *Genome Res.* **19**, 1450–1454 (2009).
84. Giovannoni, S. J., Cameron Thrash, J. & Temperton, B. Implications of streamlining theory for microbial ecology. *ISME J.* **8**, 1553–1565 (2014).
85. Galperin, M. Y. et al. COG database update: focus on microbial diversity, model organisms, and widespread pathogens. *Nucleic Acids Res.* **49**, 274–281 (2021).
86. Konstantinidis, K. T. & Tiedje, J. M. Trends between gene content and genome size in prokaryotic species with larger genomes. *Proc. Natl Acad. Sci. USA* **101**, 3160–3165 (2004).
87. Noell, S. E., Hellweger, F. L., Temperton, B. & Giovannoni, S. J. A reduction of transcriptional regulation in aquatic oligotrophic microorganisms enhances fitness in nutrient-poor environments. *Microbiol. Mol. Biol. Rev.* **30**, e0012422 (2023).
88. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Gen. Res.* **25**, 1043–1055 (2015).
89. Orakov, A. et al. GUNC: detection of chimerism and contamination in prokaryotic genomes. *Genome Biol.* **22**, 1–19 (2021).
90. Caspi, R. et al. The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Nucleic Acids Res.* **38**, 473–479 (2010).
91. Bateman, A. et al. UniProt: the universal protein knowledgebase. *Nucleic Acids Res.* **45**, 158–169 (2017).
92. Placzek, S. et al. BRENDA in 2017: new perspectives and new tools in BRENDA. *Nucleic Acids Res.* **45**, 380–388 (2017).
93. Haft, D. H. et al. TIGRFAMs and genome properties in 2013. *Nucleic Acids Res.* **41**, 387–395 (2013).
94. Mistry, J. et al. Pfam: The protein families database in 2021. *Nucleic Acids Res.* **49**, 412–419 (2021).
95. Edgar, R. C. & Bateman, A. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460–2461 (2010).
96. Eddy, S. R., Wheeler, T. J. & Development Team. HMMER User Guide. 120 (2015).
97. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* **17**, 10–12 (2011).
98. Callahan, B. J., McMurdie, P. J. & Holmes, S. P. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J.* **11**, 2639–2643 (2017).
99. Quast, C. et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* **41**, 590–596 (2013).
100. McMurdie, P. J. & Holmes, S. phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE* **8**, e61217 (2013).
101. Ramoneda, J. et al. Building a genome-based understanding of bacterial pH preferences. *Sci. Adv.* **9**, 17 (2023).
102. Ashniev, G. A., Petrov, S. N., Iablokov, S. N. & Rodionov, D. A. Genomics-based reconstruction and predictive profiling of amino acid biosynthesis in the human gut microbiome. *Microorganisms* **10**, 740 (2022).
103. Cantalapiedra, C. P., Hernandez-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J. eggNOG-mapper v2: Functional annotation, orthology assignments, and domain prediction at the metagenomic scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).
104. Weissman, J. L., Hou, S. & Fuhrman, J. A. Estimating maximal microbial growth rates from cultures, metagenomes, and single cells via codon usage patterns. *Proc. Natl Acad. Sci. USA* **118**, e2016810118 (2021).
105. R. Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.r-project.org/> (2021)
106. Ramirez, K. S. et al. Range-expansion effects on the belowground plant microbiome. *Nat. Ecol. Evol.* **3**, 604–611 (2019).
107. Gebert, M. J. et al. Ecological analyses of mycobacteria in showerhead biofilms and their relevance to human health. *MBio* **9**, e01614 (2018).
108. Oliverio, A. M. et al. The role of phosphorus limitation in shaping soil bacterial communities and their metabolic capabilities. *MBio* **11**, e01718 (2020).
109. Ortiz-Álvarez, R., Cáliz, J., Camarero, L. & Casamayor, E. O. Regional community assembly drivers and microbial environmental sources shaping bacterioplankton in an alpine lacustrine district (Pyrenees, Spain). *Environ. Microbiol.* **22**, 297–309 (2020).

110. Milici, M. et al. Bacterioplankton biogeography of the Atlantic ocean: A case study of the distance-decay relationship. *Front. Microbiol.* **7**, 590 (2016).
111. Nierychlo, M. et al. MiDAS 3: an ecosystem-specific reference database, taxonomy and knowledge platform for activated sludge and anaerobic digesters reveals species-level microbiome composition of activated sludge. *Water Res.* **182**, 115955 (2020).
112. Smets, W. et al. Leaf side determines the relative importance of dispersal versus host filtering in the phyllosphere microbiome. *bioRxiv* 2022.08.16.504148; <https://doi.org/10.1101/2022.08.16.504148> (2022).
113. Wolfe, B. E., Button, J. E., Santarelli, M. & Dutton, R. J. Cheese rind communities provide tractable systems for in situ and in vitro studies of microbial diversity. *Cell* **158**, 422–433 (2014).
114. Landis, E. A. et al. The diversity and function of sourdough starter microbiomes. *Elife* **10**, e61644 (2021).
115. Dimitriu, P. A. et al. New insights into the intrinsic and extrinsic factors that shape the human skin microbiome. *MBio* **10**, e00839 (2019).
116. Vangay, P. et al. US Immigration westernizes the human gut microbiome. *Cell* **175**, 962–972 (2018).

Acknowledgements

We thank all the people involved in the acquisition of the data compiled in this study. J.R. acknowledges funding from the Swiss National Science Foundation (Early PostDoc Mobility grant P2EZP3_199849). N.F. was supported by grants from the National Science Foundation (OPP 2133684 and AW5809-826664). M.N.P. was funded by ENIGMA—Ecosystems and Networks Integrated with Genes and Molecular Assemblies (<http://enigma.lbl.gov>), a Science Focus Area Program at Lawrence Berkeley National Laboratory is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Biological & Environmental Research under contract number DE-AC02-05CH11231. E.O.C. was supported by grants from MICIN/AEI/ERDF (INTERACTOMA RTI2018-101205-B-I00).

Author contributions

J.R. and N.F. conceived and designed the study. J.R. and T.B.N.J. performed the data analysis with the help of M.N.P. E.O.C. and M.N.P.

provided data to the study. J.R. and N.F. wrote the manuscript, with input from all co-authors.

Competing interests

The authors declare no conflicts of interest.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41467-023-43435-4>.

Correspondence and requests for materials should be addressed to Josep Ramoneda or Noah Fierer.

Peer review information *Nature Communications* thanks the anonymous reviewers for their contribution to the peer review of this work. A peer review file is available.

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023