Lawrence Berkeley National Laboratory

Recent Work

Title

Metabolic engineering strategies for sesquiterpene production in microorganism.

Permalink

<https://escholarship.org/uc/item/2zw971kg>

Journal

Critical reviews in biotechnology, 42(1)

ISSN

0738-8551

Authors

Liu, Chun-Li
Xue, Kai
Yang, Yankun
[et al.](#)

Publication Date

2022-02-01

DOI

10.1080/07388551.2021.1924112

Peer reviewed

Metabolic engineering strategies for sesquiterpene production in microorganism

Chun-Li Liu^{1, 2, 3, 4}, Kai Xue^{1, 3, 4}, Yankun Yang^{1, 3, 4}, Xiuxia Liu^{1, 3, 4}, Ye Li^{1, 3, 4}, Taek Soon Lee^{5, 6, *}, Zhonghu Bai^{1, 3, 4, *}, Tian-Wei Tan^{2, *}

¹ National Engineering Laboratory for Cereal Fermentation Technology, Jiangnan University, Wuxi, 214122, China;

² College of Life Science and Technology, Beijing University of Chemical Technology, Beijing 100029, P. R. China;

³ The Key Laboratory of Industrial Biotechnology, Ministry of Education, School of Biotechnology, Jiangnan University;

⁴ Jiangsu Provincial Research Center for Bioactive Product Processing Technology, Jiangnan University;

⁵ Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA;

⁶ Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA;

*Corresponding authors:

Dr. Tianwei Tan, College of Life Science and Technology, Beijing University of Chemical Technology, Beijing Chaoyang District North Sanhuan Road no. 15, Beijing 100029. P. R. China; Phone: +86-10-6441-6691, Fax: +86-10-6471-5443, E-mail: twtan@mail.buct.edu.cn;

Dr. Zhonghu Bai, Jiangnan University, Jiangsu Province Wuxi City Binhu District Lihu Road no. 1800, Wuxi 214122. P. R. China; Phone: +86-510-8532-9306, Fax: +86-510-8532-9306, E-mail: baizhonghu@jiangnan.edu.cn; Dr. Taek Soon Lee, Joint BioEnergy Institute, 5885

Hollis St. 4th floor, Emeryville, CA 94608, USA; Phone: +1-510-495-2470, Fax: +1-510-495-2629, E-mail: tslee@lbl.gov.

Abstract

Sesquiterpenes are a large variety of terpene natural products, widely existing in plants, fungi, marine organisms, insects, and microbes. The value-added sesquiterpenes are extensively used in industries as food, drugs, fragrances and fuels. With increase of the market demands and the price of sesquiterpenes, biosynthesis of sesquiterpenes by microbial fermentation methods from renewable feedstocks acquires increasing attention. The synthetic biology provides robust tools of sesquiterpene production in microorganisms. This review presents a summary of sesquiterpene biosynthesis and metabolic engineering strategies on the host and pathway engineering for sesquiterpene production. The diversity, native producers, and the syntheses of sesquiterpenes are presented. Advances in synthetic biology provide new strategies on creation of the desired hosts for sesquiterpene production. Especially, metabolic engineering strategies for production of sesquiterpenes such as amorphadiene, farnesene, bisabolene, and caryophyllene are emphasized in *Escherichia coli*, *Saccharomyces cerevisiae*, and some other microorganisms. Future perspectives on strain and process improvements for sesquiterpene production are also discussed.

Keywords

Sesquiterpene biosynthesis, sesquiterpene synthase, metabolic engineering strategies, MEP pathway, MVA pathway.

1. Introduction

Terpenes are a large and diverse class of organic compounds, with basic C₅ isoprene-unit hydrocarbons. According to the number of C₅ isoprene-unit, terpenes are classified as hemi- (C₅), mono- (C₁₀), sesqui- (C₁₅), di- (C₂₀), sester- (C₂₅), tri- (C₃₀) and tetra- (C₄₀) terpenes (Bohlmann and Croteau, 1999, Chen et al. , 2011). Sesquiterpene, the largest subgroup of terpenes, has several thousands of representatives and more than 300 different skeletons (Breitmaier, 2008, Connolly and Hill, 1991, Sacchettini and Poulter, 1997). Sesquiterpenes mostly consist of hydrocarbons possessing a multitude of different carbon skeletons and some unsaturated bonds, and their structures present acyclic, monocyclic, bicyclic, and tricyclic compounds (Kramer and Abraham, 2011). They present more structures after oxidation or rearrangement based on C₁₅ skeleton. Sesquiterpenes are mostly isolated from a variety of natural sources, including plant, fungi, microorganisms, and insect (Sacchettini and Poulter, 1997). In plants and microorganisms, sesquiterpenes are important components of volatile oil, existing in form of alkene, alcohol, ketone, and lactone, emitted respectively for spreading fungal spores, protecting themselves from enemies and communicating with other organisms (Rohlf's and Churchill, 2011).

In these natural sources, sesquiterpene synthases are responsible for compound biosynthesis from the common intermediate substrate farnesyl diphosphate (FPP) condensed from isopentenyl diphosphate (IPP) and dimethylallyl pyrophosphate (DMAPP) provided by mevalonate (MVA)-dependent pathway discovered in 1950s and 2-C-methyl-D-erythritol 1,4-phosphate (MEP) pathway disclosed in 1993 (Rohmer, et al. , 1993, Schwender et al. , 1996). The MVA pathway exists in some prokaryotes and almost all eukaryotes including high plants. Many, but not all eubacteria (Rohmer, 1999, Rohmer et al. , 1993, Trutko et al. , 2005), and cyanobacteria (Proteau, 1999) make exclusive use of the MEP pathway. The MVA pathway, located in cytosol, is suggested to be responsible for synthesis of sesquiterpenes (C₁₅),

73 terpenoid aldehydes (C15 and C25) and steroids/triterpenoids (C30). While all MEP-derived
74 isoprenoids are generally considered as typical and exclusive plastidial hemi- (C5), mono-
75 (C10), diterpenes (C20) and carotenoids (C40) (Bouvier et al. , 2005, Opitz et al. , 2014,
76 Pazouki and Niinemets, 2016). Although the MVA pathway and the MEP pathway operate in
77 two different subcellular compartments (cytosol and plastid, respectively), metabolite
78 exchanges or cross-regulation known as crosstalk occur between them (Hemmerlin et al. ,
79 2003).

80 Because many sesquiterpenes have complex molecular structures, inherent bioactivity,
81 and aroma, they could be applied as functional precursors of fragrances, medicines, and
82 fuels (Enserink, 2005, Little et al. , 1950, Nour et al. , 2009, Peralta-Yahya et al. , 2011).
83 Although sesquiterpenes and their derivatives display a broad range of application from
84 medicine to fuels, their production is usually at low levels in native producers (Nour et al. ,
85 2009). The small quantities and poor purities limit the scale-up of sesquiterpene production
86 through natural extraction or chemical synthesis. Given the great market demands and
87 limitations of high production of sesquiterpenes, metabolic engineering strategies are
88 attempted to produce them, such as amorphadiene (Redding-Johanson et al. , 2011, Ro et al. ,
89 2006), farnesene (Meadows et al. , 2016, Su et al. , 2015, Yang et al. , 2016c), bisabolene
90 (Kirby et al. , 2014b), and caryophyllene (Yang et al. , 2016a). Synthetic biology provides an
91 alternative approach to produce plant natural product sesquiterpenes in sufficient quantities
92 using microbial fermentation (Ro et al. , 2006), which is enabled to be an economically feasible
93 choice by recent advances.

94 Even though there have been examples of sesquiterpene biosynthesis (Block et al. , 2019,
95 Kramer and Abraham, 2011) and analytical techniques for them (Merfort, 2002), it is
96 meaningful to summarize and update the advances in sesquiterpene biosynthesis and metabolic
97 engineering strategies. Here, we review the main aspects regarding the applications of

sesquiterpenes, natural sesquiterpene biosynthesis and metabolic engineering strategies on sesquiterpene production in engineered microorganisms. First, we briefly characterize diversities and applications of sesquiterpenes. Next, we systematically discuss the natural producers of the compounds and terpene synthases (TPSs). Then, we summarize the strategies on metabolic regulation of isoprenoid pathways as well as other relating engineering networks for increasing sesquiterpene production. Finally, we illustrate the great potentials and challenges of microbial production of sesquiterpenes using metabolic engineering strategies.

2. Sesquiterpenes are widely-existing biosynthetic compounds

The spread applications of sesquiterpenes are systematically summarized here shown in Figure 1. Many sesquiterpenes have been extracted from plants by mankind for different purposes - as fragrances and flavors in cosmetic or food industries, pharmaceutical agents, insecticides and fuels (Agger et al. , 2008). For the natural fragrance of sesquiterpenes and their derivations, they could be used as blends in cosmetic.

Various plants are popular traditional Chinese medicines to prevent or treat different diseases for centuries. One of the effective constituents of the pharmacopoeia is sesquiterpene derivatives, and the bioactivities make them to exhibit useful medicinal properties. Sesquiterpene lactones make a huge group of bio-active constituents isolated from plant families. They have been reported for various potential effects of anti-microbial activities, antioxidant and anticancer potentials, served as antifeedants (Knight, 1995, Mori and Matsushima, 1995, Mullin et al. , 1991), antimigraine, anti-inflammatory (Pfaffenrath et al. , 2002, Tassorelli et al. , 2005), antitumor, and antiulcer (Cho et al. , 2000) for treatment of stomach-ache and skin infection (Heinrich et al. , 1998). From sesquiterpenes and their oxides, considerable interests are gained, such as well-known amorphadiene-derived artemisinin for antimalarial activity (Ecksteinludwig et al. , 2003), xanthatin for antitumor activity (Nibret et al. , 2011), parthenolide for curing migraine (Kwok et al. , 2001), and thapsigargin for treating

prostate cancer (Drew et al. , 2013). Furanoeudesma-1,3-diene and curzarene are responsible for the analgesic effects of myrrh by interacting with brain opioid receptors (Massoud et al. , 2010). β -2-Himachalen-6-ol, a novel sesquiterpene alcohol unique to Lebanese wild carrot, was demonstrated with potent anti-cancer activity (Taleb et al. , 2016). *Trans*-caryophyllene is a bicyclic sesquiterpene produced by plants in response to herbivore damage (Köllner et al. , 2008). Valerenadiene and valerenic acid have been suggested as the active ingredients responsible for sedative effect in *Valeriana officinalis* (*V. officinalis*) (Pyle et al. , 2012). Some plants synthesize mixture of sesquiterpenes which are believed to be ecologically more effective than single compound, such as the case of attraction of pollinators or parasites of insects (Tholl, 2006). Additionally, some sesquiterpenes are also ingredients of pheromones and juvenile hormones or their precursors (Izquierdo-Bueno et al. , 2018, Lamers, 2003).

Sesquiterpenes exist in many natural foods, such as fruits, drinks, and oil. They are comprehensively investigated in essential oil producing plants, such as peppermint, conifers, and citrus fruits (Croteau, 2001, Croteau et al. , 2005, Voo et al. , 2012). Sesquiterpenes accumulate in specialized tissues such as glandular trichomes, oil ducts or secretory cavities (Lange and Turner, 2013). Over 60 sesquiterpenes have been identified in whole grapes or skins (Petronilho et al. , 2014); α -Farnesene was found on natural coating of apples (FE and KE, 1966); (+)-valencene is an aroma compound of citrus fruits, used in flavor foods and drinks (Frohwitter et al. , 2014). Through BLAST analysis, 19 TPSs and 13 sesquiterpenes were identified in unripe fruits of black pepper (*Piper nigrum*), including β -caryophyllene, δ -elemene, α -copaene, cubebene, α -humulene, δ -cadinol, γ -cadinene and δ -cadinene (Jin et al. , 2018). Insects are repelled by subjecting clove oil and patchouli oil, and the functional ingredients are sesquiterpenes and their oxides: seychellene, α -guaiene, α -bulnesene, β - and iso-caryophyllene, β -caryophyllene oxide and ketone, α -humulene, β -patchoulene, α -patchoulene, β -elemene, (+)-longipinene, (-)-isolongifolene, (+)-longifolene, and so on

(Zhang et al. , 2011). Therefore, sesquiterpenes may have potential interests as preservative in food. Many sesquiterpenes, β -elemene, γ -elemene, α -caryophyllene, germacrene D and δ -cadinene, exist in fruit of *Zanthoxylum schinifolium*, a kind of flavourful spice and Chinese medicine (Wang et al. , 2011b). α -Humulene and α -caryophyllene are contained in the essential oil of *Humulus lupulus*, an inevitable ingredient for beer brewery (Connolly and Hill, 1991). Interestingly, potent odorants in wine are from some mixed terpenoids, of which the formation mechanism encompasses enzymatic and non-enzymatic steps (Schwab and Wust, 2015).

In addition, due to high energy density, comparable cold properties and similar carbon length with cetane, sesquiterpenes reveal great interests as next-generation jet fuel substitutes (Harrison and Harvey, 2017). Especially, the ring and branching portion of sesquiterpenes could increase the gravimetric density and the volumetric energy density of fuel. Some sesquiterpenes have been proposed as jet fuel precursors due to their low hygroscopicity and high energy density. Yoshikuni and coworkers introduced TPS mutants into *E. coli* for terpene production, obtaining eight different terpenes (α -farnesene, sabinene, γ -humulene, sibirene, longifolene, α -longipinene and β -bisabolene), most of which have potential as biofuels (Yoshikuni et al. , 2006). The biosynthesized bisabolene was chemically hydrogenated to generate bisabolane, which was identified as an alternative to D2 diesel by comparing its fuel properties with those of diesel fuels (Peralta-Yahya et al. , 2011). Sesquiterpene thujopsene, α -cedrene and β -cedrene in cedarwood oil, were hydrogenated to generate fuel blends with 12% higher volumetric net heat of combustion than conventional jet fuel (Harrison and Harvey, 2017). Sesquiterpene epi-isozizaene, pentalenene and α -isocomene are even proven to have excellent fuel properties by heat of combustion calculation (Liu et al. , 2018).

Figure 1

3. Sesquiterpene biosynthesis

3.1 Sesquiterpene producers

In nature, sesquiterpenes widely exist in plants, fungi, marine organisms, and insects (Figure 2) (Connolly and Hill, 1991, Finefield et al. , 2012). Most of sesquiterpenes are essential for the life of their producers, such as steroid hormones in mammals, carotenoids and chlorophylls in plants, and ubiquinone or menaquinone in bacteria (Sacchettini and Poulter, 1997, Thomas, 1995). In plants, sesquiterpenes and monoterpenes are important constituents of essential oils. As summarized in Figure 2, they are components of volatile oil in form of alcohol, ketone, lactone or glycoside in Asteraceae, Umbelliferae, Euphorbiaceae, Wedaceae, Leguminous, Cucurbitaceae and Ranunculaceae (Connolly and Hill, 1991). Volatile oil from plants is usually consistent of sesquiterpenes: in oil of *Xanthium strumarium*, germacrene D, guaia-4, -6-diene, β -elemene, β -caryophyllene and β -copaene were detected (Li et al. , 2016a). Sesquiterpenes are modulated by enzymes or other enzymes being induced by environmental stress factors. For example, UV-radiation (Back et al. , 1998), infection (Poole and Llewellyn, 2005) and herbivore attack (Yuan et al. , 2008) have been identified to enhance volatile sesquiterpene production. It was reported that H₂O₂ could induce sesquiterpene production by increasing expression of gene *AsTPS10*, *AsTPS16* and *AsTPS19* responsible for sesquiterpene biosynthesis in wounded *Aquilaria sinensis* (Lv et al. , 2019). *Solidago canadensis* majorly produces germacrene D, accounting for more than 90% of the total compounds (Niwa et al. , 2008, Schmidt et al. , 1998, Weyerstahl et al. , 1993). Asteraceae family plant *Parthenium hysterophorus* produces guaia-4,6-diene, whose chemical structure was previously confirmed by nuclear magnetic resonance (NMR) studies (Bohlmann et al. , 1977). Sesquiterpene 7-epizingiberene produced and stored in glandular trichomes is toxic and repellent properties (Bleeker et al. , 2012). Through gas-chromatographic-mass spectrometric (GC-MS) analysis of dynamic gas from the native scent of *Solanum tuberosum* L. leaves, the following sesquiterpenes have been identified: *trans*- and *cis*-caryophyllenes, α -ylangene, α -copaene, β -bourbonene, β -elemene, Z- β -farnesene, aromadendrene, α -humulene, α -elemene, β -bisabolene,

γ -cadinene and δ -cadinene (Khalilov et al. , 1999). In flowers of *Cananga odorata* var. *fruticosa*, bulk of the 49 VOCs were identified as sesquiterpenes: β -caryophyllene, farnesene, γ -muurolene, humulene, β -germacrene D, cedrene, δ -cadinene, α -patchoulene, α -bergamotene, β -ylangene, β -copaene, and β -cubebene (Jin et al. , 2015).

Fungi is another rich resource of terpenes, mostly existing in the format of VOCs (Figure 2). Two producers *Ascomycota* and *Basidiomycota* usually produce sesquiterpenes in late growth phase. Many species of the phylum *Ascomycota* have been reported to produce volatile sesquiterpenes (Kramer and Abraham, 2011). Caryophyllene was emitted from *Phialophora fastigata*, and α -curcumene was biosynthesized by *Penicillium commune* and *Paecilomyces variotii* (Sunesson et al. , 1995). Six TPSs and fifteen sesquiterpenes were respectively characterized and identified in basidiomycete *Coprinus cinereus* (Agger et al. , 2009). Genus *Penicillium*, belonging to phylum *Ascomycota*, is a kind of rich sesquiterpene producer: germacrene A and germacrene B were respectively produced by *P. cyclopium* and *P. expansum*, and β -elemene and β -guaiene were synthesized by *P. clavigerum* (Larsen, 1998). *P. roqueforti* could emit multiply sesquiterpenes β -patchoulene, β -elemene, β -elemene-isomer, diepi-a-cedrene, β -gurjunene, β -patchoulene-isomer, aristolochene, valencene, α -selinene, β -himachalene, α -chamigrene, β -bisabolene, and α -panasinsene (Jeleń, 2002). The fungal phylum *Basidiomycota* produces multiply sesquiterpenes via transforming and rearranging humulene (Abraham, 2001).

Many terpenes from plants and fungi have been described, but relatively fewer microbial sesquiterpenes and TPSs have been functionally summarized. As we summarized here, prokaryote producers of sesquiterpene are mainly identified belonging to Actinomyces, marine bacteria and Cyanobacteria (Figure 2). *Streptomyces* is a rich source of bioactive secondary metabolites, such as antiparasitic avermectins and sesquiterpene-derived antibiotic pentalenolactone (Chou et al. , 2010, Omura et al. , 2001, Quaderer et al. , 2006).

Sesquiterpenoid antibiotic pentalenolactone is a commonly occurring metabolite, isolated from more than 30 species of *Streptomyces* (Koe et al. , 1957, Takahashi et al. , 1983). Bioinformatics analysis of *Streptomyces* and *Kitasatospora* genomes identified 20 terpenes, including sesquiterpene β -elemene, α -muurolene, cadinadiene, epi-isozizaene, germacrene A and pentalenene (Yamada et al. , 2015a, Yamada et al. , 2015b). Many marine bacteria are found to produce many sesquiterpenes with great medical values. Four sesquiterpenes—euplotin A, euplotin B, euplotin C and their biogenetic precursor preuplotin—have been isolated from a eukaryotic unicellular marine ciliate *Euplotes crassus* (Dini et al. , 1993). Sesquiterpenes exclusively produced by marine bacteria ciliate protozoon *E. crassus* have the bioactivity of inhibiting growth and killing protozoa *L. major* and *L. infantum* in vitro (Savoia et al. , 2004).

In addition, insects were reported to synthesize sesquiterpenes (Finefield et al., 2012, Naya et al. , 1978). Some investigation suggests that insects possess a specific set of C-C cyclases for sesquiterpenoids. Volatile oil from the secretion of a scale insect *Ceroplastes ceriferus*, is consisting of numbers of sesquiterpenes: germacrene-D, β -bourbonene, α -copaene, β -bourbonene and β -elemene (Naya et al. , 1978). The presence of some cedrene-type sesquiterpenoids was reported in scale insect *Laeifer lacea* Kerr (Singh et al. , 1974).

Figure 2

3.2 Sesquiterpene synthases

In nature, terpene synthases and their carbocationic reaction mechanisms are responsible for huge varieties of terpene structures and biosynthesis, and they have garnered attentions of chemists and biologists for many years. Starting from farnesyl diphosphate (FPP), sesquiterpene skeleton is cyclized by different TPSs (Benedict et al. , 2001, Cane, 1990). From analysis of TPSs, DDXXD motifs were found to serve as a crucial site for divalent action and substrate binding (Marrero et al. , 1992, Prosser et al. , 2004). Many TPSs, such as farnesene

synthase from *C. junos*, δ -selinene synthase and λ -humulene synthase, were discovered through searching the motifs (Little and Croteau, 2002, Maruyama et al. , 2001).

Large amount of natural TPSs are extracted from plants. A TPS forming carbocyclic skeleton of δ -guaiene was isolated from *Aquilaria* cells (Kumeta and Ito, 2010). α -Farnesene synthase cDNA from peel tissue of apple fruit was cloned and characterized in *E. coli* (Pechous and Whitaker, 2004). Germacrene A synthase from *Lactuca sativa* L. was expressed in engineered *E. coli*, and β -elemene was extracted by thermally rearrangement (Bennett et al. , 2002, Kraker et al. , 1998). The caryophyllene synthase was characterized from *Artemisia annua* (Cai et al. , 2002) and *Mikania micrantha* (Wang et al. , 2009). Multiple TPSs are often expressed in different organs of one organism, and several sesquiterpenes are synthesized through one TPS catalysis. Therefore, several sesquiterpenes are often produced by one organism or one TPS catalysis of substrate FPP. For example, β -humulene synthase and δ -selinene synthase are expressed in *Abies grandis*, each catalyzing at least 52 and 36 sesquiterpenes, respectively (Steele et al. , 1998). Chamomile, with great pharmaceutical values and economic essential oil, was excavated five TPSs, four of which MrTPS1, MrTPS3, MrTPS4 and MrTPS5 were exclusively expressed in above-ground organs and respectively catalyzed (-)-(E)- β -caryophyllene, (+)-germacrene A, (E)- β -ocimene and (-)-germacrene D. Enzyme MrTPS2, mainly expressed in roots, catalyzed several Asteraceae-specific sesquiterpenes: (-)-(E)- β -caryophyllene, α -humulene, silphinene, modeph-2-ene, β -isocomene and the major product α -isocomene (Irmisch et al. , 2012). Based on transcriptome sequencing of *Santalum album*, five TPSs were characterized, including two sesquisabinene synthases (SaSQS1, SaSQS2), bisabolene synthase (SaBS), santalene synthase (SaSS) and farnesyl diphosphate synthase (SaFDS) (Srivastava et al. , 2015). In grand fir, δ -selinene synthase and λ -humulene synthase each enzyme synthesizes three major products respectively, and a third TPS catalyze to form only (E)- α -bisabolene (Bohlmann et al. , 1998, Steele et al. , 1998). Six

TPSs from *C. cinereus* were described: Cop1 and Cop2 each synthesizes germacrene A as their major product; Cop3 was identified as an α -muurolene synthase but producing β -elemene, δ -muurolene, germacrene D, and δ -cadinene; Cop4 synthesizes δ -cadinene as its major product; Cop5 has not been characterized any compounds; Cop6 catalyzes specific α -cuprenene (Agger et al. , 2009). Two TPSs were identified to synthesize a-farnesene and germacrene D from kiwifruit flowers (Nieuwenhuizen et al. , 2009). In root of *V. officinalis*, two TPSs VoTPS1 and VoTPS2 were identified to produce germacrene C/germacrene D and valerenadiene, respectively (Pyle et al. , 2012). In flowers of grapevine *Vitis vinifera* L., 69 putative TPSs were discovered through probe cross-hybridizing, and numbers of sesquiterpenes were identified in heterologous yeast and the plant flowers (Smit et al. , 2019).

Recent years, the availability of genome sequences led to the discoveries of some TPSs from actinomycetes and cyanobacteria (Agger et al. , 2008, Cane and Watt, 2003, Giglio et al. , 2008, Gust et al. , 2003, Komatsu et al. , 2008, Lin et al. , 2006a, Zhao et al. , 2008). Based on bioinformatics analysis, a biosynthetic protein (AncCL1) was reconstructed and identified to catalyze major product epi-zizaene and minor product epi-isozizaene (Guzzetti et al. , 2016). In *Streptomyces coelicolor*, epi-isozizaene synthase (EIZS) catalyzes cyclization of FPP to a novel tricyclic epi-isozizaene (Lin et al. , 2006b). Pentalenene synthase (PentS) from *Streptomyces* UC5319 catalyzes to pentalenene (Cane et al. , 1994), and a mixture of pentalenene and caryophyllene is synthesized by site-directed mutagenesis PentS N219D (Seemann et al. , 2002). Similarly, the types or the product distribution of sesquiterpenes could be changed when mutating TPSs, such as MrTPS2 (Irmisch et al. , 2012), bisabolene synthase (*BIS*) (Kirby et al. , 2014a) and amorphadiene synthase (*ADS*) (Li et al. , 2016b).

Figure 3

4. Metabolic engineering strategies on sesquiterpene production in microorganisms

To explore the potential interest of sesquiterpenes, fermentation-based production process is used to produce diverse sesquiterpenes through engineering metabolic pathways to efficiently drive conversion of the renewable resources. Some industrial model microorganisms are metabolically engineered from isoprenoid pathways, TPSs and other relative metabolic networks for sesquiterpene production. Herein, we describe the general strategies on host and pathway engineering for sesquiterpene production in engineered microorganisms.

4.1. Metabolic engineering strategies on sesquiterpene production in *E. coli*

4.1.1. Metabolic regulation on MEP pathway for sesquiterpene production in *E. coli*

E. coli employing the MEP pathway to accumulate FPP is a widely used host for sesquiterpene production. Metabolic engineering and strategies for sesquiterpene production in *E. coli* are summarized in Table 1 and Figure 4. Via the *E. coli* native MEP pathway, sesquiterpene production is generally considerable low when only over-expressing TPSs. For example, *E. coli* BL21 (DE3) harboring only gene farnesene synthase produced 0.325 mg/L α -farnesene (Zhu et al. , 2014). When codon-optimized gene *coEIZS*, *coPentS* and *coMrTPS2* were simply over-expressed in *E. coli* DH1, the titers of tricycle sesquiterpene epi-isozizaene, pentalenene and α -isocomene were 0.54 mg/L, 0.19 mg/L and 0.01 mg/L, respectively (Liu et al. , 2018). In *E. coli* BL21 (DE3), over-expressing wildtype *valerenadiene synthase* (*wvds*) gene rendered 12 μ g/L valerenadiene (Nybo et al. , 2017).

For metabolic regulation on the MEP pathway, over-expression of rate-limiting genes and TPSs is the most popular strategy to increase sesquiterpene production. Upon co-expressing gene *Dxs*, *isopentenyl-diphosphate isomerase* (*Idi*) and *ispA* in SOE4 operon, amorphadiene titer increased 3.6-fold comparing to that achieved by only *ADS* gene expression strain (Martin et al. , 2003). Medium optimization and over-expression of gene *Dxs* from *Streptomyces avermitilis*, *Idi* from *Bacillus subtilis*, *ispDF* and *ADS* increased amorphadiene titer to 331.70

mg/L, 15.5-fold of the initial titer in *E. coli* MG1655 (Wang et al. , 2015). The aim of over-expressing and deleting genes is to enrich the vital MEP pathway precursors. To enrich precursor phosphoenolpyruvate and pyruvate, carbohydrate phosphotransferase system (PTS) genes were deleted, resulting in amorphaadiene titer increased to 182 mg/L in OPT2 medium (Zhang et al. , 2013, Zhang et al. , 2015). To enrich the vital precursor DXP, a novel route of pentose phosphate to it annotated as putative xylose reductase gene *yajO* and *ribB* mutant was screened. Through fusing *ribB* mutant to *Dxr* gene and over-expressing *ispA*, *ispDF* and *Idi* gene, bisabolene titer increased above 4-fold, about 8.8 mg/g DCW (dry cell weight) (Kirby et al. , 2015). Engineering the MEP pathway by cross-lapping in vitro assembly (CLIVA) method and expressing iron sulfur (Fe–S) cluster operon increased amorphaadiene titer to ~300 mg/L (Zou et al. , 2013). Additionally, to increase sesquiterpene production, efflux transporter engineering was carried out: Over-expressing the screened outer membrane protein *tolC*, together with ABC family transporters (*macAB*) or MFS family transporters (*emrAB* or *emrKY*) increased amorphaadiene titer by more than 3-fold, up to 180 mg/L (Zhang et al. , 2016); Over-expressing two copies of *TolC* and *AcrB* improved amorphaadiene yield 118% with 404.83 mg/L (Wang et al. , 2013). For valerenadiene titer, besides over-expressing gene *Dxs*, *Idi*, FPP synthase (*FFPS*) and *cvds*, supplying glycerol into fermentation medium increased it to 11.0 mg/L (Nybo et al. , 2017).

4.1.2. Metabolic regulation on MVA pathway in *E. coli*

The low sesquiterpene production may be resulted from the insufficient FPP supplementary via native MEP pathway which is tightly controlled by endogenous regulation (Phulara et al. , 2016). The heterogenous MVA pathway was introduced to *E. coli* for dramatically improving flux to FPP and terpene production (Martin, Pitera, 2003). In *E. coli* DYM1, the synthetic *ADS* gene from *A. annua* and MVA pathway from *S. cerevisiae* were expressed, achieving approximately 112.2 mg/L of amorphaadiene (see Table 1) from LB + 0.8%

glycerol culture (Martin, Pitera, 2003). Afterwards, employing a two-phase partitioning bioreactor with dodecane led this engineered strain to produce 0.5 g/L amorphaadiene (Newman et al. , 2006). Although introduction of the MVA pathway dramatically increased amorphaadiene production, bottlenecks limiting the yield still exist. To investigate them, balancing the heterologous MVA pathway flux was proposed. All the actions of supplementing MVA into medium, increasing copies of mevalonate kinase gene (*mvk*) or plasmid with *ADS* gene and global sensitivity analysis with an equation model for the MVA pathway using kinetic parameters identified the rate-limiting genes *HMGR*, *mvk* and *ADS* (Anthony et al. , 2009, Pitera et al. , 2007, Weaver et al. , 2015). Gene titration studies and metabolomics using liquid chromatography–mass spectrometry (LC-MS) excavated a bottleneck - insufficient *HMGR* leading to HMG-CoA accumulation, being toxic to cells by inhibiting fatty acid biosynthesis and generalizing membrane stress (Kizer et al. , 2008, Pitera et al. , 2007). Sequentially, three strategies were carried out: upregulating *tHMGR* gene, over-expressing *HMGS* and *HMGR* gene from *Staphylococcus aureus* (Tsuruta et al. , 2009), and cofactor engineering using NADH-dependent *HMGR* variants from *Delftia acidovorans* (Ma et al. , 2011). Then, a commercially relevant amorphaadiene titer of 27.4 g/L was achieved through a superior fed-batch fermentation process of restricted glucose and nitrogen (Tsuruta et al. , 2009). By supplying more ATP, polyphosphate kinase (PPK)-catalyzed synthesis increased amorphaadiene titer 2.5-fold (Shimane et al. , 2012). To characterize strain variation in engineered *E. coli* for increasing the productivity, a multi-omics-based workflow was developed (Brunk et al. , 2016). Especially, targeted proteomics helped to identify *mvk* and *phosphomevalonate kinase (pmk)* as potential bottleneck genes in amorphaadiene production. Based on this, codon-optimizing *mvk* and *pmk* gene, and replacing P_{lac} with a stronger P_{trc} were conducted, improving amorphaadiene titer 3-fold (>500 mg/L) (Redding-Johanson, Batth, 2011). Assisted by metabolomics and targeted proteomics, metabolic pathway optimization with RBS

variants and combinatorial gene assembly reduced toxic intermediate accumulation and increased amorphaadiene production about 5-fold (Nowroozi et al. , 2014). Accumulation of toxic intermediates (HMG-CoA, IPP/DMAPP and FPP) of the MVA pathway resulted in cell growth issues. Through toxicity stress and whole-genome transcriptional analysis identifying FPP-responding promoter P_{rstA} and P_{gadE} , dynamic control of the MVA pathway was established, and it doubled amorphaadiene titer to 1.4 g/L (Dahl et al. , 2013). Afterwards, this dynamic control strategy was used to produce other sesquiterpenes: dynamic control of the MVA pathway expression and comparative proteome analysis made engineered *E. coli* BW25113 to produce 722.46 mg/L zeaxanthin (Shen et al. , 2016); Together promoter engineering with this dynamic control or inducible control of the MVA pathway, 727.9 mg/L epi-isozizaene, 780.3 mg/L pentalenene and 77.5 mg/L α -isocomene were produced when gene *coEIZS*, *coPentS* and *coMrTPS2* were over-expressed in *E. coli* DH1, respectively (Liu et al. , 2018).

α -Farnesene and β -farnesene are isomers, differing by the location of one double-bond. The commercially available farnesene used as perfume and fuel is a mixture of the isomers. Expressing a codon-optimized farnesene synthase gene made *E. coli* to produce 1.2 mg/L α -farnesene. Augmenting the metabolic flux for FPP synthesis through expressing the MVA pathway and over-expressing *Dxs*, *Idi*, and *ispA* gene conferred a 48-fold improvement. Together with protein fusion of FPPS and farnesene synthase, 380.0 mg/L of α -farnesene was produced (Wang et al. , 2011a). Through engineering the heterologous MVA pathway, 8.74 g/L β -farnesene was produced in a lab bioreactor (You et al. , 2017). Although up-regulating bottleneck genes and down-regulating competitive pathway improved sesquiterpene production, the ratios of enzymes in the reconstructed pathway are hardly precisely evaluated. Therefore, targeted engineering analysis was employed: the MVA pathway was reconstructed *in vitro* to produce farnesene using purified enzymes, and the molar ratio of

AtoB:ERG13:tHMG1:ERG12:ERG8:MVD1:Idi:ispA:AFS was slightly adjusted to 1:10:2:5:5:2:5:2:2. Based on this information, a rationally designed strain was constructed *in vivo* for a high titer of farnesene, achieving 1.1 g/L after 96 h post induction in *E. coli* BL21(DE3) (Zhu, Zhong, 2014). The *in vitro* biosynthesis of sesquiterpenes starting from acetic acid using the MVA pathway was also investigated: a ten-step cascade plus added cofactor regeneration systems led conversion of farnesyl pyrophosphate >40% and conversion of sesquiterpenes >90% (Dirkmann et al. , 2018).

Bisabolene was identified as a biosynthetic precursor of a novel alternative to D2 diesel fuel. *E. coli* expressing the MVA pathway from *S. cerevisiae* and plant-derived bisabolene synthase produced 388 mg/L bisabolene. Codon-optimization of *tHMGR*, *HMGS*, *mvk* and *pmk* genes, and insertion of a P_{trc} in front of *mvk* gene increased bisabolene titer to 912 mg/L (Peralta-Yahya et al. , 2011). Later, this *E. coli* variant was re-engineered to produce 363 µg/L cedrol under shaking flask culture conditions for 72 h (Luo et al. , 2019). To use the industrial biomass pretreated by ionic liquid (IL) for biofuel production, *E. coli* containing plasmid with bisabolene pathway genes was engineered to be resist tolerant to the IL with an inner membrane transporter regulated by an IL-inducible repressor (Ruegg et al. , 2014). Sequentially, principal component analysis of proteomics (PCAP) was proposed to increase bisabolene titer to 1,150 mg/L in strain CB with an extra copy of *BIS* (Alonso-Gutierrez et al. , 2015). These above MVA pathway genes and sucrose utilization genes were also integrated into chromosome to address issues associated with *E. coli* as an industrial platform for isoprenoids and growth ability on sucrose. Although a CRISPR-Cas9 system for replacing promoter sequences increased the bisabolene titer 5-fold, low titer of ~435 µg/L was finally resulted probably due to lower enzymes levels from chromosomally integrated pathway than the strains using plasmids to express pathway proteins (Alonso-Gutierrez et al. , 2018). Similar with the dynamic control, quorum sensing (QS) system controlling gene expression in response to

population density, was applied to establish an inducer-free system of bisabolene production. The LuxI/R system from *Vibrio fischeri* was employed to construct sensor and response plasmids and variants, of which a QS chromosome-integrated strain produced 1.1 g/L bisabolene without inducers added, increasing 44% comparing to the previous inducible system (Kim et al. , 2017). In addition, simulation including of molecular simulation to study the membrane permeability of terpenoids (Vermaas et al. , 2018), CFD and kinetic-based modeling to optimize the sparger design of a large-scale photobioreactor (Ali et al. , 2019) were also employed to improve bisabolene production.

β -Caryophyllene is a bio-based product with economic potentials and a promising precursor candidate for petroleum-based products. Glucose-6-phosphate dehydrogenase (*G6PH*) for increasing NADPH, geranyl diphosphate synthase (*GPPS2*) and β -caryophyllene synthase (*QHS1*) genes, were co-overexpressed with the MVA pathway in engineered *E. coli* YJM59, producing 1,520 mg/L β -caryophyllene in fed-batch fermentation (Yang et al. , 2016a). To convert acetic acid to β -caryophyllene, genes *QHS1*, *GPPS2*, acetyl-CoA synthase (*ACS*) and acetoacetyl-CoA synthase (*AACS*) were co-expressed along with the heterologous MVA pathway in *E. coli* YJM67 and produced up to 1.05 g/L of β -caryophyllene accumulated after 72 h (Yang and Nie, 2016). By expressing the MVA pathway and gene *QHS1* in *E. coli*, 406 mg/L terpene mix enriched in 100 mg/L caryophyllene was produced (Wu et al. , 2018).

For α -humulene production, the MVA pathway genes from *Streptomyces* (from *HMGS* to *Idi2*) and yeast (*Idi1*), and rat gene acetoacetate-CoA ligase (*Aacl*) and α -humulene synthase of shampoo ginger were co-expressed, and the resultant *E. coli* strain produced 958 mg/L α -humulene when supplemented with a lithium acetoacetate (LAA), a 13.6-fold increase compared with strain expressing only humulene synthase (Harada et al. , 2009). These resulted from that exogenous LAA is efficiently converted into acetoacetyl-CoA by mammalian *Aacl* and used as a dominant precursor for isoprenoid biosynthesis. Strategy of a probable

mechanism of adaptive evolution was used to engineer *tHMGR* and terpene synthases (λ -humulene synthase, bisabolene synthase, sibirene synthase, longifolene synthase, and α -longipinene synthase) in *E. coli*, and thereby eliminate the bottleneck from inefficient enzymes, increasing the productivity by ~1,000 fold (Yoshikuni et al. , 2008). Protoilludene was successfully produced by overexpressing a hybrid exogenous MVA pathway, endogenous *ispA*, and protoilludene synthase of *Omphalotus olearius*, and was improved to 1199 mg/L by altering promoters and copy numbers (Yang et al. , 2016b). Additionally, 62.0 mg/L valerenadiene was produced via expressing the MVA pathway and *cvds* gene respectively promoted by P_{trc} and P_{T7} (Nybo et al. , 2017). Co-expression of codon-optimized longifolene synthase from *Picea abies*, *FPPS*, and the MVA pathway genes in *E. coli* BL21 (DE3) rendered 382 mg/L longifolene in a 5-L bioreactor (Cao et al. , 2019).

Figure 4

Table 1

4.2. Metabolic engineering strategies for sesquiterpene production in *S. cerevisiae*

S. cerevisiae harboring the MVA pathway is one of the most favorite microorganisms for industrial production of isoprenoids. The low productivity limits commodity scale production of sesquiterpene. The metabolic engineering strategies, synthetic biology technologies and advanced tools (summarized in Figure 5 and Table 2) have exhibited potential to convert *S. cerevisiae* to a high sesquiterpene-producing host.

In *S. cerevisiae*, up-regulating the upstream MVA pathway for increasing a precursor FPP synthesis is the most common strategy to increase sesquiterpene production. Over-expressing *tHMGR* gene increased amorphadiene titer approximately 5-fold, and down-regulating *ERG9* encoding squalene synthase using a methionine repressible promoter (P_{MET3}) increased it an additional 2-fold by increasing FPP pool (Paradise et al. , 2008, Ro et al. , 2006). In *S. cerevisiae*, a global transcription factor *UPC2* was mutated to regulate biosynthesis of sterols,

an additional copy of *tHMGR* was integrated into chromosome, and *ERG20* encoding FPP synthase was over-expressed, generating FPP-biosynthesis platform EPY224 which produced 153 mg/L amorphaadiene (Ro et al. , 2006). Over-expressing heterologous ADS, replacing promoter by P_{MET3} for repressing *ERG9* gene and enzyme fusion made amorphaadiene titer to 25.02 mg/L in *S. cerevisiae* MTCC 3157 (Baadhe et al. , 2013). P_{CTR3} repressed by cheap CuSO₄ was slight better to suppress *ERG9* expression to reduce FPP flux to squalene than P_{MET3} repressed by methionine (Paddon et al. , 2013, Peng et al. , 2017). To achieve high sesquiterpene, fermentation process needs to be optimized via such as response surface methodology (Baadhe et al. , 2014) and feeding various carbon sources. Feedback-controlled ethanol pulse feed has several significant advantages over restricted glucose feed. This strategy is successfully applied to amorphaadiene with titer of 41 g/L and artemisinic acid production in *S. cerevisiae* CEN.PK2 Y293 (Paddon et al. , 2013, Westfall et al. , 2012). With development of genomics and relevant technologies, comprehensive metabolic models were developed. Model prediction via network models has been applied to sesquiterpene production: OptGene (Burgard et al. , 2003, Patil et al. , 2005), as an extension modeling framework of OptKnock, was subjected to minimization of metabolic adjustment (MOMA) method to identify *GDH1* coding NADPH-dependent glutamate dehydrogenase as the best knockout target. Then, through over-expressing *tHMG1* and replacing NADH-dependent *GDH2* with *GDH1* gene, sesquiterpene titer increased to 30.1 mg/L from 11.3 mg/L (Asadollahi et al. , 2009). Flux balance analysis (FBA) and MOMA methods were used for predicting knockout targets, resulting in 54. 55 mg/L amorphaadiene in the highest strain *hvk2D:: his3::ADS* (Sun et al. , 2014). Assisted by 3D model of ADS protein, mutability landscape guided enzyme engineering was performed to improve the catalytic activity, increasing catalytic efficiency ~ 4-fold in mutant H448A and k_{cat} by 5-fold in mutant T399S/H448A (Abdallah et al. , 2018).

Based on FPP-biosynthesis platform EPY224, many studies were carried out: through replacing *ADS* gene with *BIS* gene, more than 900 mg/L bisabolene was produced (Peralta-Yahya et al. , 2011). Then, this EPY224-derived strain was applied to evolving *BIS*, enriching a mixed population and removing little bisabolene producing strains in presence of selective pressure nonionic surfactants Tween 20 (T20), for the protection of bisabolene against the disruptive action of T20 (Kirby et al. , 2014a). Recently, *S. cerevisiae* EPY224 was re-engineered to produce 344 mg/L pentalenene by replacing gene *BIS* with gene *coPentS* (Liu et al. , 2018). In terpene production, screening color distribution of strains harboring carotenoid-producing pathway is a tool of screening the targeted mutants (Furubayashi et al. , 2014, Wadhwa and Bachhawat, 2019, Yuan and Ching, 2014) and identifying gene deletions contributing to improving isoprenoid production (Ozaydin et al. , 2013). Combination of the deletions and MVA pathway modifications increased bisabolene titer to 5.2 g/L in a fermentation process (Ozaydin et al. , 2013).

In yeast, *FS* gene from *Malus domestica* led to higher α -farnesene titer (4 mg/L) than plant-origins of *C. junos* and *A. annua*. Gene *FS* and *tHMG1* were over-expressed in a high-copy plasmid in *S. cerevisiae* CEN.PK113-5D, which produced 170 mg/L farnesene in fed-batch fermentations with a respiratory quotient (RQ)-controlled exponential feeding (Tippmann et al. , 2016). After *FS* gene was replaced by santalene synthase from *Clausena lansium*, this engineered *S. cerevisiae* produced 163 mg/L santalene under RQ-controlled exponential feeding (Tippmann et al. , 2016). Before this, dynamic control of gene expression including using P_{HXT1} promoter to control *ERG9* expression, overexpression of gene *tHMGR* and deletion of gene *LPPI* and *DPPI* led strain to produce 92 mg/L α -santalene by diverting the carbon flux FPP towards α -santalene (Scalcinati et al. , 2012a). In yeast, protein fusion strategy could also make multi-enzyme complexes for increasing metabolic flux and compound production. Comparing with that observed in a non-fused control, over-expression of Tya-

fused (Tya: Ty1 retrotransposon element) enzymes tHmg1, ispA and FS increased farnesene production 4-fold, yielding 930 mg/L after 7 days in *S. cerevisiae* ATCC200589 (Han et al. , 2018). Specifically, scaffold of FFPS and FS protein through recognizing affibodies to their anti-idiotypic partners, the farnesene yield on glucose was improved by 135% (Tippmann et al. , 2017a). Sequentially, acetoacetyl-CoA synthase (*nphT7*) identified to catalyze the irreversible condensation of malonyl-CoA and acetyl-CoA to acetoacetyl-CoA from *Streptomyces* sp. CL190, was expressed to increase flux, however, it was detrimental for growth and farnesene production (Tippmann et al. , 2017b). Many important insights into microbial metabolism have been achieved using metabolic nodes to disturb metabolic flux and efficiency. Pantothenate (vitamin B5) was utilized as a “metabolic switch” for tuning synthesis rates of molecules relying on CoA intermediates β -farnesene (Sandoval et al. , 2014). To overcome the metabolic inefficiencies including a low ATP/oxygen ratio and challenges of commodity-scale production of anabolic molecules in *S. cerevisiae*, gene aldehyde dehydrogenase acylating (*ADA*), xylulose-5-phosphate specific phosphoketolase (*xPK*), phosphotransacetylase (*PTA*) and NADH-*HMGR* were over-expressed, and DL-glycerol-3-phosphatase (*Rhr2*) was deleted. They rewired the central carbon metabolism, reduced ATP requirement and carbon loss, and improved pathway redox balance of cytosolic acetyl-CoA biosynthesis, leading to farnesene titer exceeded 130 g/L using ~25% less glucose and ~75% less O₂ than previous strains in a fed-batch fermentation (Meadows et al. , 2016, Yu and Chang, 2016). Additionally, evolutionary engineering discovered that truncated tTcb3p protein could increase tolerance of replacement jet fuels blend AMJ-700t (10% cymene, 50% limonene, 40% farnesene) in *S. cerevisiae* (Brennan et al. , 2015).

Compared with *S. cerevisiae* BY4741, industrial *S. cerevisiae* EG60 showed more advantages on cineole and sesquiterpene production levels (Ignea et al. , 2011). In *S. cerevisiae* EG60, through over-expressing gene *IDII*, *HMG2* (K6R) variant, *ERG20* and *SF* homologue

546 P330 from *Salvia pomifera*, and deleting a single copy of *ERG9* gene, the titers of *trans*- β -
547 caryophyllene, α -cubebene, α -copaene and δ -cadinene were respectively ~10 mg/L, ~7 mg/L,
548 ~12 mg/L and ~38 mg/L (Ignea et al. , 2011). This strain was further engineered to produce
549 125 mg/L caryophyllene through replacing *Salvia fruticosa* caryophyllene synthase Sf126 with
550 *SF*, as well as deleting the ubiquitin ligases *ubc7* and *ssm4/doa10* and endoplasmic reticulum
551 associated degradation resident protein *pho86* (Ignea et al. , 2012). In *S. cerevisiae*, over-
552 expressing gene *tHMG1* and down-regulating gene *ERG9* are two widely-used strategies, and
553 the effect on sesquiterpene biosynthesis was specially studied (Asadollahi et al. , 2008,
554 Asadollahi et al. , 2010). *S. cerevisiae* YZL141 was metabolically engineered to be a
555 sesquiterpene production platform by over-expressing gene *tHMG1*, *ERG20* and *FgJ03939*.
556 Based on this platform, sesquiterpene cyclase and skeletons of fusariumdiene and
557 fusagramineol from *Fusarium graminearum* were characterized, identifying 8 compounds -
558 nerolidol, (-)- α -acorenol, (E)- β -farnesene, (+)- α -bisabolol, (-)-acoradiene, fusariumdiene,
559 epi-fusagramineol and fusagramineol (Bian et al. , 2018). Mutation of *ERG9* gene and uptake
560 of exogenous ergosterol (sue) made *S. cerevisiae* WAT11 to produce about 90 mg/L 5-epi-
561 aristolochene, 90 mg/L premnaspirodien and 20 mg/L valencene when respectively
562 introducing gene *TEAS* (tobacco epi-aristolochene synthase), *HPS* (*Hyoscyamus muticus*
563 premnaspirodien synthase) and *CVS* (citrus valencene synthase) (Takahashi et al. , 2007).
564 Thereafter, applying this strategy in *S. cerevisiae* BY4741, co-expressing *HPS* gene with an
565 amino-terminal truncated *tHMGR* gene made it to produce 170.23 mg/L premnaspirodien
566 (Zhuang and Chappell, 2015). Protein fusion engineering is an efficient strategy in
567 sesquiterpene production both in *E. coli* and yeast. Besides farnesene, it was also applied to
568 germacrene A production increasing 6-fold to 190.7 mg/L by over-expressing *tHMGR* gene
569 and fusing FPPS and germacrene A synthase (Hu et al. , 2017), and hirsutene and Δ 6-
570 protoilludene production by fusion of hirsutene synthase (*HS*) from *Stereum hirsutum* to an

HMGS domain (Flynn and Schmidt-Dannert, 2018). Optimizing FPP branch point, modulating MVA pathway, modifying ammonium assimilation pathway and enhancing the transcriptional activator activity as well as a specifically developed continuous fermentation process were combined to produce 0.036 mg/g DCW of α -santalene (Scalcinati et al. , 2012b). Engineering protein degradation of FPPS protein is an effective regulatory strategy to regulate monoterpene and sesquiterpene production in *S. cerevisiae* (Peng et al. , 2018).

Figure 5

Table 2

4.3. Metabolic engineering strategies for sesquiterpene production in other microorganisms

Besides *E. coli* and *S. cerevisiae*, other microorganisms are also engineered to produce sesquiterpenes, such as *Corynebacterium glutamicum*, *B. Subtilis*, *Lactococcus lactis*, cyanobacterium and some eucaryons. The metabolic engineering and strategies for sesquiterpene production in these microorganisms are summarized in Table 3. In *C. glutamicum*, due to the low provision of FPP, heterologous expression of CVS gene from *Citrus sinensis* could not enable (+)-valencene production. Upon deleting two endogenous prenyltransferase genes and expressing either gene *ispA* from *E. coli* or gene *ERG20* from *S. cerevisiae*, (+)-valencene production was observed. Employing CVS gene from *Nootka cypress* further improved (+)-valencene titer to 2.41 mg/L, increasing 10-fold comparing to VS gene from *C. sinensis* (Frohwitter et al. , 2014).

B. subtilis and *Lactococcus lactis* are two Gram-positive prokaryotes reported to be engineered for sesquiterpene production. *B. subtilis* is a promising microbial host for chemical production purpose because of its fast growth rate and generally regarded as safe status. In *B. subtilis*, protein translation engineering was conducted by fusing the highly positive charged six-arginine tag to ADS to increase protein expression. Coupled with expressing gene *Dxs* and *Idi* under P_{xyIA} and systematic media optimization, ~20 mg/L amorphadiene was produced in

shake flask scale (Zhou et al. , 2013). *L. lactis*, harboring MVA pathway, is also an attractive host for characterizing sesquiterpene production: TPS *VMPSTS* from *Vanda Mimi Palmer* was characterized multiple sesquiterpenes with the dominating germacrene D (Song et al. , 2012). Comparing with the β -sesquiphellandrene titer when only expressing β -sesquiphellandrene synthase, co-expression of *mvk* and *mvaA* gene doubled the amount to 15.5 mg/L (Song et al. , 2014). An industrial bacterium *Rhodobacter sphaeroides* engineered by co-expressing MVA operon from *Paracoccus zeaxanthinifaciens* and *CVS* from *Callitropsis nootkatensis* produced 352 mg/L valencene in shaker flasks, 14-fold higher than that of only *CVS* expressing in *S. cerevisiae* (Beekwilder et al. , 2014).

Cyanobacterium, as a photosynthetic prokaryote, was engineered to convert CO₂ and H₂O directly into high-valued biofuel sesquiterpenes (Lin and Pakrasi, 2019). Filamentous cyanobacterium *Anabaena sp.* PCC 7120 produced a farnesene photosynthetic productivity of 69.1 $\mu\text{g}\cdot\text{L}^{-1}\cdot\text{OD}^{-1}\cdot\text{d}^{-1}$ through its endogenous MEP pathway and *coFS* gene using CO₂, mineralized water and light (Halfmann et al. , 2014). Unicellular cyanobacterium *Synechococcus elongatus* PCC 7942 produced 4.6 mg/L farnesene or 19.8 mg/L amorphadiene when over-expressing MEP pathway gene *Dxs*, *Idi* and *ispA*, with *FS* gene from apple or *ADS* gene from *A. annua* with a dodecane overlay in 7 days (Choi et al. , 2016, Lee et al. , 2017). With *QHS1* gene from *A. annua* integrated into cyanobacterium *Synechocystis sp.* PCC6803 genome, $\sim 4 \text{ mg gDCW}^{-1}$ β -caryophyllene was measured by a methanol: chloroform extraction method (Reinsvold et al. , 2011). *Synechococcus sp.* PCC 7002 engineered by expressing *A. grandis* *BIS* gene and deleting *glgC* gene produced more than 0.6 mg/L bisabolene with a dodecane overlay (Davies et al. , 2014). Recently, through genetic engineering, serial enzyme loading and artificial micro RNA-based repression of competing pathways, green microalga *Chlamydomonas reinhardtii* produced 10.3 mg bisabolene $\cdot \text{g}^{-1}$ CDW, yielding up to 11.0 mg/L by mixotrophic cultivation in 7 days (Wichmann et al. , 2018).

For eukaryote microorganism, *Yarrowia lipolytica*, *Rhodospiridium toruloides*, *Streptomyces venezuelae* and *Xanthophyllomyces dendrorhous* are reported to produce sesquiterpenes for their simple industrial scale-up operations. *Y. lipolytica* Po1h, with *tHMG1*, *Idi*, *ERG20* and *coFS* gene integrated into the genome, produced 57.08 mg/L of α -farnesene, a 20.8-fold increase over the initial titer in shake flasks with YPD medium. Bioreactor scale-up in PM medium led α -farnesene concentration up to 259.98 mg/L (Yang et al. , 2016c). The carotenogenic yeast *R. toruloides* was reported to utilize lignocellulose hydrolysate to produce bisabolene and amorphadiene. *R. toruloides* showed superior growth in corn stover hydrolysates prepared by two pretreatment methods: biocompatible IL choline α -ketoglutarate resulting in 261 mg/L bisabolene at bench scale; and an alkaline resulting in 680 mg/L bisabolene in a high-gravity fed-batch bioreactor (Yaegashi et al. , 2017). In *S. venezuelae*, 10.52 mg/L bisabolene was produced after deleting FPP and GPP consumption gene *SVEN_0269* and *SVEN_7111*, and inserting P_{SV} -coAgBis and P_{ermEp} -FPPS1 at the *aatP* chromosomal site (Phelan et al. , 2015). Comparing α -cuprenene production in *E. coli*, *S. cerevisiae* and *X. dendrorhous* expressing *Cop6* gene from *C. cinereus*, α -cuprenene titer in *X. dendrorhous* reached to 80 mg/L, far higher than the other mutants (Melillo et al. , 2013).

Table 3

5. Conclusions and perspective

Biosynthesizing sesquiterpene by engineered microorganisms is a worthy-studied theme for their widely promising applications and values. More and more sesquiterpenes and their synthases are discovered from all kinds of organisms. Although great values contained in sesquiterpenes, minute amount of sesquiterpenes are biosynthesized by native producers. Benefit from the development of metabolic engineering and synthetic biology, microorganisms have become tractable cell factories for sustainable biomanufacturing of the high-value sesquiterpenes. Recent advances in cells and process engineering have made fermentation to

be an attractive route for sesquiterpene production from renewable, low-cost biomass feedstocks.

Sesquiterpene production depends on metabolic strategies, and metabolic strategies also need biotechnology innovation. Recent progress in synthetic biology of terpenoid synthases has opened new avenues of research by providing diversities of strategies. Biotechnological applications made possible by recent molecular advances including the engineering of industrial microbiology and regulation of two isoprenoid pathways to increase the desirable sesquiterpenes. The designing, engineering and optimizing processes are served to establish a high efficiency fermentation-based process and stable sesquiterpene production hosts. Although there have been successful applications of sesquiterpenes, like amorphadiene (Westfall et al. , 2012) and farnesene (Harvey et al. , 2014), more studies and challenges need to be performed and overcome to expand its transformational efficiency, stability and recovery. For example, the use of solvents reduced sesquiterpene evaporation during fermentation and improved product recovery, but it resulted in higher costs, due to the additional equipment cost for solvent-product separation (Cuesta et al. , 2019). Further understanding of the enzymatic mechanism and optimal expression of isoprenoid pathway enzymes will be necessitated. Challenges on making bio-based sesquiterpene more competitive in applications come from aspects of searching high-activity enzymes, supplying energy and cofactor, and balancing metabolic flux, as well as efficient recovery and purification of sesquiterpenes from fermentation broth. Furthermore, issues concerning microbial biosynthesis for achieving higher production help to develop approaches of identifying regulatory effects in metabolic pathways for redirecting carbon fluxes to terpenoid pathways.

Ethical Statement

This article does not contain any studies with human participants performed by any of the authors.

Funding

This work was supported by the Natural Science Foundation of Jiangsu Province, the Fundamental Research Funds for the Central Universities (no. JUSRP11963), China Postdoctoral Foundation (2019M651696), the 111 Project (111-2-06) and national first-class discipline program of Light Industry Technology and Engineering (LITE2018-24). This work also was part of the DOE Joint BioEnergy Institute (<http://www.jbei.org>) supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U.S. Department of Energy.

Competing interests

The authors declare that they have no competing interests.

References

- Abdallah, II., van Merkerk R., Klumpenaar E., Quax W.J., 2018. Catalysis of amorpho-4,11-diene synthase unraveled and improved by mutability landscape guided engineering. *Sci. rep.* 8:9961.
- Abraham W.R., 2001. Bioactive sesquiterpenes produced by fungi: are they useful for humans as well? *Current Med. Chem.* 8
- Agger S., Lopez-Gallego F., Schmidt-Dannert C., 2009. Diversity of sesquiterpene synthases in the basidiomycete *Coprinus cinereus*. *Molecul. Microbiol.* 72:1181-95.
- Agger S.A., Lopez-Gallego F., Hoyer T.R., Schmidt-Dannert C., 2008. Identification of sesquiterpene synthases from *Nostoc punctiforme* PCC 73102 and *Nostoc sp.* strain PCC 7120. *J. bacteriol.* 190:6084-96.
- Ali H., Solsvik J., Wagner J.L., Zhang D., Hellgardt K., Park C.W., 2019. CFD and kinetic-based modeling to optimize the sparger design of a large-scale photobioreactor for scaling up of biofuel production. *Biotechnol bioeng.*
- Alonso-Gutierrez J., Kim E.M., Batth T.S., Cho N., Hu Q., Chan L.J.G., Petzold C.J., Hillson N.J., Adams P.D., Keasling J.D., Martin H.G., Lee T.S., 2015. Principal component analysis of proteomics (PCAP) as a tool to direct metabolic engineering. *Metab eng.* 28:123-33.
- Alonso-Gutierrez J., Koma D., Hu Q., Yang Y., Chan LJG, Petzold CJ, Adams P.D., Vickers C.E., Nielsen L.K., Keasling J.D., Lee T.S., 2018. Toward industrial production of isoprenoids in *Escherichia coli*: Lessons learned from CRISPR-Cas9 based optimization of a chromosomally integrated mevalonate pathway. *Biotechnol bioeng.* 115:1000-13.
- Anthony J.R., Anthony L.C., Nowroozi F., Kwon G., Newman J.D., Keasling J.D., 2009. Optimization of the mevalonate-based isoprenoid biosynthetic pathway in *Escherichia*

coli for production of the anti-malarial drug precursor amorpha-4,11-diene. Metab eng.,11:13-9.

Asadollahi M.A., Maury J., Moller K., Nielsen K.F., Schalk M., Clark A., Nielsen J., 2008. Production of plant sesquiterpenes in *Saccharomyces cerevisiae*: effect of ERG9 repression on sesquiterpene biosynthesis. Biotech bioeng. 99:666-77.

Asadollahi M.A., Maury J., Patil K.R., Schalk M., Clark A., Nielsen J., 2009. Enhancing sesquiterpene production in *Saccharomyces cerevisiae* through in silico driven metabolic engineering. Metab eng, 11:328-34.

Asadollahi M.A., Maury J., Schalk M., Clark A., Nielsen J., 2010. Enhancement of farnesyl diphosphate pool as direct precursor of sesquiterpenes through metabolic engineering of the mevalonate pathway in *Saccharomyces cerevisiae*. Biotechnol bioeng., 106:86-96.

Baadhe R.R., Mekala N.K., Parcha S.R., Prameela DY., 2013. Combination of ERG9 repression and enzyme fusion technology for improved production of amorphadiene in *Saccharomyces cerevisiae*. J Anal Methods Chem,140469.

Baadhe R.R., Mekala N.K., Rao P. S., Prameela D. Y., 2014. Optimization of amorphadiene production in engineered yeast by response surface methodology. Biotech., 4:317-24.

Back K., He S., Kim K.U., Dong H.S., 1998. Cloning and bacterial expression of sesquiterpene cyclase, a key branch point enzyme for the synthesis of sesquiterpenoid phytoalexin capsidiol in uv-challenged leaves of *Capsicum annuum*. Plant Cell Physiol., 39:899-904.

Beekwilder J., van Houwelingen A., Cankar K., van Dijk A.D., de Jong R.M., Stoop G., Bosch D., 2014. Valencene synthase from the heartwood of *Nootka cypress* (*Callitropsis nootkatensis*) for biotechnological production of valencene. Plant Biotechnol J., 12:174-82.

730 Benedict C.R., Lu J.L., Pettigrew D.W., Liu J., Stipanovic R.D., Williams H.J., 2001. The
 731 cyclization of farnesyl diphosphate and nerolidyl diphosphate by a purified recombinant
 732 delta-cadinene synthase. *Plant Physiol.* 125:1754-65.

733 Bennett M.H., Mansfield J.W., Lewis M.J., Beale M.H., 2002. Cloning and expression of
 734 sesquiterpene synthase genes from lettuce (*Lactuca sativa* L.). *Phytochem.* 60:255-61.

735 Bian G., Hou A., Yuan Y., Hu B., Cheng S., Ye Z., Di Y., Deng Z., Liu T., 2018. Metabolic
 736 engineering-based rapid characterization of a sesquiterpene cyclase and the skeletons of
 737 fusariumdiene and fusagramineol from *fusarium graminearum*. *Org Lett.* 20:1626-9.

738 Bleeker P.M., Rossana M., Diergaarde P.J., Arjen V.D., Alain T., Kant M.R., Prinsb M., de
 739 Vosb M., Haringa M.A., Schuurinka R.C., 2012. Improved herbivore resistance in
 740 cultivated tomato with the sesquiterpene biosynthetic pathway from a wild relative. *PNAS*
 741 109:20124-9.

742 Block A.K., Vaughan M.M., Schmelz E.A., Christensen S.A., 2019. Biosynthesis and function
 743 of terpenoid defense compounds in maize (*Zea mays*). *Planta*, 249:21-30.

744 Bohlmann F., Zdero C., Lonitz M., 1977. Neue guajen-derivate aus *Parthenium hysterophorus*
 745 und ein weiteres pseudoguajanolid aus *Ambrosia cumanensis*. *Phytochem.*,16:575-7.

746 Bohlmann J., Croteau R., 1999. Diversity and variability of terpenoid defences in conifers:
 747 molecular genetics, biochemistry and evolution of the terpene synthase gene family in
 748 grand fir (*Abies grandis*). *Novartis Found Symp.* 223:132-45; discussion 46-9.

749 Bohlmann J., Meyer-Gauen G., Croteau R., 1998. Plant terpenoid synthases: molecular biology
 750 and phylogenetic analysis. *PNAS.* 95:4126-33.

751 Bouvier F., Rahier A., Camara B., 2005. Biogenesis, molecular regulation and function of plant
 752 isoprenoids. *Prog. Lipid Res.* 44:357-429.

753 Breitmaier E., 2008. Chapter 3. Sesquiterpene: Wiley-VCH Verlag GmbH & Co. KGaA,;

754 Brennan T.C., Williams T.C., Schulz B.L., Palfreyman R.W., Kromer J.O., Nielsen L.K., 2015.
 755 Evolutionary engineering improves tolerance for replacement jet fuels in *Saccharomyces*
 756 *cerevisiae*. Appl Environ Microbiol.;81:3316-25.

757 Brunk E., George K.W., Alonso-Gutierrez J., Thompson M., Baidoo E., Wang G., Petzolda C.
 758 J., McCloskey D., Monk J., Yang L., Battha T. S., Martina H. G., Feistb A., Adams P. D.,
 759 Keasling J.D., Palsson B., Lee T.S., 2016. Characterizing strain variation in engineered *E.*
 760 *coli* using a multi-omics-based workflow. Cell Syst. 2:335-46.

761 Burgard A.P., Pharkya P., Maranas C.D., 2003. Optknock: A bilevel programming framework
 762 for identifying gene knockout strategies for microbial strain optimization.

763 Cai Y., Jia J.W., Crock J., Lin Z.X., Chen X.Y., Croteau R., 2002. A cDNA clone for β -
 764 caryophyllene synthase from *Artemisia annua*. Phytochem, 61:523-9.

765 Cane D.E., 1990. Enzymic formation of sesquiterpenes. Chemical Reviews, 90:26-30.

766 Cane D.E., Watt R.M., 2003. Expression and mechanistic analysis of a germacradienol
 767 synthase from *Streptomyces coelicolor* implicated in geosmin biosynthesis. PNAS
 768 100:1547-51.

769 Cane D. E., Sohng J.-K., Carol L. R., Suzanne R. M., Wu Z., Lloyd M.D., Oliver S. J., Hubbard
 770 B. R., 1994. Pentalenene synthase. purification, molecular cloning, sequencing, and high-
 771 level expression in *Escherichia coli* of a terpenoid cyclase from *Streptomyces* UC5319.
 772 Biochem. 33:12.

773 Cao Y, Zhang R, Liu W, Zhao G, Niu W, Guo J, Xian M., Liu H., 2019. Manipulation of the
 774 precursor supply for high-level production of longifolene by metabolically engineered
 775 *Escherichia coli*. Sci. rep. 9:95.

776 Chen F., Tholl D., Bohlmann J., Pichersky E., 2011. The family of terpene synthases in plants:
 777 a mid-size family of genes for specialized metabolism that is highly diversified throughout
 778 the kingdom. The Plant journal : for cell and molecular biology, 66:212-29.

779 Cho J.Y., Baik K.U., Jung J.H., Park M.H., 2000. In vitro anti-inflammatory effects of
780 cynaropicrin, a sesquiterpene lactone, from *Saussurea lappa*. Eur. J. Pharmacol, 398:399.

781 Choi S.Y., Lee HJ, Choi J, Kim J, Sim SJ, Um Y, Kim Y., Lee T.S., Keasling J.D., Woo H.
782 M., 2016. Photosynthetic conversion of CO₂ to farnesyl diphosphate-derived
783 phytochemicals (amorpha-4,11-diene and squalene) by engineered cyanobacteria.
784 Biotechnol. biofuels. 9:202.

785 Chou W.K., Fanizza I., Uchiyama T., Komatsu M., Ikeda H., Cane D.E., 2010. Genome mining
786 in *Streptomyces avermitilis*: cloning and characterization of SAV_76, the synthase for a
787 new sesquiterpene, avermitilol. J Am Chem Soc. 132:8850.

788 Connolly J.D., Hill R.A., 1991. Dictionary of terpenoids: Chapman & Hall.

789 Croteau R., 2001. Defensive resin biosynthesis in conifers. Annu Rev Plant Physiol Plant Mol
790 Biol.;39:689-724.

791 Croteau R.B., Davis E.M., Ringer K.L., Wildung M.R., 2005. Menthol biosynthesis and
792 molecular genetics. Naturwissenschaften, 92:562-77.

793 Cuesta S.P.-dl, Knopper L., van der Wielen L.A.M., Cuellar M.C., 2019. Techno-economic
794 assessment of the use of solvents in the scale-up of microbial sesquiterpene production
795 for fuels and fine chemicals. Biofuel Bioprod Bior. 13:140-52.

796 Dahl R.H., Zhang F., Alonso-Gutierrez J, Baidoo E, Bath TS, Redding-Johanson AM, Petzold
797 C. J., Mukhopadhyay A., Lee T. S., Adams P. D. Keasling J.D., 2013. Engineering
798 dynamic pathway regulation using stress-response promoters. Nat Biotechnol, 31:1039-
799 46.

800 Davies F.K., Work V.H., Beliaev A.S., Posewitz M.C., 2014. Engineering limonene and
801 bisabolene production in wild type and a glycogen-deficient mutant of *Synechococcus sp.*
802 PCC 7002. Front Bioeng Biotechnol., 2:21.

803 Dini E. G.G., Giubbilini P., Mancini I., Pietra E., 1993. Control of interspecific relationships
804 in marine ciliate protists by most evolved natural products. *Naturwissenschaften*.;80:3.

805 Dirkmann M., Nowack J., Schulz F., 2018. An in Vitro Biosynthesis of Sesquiterpenes Starting
806 from Acetic Acid. *Chem Bio Chem*. 19:6.

807 Drew D., Dueholm B., Weitzel C., Zhang Y., Sensen C., Simonsen H., 2013. Transcriptome
808 analysis of *thapsia laciniata* rouy provides insights into terpenoid biosynthesis and
809 diversity in apiaceae. *Int. J. Mol. Sci*. 14:9080-98.

810 Ecksteinludwig U., Webb R.J, Van Goethem I.D., East J.M., Lee A.G., Kimura M., Neill
811 P.M.O., Bray P.G., Ward S.A., Krishna S., 2003. Artemisinin target the SERCA of
812 *Plasmodium falciparum*. *Nat*. 424:957.

813 Enserink M. 2005. Infectious diseases. Source of new hope against malaria is in short supply.
814 *Sci*. 307:33.

815 FE H., KE M., 1966. Alpha-farnesene in the natural coating of apples. *Nat*. 210:1260.

816 Finefield J.M., Sherman D.H., Kreitman M., Williams R.M., 2012. Enantiomeric natural
817 products: occurrence and biogenesis. *Angew Chem Int Ed Engl*. 51:4802-36.

818 Flynn M.C., Schmidt-Dannert C., 2018. Sesquiterpene synthase–3-hydroxy-3-methylglutaryl
819 coenzyme a synthase fusion protein responsible for hirsutene biosynthesis in *stereum*
820 *hirsutum*. *App. Environ. Microbiol*. 84:18.

821 Frohwitter J., Heider S.A., Peters-Wendisch P., Beekwilder J., Wendisch V.F., 2014.
822 Production of the sesquiterpene (+)-valencene by metabolically engineered
823 *Corynebacterium glutamicum*. *J Biotechnol*. 191:205-13.

824 Furubayashi M., Ikezumi M., Kajiwarra J., Iwasaki M., Fujii A., Li L., Saito K., Umeno D.,
825 2014. A high-throughput colorimetric screening assay for terpene synthase activity based
826 on substrate consumption. *PLoS One* 9:e93317.

827 Giglio S., Jiang J., Saint C.P., Cane D., Monis P.T., 2008. Isolation and characterization of the
828 gene associated with geosmin production in Cyanobacteria. Environ. Sci. Technol.
829 42:8027-32.

830 Gust B., Challis G.L., Fowler K., Kieser T., Chater K.F., 2003. PCR-targeted Streptomyces
831 gene replacement identifies a protein domain needed for biosynthesis of the sesquiterpene
832 soil odor geosmin. PNAS. 100:1541-6.

833 Guzzetti D., Lebrun A., Subileau M., Grousseau E., Dubreucq E., Drone J., 2016. A
834 catalytically competent terpene synthase inferred using ancestral sequence reconstruction
835 strategy. ACS Catalysis, 6:5345-9.

836 Halfmann C., Gu L., Gibbons W., Zhou R., 2014. Genetically engineering cyanobacteria to
837 convert CO₂, water, and light into the long-chain hydrocarbon farnesene. Appl Microbiol
838 Biotechnol. 98:9869-77.

839 Han J.Y., Song J.M., Seo S.H., Wang C., Lee S.G., Lee H., Kim S.W., Choi E.S., 2018. Ty1-
840 fused protein-body formation for spatial organization of metabolic pathways in
841 *Saccharomyces cerevisiae*. Biotechnol Bioeng. 115:694-704.

842 Harada H., Yu F., Okamoto S., Kuzuyama T., Utsumi R., Misawa N., 2009. Efficient synthesis
843 of functional isoprenoids from acetoacetate through metabolic pathway-engineered
844 *Escherichia coli*. Appl Microbiol Biotechnol. 81:915-25.

845 Harrison K.W., Harvey B.G., 2017. Renewable high density fuels containing tricyclic
846 sesquiterpanes and alkyl diamondoids. Sustain Energ Fuels, 1:467-73.

847 Harvey B.G., Meylemans H.A., Gough R.V., Quintana R.L., Garrison M.D., Bruno T.J., 2014.
848 High-density biosynthetic fuels: the intersection of heterogeneous catalysis and metabolic
849 engineering. Physical Chemistry Chemical Physics Pccp. 16:9448-57.

850 Heinrich M., Robles M., West J.E., Br O.D.M., Rodriguez E., 1998. Ethnopharmacology of
851 Mexican asteraceae (Compositae). Annu Rev Pharmacol. 38:539.

852 Hemmerlin A., Hoeffler J.F., Meyer O., Tritsch D., Kagan I.A., Grosdemange-Billiard C.,
853 Rohmer M., Bach T.J., 2003. Cross-talk between the cytosolic mevalonate and the
854 plastidial methylerythritol phosphate pathways in tobacco bright yellow-2 cells. J.
855 Biological Chem. 278:26666.

856 Hu Y., Zhou Y.J., Bao J., Huang L., Nielsen J., Krivoruchko A., 2017. Metabolic engineering
857 of *Saccharomyces cerevisiae* for production of germacrene A, a precursor of beta-elemene.
858 J Ind Microbiol Biotechnol. 44:1065-72.

859 Ignea C., Cvetkovic I., Loupassaki S., Kefalas P., Johnson C.B., Kampranis S.C., Makris A.M.,
860 2011. Improving yeast strains using recyclable integration cassettes, for the production of
861 plant terpenoids. Microbial Cell Factories 10:4.

862 Ignea C., Trika F.A., Kourtzelis I., Argiriou A., Kanellis A.K., Kampranis S.C., Makris A.M.,
863 2012. Positive genetic interactors of HMG2 identify a new set of genetic perturbations for
864 improving sesquiterpene production in *Saccharomyces cerevisiae*. Microbial Cell
865 Factories 11:162-.

866 Irmisch S., Krause S.T., Kunert G., Gershenzon J., Degenhardt J., Kollner T.G., 2012. The
867 organ-specific expression of terpene synthase genes contributes to the terpene
868 hydrocarbon composition of chamomile essential oils. BMC plant biology 12:84.

869 Izquierdo-Bueno I., González-Rodríguez V.E., Simon A., Dalmais B., Pradier J.M., Le P.P.,
870 Mercier A., Walker A.S., Garrido C., Collado I.G., Viaud M., 2018. Biosynthesis of
871 abscisic acid in fungi: Identification of a sesquiterpene cyclase as the key enzyme in
872 *Botrytis cinerea*. Environ Microbiol. 20(7):2469-2482.

873 Jeleń H.H. 2002. Volatile sesquiterpene hydrocarbons characteristic for penicillium roqueforti
874 strains producing prtoxin. J Agr Food Chem. 50:6569-74.

875 Jin J., Kim M.J., Dhandapani S., Tjhang J.G., Yin J.L., Wong L., Sarojam R., Chua N.H., Jang
876 I.C., 2015. The floral transcriptome of ylang ylang (*Cananga odorata* var. *fruticosa*)

877 uncovers biosynthetic pathways for volatile organic compounds and a multifunctional and
878 novel sesquiterpene synthase. *J Exp Bot.* 66:3959-75.

879 Jin Z., Kwon M., Lee A.R., Ro D.K., Wungsintaweekul J., Kim S.U., 2018. Molecular cloning
880 and functional characterization of three terpene synthases from unripe fruit of black
881 pepper (*Piper nigrum*). *Arch Biochem Biophys* 638:35-40.

882 Köllner T.G., Held M., Lenk C., Hiltbold I., Turlings T.C.J., Gershenzon J., Degenhardt J.,
883 2008. A maize (e)- β -caryophyllene synthase implicated in indirect defense responses
884 against herbivores is not expressed in most american maize varieties. *Plant Cell* 20:482-
885 94.

886 Khalilov L.M., Khalilova A.Z., Odinkov V.N., Baltaev U.A., Paramonov E.A., Dzhemilev
887 U.M., 1999. Identification and biological activity of the volatile organic substances
888 emitted by plants and insects II. Sesquiterpene composition of the native scent of leaves
889 of the potato *Solanum tuberosum*. *Chem Nat Compd.* 35:422-6.

890 Kim E.M., Min H.W., Tian T., Yilmaz S., Javidpour P., Keasling J.D., Lee T.S., 2017.
891 Autonomous control of metabolic state by a quorum sensing (QS)-mediated regulator for
892 bisabolene production in engineered *E. coli*. *Metab eng* 44:325-336.

893 Kirby J., Nishimoto M., Chow R.W., Baidoo E.E., Wang G., Martin J., Schackwitz W., Chan
894 R., Fortman J.L., Keasling J.D., 2015. Enhancing Terpene yield from sugars via novel
895 routes to 1-deoxy-d-xylulose 5-phosphate. *Appl Environ Microbiol* 81:130-8.

896 Kirby J., Nishimoto M., Chow R.W.N., Pasumarthi V.N., Chan R., Chan L.J.G., Petzold C. J.,
897 Keasling J. D., 2014. Use of Nonionic Surfactants for Improvement of Terpene Production
898 in *Saccharomyces cerevisiae*. *Appl Environ Microbiol.* 6685–6693.

899 Kizer L., Pitera D.J., Pfleger B.F., Keasling J.D., 2008. Application of functional genomics to
900 pathway optimization for increased isoprenoid production. *Appl Environ Microbiol.*
901 74:3229-41.

902 Knight D.W., 1995. Feverfew: chemistry and biological activity. *Nat Prod Rep.* 12:271-6.
 903 Koe B.K., Sobin B.A., Celmer W.D., 1957. PA 132, a new antibiotic. I. Isolation and chemical
 904 properties. *Antibiot Annu.* 672-5.
 905 Komatsu M., Tsuda M., Ōmura S., Oikawa H., Ikeda H., 2008. Identification and functional
 906 analysis of genes controlling biosynthesis of 2-methylisoborneol. *PNAS* 105:7422.
 907 Kraker J.W.D., Franssen M.C.R., Groot A.D., König W.A., Bouwmeester H.J., 1998. (+)-
 908 Germacrene A Biosynthesis. *Plant Physiol.* 117.
 909 Kramer R., Abraham W.-R., 2011. Volatile sesquiterpenes from fungi: what are they good for?
 910 *Phytochem Rev.* 11:15-37.
 911 Kumeta Y., Ito M., 2010. Characterization of δ -guaiene synthases from cultured cells of
 912 *aquilaria*, responsible for the formation of the sesquiterpenes in agarwood. *Plant Physiol.*
 913 154:1998.
 914 Kwok B.H.B., Koh B., Ndubuisi M.K.I., Eloffson M., Crews CM., 2001. The anti-
 915 inflammatory natural product parthenolide from the medicinal herb Feverfew directly
 916 binds to and inhibits I κ B kinase. *Chem Biol.* 8:759-66.
 917 Lamers Y.M.A.W., 2003. (+)-Aromadendrene as chiral starting material for the synthesis of
 918 fragrances and pheromones. *Modern Educational Technology.*
 919 Lange B.M., Turner G.W., 2013. Terpenoid biosynthesis in trichomes—current status and
 920 future opportunities. *Plant Biotechnol J.* 11:2–22.
 921 Larsen T.O., 1998. Volatile flavour production by *Penicillium caseifulvum*. *Int Dairy J.* 8:883-
 922 7.
 923 Lee H.J., Lee J., Lee S.M., Um Y., Kim Y., Sim S.J., Choi J.I., Woo H.M., 2017. Direct
 924 conversion of co₂ to alpha-farnesene using metabolically engineered *Synechococcus*
 925 *elongatus* PCC 7942. *J Agric Food Chem.* 65:10424-8.

926 Li Y., Chen F., Li Z., Li C., Zhang Y., 2016a. Identification and functional characterization of
 927 sesquiterpene synthases from *Xanthium strumarium*. *Plant Cell Physiol.* 57:630.
 928 Li Z., Gao R., Hao Q., Zhao H., Cheng L., He F., Liu L., Liu X., Chou W.K., Zhu H., Cane
 929 D.E., 2016b. The T296V mutant of amorpho-4,11-diene synthase is defective in allylic
 930 diphosphate isomerization but retains the ability to cyclize the intermediate (3r)-nerolidyl
 931 diphosphate to amorpho-4,11-diene. *Biochem.* 55:6599-604.
 932 Lin P.C., Pakrasi H.B., 2019. Engineering cyanobacteria for production of terpenoids. *Planta*
 933 249:145-54.
 934 Lin X., Hopson R., Cane D.E., 2006a. Genome mining in *Streptomyces coelicolor* molecular
 935 cloning and characterization of a new sesquiterpene synthase. *J Am Chem Soc.*128:6022-
 936 3.
 937 Lin X., Hopson R., Cane D.E., 2006b. Genome mining in *Streptomyces coelicolor*: molecular
 938 cloning and characterization of a new sesquiterpene synthase. *J Am Chem Soc.* 128:6022-
 939 3.
 940 Little D.B., Croteau R.B., 2002. Alteration of product formation by directed mutagenesis and
 941 truncation of the multiple-product sesquiterpene synthases delta-selinene synthase and
 942 gamma-humulene synthase. *Arch Biochem Biophys.* 402:120-35.
 943 Little J.E., Foote M.W., Johnstone D.B., Xanthatin: an antimicrobial agent from *Xanthium*
 944 pennsylvanicum. *Arch Biochem.* 1950;27:247-54.
 945 Liu C.-L., Tian T., Alonso-Gutierrez J., Garabedian B., Wang S., Baidoo E.E.K., Benites V.,
 946 Chen Y., Petzold C.J., Adams P.D., Keasling J.D., Tan T., Lee T.S., 2018. Renewable
 947 production of high density jet fuel precursor sesquiterpenes from *Escherichia coli*.
 948 *Biotechnol biofuels.*; 11.
 949 Luo F., Ling Y., Li D.S., Tang T., Liu Y.C., Liu Y., Li S.H., 2019. Characterization of a
 950 sesquiterpene cyclase from the glandular trichomes of *Leucosceptrum canum* for sole

951 production of cedrol in *Escherichia coli* and *Nicotiana benthamiana*. *Phytochem* 162:121-
 952 8.

953 Lv F., Li S., Feng J., Liu P., Gao Z., Yang Y., Xu Y., Wei J., 2019. Hydrogen peroxide burst
 954 triggers accumulation of jasmonates and salicylic acid inducing sesquiterpene
 955 biosynthesis in wounded *Aquilaria sinensis*. *J Plant Physiol* 234-235:167-75.

956 Ma S.M., Garcia D.E., Redding-Johanson A.M., Friedland G.D., Chan R., Batth T.S.,
 957 Haliburton J.R., Chivian D., Keasling J.D., Petzold C.J., Lee T.S., Chhabra S.R., 2011.
 958 Optimization of a heterologous mevalonate pathway through the use of variant HMG-
 959 CoA reductases. *Metab eng* 13:588-97.

960 Marrero P.F., Poulter C.D., Edwards P.A., 1992. Effects of site-directed mutagenesis of the
 961 highly conserved aspartate residues in domain II of farnesyl diphosphate synthase activity.
 962 *J Biol Chem.* 267:21873.

963 Martin V.J., Pitera D.J., Withers S.T., Newman J.D., Keasling J.D., 2003. Engineering a
 964 mevalonate pathway in *Escherichia coli* for production of terpenoids. *Nat Biotechnol.*
 965 21:796-802.

966 Maruyama T., Ito M., Honda G., 2001. Molecular cloning, functional expression and
 967 characterization of (E)-beta farnesene synthase from *Citrus junos*. *Biol Pharm Bull.*
 968 24:1171-5.

969 Massoud A.M., Khateeb R.M.E., Shalaby H.A., Kutkat M.A., 2010. Efficacy of volatile oils
 970 (curzerene, furanoeudesma-1, 3-diene and lindestrene) on avian coccidiosis under
 971 laboratory conditions. *J Egypt Soc Parasitol.* 40:699-706.

972 Meadows A.L., Hawkins K.M., Tsegaye Y., Antipov E., Kim Y., Raetz L., Dahl R.H., Tai A.,
 973 Mahatdejkul-Meadows T., Xu L., Zhao L., Dasika M.S., Murarka A., Lenihan J., Eng D.,
 974 Leng J.S., Liu C.L., Wenger J.W., Jiang H., Chao L., Westfall P., Lai J., Ganesan S.,
 975 Jackson P., Mans R., Platt D., Reeves C.D., Saija P.R., Wichmann G., Holmes V.F.,

Benjamin K., Hill P.W., Gardner T.S., Tsong A.E., 2016. Rewriting yeast central carbon
 metabolism for industrial isoprenoid production. *Nat.* 537:694-7.

Melillo E., Setroikromo R., Quax W.J., Kayser O., 2013. Production of alpha-cuprenene in
Xanthophyllomyces dendrorhous: a step closer to a potent terpene biofactory. *Microbial*
Cell Factories 12:13.

Merfort I., 2002. Review of the analytical techniques for sesquiterpenes and sesquiterpene
 lactones. *J Chromatogr A.* 967:115-30.

Mori K., Matsushima Y., 1995. Synthesis of mono and sesquiterpenoids; xxiv: (-)-
 homogynolide a, an insect antifeedant isolated from *Homogyne alpina*. *Synthesis* 845-50.

Mullin C.A., Alfatafta A.A., Harman J.L., Everett S.L., Serino A.A., 1991. Feeding and toxic
 effects of floral sesquiterpene lactones, diterpenes, and phenolics from sunflower
 (*Helianthus annuus* L.) on western corn rootworm. *Jagricfood Chem* 39:2293-9.

Naya Y., Miyamoto F., Takemoto T., 1978. Formation of enantiomeric sesquiterpenes in the
 secretions of scale insects. *Experientia* 34:984-6.

Newman J.D., Marshall J., Chang M., Nowroozi F., Paradise E., Pitera D., Newman K.L.,
 Keasling J.D., 2006. High-level production of amorpho-4,11-diene in a two-phase
 partitioning bioreactor of metabolically engineered *Escherichia coli*. *Biotechnol bioeng.*
 95:684-91.

Nibret E., Youns M., Krauthsiegel R.L., Wink M., 2011. Biological activities of xanthatin from
Xanthium strumarium leaves. *Phytother Res* 25:1883-90.

Nieuwenhuizen N.J., Wang M.Y., Matich A.J., Green S.A., Chen X., Yauk Y.K., Beuning L.L.,
 Nagegowda D.A., Dudareva N., Atkinson R.G., 2009. Two terpene synthases are
 responsible for the major sesquiterpenes emitted from the flowers of kiwifruit (*Actinidia*
deliciosa). *J Exp Bot.* 60:3203-19.

1000 Niwa M., Iguchi M., Yamamura S., 2008. Co-occurrence of (-) and (+)-germacrene-d in
 1001 *Solidago altissima* L. : determination of the optical rotation of optically pure germacrene-
 1002 D. Chempharmbull 28:997-9.

1003 Nour A.M., Khalid S.A., Kaiser M., Brun R., Abdallah W.E., Schmidt T.J., 2009. The
 1004 antiprotozoal activity of sixteen asteraceae species native to Sudan and bioactivity-guided
 1005 isolation of xanthanolides from *Xanthium brasiliicum*. Planta Medica 75:1363-8.

1006 Nowroozi F.F., Baidoo E.E.K., Simon E., Redding-Johanson A.M., Batth T.S., Petzold C.J.,
 1007 Keasling J.D., 2014. Metabolic pathway optimization using ribosome binding site variants
 1008 and combinatorial gene assembly. Appl Microbiol Biotechnol. 98:1567-81.

1009 Nybo S.E., Saunders J., McCormick S.P., 2017. Metabolic engineering of *Escherichia coli* for
 1010 production of valerenadiene. J Biotechnol. 262:60-6.

1011 Omura S., Ikeda H., Ishikawa J., Hanamoto A., Takahashi C., Shinose M., Takahashi Y.,
 1012 Horikawa H., Nakazawa H., Osonoe T., Kikuchi H., Shiba T., Sakaki Y., Hattori M., 2001.
 1013 Genome sequence of an industrial microorganism *Streptomyces avermitilis*: deducing the
 1014 ability of producing secondary metabolites. PNAS 98:12215-20.

1015 Opitz S., Nes W.D., Gershenzon J., 2014. Both methylerythritol phosphate and mevalonate
 1016 pathways contribute to biosynthesis of each of the major isoprenoid classes in young
 1017 cotton seedlings. Phytochemistry, 98:110-9.

1018 Ozaydin B., Burd H., Lee T.S., Keasling J.D., 2013. Carotenoid-based phenotypic screen of
 1019 the yeast deletion collection reveals new genes with roles in isoprenoid production. Metab
 1020 eng. 15:174-83.

1021 Paddon C.J., Westfall P.J., Pitera D.J., Benjamin K., Fisher K., McPhee D., Leavell M.D., Tai
 1022 A., Main A., Eng D., Polichuk D.R., Teoh K.H., Reed D.W., Treynor T., Lenihan J., Fleck
 1023 M., Bajad S., Dang G., Dengrove D., Diola D., Dorin G., Ellens K.W., Fickes S., Galazzo
 1024 J., Gaucher S.P., Geistlinger T., Henry R., Hepp M., Horning T., Iqbal T., Jiang H., Kizer

1025 L., Lieu B., Melis D., Moss N., Regentin R., Secrest S., Tsuruta H., Vazquez R.,
 1026 Westblade L.F., Xu L., Yu M., Zhang Y., Zhao L., Lievens J., Covello P.S., Keasling
 1027 J.D., Reiling K.K., Renninger N.S., Newman J.D., 2013. High-level semi-synthetic
 1028 production of the potent antimalarial artemisinin. *Nat.* 496:528-32.

1029 Paradise E.M., Kirby J., Chan R., Keasling J.D., 2008. Redirection of flux through the FPP
 1030 branch-point in *Saccharomyces cerevisiae* by down-regulating squalene synthase.
 1031 *Biotechnol bioeng.* 100:371-8.

1032 Patil K.R., Rocha I., Förster J., Nielsen J., 2005. Evolutionary programming as a platform for
 1033 in silico metabolic engineering. *Bmc Bioinformatics* 6:1-12.

1034 Pazouki L., Niinemets U., 2016. Multi-substrate terpene synthases: their occurrence and
 1035 physiological significance. *Frontiers in plant science* 7:1019.

1036 Pechous S.W., Whitaker B.D., 2004. Cloning and functional expression of an (E, E)-alpha-
 1037 farnesene synthase cDNA from peel tissue of apple fruit. *Planta* 219:84-94.

1038 Peng B., Nielsen L.K., Kampranis S.C., Vickers C.E., 2018. Engineered protein degradation
 1039 of farnesyl pyrophosphate synthase is an effective regulatory mechanism to increase
 1040 monoterpene production in *Saccharomyces cerevisiae*. *Metab eng.* 47:83-93.

1041 Peng B., Plan M.R., Carpenter A., Nielsen L.K., Vickers C.E., 2017. Coupling gene regulatory
 1042 patterns to bioprocess conditions to optimize synthetic metabolic modules for improved
 1043 sesquiterpene production in yeast. *Biotechnol biofuels* 10:43.

1044 Peralta-Yahya P.P., Ouellet M., Chan R., Mukhopadhyay A., Keasling J.D., Lee T.S., 2011.
 1045 Identification and microbial production of a terpene-based advanced biofuel. *Nat comm.*
 1046 2:483.

1047 Petronilho S., Coimbra M.A., Rocha S.M., 2014. A critical review on extraction techniques
 1048 and gas chromatography based determination of grapevine derived sesquiterpenes. *Anal*
 1049 *Chim Acta.* 846:8-35.

1050 Pfaffenrath V., Diener H.C., Fischer M., Friede M., Henneicke-von Zepelin H.H., 2002. The
 1051 efficacy and safety of *Tanacetum parthenium* (feverfew) in migraine prophylaxis--a
 1052 double-blind, multicentre, randomized placebo-controlled dose-response study.
 1053 *Cephalalgia* 22:523-32.

1054 Phelan R.M., Sekurova O.N., Keasling J.D., Zotchev S.B., 2015. Engineering terpene
 1055 biosynthesis in *Streptomyces* for production of the advanced biofuel precursor bisabolene.
 1056 *ACS syn biol.* 4:393-9.

1057 Phulara S.C., Chaturvedi P., Gupta P., 2016. Isoprenoid-based biofuels: homologous
 1058 expression and heterologous expression in prokaryotes. *Appl Environ Microbiol.*
 1059 82:5730-40.

1060 Pitera D.J., Paddon C.J., Newman J.D., Keasling J.D., 2007. Balancing a heterologous
 1061 mevalonate pathway for improved isoprenoid production in *Escherichia coli*. *Metab eng.*
 1062 9:193-207.

1063 Poole A., Llewellyn D.J., 2005. Antisense suppression of a (+)- δ -cadinene synthase gene in
 1064 cotton prevents the induction of this defense response gene during bacterial blight
 1065 infection but not its constitutive expression. *Plant Physiology* 138:516.

1066 Prosser I., Altug I.G., Phillips A.L., König W.A., Bouwmeester H.J., Beale M.H., 2004.
 1067 Enantiospecific (+)- and (–)-germacrene D synthases, cloned from goldenrod, reveal a
 1068 functionally active variant of the universal isoprenoid-biosynthesis aspartate-rich motif.
 1069 *Arch Biochem Biophys* 432:136-44.

1070 Proteau P.J., 1999. ChemInform abstract: probing the non-mevalonate pathway to phytol in
 1071 the cyanobacterium *synechocystis* sp. utex 2470 using deuterium - labeled glucose.
 1072 *Cheminform.* 30.

1073 Pyle B.W., Tran H.T., Pickel B., Haslam T.M., Gao Z., MacNevin G., Vederas J.C., Kim S.U.,
 1074 Ro D.K., 2012. Enzymatic synthesis of valerena-4,7(11)-diene by a unique sesquiterpene
 1075 synthase from the valerian plant (*Valeriana officinalis*). The FEBS J. 279:3136-46.
 1076 Quaderer R., Omura S., Ikeda H., Cane D.E., 2006. Pentalenolactone biosynthesis. molecular
 1077 cloning and assignment of biochemical function to PtlI, a Cytochrome P450 of
 1078 *Streptomyces avermitilis*. J Am Chem Soc. 128:13036.
 1079 Redding-Johanson A.M., Batth T.S., Chan R., Krupa R., Szmidt H.L., Adams P.D., Keasling
 1080 J.D., Lee T.S., Mukhopadhyay A., Petzold C.J., 2011. Targeted proteomics for metabolic
 1081 pathway optimization: application to terpene production. Metab eng. 13:194-203.
 1082 Reinsvold R.E., Jinkerson R.E., Radakovits R., Posewitz M.C., Basu C., 2011. The production
 1083 of the sesquiterpene β -caryophyllene in a transgenic strain of the cyanobacterium
 1084 *Synechocystis*. J. Plant Physiol. 168:848-52.
 1085 Ro D.K., Paradise E.M., Ouellet M., Fisher K.J., Newman K.L., Ndungu J.M., Ho K.A., Eachus
 1086 R.A., Ham T.S., Kirby J., Chang M.C., Withers S.T., Shiba Y., Sarpong R., Keasling J.D.,
 1087 2006. Production of the antimalarial drug precursor artemisinic acid in engineered yeast.
 1088 Nat. 440:940-3.
 1089 Rohlf M., Churchill A., 2011. Fungal secondary metabolites as modulators of interactions
 1090 with insects and other arthropods. Fungal Genet Biol. 48:23-34.
 1091 Rohmer M. A., 1999. Mevalonate-independent route to isopentenyl diphosphate.
 1092 Comprehensive Natural Products Chemistry. 45-67.
 1093 Rohmer M., Knani M., Simonin P., Sutter B., Sahm H., 1993. Isoprenoid biosynthesis in
 1094 bacteria: a novel pathway for the early steps leading to isopentenyl diphosphate. Biochem
 1095 J. 295:8.

1096 Ruegg T.L., Kim E.M., Simmons B.A., Keasling J.D., Singer S.W., Lee T.S., Thelen M.P.,
 1097 2014. An auto-inducible mechanism for ionic liquid resistance in microbial biofuel
 1098 production. Nat comm. 5:3490.

1099 Sacchettini J.C., and Poulter C.D., 1997. Creating isoprenoid diversity. Sci. 277.

1100 Sandoval C.M., Ayson M., Moss N., Lieu B., Jackson P., Gaucher S.P., Horning T., Dahl R.H.,
 1101 Denery J.R., Abbott D.A., Meadows A.L., 2014. Use of pantothenate as a metabolic
 1102 switch increases the genetic stability of farnesene producing *Saccharomyces cerevisiae*.
 1103 Metab eng. 25:215-26.

1104 Savoia D., Avanzini C., Allice T., Callone E., Guella G., Dini F., 2004. Antimicrobial activity
 1105 of euplotin C, the sesquiterpene taxonomic marker from the marine ciliate *Euplotes*
 1106 *crassus*. Antimicrob Agents Chemother 48:3828-33.

1107 Scalcinati G., Knuf C., Partow S., Chen Y., Maury J., Schalk M., Daviet L., Nielsen J., Siewers
 1108 V., 2012a. Dynamic control of gene expression in *Saccharomyces cerevisiae* engineered
 1109 for the production of plant sesquiterpene alpha-santalene in a fed-batch mode. Metab eng.
 1110 14:91-103.

1111 Scalcinati G., Partow S., Siewers V., Schalk M., Daviet L., Nielsen J., 2012b. Combined
 1112 metabolic engineering of precursor and co-factor supply to increase α -santalene.
 1113 Microbial Cell Factories. 11:16.

1114 Schmidt C.O., Bouwmeester H.J., De J.K., König W.A., 1998. Biosynthesis of (+)- and (-)-
 1115 germacrene D in *Solidago canadensis*: isolation and characterization of two
 1116 enantioselective germacrene D synthases. Angew Chem Int Edit. 37:1400–2.

1117 Schwab W., Wust M., 2015. Understanding the constitutive and induced biosynthesis of mono-
 1118 and sesquiterpenes in grapes (*Vitis vinifera*): a key to unlocking the biochemical secrets
 1119 of unique grape aroma profiles. J Agric Food Chem. 63:10591-603.

1120 Schwender, J., Seemann M., Lichtenthaler H.K., Rohmer M., 1996. Biosynthesis of
 1121 isoprenoids (carotenoids, sterols, prenyl side-chains of chlorophylls and plastoquinone)
 1122 via a novel pyruvate/glyceraldehyde 3-phosphate non-mevalonate pathway in the green
 1123 alga *Scenedesmus obliquus*. Biochem J. 316.
 1124 Seemann M., Zhai G., Kraker de J.-W., Paschall M. Chiana, Christianson W.D., Cane D., 2002.
 1125 Pentalenene synthase. analysis of active site residues by site-directed mutagenesis. J Am
 1126 Chem Soc. 124:12.
 1127 Shen H.J., Cheng B.Y., Zhang Y.M., Tang L., Li Z., Bu Y.F., Li X.R., Tian G.Q., Liu J.Z.,
 1128 2016. Dynamic control of the mevalonate pathway expression for improved zeaxanthin
 1129 production in *Escherichia coli* and comparative proteome analysis. Metab eng. 38:180-
 1130 90.
 1131 Shimane M., Sugai Y., Kainuma R., Natsume M., Kawaide H., 2012. Mevalonate-dependent
 1132 enzymatic synthesis of amorphaadiene driven by an ATP-regeneration system using
 1133 polyphosphate kinase. Biosci biotechnol biochem. 76:1558-60.
 1134 Singh A.N., Upadhye A.B., Mhaskar V.V., Dev S., Pol A.V., Naik V.G., 1974. Chemistry of
 1135 lac resin—VII : Pure lac resin—3: Structure. Tetrahedron. 30:3689-93.
 1136 Smit S.J., Vivier M.A., Young P.R., 2019. Linking terpene synthases to sesquiterpene
 1137 metabolism in grapevine flowers. Front Plant Sci 10:177.
 1138 Song A.A., Abdullah J.O., Abdullah M.P., Shafee N., Othman R., Noor N.M., Rahim R.A.,
 1139 2014. Engineering the lactococcal mevalonate pathway for increased sesquiterpene
 1140 production. FEMS microbiol lett. 355:177-84.
 1141 Song A.A., Abdullah J.O., Abdullah M.P., Shafee N., Rahim R.A., 2012. Functional expression
 1142 of an orchid fragrance gene in *Lactococcus lactis*. Int J Mol Sci. 13:1582-97.

1143 Srivastava P.L., Daramwar P.P., Krithika R., Pandreka A., Shankar S.S., Thulasiram H.V.,
 1144 2015. Functional characterization of novel sesquiterpene synthases from indian
 1145 sandalwood, *Santalum album*. Sci rep. 5:10095.

1146 Steele C.L., Crock J., Bohlmann J., Croteau R., 1998. Sesquiterpene synthases from grand fir
 1147 (*Abies grandis*). Comparison of constitutive and wound-induced activities, and cDNA
 1148 isolation, characterization, and bacterial expression of delta-selinene synthase and
 1149 gamma-humulene synthase. J Biol Chem. 273:2078-89.

1150 Su S., Liu X., Pan G., Hou X., Zhang H., Yuan Y., 2015. In vitro characterization of a (E)-
 1151 beta-farnesene synthase from *Matricaria recutita* L. and its up-regulation by methyl
 1152 jasmonate. Gene 571:58-64.

1153 Sun Z., Meng H., Li J., Wang J., Li Q., Wang Y., Zhang Y., 2014. Identification of novel
 1154 knockout targets for improving terpenoids biosynthesis in *Saccharomyces cerevisiae*.
 1155 PLoS One 9:e112615.

1156 Sunesson A., Vaes W., Nilsson C., Blomquist G., Andersson B., Carlson R., 1995.
 1157 Identification of volatile metabolites from five fungal species cultivated on two media.
 1158 Appl Environ Microbiol. 61:2911-8.

1159 Takahashi S., Takeuchi M., Arai M., Seto H., Otake N., 1983. Studies on biosynthesis of
 1160 pentalenolactone. V isolation of deoxypentalenylglucuron. J Antibiot. 36:226.

1161 Takahashi S., Yeo Y., Greenhagen T.B., McMullin T., Song L., Maurina-Brunker J., Rosson
 1162 R., Noel J.P., Chappell J., 2007. Metabolic engineering of sesquiterpene metabolism in
 1163 yeast. Biotechnol Bioeng. 97:11.

1164 Taleb R.I., Najm P., Shebaby W., Boulos J.C., Demirdjian S., Hariri E., El-Sibai M., Daher C.,
 1165 Mroueh M., 2016. β -2-himachalen-6-ol: A novel anticancer sesquiterpene unique to the
 1166 Lebanese wild carrot. J Ethnopharmacol. 190:59-67.

1167 Tassorelli C., Greco R., Morazzoni P., Riva A., Sandrini G., Nappi G., 2005. Parthenolide is
 1168 the component of *tanacetum parthenium* that inhibits nitroglycerin-induced Fos activation:
 1169 studies in an animal model of migraine. *Cephalalgia* 25:612-21.

1170 Tholl D., 2006. Terpene synthases and the regulation, diversity and biological roles of terpene
 1171 metabolism. *Curr Opin Plant Biol.* 9:297-304.

1172 Thomas B.J., 1995. Some new aspects of isoprenoid biosynthesis in plants a review. *Lipids*
 1173 30:12.

1174 Tippmann S., Anfelt J, David F, Rand JM, Siewers V, Uhlen M, Nielsen J., Hudson E.P., 2017a.
 1175 Affibody scaffolds improve sesquiterpene production in *Saccharomyces cerevisiae*. *ACS*
 1176 *syn biol.* 6:19-28.

1177 Tippmann S., Ferreira R., Siewers V., Nielsen J., Chen Y., 2017b. Effects of acetoacetyl-CoA
 1178 synthase expression on production of farnesene in *Saccharomyces cerevisiae*. *J Ind*
 1179 *Microbiol Biotechnol.* 44:911-22.

1180 Tippmann S., Scalcinati G., Siewers V., Nielsen J., 2016. Production of farnesene and santalene
 1181 by *Saccharomyces cerevisiae* using fed-batch cultivations with RQ-controlled feed.
 1182 *Biotechnol Bioeng.* 113:72-81.

1183 Trutko S.M., Dorofeeva L.V., Shcherbakova V.A., Chuvil'Skaia N.A., Laurinavichus K.S.,
 1184 Biniukov V.I., Ostrovskii D.N., Hintz M., Wiesner J., Jomaa H., Akimenko V.K., 2005.
 1185 Prevalence of nonmevalonate and mevalonate pathways for isoprenoid biosynthesis
 1186 among bacteria of different systematic groups. *Microbiol.* 74:153-8.

1187 Tsuruta H., Paddon C.J., Eng D., Lenihan J.R., Horning T., Anthony L.C., Regentin R.,
 1188 Keasling J.D., Renninger N.S., Newman J.D., 2009. High-level production of amorpho-
 1189 4,11-diene, a precursor of the antimalarial agent artemisinin, in *Escherichia coli*. *PLoS*
 1190 *One* 4:e4489.

1191 Vermaas J.V., Bentley G.J., Beckham G.T., Crowley M.F., 2018. Membrane permeability of
 1192 terpenoids explored with molecular simulation. *J Phys Chem B*. 122:10349-61.

1193 Voo S.S., Grimes H.D., Lange B.M., 2012. Assessing the biosynthetic capabilities of secretory
 1194 glands in Citrus peel. *Plant Physiology* 159:81-94.

1195 Wadhwa M., Bachhawat K.A., 2019. A genetic screen for the isolation of mutants with
 1196 increased flux in the isoprenoid pathway of yeast. *Microb Metab Eng*. 1927:16.

1197 Wang C., Yoon S.H., Jang H.J., Chung Y.R., Kim J.Y., Choi E.S., Kim S.W., 2011a. Metabolic
 1198 engineering of *Escherichia coli* for alpha-farnesene production. *Metab eng*. 13:648-55.

1199 Wang C.F., Yang K., Zhang H.M., Cao J., Fang R., Liu Z.L., Du S.S., Wang Y.Y., Deng Z.W.,
 1200 Zhou L., 2011b. Components and insecticidal activity against the maize weevils of
 1201 *Zanthoxylum schinifolium* fruits and leaves. *Molecules*. 16:3077-88.

1202 Wang J., Xiong Z., Li S., Wang Y., 2015. Exploiting exogenous MEP pathway genes to
 1203 improve the downstream isoprenoid pathway effects and enhance isoprenoid production
 1204 in *Escherichia coli*. *Proc Biochem*. 50:24-32.

1205 Wang J.F., Xiong Z.Q., Li S.Y., Wang Y., 2013. Enhancing isoprenoid production through
 1206 systematically assembling and modulating efflux pumps in *Escherichia coli*. *Appl*
 1207 *Microbiol Biotechnol*. 97:8057-67.

1208 Wang R.L., Peng S.L., Zeng R.S., Ding L.W., Xu Z.F., 2009. Cloning, expression and
 1209 wounding induction of β -caryophyllene synthase gene from *Mikania micrantha* HBK and
 1210 allelopathic potential of β -caryophyllene. *Allelopathy J*. 24:35-44.

1211 Weaver L.J., Sousa M.M., Wang G., Baidoo E., Petzold C.J., Keasling J.D., 2015. A kinetic-
 1212 based approach to understanding heterologous mevalonate pathway function in *E. coli*.
 1213 *Biotechnol Bioeng*. 112:111-9.

1214 Westfall P.J., Pitera D.J., Lenihan J.R., Eng D., Woollard F.X., Regentin R., Horning T.,
 1215 Tsuruta H., Melis D.J., Owens A., Fickes S., Diola D., Benjamin K.R., Keasling J.D.,

1216 Leavell M.D., McPhee D.J., Renninger N.S., Newman J.D., Paddon C.J., 2012.
 1217 Production of amorphadiene in yeast, and its conversion to dihydroartemisinic acid,
 1218 precursor to the antimalarial agent artemisinin. PNAS 109:E111-8.
 1219 Weyerstahl P., Marschall H., Christiansen C., Kalembe D., Góra J., 1993. Constituents of the
 1220 essential oil of *Solidago canadensis* ("Goldenrod") from Poland - A Correction. Planta
 1221 Medica. 59:281-2.
 1222 Wichmann J., Baier T., Wentnagel E., Lauersen K.J., Kruse O., 2018. Tailored carbon
 1223 partitioning for phototrophic production of (E)-alpha-bisabolene from the green microalga
 1224 *Chlamydomonas reinhardtii*. Metab eng. 45:211-22.
 1225 Wu W., Fang L., Davis R.W., 2018. Engineering *Escherichia coli* for the production of terpene
 1226 mixture enriched in caryophyllene and caryophyllene alcohol as potential aviation fuel
 1227 compounds. Metab eng Comm. 6:13-21.
 1228 Yaegashi J., Kirby J., Ito M., Sun J., Dutta T., Mirsiaghi M., Sundstrom E.R., Rodriguez A.,
 1229 Baidoo E.,anjore D., Pray T., Sale K., Singh S., Keasling J.D., Simmons B.A., Singer
 1230 S.W., Magnuson J.K., Arkin A.P., Skerker J.M., Gladden J.M., 2017. *Rhodospiridium*
 1231 *toruloides*: a new platform organism for conversion of lignocellulose into terpene biofuels
 1232 and bioproducts. Biotechnol Biofuels. 10:241.
 1233 Yamada Y., Arima S., Nagamitsu T., Johmoto K., Uekusa H., Eguchi T., Shin-ya K., Cane
 1234 D.E., Ikeda H., 2015a. Novel terpenes generated by heterologous expression of bacterial
 1235 terpene synthase genes in an engineered *Streptomyces* host. J Antibiot. 68:385-94.
 1236 Yamada Y., Kuzuyama T., Komatsu M., Shin-Ya K., Omura S., Cane D.E., Ikeda H., 2015b.
 1237 Terpene synthases are widely distributed in bacteria. PNAS 112:857-62.
 1238 Yang J., Li Z., Guo L., Du J., Bae H.-J., 2016a. Biosynthesis of β -caryophyllene, a novel
 1239 terpene-based high-density biofuel precursor, using engineered *Escherichia coli*. Renew
 1240 Energ 99:216-23.

1241 Yang J., Nie Q., 2016. Engineering *Escherichia coli* to convert acetic acid to β -caryophyllene.
1242 Microbial Cell Factories. 15:1-9.

1243 Yang L., Wang C., Zhou J., Kim S.W., 2016b. Combinatorial engineering of hybrid mevalonate
1244 pathways in *Escherichia coli* for protoilludene production. Microb Cell Fact. 15:14.

1245 Yang X., Nambou K., Wei L., Hua Q., 2016c. Heterologous production of alpha-farnesene in
1246 metabolically engineered strains of *Yarrowia lipolytica*. Bioresour technol. 216:1040-8.

1247 Yoshikuni Y., Dietrich J.A., Nowroozi F.F., Babbitt P.C., Keasling J.D., 2008. Redesigning
1248 enzymes based on adaptive evolution for optimal function in synthetic metabolic
1249 pathways. Chem Biol. 15:607-18.

1250 Yoshikuni Y., Ferrin T.E., Keasling J.D., 2006. Designed divergent evolution of enzyme
1251 function. Nat. 440:1078.

1252 You S., Yin Q., Zhang J., Zhang C., Qi W., Gao L., Tao Z., Su R., He Z., 2017. Utilization of
1253 biodiesel by-product as substrate for high-production of β -farnesene via relatively
1254 balanced mevalonate pathway in *Escherichia coli*. Bioresour Technol. 243:228-36.

1255 Yu V.Y., Chang M.C., 2016. High-yield chemical synthesis by reprogramming central
1256 metabolism. Nat Biotechnol. 34:1128-9.

1257 Yuan J., Ching C.B., 2014. Combinatorial engineering of mevalonate pathway for improved
1258 amorpho-4,11-diene production in budding yeast. Biotechnol Bioeng. 111:608-17.

1259 Yuan J.S., Köllner T.G., Wiggins G., Grant J., Degenhardt J., Chen F., 2008. Molecular and
1260 genomic basis of volatile-mediated indirect defense against insects in rice.

1261 Zhang C., Chen X., Stephanopoulos G., Too H.P., 2016. Efflux transporter engineering
1262 markedly improves amorphadiene production in *Escherichia coli*. Biotechnol Bioeng.
1263 113:1755-63.

1264 Zhang C., Chen X., Zou R., Zhou K., Stephanopoulos G., Too H.P., 2013. Combining genotype
 1265 improvement and statistical media optimization for isoprenoid production in *E. coli*. Plos
 1266 One 8:e75164.

1267 Zhang C., Zou R., Chen X., Stephanopoulos G., Too H.-P., 2015. Experimental design-aided
 1268 systematic pathway optimization of glucose uptake and deoxyxylulose phosphate
 1269 pathway for improved amorphaadiene production. Appl Microbiol Biotechnol. 99:13.

1270 Zhang Q., Margaryan A., Schneidmiller R.G., 2011. Methods for repelling insects using
 1271 sesquiterpene hydrocarbons and their derivatives. US.

1272 Zhao B., Lin X.L., Lamb D., Kelly S., Waterman M., Cane D., 2008. Biosynthesis of the
 1273 sesquiterpene antibiotic albaflavenone in *Streptomyces coelicolor* A3(2). J Biol Chem.
 1274 283:8183.

1275 Zhou K., Zou R., Zhang C., Stephanopoulos G., Too H.P., 2013. Optimization of amorphaadiene
 1276 synthesis in *Bacillus subtilis* via transcriptional, translational, and media modulation.
 1277 Biotechnol Bioeng. 110:2556-61.

1278 Zhu F., Zhong X., Hu M., Lu L., Deng Z., Liu T., 2014. In vitro reconstitution of mevalonate
 1279 pathway and targeted engineering of farnesene overproduction in *Escherichia coli*.
 1280 Biotechnol Bioeng. 111:1396-405.

1281 Zhuang X., Chappell J., 2015. Building terpene production platforms in yeast. Biotechnol
 1282 Bioeng. 112:1854-64.

1283 Zou R., Zhou K., Stephanopoulos G., Too H.P., 2013. Combinatorial engineering of 1-deoxy-
 1284 D-xylulose 5-phosphate pathway using cross-lapping in vitro assembly (CLIVA) method.
 1285 PLoS One 8:e79557.

Table 1. Metabolic engineering of *E. coli* for sesquiterpene biosynthesis.

Sesquiterpene	Strain	Description	Titer	Reference
Amorphadiene	<i>E. coli</i> DYM1	Over-expressing <i>ADS</i> gene from <i>A. annua</i> and MVA pathway genes from <i>S. cerevisiae</i> , with <i>mvk</i> , <i>pmk</i> , <i>mvaD</i> , <i>idi</i> and <i>ispA</i> gene under control of P _{LAC} (plasmid pMBIS); <i>atoB</i> , <i>HMGS</i> and <i>tHMGR</i> gene under control of P _{LAC} (plasmid MevT)	112.2 mg/L	(Martin et al. , 2003)
	<i>E. coli</i> K12 MG1655 Δ recA, Δ endA	Containing plasmid P _{BAD} -Dxs-Idi-ispDF and P _{ACM} -ADS, over-expressing multidrug-resistant membrane efflux transporter protein TolC and ABC family transporters (macAB) or MFS family transporters (emrAB or emrKY)	~180 mg/L	(Zhang et al. , 2016)
	<i>E. coli</i> K12 MG1655	Containing plasmid P _{BAD} -Dxs-Idi-ispDF and P _{ACM} -ADS, deleting PTS genes, using optimized OPT2 medium	182 mg/L	(Zhang et al. , 2013)
	<i>E. coli</i> MG1655 DE3	Δ pstHIerr::FRT, containing plasmid pETA-TM2-galP-glk and pACM-P _{T7} -dxs-P _{T7} -idi-ADS-ispA	201 mg/L	(Zhang et al. , 2015)
	<i>E. coli</i> DP10	Containing plasmid pMBIS, supplementing 20 mM MVA to medium	~290 mg /L	(Pitera et al. , 2007)
	<i>E. coli</i> DH1	Plasmid pAM4: lacUV5 promoters for codon-optimized MevT operon and MBIS operon, Cm ^R , Plasmid pAM94: pTrc99A derivative containing <i>ADS</i> and codon-optimized <i>mvk</i> , Ap ^R ; fermentation for 75 h.	293 mg/L/OD ₆₀₀	(Anthony et al. , 2009)
	<i>E. coli</i> BL21-Gold DE3	Over-expressing <i>Dxs</i> , <i>Idi</i> , <i>ispA</i> and <i>ADS</i> gene	~300 mg/L	(Yuan and Ching, 2014)
	<i>E. coli</i> BL21 (DE3)	Co-expressing <i>Dxs2</i> from <i>S. avermitilis</i> and <i>Idi</i> from <i>B. subtilis</i>	331.7 mg/L	(Wang et al. , 2015)
	<i>E. coli</i> BL21 (DE3)	Harboring pET21c-ADS, P _{T7} -tolC-tolC-acrB operon	404.8 mg/L	(Wang et al. , 2013)
	<i>E. coli</i> W3110 (ATCC 27325)	Containing plasmid pMBIS and MevT, employing a two-phase partitioning bioreactor with a dodecane organic phase	500 mg/L	(Newman et al. , 2006)
	<i>E. coli</i> DH1	Codon-optimizing <i>mvk</i> and <i>pmk</i> gene, and replacing P _{lac} with a stronger P _{trc}	>500 mg/L	(Redding-Johanson et al. , 2011)

	<i>E. coli</i> DH1	Bearing <i>D. acidovorans</i> <i>HMGR</i> by increasing intracellular NADH level using a NADP-dependent formate dehydrogenase (<i>fdhI</i>) from <i>Candida boidinii</i> , along with formate supplementation after 48 h of fermentation.	700 mg/L	(Ma et al. , 2011)
	<i>E. coli</i> DH1	Dynamic control the MVA pathway expression using the identified FPP-responding P_{rstA} and P_{gadE}	1400 mg/L	(Dahl et al. , 2013)
	<i>E. coli</i> DH1	Metabolic pathway optimization using rbs variants and combinatorial gene assembly, pTrc-RBS.ADS-MBIS, using 3% glycerol	3550 mg/L	(Nowroozi et al. , 2014)
	<i>E. coli</i> DH1	Heterologous MVA pathway with <i>HMGS</i> and <i>HMGR</i> from <i>S. aureus</i> , using a fermentation process of restricted glucose and nitrogen with NaOH pH control	29.7 g/L	(Tsuruta et al. , 2009)
α -Farnesene	<i>E. coli</i> DH5a	Codon-optimizing <i>FS</i> gene, augmenting metabolic flux for FPP synthesis by introducing MVA pathway and over-expressing <i>Dxs</i> , <i>Idi</i> and <i>ispA</i> gene, fusing <i>FPPS</i> and <i>FS</i>	380.0 mg/L	(Wang et al. , 2011a)
	<i>E. coli</i> BL21 (DE3)	Developing a reconstituted in vitro farnesene biosynthetic system, rationally designing strain F4 with P_{lac} for <i>ERG12</i> , <i>ERG8</i> , <i>MVD1</i> and <i>Idi</i> gene with pBBR1MCS ori, and P_{T7} for <i>AFS</i> , <i>ispA</i> and <i>Idi</i> gene with pBBR322 ori	1100 mg/L	(Zhu et al. , 2014)
	<i>E. coli</i> BL21 (DE3)	Expressing heterologous MVA pathway, and high-expressing <i>Idi</i> and <i>ispA</i> gene in a lab bioreactor.	8740 mg/L	(You et al. , 2017)
Bisabolene	<i>E. coli</i> BL21 (DE3)	Harboring plasmid pBbA1k-AgBIS-ispA-ispDF-idi, fusing nDXP gene <i>ribB</i> to <i>Dxr</i> gene,	8.8 mg/g DCW	(Kirby et al. , 2015)
	<i>E. coli</i> DH1	F-, λ -, endA1, recA1, gyrA96, thi-1, glnV44, relA1, hsdR17(rK- mK+), DSMBT7, rbsR:: P_{T7} -BS-idi-ispA, poxB:: P_{T7} -MevT	~435 μ g/L	(Alonso-Gutierrez et al. , 2018)
	<i>E. coli</i> DH1	Screening BIS enzyme, codon-optimizing <i>HMGS</i> , <i>tHMGR</i> , <i>mvk</i> and <i>pmk</i> gene, and inserting a strong P_{trc} before <i>mvk</i> gene (pJBEI-2999)	> 900 mg/L	(Peralta-Yahya et al. , 2011)
	<i>E. coli</i> DH1	Using quorum sensing LuxI/R system from <i>Vibrio fischeri</i> to construct sensor plasmid carrying <i>luxI-luxR</i> genes, and response plasmid carrying bisabolene producing pathway genes under control of P_{luxI}	1100 g/L	(Kim et al. , 2017)

	<i>E. coli</i> DH1	Harboring plasmid pBbA5c-CL1-T61-CL2-T21-AgBISppTrc-AgBIS	1150 mg/L	(Alonso-Gutierrez et al. , 2015)
β -Caryophyllene	<i>E. coli</i> DH1	Expressing MVA pathway and <i>QHS1</i> gene	100 mg/L	(Wu et al. , 2018)
	<i>E. coli</i> BL21 (DE3)	Co-expressing <i>ACS</i> , <i>QHS1</i> , <i>GPPS2</i> , <i>AACS</i> gene and heterologous MVA pathway	1050 mg/L	(Yang and Nie, 2016)
	<i>E. coli</i> BL21 (DE3)	Over-expressing MVA pathway and <i>GPPS</i> , <i>G6PH</i> , <i>QHS1</i> genes, fed-batch fermentation for 60 h	1520 mg/L	(Yang et al. , 2016a)
α -Humulene	<i>E. coli</i> BL21 (DE3)	Expressing <i>Streptomyces</i> MVA pathway genes from <i>HMGS</i> to <i>Idi2</i> , yeast <i>Idi1</i> and rat <i>Aacl</i> gene, α -humulene synthase gene of shampoo ginger, supplying with LAA	958 mg/L	(Harada et al. , 2009)
5-Epi-aristolochene	<i>E. coli</i> XL1-Blue	P _{lac} -crtM-crtN, <i>TEAS</i> from <i>Nicotiana tabacum</i> , using the screened TEASmut7 Q481R by scoring their substrate consumption based on color loss of cell harboring carotenoid pathways	160 mg/L	(Furubayashi et al. , 2014)
Protoilludene	<i>E. coli</i> DH5 α	Over-expressing a exogenous MVA pathway, endogenous <i>ispA</i> , and protoilludene synthase of <i>Omphalotus olearius</i>	1199 mg/L	(Yang et al. , 2016b)
Valerenadiene	<i>E. coli</i> BL21 (DE3)	Codon-optimizing gene <i>cvds</i> , over-expressing <i>Dxs</i> , <i>Idi</i> and <i>FPPS</i> , optimizing fermentation medium with glycerol as supplementary	11 mg/L	(Nybo et al. , 2017)
Longifolene	<i>E. coli</i> BL21 (DE3)	pTrcLower: pTrcHis2B harboring <i>S. cerevisiae</i> <i>erg19</i> , <i>erg8</i> , <i>erg12</i> and <i>idi1</i> ; pA-mvaESispALFS: pACYCDuet-1 harboring <i>E. faecalis</i> <i>mvaE</i> , <i>mvaS</i> , <i>E. coli</i> <i>ispA</i> and <i>P. abies</i> longifolene synthase	382 mg/L	(Cao et al. , 2019)

1287

1288

Table 2. Metabolic engineering of *S. cerevisiae* for sesquiterpene biosynthesis.

Sesquiterpene	Strain	Description	Titer	Reference
Amorphadiene	<i>S. cerevisiae</i> WAT11	Using flux balance analysis (FBA) and MOMA methods for predicting knockout targets, strain MATa (leu2-3,112 trp1-1 can1-100 ura3-1 ade2-1 his3-11,15, <i>hvk2D::his3::ADS</i>)	54.6 mg/L	(Sun et al. , 2014)
	<i>S. cerevisiae</i> EPY224	Upregulating <i>HMGS</i> , <i>mvk</i> and <i>pmk</i> gene by <i>upc2-1</i> expression; over-expressing a truncated <i>tHMGR</i> , <i>ispA</i> and <i>ADS</i> gene; <i>erg9</i> : P _{met3} -ERG9	153 mg/L	(Ro et al. , 2006)
	<i>S. cerevisiae</i> CEN.PK2 Y293	Genotype: MATa, <i>erg9Δ::kan^r</i> P _{MET3} -ERG9, leu2-3,112::HIS P _{GAL1} -MVD1, P _{GAL10} -ERG8, his3Δ1::HIS, P _{GAL1} -ERG12, P _{GAL10} -ERG10, <i>ade1Δ::P_{GAL1}-tHMGI</i> , P _{GAL10} -IDI1 ADE1, ura3-52::P _{GAL1} -tHMGI, P _{GAL10} -ERG13, URA3trp1-289::P _{GAL1} -tHMGI, P _{GAL10} -ERG20 TRP1; gal80Δ::nat ^r ; plasmid pAM426: P _{GAL1} -tHMGI. Fermentation with unrestricted ethanol pulse feed with lower oxygen uptake rate (OUR)	41 g/L	(Westfall et al. , 2012)
Bisabolene	<i>S. cerevisiae</i> EPY224	EPY224 platform through replacing <i>ADS</i> gene with <i>BIS</i> gene	900 mg/L	(Peralta-Yahya et al. , 2011)
	<i>S. cerevisiae</i> EPY224	EPY224 platform through replacing <i>ADS</i> gene with <i>BIS</i> gene, enriching a mixed population and removing little or no bisabolene producing strains using selective pressure nonionic surfactants (T20)	375 mg/L	(Kirby et al., 2014a)
	<i>S. cerevisiae</i> CEN.PK2-1C	Genotype: MATa ura3-52 trp1-289 leu2-3,112 his3Δ 1 MAL2-8C SUC2, YPRC15::KANMX-PGAL1-C.O.BisSyn~ERG20/PGAL10-tHMG, gal80::HIS5, ypl064w::hph yjl064w::Ca.URA3, harboring pRS-AgBIS	5.2 g/L	(Ozaydin et al. , 2013)
β-Farnesene	<i>S. cerevisiae</i> CEN.PK113-5D	Genotype: MATa, MAL2-8c, SUC2, ura3-52, <i>lpp1Δ::loxP</i> , <i>dpp1Δ::loxP</i> , PERG9Δ::loxP-P _{HXT1} , P _{PGK1} -Z _{IgA} -ERG20, P _{TEF1} -Z _{Taq} -FarnSyn-Cj, P _{TEF1} -anti-Z _{Taq} -anti-Z _{IgA} with (SSSSG) ₄ linker	0.813 mg/g	(Tippmann et al. , 2017a)
	<i>S. cerevisiae</i> CEN.PK113-5D	Genotype: MATa, MAL2-8 ^c , SUC2, ura3-52, <i>lpp1Δ::loxP</i> , <i>dpp1Δ::loxP</i> , PERG9Δ::loxP-P _{HXT1} , <i>gdh1Δ::loxP</i> , P _{TEF1} -ERG20, P _{PGK1} -GDH2, over-expressing <i>FS</i> and <i>tHMG1</i> gene in a 2-μm plasmid with respiratory quotient-controlled exponential feeding	170 mg/L	(Tippmann et al. , 2016)

	<i>S. cerevisiae</i> ATCC200589	Over-expressing structural component Ty1 retrotransposon element Tya-fused enzyme tHmg1, ispA and FS in a two-phase partitioning fed-batch fermentation for 7 days	930 mg/L	(Han et al. , 2018)
	<i>S. cerevisiae</i> CEN.PK2-1C	Genotype: MAT α , erg9 Δ ::P _{GAL4} (OC)-GAL4_syn_P _{MET3} -ERG9, trp1-289::P _{GAL1} -tHMG1_P _{GAL10} -ERG20_T _{RP1} , leu2-3,112::HISM _X _P _{GAL1} -MVD1_P _{GAL10} -ERG8, ade1 Δ ::P _{GAL1} -tHMGR_P _{GAL10} -IDI1_ADE1, his3 Δ 1::HISM _X _P _{GAL1} -ERG12_P _{GAL10} -ERG10_HIS3, ura3-52:: P _{GAL1} -tHMG1_P _{GAL10} -ERG13_URA3, gal80 Δ ::syn_P _{GAL1} -ERG12, erg9 Δ ::P _{GAL4} (OC)-GAL4_syn_P _{MET3} -ERG9, hxt3 Δ ::syn_P _{TDH3} -ERG10-T _{AHP1} -P _{YPD1} -Bj.HMGs-T _{CCW12} -P _{TUB2} -Ca.THL-T _{HXT3} , dit1 Δ ::syn_P _{GAL1} -Aa.FS_A-T _{CYC1} , ndt80 Δ :: P _{GAL1} -Aa.FS_A-T _{CYC1} -P _{HXT3} -ACS2-T _{PGK1} -P _{GAL7} -Zm.PDC-T _{TDH3} , ho Δ ::syn, gal2 Δ ::syn, adh5 Δ :: P _{GAL1} -Aa.FS_B-T _{ADH1} -P _{GAL10} -Aa.FS_A-T _{CYC1} -LEU2_P _{GAL1} -Aa.FS_B-T _{ADH1} -P _{GAL10} -Aa.FS_A-T _{CYC1} ; ald4 Δ ::P _{GAL1} -Dz.eutE-T _{PGK1} -syn_P _{GAL1} -Dz.eutE-T _{PGK1} , ald6 Δ ::syn, rhr2 Δ ::syn_P _{TDH3} -Lm.PK-T _{TDH3} , erg13 Δ ::syn_P _{HXT1} -GAL4*, ygr250c Δ ::syn_P _{ACT3} -PDR3, bdh1 Δ ::P _{TDH3} -Lm.PK-T _{TDH3} -syn_P _{ACT1} -Ck.PTA-T _{PGK1} , adh1 Δ ::P _{FBA1} -ACS2_syn_P _{TDH3} -Sp.HMGr-T _{HMG1} , P _{TDH3} -ERG13*_P _{GAL10} -Sp.HMGr-T _{HMG1} ; 200,000-L industrial bioreactors.	130 g/L	(Meadows et al. , 2016)
α -Santalene	<i>S. cerevisiae</i> CEN.PK113-5D	Genotype: MAT α MAL2-8c SUC2 ura3-52 lpp1 Δ ::loxP dpp1 Δ ::loxP P _{ERG9} Δ ::loxP-P _{HXT1} , plasmid pICK01 containing a copy of tHMG1 and santalene synthase	92 mg/L	(Scalcinati et al. , 2012a)
	<i>S. cerevisiae</i> CEN.PK113-5D	Genotype: MAT α , MAL2-8 c , SUC2, ura3-52, lpp1 Δ ::loxP, dpp1 Δ ::loxP, P _{ERG9} Δ ::loxP-P _{HXT1} , gdh1 Δ ::loxP, P _{TEF1} -ERG20, P _{PGK1} -GDH2, over-expressing santalene synthase and <i>tHMG1</i> gene in a 2- μ m plasmid with respiratory quotient-controlled exponential feeding	163 mg/L	(Tippmann et al. , 2016)
Caryophyllene	<i>S. cerevisiae</i> EG60	Genotype: Mat a/ α , P _{Gal1} -HMG2-(K6R) x2, P _{Gal1} -HA-ERG20, ura3, trp1, his3, leu2, Δ erg9:: HIS5, pYESmyc/P330, 2 μ , URA3, P _{GAL1} -myc-P330	~ 10 mg/L	(Ignea et al. , 2011)
	<i>S. cerevisiae</i> EG60	Genotype: Mat a/ α , P _{Gal1} -HMG2(K6R):: HOX2, ura3, trp1, his3, P _{TDH3} -HMG2(K6R)X2-::leu2 ERG9/erg9, UBC7/ubc7, SSM4/ssm4, PHO86/pho86	125 mg/L	(Ignea et al. , 2012)
α -Cubebene	<i>S. cerevisiae</i> EG60	Genotype: Mat a/ α , P _{Gal1} -HMG2-(K6R) x2, P _{Gal1} -HA-ERG20, ura3, trp1, his3, leu2, Δ erg9:: HIS5, pYESmyc/P330, 2 μ , URA3, P _{GAL1} -myc-P330	~ 7 mg/L	(Ignea et al. , 2011)

α -Copaene	<i>S. cerevisiae</i> EG60	Genotype: Mat a/ α , P _{Gal1} -HMG2-(K6R) x2, P _{Gal1} -HA-ERG20, ura3, trp1, his3, leu2, Δ erg9:: HIS5, pYESmyc/P330, 2 μ , URA3, P _{GAL1} -myc-P330	~12 mg/L	(Ignea et al. , 2011)
α -Cadinene	<i>S. cerevisiae</i> EG60	Genotype: Mat a/ α , P _{Gal1} -HMG2-(K6R) x2, P _{Gal1} -HA-ERG20, ura3, trp1, his3, leu2, Δ erg9:: HIS5, pYESmyc/P330, 2 μ , URA3, P _{GAL1} -myc-P330	~38 mg/L	(Ignea et al. , 2011)
Pentalenene	<i>S. cerevisiae</i> EPY	Upregulating <i>HMGS</i> , <i>MK</i> and <i>PMK</i> gene by upc2-1 expression; over-expressing <i>tHMGR</i> , <i>ispA</i> and <i>PentS</i> gene; <i>erg9</i> : P _{met3} -ERG9	344 mg/L	(Liu et al. , 2018)
5-Epi-aristolochene	<i>S. cerevisiae</i> WAT11	Mutating gene <i>ERG9</i> , uptake of exogenous ergosterol (sue), introducing gene <i>TEAS</i> in plasmid pESC-HIS3-GPD-TEAS	90 mg/L	(Takahashi et al. , 2007)
Premnaspirodienone	<i>S. cerevisiae</i> WAT11	Mutating gene <i>ERG9</i> , uptake of exogenous ergosterol (sue), introducing gene <i>HPS</i> in plasmid pESC-HIS3-GPD-HPS	90 mg/L	(Takahashi et al. , 2007)
	<i>S. cerevisiae</i> BY4741	Expressing <i>HPS</i> gene with an amino-terminal truncated in plasmid pESC-HIS3-GPD-HPS, catalytic form of tHMGR gene, uptake of exogenous ergosterol (sue)	170.23 mg/L	(Zhuang and Chappell, 2015)
germacrene A	<i>S. cerevisiae</i> SCIGS22a	MATa MAL2-8c SUC2 ura3-52 lpp1 Δ ::loxP dpp1 Δ ::loxP PERG9 Δ ::loxP-P _{HXT1} gdh1 Δ ::loxP P _{TEF1} -ERG20, P _{GK1} -GDH2 P _{TEF1} -tHMG1, P _{TEF1} -ERG20-GGGGS-LTC2 _{opt} aaaaaaaaaa P _{PGK1} -tHMG1	190.7 mg/L	(Hu et al. , 2017)
Valencene	<i>S. cerevisiae</i> YIP-00-03 (CEN.PK113-5D)	MATa MAL2-8c SUC2 ura3-52, <i>erg9</i> ::P _{MET3} -ERG9, pESC-URA 2 μ URA3 P _{GAL1} - GFTpSD	2 mg/L	(Asadollahi et al. , 2008)
	<i>S. cerevisiae</i> WAT11	Mutating <i>ERG9</i> gene, uptake of exogenous ergosterol (sue), introducing <i>CVS</i> gene	20 mg/L	(Takahashi et al. , 2007)

1290

1291

Table 3. Metabolic engineering of other organisms for sesquiterpene biosynthesis.

Sesquiterpene	Strain	Description	Titer	Reference
(+)-Valencene	<i>C. glutamicum</i> ATCC 13032	Deleting two endogenous prenyltransferase genes, expressing heterologous gene <i>ispA</i> from <i>E. coli</i> or gene <i>ERG20</i> from <i>S. cerevisiae</i>	2.41 mg/L	(Frohwitter et al. , 2014)
Amorphadiene	<i>B. subtilis</i> 1A1 (BGSC)	Expressing gene <i>ADS</i> under P_{grac} , gene <i>Dxs</i> and <i>Idi</i> under P_{xyIA} , protein translation engineering fused six-arginine tag to N terminus of gene <i>ADS</i> , optimizing media by adding pyruvate and dipotassium phosphate	~20 mg/L	(Zhou et al. , 2013)
β -Sesquiphellandrene	<i>L. lactis</i> NZ9000	Introducing β -sesquiphellandrene synthase, co-expressing <i>mvk</i> and <i>mvaA</i> gene	15.5 mg/L	(Song et al. , 2014)
Valencene	<i>R. sphaeroides</i>	Co-expressing MVA operon from <i>Paracoccus zeaxanthinifaciens</i> and <i>CVS</i> from <i>Callitropsis nootkatensis</i> on plasmid (pBBR-MEV) in shaker flasks	352 mg/L	(Beekwilder et al. , 2014)
Farnesene	<i>Anabaena</i> sp. PCC 7120	Expressing a <i>coFS</i> gene, through its endogenous MEP pathway using CO ₂ , mineralized water, and light	69.1 $\mu\text{g} \cdot \text{L}^{-1} \cdot \text{O}_2 \cdot \text{d}^{-1}$	(Halfmann et al. , 2014)
	<i>S. elongatus</i> PCC 7942	<i>NSI::Bb1s-dxs-idi-ispA</i> , <i>NSII::Bb1k-AFS</i> direct conversion of CO ₂ , with a dodecane overlay in 7 days	4.6 mg/L	(Lee et al. , 2017)
Amorphadiene	<i>S. elongatus</i> PCC 7942	<i>NSI::Bb1 s-dxs-idi-ispA</i> <i>NSII::Bb2 k-ADS</i> , direct conversion of CO ₂ , with a dodecane overlay in 7 days	19.8 mg/L	(Choi et al. , 2016)
Caryophyllene	<i>Synechocystis</i> sp. PCC6803	Inserting <i>QHSI</i> gene from <i>A. annua</i> via double homologous recombination into genome, converting CO ₂ and H ₂ O, measured by a methanol: chloroform extraction method	~ 4 mg/g DCW	(Reinsvold et al. , 2011)
Bisabolene	<i>Synechococcus</i> sp. PCC 7002	Expressing heterologous <i>A. grandis</i> <i>BIS</i> gene, deleting <i>glgC</i> gene and applying a dodecane overlay	>0.6 mg L ⁻¹	(Davies et al. , 2014)
	<i>C. reinhardtii</i>	Through genetic engineering, serial enzyme loading and artificial micro RNA-based repression of competing pathways	11.0 mg/L	(Wichmann et al. , 2018)
Farnesene	<i>Y. lipolytica</i> Polh	Integrating <i>tHMG1</i> , <i>Idi</i> , <i>ERG20</i> and <i>coFS</i> gene into genome in a bioreactor scale-up with PM medium	259.98 mg/L	(Yang et al. , 2016c)

Amorphadiene	<i>R. toruloides</i> IFO0880	Mating type A2, P _{GAPDH} -BIS-TNOS, in corn stover hydrolysates prepared by a biocompatible ionic liquid (IL) choline α -ketoglutarate pretreatment	36 mg/L	(Yaegashi et al. , 2017)
Bisabolene	<i>R. toruloides</i> IFO0880	Mating type A2, plasmid P _{GAPDH} -BIS-TNOS, with corn stover hydrolysates prepared by an alkaline pretreatment, in high-gravity fed-batch bioreactor	680 mg/L	(Yaegashi et al. , 2017)
Bisabolene	<i>S. venezuelae</i> ATCC 10712	Deleting gene <i>ΔSVEN_0269</i> and <i>ΔSVEN_7111</i> , inserting P _{sv} -coAgBis and P _{ermEp} -FPPS1 at the <i>aatP</i> chromosomal site	10.52 mg/L	(Phelan et al. , 2015)
α -cuprenene	<i>X. dendrorhous</i>	Integrating gene <i>Cop6</i> in the rDNA from <i>C. cinereus</i>	80 mg/L	(Melillo et al. , 2013)

1293

1294