

Submitted to the JOURNAL OF APPLIED PHYSICS

LBL-8076  
Preprint

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

OCT 26 1978

LIBRARY AND  
DOCUMENTS SECTION

THERMODYNAMIC PROPERTIES OF SeS GAS

Ming-Der Huang, David J. Meschi and Alan W. Searcy


September, 1978

Prepared for the U. S. Department of Energy  
under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy  
which may be borrowed for two weeks.

For a personal retention copy, call  
Tech. Info. Division, Ext. 6782



LBL-8076  
c.2

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THERMODYNAMIC PROPERTIES OF SeS GAS

Ming-Der Huang, David J. Meschi and Alan W. Searcy

Materials and Molecular Research Division, Lawrence Berkeley Laboratory  
and Department of Materials Science and Mineral Engineering,  
University of California, Berkeley, California

ABSTRACT

Mass-spectrometric Knudsen effusion measurements show the heat of formation of SeS to be 1.4 ( $\pm 1.7$ ) kcal per mole at 1000°K. SeS can be an important vapor species when selenium is an impurity in metal sulfides.

In the vacuum deposition of sulfide and selenide coatings, the presence of SeS gas, if its concentration is high enough, might affect the coating compositions. Thus thermodynamic properties of SeS are of practical interest in semiconductor technology.

The SeS gas molecule was observed by Umilin et al.<sup>1</sup> and by Chernozubov and Selivanov.<sup>2</sup> Ahmed and Barrow<sup>3</sup> have reported spectroscopic data for it. While the present work was in progress Drowart and Smoes published a mass-spectrometric study of the stability of SeS.<sup>4</sup>

We used a quadrupole mass-filter to study the gas-phase reaction,



These gases were generated by vaporizing a mixture of  $In_2S_3$  and  $In_2Se_3$  in a 99.5% alumina Knudsen cell. Orifice diameters of 1.03 and 0.52 mm were used. Measurements were made of the  $S_2^+$ ,  $Se_2^+$  and  $SeS^+$  ion currents at 64, 160 and 112 atomic mass units, respectively, at a series of temperatures between 1117° and 1195°K (Table I).

In mass-spectrometric measurements the pressure  $P_m$  of a given vapor species in a Knudsen cell can be determined from the corresponding total ion current  $I_m^+$  by means of the relation,

$$P_m \sigma_m g_m = I_m^+ T. \quad (2)$$

In this expression  $\sigma_m$  is the ionization cross-section for the formation of the ion in question,  $g_m$  is the machine constant and T is the absolute temperature. For a quadrupole mass-filter,  $g_m$  depends on the ion mass. In this study the coincidences of masses 64 for  $^{32}S^{32}S$  and  $^{64}Zn$  and of mass 112 for  $^{78}Se^{34}S$ ,  $^{80}Se^{32}S$ , and  $^{112}Cd$  were exploited to obtain relative values of  $g_m$  needed for calibration.

$CdS^5$  and  $ZnSe^6$  vaporize congruently in a Knudsen cell. Taking CdS as an example, this means that,

$$P_{\text{Cd}}/M_{\text{Cd}}^{1/2} = 2 P_{\text{S}_2}/M_{\text{S}_2}^{1/2} \quad (3)$$

where P is the pressure of the species indicated in the subscript, and M the molecular weight. Combining (2) and (3) gives

$$\frac{I_{64}^+}{I_{112}^+} = \frac{a_{\text{S}_2}^{(64)} g_{64} \sigma_{\text{S}_2} P_{\text{S}_2}}{a_{\text{Cd}}^{(112)} g_{112} \sigma_{\text{Cd}} P_{\text{Cd}}} = 1/2 \left( \frac{64}{112} \right)^{1/2} \frac{a_{\text{S}_2}^{(64)} g_{64} \sigma_{\text{S}_2}}{a_{\text{Cd}}^{(112)} g_{112} \sigma_{\text{Cd}}}$$

In this example the ion currents are those for the  $^{112}\text{Cd}^+$  and  $^{32}\text{S}^{32}\text{S}^+$  ions, with the corresponding abundances  $a_{\text{Cd}}^{(112)} = 0.24$  and  $a_{\text{S}_2}^{(64)} = 0.90$ . For these calibrations the potential of the ionizing electrons was 40 V, whereas the listed values of  $\sigma^7$  are given for  $V_{\text{max}}$ , the potential at which  $\sigma$  is a maximum. Values of  $\sigma$  at 40V. were calculated by assuming  $\sigma$  to be a linear function of the potential, equal to zero at the appearance potential and to the listed value at  $V_{\text{max}}$ . The calculated values were  $\sigma_{\text{Zn}} = 4.11$ ,  $\sigma_{\text{Se}_2} = 6.87$ ,  $\sigma_{\text{Cd}} = 5.13$  and  $\sigma_{\text{S}_2} = 4.78$ .

In this manner it was possible to determine the ratios,  $g_{64}/g_{160} = 1.05$  and  $g_{64}/g_{112} = 0.76$  which were then used to calculate the equilibrium constants for reaction (1) from the ion currents listed in Table 1:

$$K = \frac{P_{\text{SeS}}}{(P_{\text{S}_2} P_{\text{Se}_2})^{1/2}} = \alpha \frac{I_{112}^+}{(I_{64}^+ I_{160}^+)^{1/2}}$$

$$\text{where } \alpha = \frac{g_{64}}{g_{112}} \left( \frac{g_{160}}{g_{64}} \right)^{1/2} \frac{(\sigma_{\text{S}_2} \sigma_{\text{Se}_2})^{1/2}}{\sigma_{\text{SeS}}} \frac{[a_{\text{S}_2}^{(64)} a_{\text{Se}_2}^{(160)}]^{1/2}}{a_{\text{SeS}}^{(112)}} = 0.774$$

These measurements were made using an ionizing potential of 36 V, and the ionization cross-sections were calculated from the listed values<sup>7</sup> by the method discussed above. The calculated values were,  $\sigma_{\text{S}_2} = 4.12$ ,  $\sigma_{\text{Se}_2} = 5.98$ ,  $\sigma_{\text{SeS}} = 5.05$ .

Plots of  $\log K$  vs.  $1/T$  yielded from the slopes  $\Delta H^\circ = -2100$  ( $\pm 2600$ ) cal. for reaction (1) from measurements with the 1.03 mm orifice and  $\Delta H^\circ = 2500$  ( $\pm 3600$ ) cal. with the 0.52 mm orifice.

From the measurements of Ahmed and Barrow,<sup>3</sup> we calculated the entropy of SeS, and found  $\Delta S^\circ$  for reaction (1) to be 1.42 cal deg.<sup>-1</sup> at 1100°K and 1.41 cal deg.<sup>-1</sup> at 1200°K. Values of  $\Delta H^\circ$  for reaction (1) can thus be calculated from  $\Delta S^\circ$  and each measured  $K$ . For the 1.03 mm orifice, the mean  $\Delta H^\circ$  was 2080 cal per mole of SeS, and for the 0.52 mm orifice, 380 cal per mole. The average of all twenty values for both orifices is 1400 ( $\pm 1700$ ) cal. We prefer this value to those obtained from the slope because the latter are unduly sensitive to measurements at the temperature extremes. For reaction (1)  $\Delta C_p$  is essentially zero between room temperature and 1200°K, so  $\Delta H_{298}^\circ$  is also 1400 ( $\pm 1700$ ) cal. Drowart and Smoes<sup>4</sup> report a  $\Delta H^\circ$  value of 1900 ( $\pm 1600$ ) cal corresponding to the above, and a value of 1300 ( $\pm 700$ ) cal derived from the slope.

The importance of SeS as a vapor species can be illustrated for a solution (assumed ideal) of 1% CdSe dissolved in CdS at 1000°K. From known data for CdS<sup>5</sup> and CdSe<sup>6</sup> and  $K = 1.01$  for reaction (1) we find  $P_{S_2} = 1.9 \times 10^{-5}$  atm,  $P_{SeS} = 1.0 \times 10^{-5}$  atm, and  $P_{Se_2} = 5.6 \times 10^{-6}$  atm. Thus, SeS is predicted to be a major Se impurity carrier in vacuum deposition processes.

#### ACKNOWLEDGMENT

This work was supported by the Division of Materials Sciences, Office of Basic Energy Sciences, U.S. Department of Energy.

REFERENCES

1. V. A. Umilin, I. L. Agafonov, L. N. Kornev, and G. G. Devyatykh, Russian J. Inorg. Chem. 9, 1345 (1964).
2. Y. S. Chernozubov and G. K. Selivanov, Russian J. Phys. Chem. 44, 465 (1970).
3. F. Ahmed and R. Barrow, J. Phys. B., 7, 2256 (1974).
4. J. Drowart and S. Smoes, Trans. Faraday Soc. II 73, 1755 (1977).
5. Z. A. Munir, High Temp. Sci. 2, 58 (1970).
6. J. Berkowitz and W. A. Chupka, J. Chem. Phys. 45, 4289 (1966).
7. J. Mann, in "Recent Developments in Mass Spectrometry," ed. by K. Ogata and T. Hayakawa, p. 814 (1969).
8. S. Smoes, J. Drowart and J. M. Welter, J. Chem. Thermodynamics 9, 275 (1977).
9. S. Smoes, W. R. Pattje and J. Drowart, to be published.

Table I. Ion Intensities and Calculated Equilibrium Constants for Reaction (1).

T(°K)	$I_{64}^+(S_2^+)$	$I_{112}^+(SeS^+)$	$I_{160}^+(Se_2^+)$	$K = \alpha \frac{I_{112}^+}{(I_{64}^+ I_{160}^+)^{1/2}}$
d= 1.03mm				
1117	4.91	1.38	0.283	0.91
1137	9.30	2.47	0.435	0.95
1166	12.00	3.87	0.977	0.88
1138	24.3	12.27	5.55	0.82
1154	28.2	14.52	8.48	0.73
1165	27.6	18.15	10.08	0.84
1188	37.2	23.40	12.51	0.84
1130	9.68	4.80	2.25	0.80
1156	11.4	6.87	3.36	0.86
1195	21.6	9.09	3.56	0.80
1182	15.7	7.01	3.66	0.72
1119	4.48	2.34	1.23	0.77
d= 0.52mm				
1125	1.05	0.71	0.087	1.82
1125	1.25	0.63	0.090	1.45
1125	1.25	0.82	0.093	1.86
1146	1.39	0.81	0.108	1.62
1146	1.41	0.87	0.114	1.68
1164	1.8	1.41	0.186	1.89
1188	2.75	1.56	0.192	1.66
1188	2.50	1.70	0.195	1.88

d is the orifice diameter.



This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

TECHNICAL INFORMATION DEPARTMENT  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720