

# UC Irvine

## UC Irvine Previously Published Works

### Title

Chronic kidney disease

### Permalink

<https://escholarship.org/uc/item/3051x4gr>

### Journal

The Lancet, 398(10302)

### ISSN

0140-6736

### Authors

Kalantar-Zadeh, Kamyar  
Jafar, Tazeen H  
Nitsch, Dorothea  
[et al.](#)

### Publication Date

2021-08-01

### DOI

10.1016/s0140-6736(21)00519-5

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



## Chronic kidney disease

Kamyar Kalantar-Zadeh, Tazeen H Jafar, Dorothea Nitsch, Brendon L Neuen, Vlado Perkovic

Lancet 2021; 398: 786–802

Published Online

June 24, 2021

[https://doi.org/10.1016/S0140-6736\(21\)00519-5](https://doi.org/10.1016/S0140-6736(21)00519-5)

50140-6736(21)00519-5

Division of Nephrology,  
Hypertension and  
Kidney Transplantation,  
University of California Irvine,  
Orange, CA, USA

(Prof K Kalantar-Zadeh MD);  
Tibor Rubin Veterans Affairs  
Medical Center, Long Beach,  
CA, USA (Prof K Kalantar-Zadeh);

Duke-NUS Graduate Medical  
School, Singapore  
(Prof T H Jafar MD); Department  
of Renal Medicine, Singapore  
General Hospital, Singapore

(Prof T H Jafar); Duke Global  
Health Institute, Durham, NC,  
USA (Prof T H Jafar); Faculty of  
Epidemiology and Population

Health, London School of  
Hygiene & Tropical Medicine,  
London, UK (Prof D Nitsch MD);

United Kingdom Renal  
Registry, Bristol, UK  
(Prof D Nitsch); Department of  
Nephrology, Royal Free London  
NHS Foundation Trust,

London, UK (Prof D Nitsch);  
The George Institute for Global  
Health (B L Neuen MD), and

Faculty of Medicine  
(Prof V Perkovic MD), University  
of New South Wales Sydney,  
Sydney, NSW, Australia

Correspondence to:

Prof Kamyar Kalantar-Zadeh,  
Division of Nephrology,  
Hypertension and Kidney  
Transplantation, University of  
California Irvine,  
Orange 92868-3217, CA, USA  
[kkz@uci.edu](mailto:kkz@uci.edu)

See Online for appendix

Chronic kidney disease is a progressive disease with no cure and high morbidity and mortality that occurs commonly in the general adult population, especially in people with diabetes and hypertension. Preservation of kidney function can improve outcomes and can be achieved through non-pharmacological strategies (eg, dietary and lifestyle adjustments) and chronic kidney disease-targeted and kidney disease-specific pharmacological interventions. A plant-dominant, low-protein, and low-salt diet might help to mitigate glomerular hyperfiltration and preserve renal function for longer, possibly while also leading to favourable alterations in acid-base homeostasis and in the gut microbiome. Pharmacotherapies that alter intrarenal haemodynamics (eg, renin–angiotensin–aldosterone pathway modulators and SGLT2 [SLC5A2] inhibitors) can preserve kidney function by reducing intraglomerular pressure independently of blood pressure and glucose control, whereas other novel agents (eg, non-steroidal mineralocorticoid receptor antagonists) might protect the kidney through anti-inflammatory or antifibrotic mechanisms. Some glomerular and cystic kidney diseases might benefit from disease-specific therapies. Managing chronic kidney disease-associated cardiovascular risk, minimising the risk of infection, and preventing acute kidney injury are crucial interventions for these patients, given the high burden of complications, associated morbidity and mortality, and the role of non-conventional risk factors in chronic kidney disease. When renal replacement therapy becomes inevitable, an incremental transition to dialysis can be considered and has been proposed to possibly preserve residual kidney function longer. There are similarities and distinctions between kidney-preserving care and supportive care. Additional studies of dietary and pharmacological interventions and development of innovative strategies are necessary to ensure optimal kidney-preserving care and to achieve greater longevity and better health-related quality of life for these patients.

### Introduction

Chronic kidney disease is a progressive condition characterised by structural and functional changes to the kidney due to various causes. Chronic kidney disease is typically defined as a reduction in kidney function, an estimated glomerular filtration rate (eGFR) of less than 60 mL/min per 1.73 m<sup>2</sup>, or markers of kidney damage, such as albuminuria, haematuria, or abnormalities detected through laboratory testing or imaging and that are present for at least 3 months (appendix p 5).<sup>1</sup> The global burden of chronic kidney disease is substantial and growing: approximately 10% of adults worldwide are affected by some form of chronic kidney disease, which results in 1.2 million deaths and 28.0 million years of life lost each year.<sup>2,3</sup> By 2040, chronic kidney disease is estimated to become the fifth


leading cause of death globally—one of the largest projected increases of any major cause of death.<sup>4</sup>

The prevalence of different aetiologies of chronic kidney disease varies considerably by region. There are many causes of chronic kidney disease, including those that are common and well researched, such as diabetes, glomerulonephritis, and cystic kidney diseases, but causation in chronic kidney disease is not yet fully understood. For instance, despite a close association between chronic kidney disease and hypertension, whether hypertension is a cause or a consequence of chronic kidney disease is controversial.<sup>5</sup> As another example, chronic kidney disease of unknown aetiology and for which there is no known treatment is found in some agricultural communities in south Asia and central America; recurrent volume depletion has been speculated to be a cause, especially with the increasing frequency of climate change-related heat waves.<sup>6</sup> This possible cause of chronic kidney disease of unknown aetiology highlights the potential role of adequate hydration as a kidney-preserving strategy. The global burden of chronic kidney disease has also been attributed to air pollution, and is disproportionately borne by some world regions.<sup>7</sup> Chronic kidney disease severity also varies from kidney damage with normal function to kidney failure (or end-stage renal disease), which typically occurs when eGFR decreases to less than 15 mL/min per 1.73 m<sup>2</sup>. In general, the prevalence of chronic kidney disease increases with age and, in high-income countries, is more common in people with obesity, diabetes, and hypertension.<sup>8,9</sup>

Chronic kidney disease is usually insidious, and most affected individuals are asymptomatic until the

### Search strategy and selection criteria

We reviewed biomedical literature to identify all studies in which the effects of diet and lifestyle modifications and pharmacotherapeutic interventions were examined for conservative or preservative management of chronic kidney disease. Both the PubMed and Google Scholar databases were searched for studies published in English or with an English abstract between Jan 1, 1970, and Feb 1, 2021, using the terms “chronic kidney disease”, “kidney-preserving therapy”, “conservative management”, “diet”, “lifestyle modifications”, “pharmacological strategies”, and “renal hyperfiltration”. Full-text articles deemed pertinent were selected and the reference lists of the identified reports and articles were searched for further material.



**Figure 1: Conservative and preservative management of chronic kidney disease without dialysis or renal transplantation**

This chart highlights the role of preservative management and its goals (green domain) within the overall conservative management of chronic kidney disease without dialysis (blue zone), juxtaposing renal replacement therapy including dialysis and kidney transplantation (yellow zone). The X axis (showing chronic kidney disease progression) should be read exclusively from left to right. The bottom half of the chart represents conventional (life-prolonging and kidney-prolonging) strategies, whereas the top half represents supportive care, including palliative and hospice care, in which dialysis is often avoided or withdrawn (violet domain).<sup>17</sup> The oblique dotted line between the two main zones (conservative management vs renal replacement therapy) suggests that there is variability in transitioning to dialysis therapy (moving from bottom left to top right), including timing (early vs late vs never), level of care (life-prolonging vs supportive care), and type of dialysis (conventional vs incremental). The symptom management (purple) domain provides wide ranges of interventions to encompass the goals of care under both kidney preserving care and palliative and hospice care. Preservative management can preserve residual kidney function for longer, especially after incremental transition to dialysis. See appendix p 6 for an overview of these strategies over the course of chronic kidney disease progression. BP= blood pressure. GFR=glomerular filtration rate. MR=mineralocorticoid receptor. RAAS=renin-angiotensin-aldosterone system.

disease becomes advanced (ie, eGFR of less than 30 mL/min per 1.73 m<sup>2</sup>). The rate of loss of kidney function varies by aetiology, exposures, and interventions but, in most cases, progression to kidney failure typically takes between months and decades to develop. Signs and symptoms of kidney failure result from progressive uraemia, anaemia, volume overload, electrolyte abnormalities, mineral and bone disorders, and acidaemia, and inevitably lead to death if left untreated.<sup>10</sup>

Renal replacement therapy, either in the form of chronic dialysis or kidney transplantation, is a life-sustaining treatment for people with kidney failure. Because of the shortage of kidney donors and of the comorbidities that develop with age and often preclude kidney transplantation,<sup>11,12</sup> dialysis remains the prevailing treatment option for most people with kidney failure.<sup>13</sup> Kidney failure requiring dialysis is often associated with substantially reduced quality of life and high mortality rates,<sup>14</sup> especially in the first year after transition to dialysis,<sup>15</sup> underscoring the importance of preserving kidney function in people with, or at high risk of, chronic kidney disease.

### Approaches to preserving kidney function

There has been growing recognition that conservative management without dialysis is a viable, patient-centred treatment option for a substantial proportion of patients with chronic kidney disease.<sup>16</sup> Within conservative management strategies, there are several overlapping intervention domains, with similarities and differences, that can be offered to patients with chronic kidney disease (figure 1).<sup>17</sup> Kidney-preserving care is a life-sustaining conservative management therapy with the primary goal of slowing chronic kidney disease progression and preserving kidney function to avoid dialysis for as long as possible or, ideally, altogether. This approach strives to achieve the greatest possible survival, improved cardiovascular health, and superior health-related quality of life through effective treatment of renal and non-renal comorbidities and their associated symptoms.<sup>18,19</sup>

Given that conservative management is defined as chronic kidney disease care without dialysis or kidney transplantation,<sup>20</sup> misconceptions of dialysis-free management as so-called no care, or misguided conflation with hospice care might have contributed to an underuse of the full spectrum of kidney-preserving management.<sup>12,18</sup>

	Advantages and rationale	Disadvantages	Additional comments
<b>Diet and lifestyle</b>			
Plant-dominant, low-protein diets*	Patient-centred, inexpensive, improves metabolic markers, and mitigates acidosis; might slow disease progression and attenuate uraemia	Risk of muscle loss and frailty; risk of diet-induced hyperkalaemia	Effects on patient-level kidney outcomes not yet confirmed in randomised controlled trials
Nutrient-focused dietary interventions (low sodium, low phosphate, and low potassium)*†	Experience in clinical practice	Uncertain effectiveness of single-nutrient approaches (eg, strict phosphate control) on patient-level outcomes	Potassium-restricted diets might cause more harm than no intervention by limiting intake of potassium-rich, healthy fruits and vegetables
Increasing physical activity, weight reduction, and smoking cessation*†	Numerous clear health benefits	Sustained goals can be challenging to achieve	See table 2 for more details
<b>Pharmacological agents to slow chronic kidney disease progression</b>			
Renin-angiotensin-aldosterone system blockade (ACEi, ARB)‡	Proven benefit in preventing kidney failure in randomised trials in people with diabetes	Risk of AKI and of hyperkalaemia	Uncertain benefits for non-diabetic kidney disease with proteinuria <0.5 g/day
SGLT2 inhibitors‡	Clear reductions in patient-level adverse cardiovascular and renal outcomes in people with type 2 diabetes	Risk of mycotic genital infections, volume depletion, and euglycaemic ketoacidosis; reported increased risk of limb amputation not substantiated	Less data for initiation at eGFR <30 mL/min per 1.73 m <sup>2</sup> ; emerging evidence of benefit in people with non-diabetic kidney disease
Non-steroidal mineralocorticoid receptor antagonists†‡	Reduction in adverse cardiovascular and renal outcomes in people with type 2 diabetes; potential anti-inflammatory and antifibrotic effects	Risk of hyperkalaemia	Uncertain effects of steroidal mineralocorticoid receptor antagonists (eg, spironolactone) on patient-level kidney outcomes; non-steroidal mineralocorticoid receptor antagonists have not yet been evaluated in non-diabetic kidney disease
Tolvaptan for polycystic kidney disease*†	Slowed decline in glomerular filtration rate	Risk of polydipsia or polyuria and deranged liver function	Tolvaptan might delay the predicted onset of kidney failure
Rituximab for primary membranous nephropathy*†	Increased likelihood of long-term remission compared with ciclosporin	Little data from randomised controlled trials directly comparing with alkylating agents	Effect of combination therapies to be determined
Steroids for IgA nephropathy*†	Experience in clinical practice	Mixed results in clinical trials; increased risk of adverse events, especially serious infection	Additional trial ongoing (NCT01560052)
<b>Pharmacological strategies to reduce cardiovascular risk</b>			
Lipid-lowering treatments‡	Reduction in adverse cardiovascular events in people with chronic kidney disease; well tolerated	No clear benefit for initiating treatment in people requiring dialysis	Few data on new lipid-lowering therapies in chronic kidney disease
Blood pressure-lowering treatments‡	Reduction in adverse cardiovascular and potentially renal outcomes	Possibly greater risk of adverse events as kidney function declines	Addressing volume overload is a crucial aspect of blood pressure-lowering in advanced chronic kidney disease
Glucose-lowering drugs‡	SGLT2 inhibitors and GLP-1 receptor agonists reduce adverse cardiovascular events in people with type 2 diabetes	Risk of hypoglycaemia and other treatment-related adverse events with intensive glucose reduction	Inconsistent benefits for cardiovascular and kidney outcomes with intensive glucose control; cardiovascular and renal benefits vary with class of glucose-lowering drug
<b>Pharmacological and other strategies to slow progression and manage uraemia and associated symptoms</b>			
Sodium bicarbonate and veverimer for acidosis management*†	Sodium bicarbonate improves acidosis, might slow progression of chronic kidney disease	Effect on long-term outcomes uncertain; sodium bicarbonate might worsen sodium and fluid retention	Veverimer might improve acidosis without causing sodium retention; randomised trial ongoing (NCT03710291)
Potassium binders (sodium polystyrene, zirconium, and patiromer)*†	Reduced risk of hyperkalaemia; enables use of ACEi and ARB	No data on patient-level outcomes or progression of kidney disease	Randomised trial ongoing (NCT03888066)
Sodium and volume management (sodium restriction, loop diuretics, and thiazide diuretics)*†	Experience in clinical practice	Uncertain effect on chronic kidney disease progression	Few data from randomised controlled trials
Symptom management (eg, for pruritus, pain, fatigue, and sleep disorders)*†	Important priority for patients with unpleasant symptoms	Unlikely to influence need for renal replacement therapy or to affect risk of chronic kidney disease progression	Few data from randomised controlled trials
Prevention of AKI*	AKI might increase future risk of chronic kidney disease	Effect of sick day advice (ie, temporary discontinuation of ACEi or ARBs) on incidence of AKI not yet evaluated in randomised trials	Some drug combinations (eg, ACEi and ARB) increase risk of AKI
Prevention of infection (eg, hepatitis C and COVID-19)*	Many infectious events might cause AKI, chronic kidney disease, or both	Direct kidney involvement unknown	Effect of infection prevention strategies not clear
Footnotes indicate the strength of the evidence for intervention efficacy (two footnotes signify that the evidence ranges between the two levels). For the potential roles of these measures in secondary and tertiary prevention of chronic kidney disease see appendix p 7. ACEi=ACE inhibitors. AKI=acute kidney injury. ARB=angiotensin receptor blockers. eGFR=estimated glomerular filtration rate. *Supported by biological or observational data but little or no evidence from randomised controlled trials. †Some evidence from small randomised trials or trials evaluating effects on surrogate outcomes. ‡Supported by data from large randomised trials assessing effects on patient-level outcomes.			
<b>Table 1: Intervention strategies to preserve kidney function in people with chronic kidney disease</b>			

	Effect on chronic kidney disease progression	Effect on cardiovascular disease and mortality	Comments	Recommendations
Physical activity	Slower decline in kidney function	Lower risk of adverse cardiovascular outcomes and mortality	Evidence on physical activity and progression of kidney disease and cardiovascular outcomes is largely based on observational studies; small trials of physical activity show improvements in kidney function and blood pressure in people with chronic kidney disease not receiving dialysis; small trials in patients receiving dialysis show improvements in physical function and health-related quality of life	Target of 150 min/week of moderate intensity physical activity for patients with chronic kidney disease; exercise should be individualised for patients according to comorbidities and functional status (mixed data in dialysis-dependent patients)
Smoking cessation or avoidance	Smoking is associated with a greater risk of incident chronic kidney disease	Smoking is associated with increased risk of all-cause mortality, including vascular causes and cancer, in people with chronic kidney disease	Smoking cessation should be prioritised in all individuals for numerous recognised health benefits	Smoking cessation for all patients with behavioural counselling and pharmacological therapies as required, with appropriate dose adjustment for patients with chronic kidney disease
Dietary sodium restriction	Reduced albuminuria and improved fluid status in people with and without chronic kidney disease	Reduces blood pressure and improves arterial stiffness in people with and without chronic kidney disease	People with chronic kidney disease are more likely to have salt-sensitive hypertension	Limit sodium intake to a maximum of 2.3 g/day (<100 mmol) according to the American Heart Association
Higher proportion of plant-based protein in diet	Higher proportion of plant-based protein and fibre intake might improve acidosis, mitigate inflammation, reduce phosphorus burden, slow progression of chronic kidney disease, and create less uraemic toxins	Higher red meat intake might be associated with atherosclerosis due to higher carnitine generation via gut microbiota	Reducing dietary protein intake can increase the risk of muscle mass loss and frailty; protein intake recommendations vary depending on chronic kidney disease stage, acute kidney injury, and need for dialysis	Higher intake of complex carbohydrates and fresh fruits and vegetables as opposed to processed carbohydrates
Weight reduction	Improved cardiometabolic health; potentially slower decline in kidney function and improved albuminuria	Improves blood pressure	Little evidence from randomised controlled trials to guide the dietetic management of people with overweight and obesity and chronic kidney disease	Multidisciplinary approach to weight loss in overweight and obese individuals with chronic kidney disease with involvement of a renal dietitian; mixed data in dialysis patients related to the obesity paradox

See also appendix p 6 for schematic depictions of these strategies over the course of chronic kidney disease progression.

**Table 2: Lifestyle modification strategies to slow the progression of chronic kidney disease and preventing adverse cardiovascular outcomes**

Notwithstanding heterogeneity in definitions, provision of (or access to) care, and patient demographics or socioeconomic status across different domains of the conservative management of chronic kidney disease, the use of conservative management is rising, with a greater focus on kidney-preserving strategies.<sup>21–33</sup>

The focus of efforts to slow the loss of kidney function varies depending on the severity of chronic kidney disease and underlying causes, encompassing a range of pharmacological and non-pharmacological approaches (figure 1; appendix pp 6–7), given that these measures are consistent with the secondary and tertiary prevention of chronic kidney disease<sup>34</sup> (table 1). In patients with chronic kidney disease (as in the general population), lifestyle and dietary modifications (table 2) should be prioritised because these can improve cardiometabolic health and are likely to have favourable long-term effects on the kidney. The focus of care in primary prevention is to achieve optimal control of risk factors for chronic kidney disease by addressing physical inactivity and obesity, smoking, high blood pressure, and high blood glucose.<sup>35</sup> Addressing these risk factors is important across the spectrum of kidney function. Although the cost-effectiveness of population-wide screening for chronic kidney disease is

controversial, targeted screening of individuals with risk factors (eg, obesity, hypertension, and diabetes) through regular assessments of eGFR and albuminuria is recommended.<sup>36</sup>

For individuals with established chronic kidney disease, addressing complications and associated comorbidities and managing symptoms in addition to protecting kidney function are important steps. Slowing the progression of chronic kidney disease can be achieved through a range of lifestyle, dietary, and pharmacological strategies, which include weight loss, moderate dietary protein restriction, blood pressure and glucose control, and renin-angiotensin-aldosterone system blockade (appendix pp 6–7). For specific aetiologies, such as primary glomerulonephritis and autosomal dominant polycystic kidney disease, newer targeted therapies also have an important role. Because cardiovascular disease is a more common cause of death than kidney failure in patients with chronic kidney disease, reducing cardiovascular risk is a fundamental aspect of care for this population.<sup>37</sup>

Large-scale, collaborative meta-analyses have shown that eGFR and albuminuria are strongly and independently associated with risk of a range of adverse outcomes, including progression to kidney failure,

cardiovascular events, and death, and both kidney markers should be used to inform prognosis and direct care priorities for people with chronic kidney disease.<sup>38–42</sup> The Kidney Disease: Improving Global Outcomes (KDIGO) classification of chronic kidney disease incorporates eGFR and the urine albumin-to-creatinine ratio into a two-dimensional framework to stratify individuals' risk, focus management priorities, and guide referral to specialist care, and is perhaps the most widely used staging system for chronic kidney disease (appendix p 5).<sup>43</sup> Other tools, such as the Kidney Failure Risk Equation<sup>44</sup> to estimate the risk of kidney failure, and the Dialysis Transition Mortality Prediction Score,<sup>45</sup> to estimate mortality in the first year of dialysis, can also be used to inform discussion, facilitate specialist referral, and contribute to shared decision making. After progression to advanced chronic kidney disease, when uraemia cannot be controlled without renal replacement therapy, incremental transition to peritoneal or haemodialysis therapy might be a preferred approach with the goal of preserving residual kidney function while reducing the frequency of dialysis, although clinical trials are needed to examine this and other alternative dialysis transition strategies.<sup>46</sup>

### Physical activity, obesity, and weight loss

Obesity is the hallmark of metabolic syndrome and associated with chronic kidney disease.<sup>47</sup> Evidence suggests that increased adiposity measures (eg, body-mass index and waist circumference) are independently associated with a decline in glomerular filtration rate.<sup>48</sup> This association might result from systemic and intraglomerular hypertension, the effect of prediabetes concentrations of blood glucose on podocyte stress, and other unrecognised factors.<sup>49,50</sup> Physical activity is the core component of lifestyle modification strategies to manage weight for a positive effect on chronic kidney disease progression.

The Look AHEAD trial,<sup>51</sup> in which 5145 people with obesity and type 2 diabetes were randomly assigned to intensive lifestyle intervention or diabetes support and education, showed that intensive lifestyle intervention reduced weight by an average of 4 kg and resulted in a relative risk reduction of onset of very high-risk chronic kidney disease (according to the KDIGO classification system) of approximately 30% compared with control ( $p=0.0016$ ).

A range of weight loss interventions can be recommended to people with chronic kidney disease. Caloric restriction, in conjunction with a plant-dominant, low-protein diet, can lead to gradual weight loss in most people with obesity and chronic kidney disease.<sup>52,53</sup> Efforts to identify pharmacological agents that can reduce bodyweight and improve clinical outcomes have yielded few or modest results.<sup>54</sup> The role of bariatric surgery in mitigating the risk of chronic kidney disease is also uncertain.<sup>55</sup> Observational studies have suggested that

bariatric surgery is associated with a lower risk of patient-level kidney outcomes.<sup>56</sup> Gastric bypass surgery increases remission of albuminuria in people with type 2 diabetes, obesity, and microalbuminuria when compared with optimal medical treatment, and might represent an important treatment option for selected individuals.<sup>57</sup>

In more advanced chronic kidney disease, prolonged survival has been paradoxically reported with larger body-mass index, a phenomenon that is known as the obesity paradox or reverse epidemiology.<sup>58</sup> Conversely, weight loss can contribute to poorer outcomes, whereas effective nutritional interventions to gain weight including muscle mass might improve longevity.<sup>59</sup> Any unintentional weight loss warrants prompt investigation and dietary interventions, and unnecessary weight loss in advanced chronic kidney disease should be avoided, unless absolutely required (eg, as a strict requirement for imminent kidney transplantation or other life-saving procedures that require a lower weight).<sup>60</sup>

### Plant-dominant, low-protein diet


In many causes of chronic kidney disease, afferent arterioles are relatively dilated and efferent arterioles are relatively contracted as a compensatory mechanism to maintain glomerular filtration rate in the short term—a process known as glomerular hyperfiltration or intraglomerular hypertension. This interaction is partly regulated via tubuloglomerular feedback.<sup>49</sup> In the long term, glomerular hyperfiltration can cause further damage to the kidney through mechanisms such as mechanical stress and activation of inflammatory mediators that promote interstitial fibrosis.<sup>61,62</sup> Dietary protein restriction, by enhancing the afferent arteriole tone, might alleviate intraglomerular hypertension, mitigate renal interstitial fibrosis,<sup>61,62</sup> and slow the progression of chronic kidney disease (figure 2). This effect acts in parallel and is complementary to the postglomerular effect of renin–angiotensin–aldosterone pathway modulators, such as angiotensin-converting enzyme (ACE) inhibitors and angiotensin receptor blockers, which reduce intraglomerular pressure by promoting efferent arteriolar vasodilatation.<sup>63</sup>

Evidence from randomised controlled trials supporting the beneficial effect of dietary protein restriction comes from the Modification of Diet in Renal Disease study, which randomly assigned 585 participants with non-diabetic kidney disease to assess the effect of typical (1.3 g/kg per day) versus low-protein diets (0.58 g/kg per day) on eGFR decline. Although the primary results of this trial were inconclusive, they did not take into account the acute effect of dietary protein restriction, which reduces short-term glomerular filtration rate through afferent arteriolar constriction, similarly to what is observed with initiation of ACE inhibitors or angiotensin receptor (AGTR1) blockade therapy. Subsequent analyses of the Modification of Diet in Renal Disease study data excluding the acute effect of the dietary intervention on glomerular

filtration rate suggested that there was a benefit of dietary protein restriction (appendix p 2).<sup>64</sup> Additional analyses also showed that dietary protein restriction might reduce blood pressure and proteinuria.<sup>65</sup> The findings from this study are supported by meta-analyses that show a reduced risk of progression to kidney failure and improvements in proteinuria and other favourable biochemical outcomes, such as higher serum concentrations of bicarbonate, lower azotaemia, and lower serum concentrations of phosphorus.<sup>66–69</sup> The benefits of dietary protein restriction need to be considered in the context of potential risks to protein-energy wasting and loss of muscle mass and strength, particularly in frailer people or older than 80 years.<sup>70</sup> Therefore, current guidelines recommend a conservatively low range of 0.6–0.8 g/kg per day of dietary protein in people with substantial albuminuria (more than 300 mg/g) to ensure safety and adequate nutritional intake (appendix p 2).<sup>71</sup>

More recent data suggest salutary effects of plant-dominant, low-protein diets,<sup>52</sup> in which more than 50% of ingested protein is derived from non-animal sources (ie, fruits, vegetables, nuts, legumes, and seeds). There are different types of plant-dominant diets, listed here with increasing amounts of foods from plant sources:<sup>72</sup> vegan or strict vegetarian diets that not only exclude meat, poultry, and seafood but also eggs and dairy products; lacto-vegetarian, ovo-vegetarian, or lacto-ovo-vegetarian diets that can include dairy products and eggs; and pescatarian or pesco-vegetarian diets that include a vegetarian diet combined with occasional intake of some or all types of seafoods, mostly fish.<sup>72</sup> Although some, but not all studies have shown that plant-dominant diets are associated with a lower risk of chronic kidney disease and glomerular filtration rate decline, less proteinuria, amelioration of acidosis, and better cardiovascular profile,<sup>73</sup> and although experimental data also suggest that such diets can reduce uraemic toxin generation and exert favourable effects on cardiovascular health in people with kidney failure,<sup>32,53,74,75</sup> these effects have not yet been definitively shown in randomised controlled trials.<sup>76</sup>

There is growing interest in the role of the gut microbiome in chronic kidney disease, although the role of any microbiome-related interventions is not yet proven.<sup>76</sup> The gut microbiome in chronic kidney disease might be altered by uraemia, natural intake of probiotics,<sup>77</sup> and the type of diet (including plant-origin vs animal-origin foods).<sup>78</sup> A plant-dominant, fibre-rich, low-protein diet can lead to favourable alterations in the gut microbiome, which might modulate uraemic toxin generation.<sup>52</sup> Several gut-derived uraemic toxins, including indoxyl sulfate, indole-3 acetic acid, p-cresyl sulfate, trimethylamine N-oxide, and phenylacetylglutamine, are associated with cardiovascular disease and mortality in chronic kidney disease.<sup>79</sup> Circulating p-cresyl sulfate and indoxyl sulfate (protein-bound uraemic retention solutes) and other catabolic by-products of protein metabolism can exert harmful effects on several organs and homeostatic pathways, such



**Figure 2: Effects of dietary protein and sodium intake and pharmacological therapies on afferent and efferent arteriolar tone, intraglomerular pressure, and glomerular structures and functions**

Dietary protein restriction results in contraction of the afferent arterioles leading to reduced intraglomerular pressure, reducing damage to glomerular structure and function in the long term. The kidney-protective effects of a plant-dominant, low protein diet act in parallel and might be complementary to the effect of SGLT2 inhibitors, RAAS blockade, mineralocorticoid receptor antagonism, and other blood pressure-lowering agents, thereby more effectively reducing intraglomerular pressure and mitigating interstitial fibrosis. GFR=glomerular filtration rate. M=mesangial cells. RAAS=renin-angiotensin-aldosterone system.

as inflammation, oxidative stress, endothelial dysfunction, muscle wasting, renal interstitial fibrosis, worsening proteinuria, accelerated chronic kidney disease progression, and insulin resistance.<sup>80–82</sup> Therefore, a high-fibre, plant-dominant, low-protein diet has been proposed, but not yet proven in clinical trials, to favourably modulate microbiome, reduce uraemic toxin generation, and help to control uraemia without dialysis, potentially while enhancing cardiovascular health, which is consistent with the goals of the conservative and preservative management of chronic kidney disease.<sup>83</sup>

### Intravascular volume and acid-base and electrolyte homeostasis

Subclinical volume overload is highly prevalent in people with chronic kidney disease and perturbations in systemic haemodynamics are strongly associated with risk of poor cardiovascular and kidney outcomes.<sup>84–86</sup> Optimisation of intravascular volume is, therefore, an important focus of care, particularly as kidney function declines, and can be achieved through dietary sodium restriction and diuretics. Many patients with chronic kidney disease exhibit a tendency for salt-sensitive hypertension.<sup>87</sup> Although loop diuretics are the mainstay pharmacological therapy for the management of excess fluid, emerging data from randomised controlled trials suggest that distal thiazide

diuretics can reduce blood pressure and extracellular volume even at lower eGFRs,<sup>88</sup> with additional trials ongoing.<sup>89</sup>

The prevalence of hyperkalaemia increases as kidney function declines, and epidemiological data show a U-shaped association between serum potassium and adverse outcomes.<sup>90</sup> Commonly used treatments such as ACE inhibitors, angiotensin receptor blockers, and mineralocorticoid receptor antagonists increase the risk of hyperkalaemia. Although dietary potassium restriction is traditionally recommended for people with advanced chronic kidney disease, there are concerns that such diets might limit the consumption of healthy plant proteins, fruit, and vegetables.<sup>91</sup> Traditional and newer potassium binders might allow more effective control of hyperkalaemia, and trials are underway to test the effect of newer potassium binders on clinical outcomes.<sup>92,93</sup>

Metabolic acidosis in chronic kidney disease results from the inability of the kidney to excrete endogenous acid, and has been shown to be associated with loss of kidney function and unfavourable effects on muscle mass and bone health.<sup>94</sup> Small trials have collectively suggested that correction of metabolic acidosis with bicarbonate might slow progression of chronic kidney disease.<sup>95</sup> In 2019, veverimer (a novel binder of hydrochloric acid in the gastrointestinal tract) was shown to increase serum bicarbonate concentrations in people with chronic kidney disease, with an ongoing trial to test the effect of this agent on clinical outcomes, including progression of kidney disease or kidney failure requiring dialysis.<sup>96</sup>

### Traditional and emerging pharmacotherapies Renin–angiotensin–aldosterone system inhibition, mineralocorticoid receptor antagonism, and other blood pressure-lowering therapies

Renin–angiotensin–aldosterone pathway modulators, specifically ACE inhibitors or angiotensin receptor blockers, have been the cornerstone of kidney-preserving pharmacological therapies for several decades. The evidence is strongest in people with type 2 diabetes and albuminuric chronic kidney disease, for whom the angiotensin receptor blockers losartan (RENAAL trial) and irbesartan (IDNT trial) have been shown to reduce the risk of kidney failure, doubling of creatinine, or death (appendix p 8).<sup>97,98</sup> Importantly, several studies<sup>99–101</sup> have shown an increased risk of adverse outcomes and no clear benefits with multi-agent renin–angiotensin–aldosterone system blockade, for which reason angiotensin-converting enzyme inhibitors and angiotensin receptor blockade combination therapy is strongly discouraged.

The FIDELIO-DKD trial of the non-steroidal mineralocorticoid receptor antagonist finerenone in diabetic kidney disease showed a significant reduction in the risk of the primary kidney outcome (sustained 40% reduction in eGFR, kidney failure, or kidney death) and cardiovascular outcome (myocardial infarction,

stroke, heart failure, or cardiovascular death).<sup>102</sup> Several studies have suggested that the older steroidal mineralocorticoid receptor antagonist spironolactone reduces proteinuria in diabetic kidney disease,<sup>103</sup> but clear data regarding its effects on hard outcomes such as kidney disease progression or need to start dialysis are not currently available. Renin–angiotensin–aldosterone pathway modulation with angiotensin-converting enzyme inhibitors or angiotensin receptor blockade therapy is also generally recommended for people with type 1 diabetes and kidney disease<sup>98</sup> and for those without diabetes but with significant proteinuria;<sup>104</sup> the effects in people with little or no proteinuria are uncertain.

Although blood pressure reduction per se probably reduces the risk of kidney failure,<sup>105,106</sup> the haemodynamic effect of intensive blood pressure lowering might paradoxically induce a more rapid decline in kidney function.<sup>107–110</sup> Notwithstanding such mixed data, the KDIGO guidelines recommend a systolic blood pressure target of less than 120 mm Hg in people with chronic kidney disease not on dialysis to reduce the risk of cardiovascular events and mortality. However, individualisation of blood pressure targets is crucial and should take into account comorbidities, frailty, and patient preferences, given the ongoing uncertainty about the risk–benefit profile of intensive blood pressure targets in patients with moderate-to-advanced chronic kidney disease.<sup>107</sup>

### Glucose-lowering therapies

Glucose-lowering therapies could potentially reverse the fundamental metabolic abnormality pathognomonic of diabetes.<sup>111</sup> Although post-hoc analyses of the ADVANCE trial<sup>112</sup> suggested that the risk of kidney failure might be smaller in people treated to lower targets of haemoglobin A<sub>1c</sub>; meta-analyses of more intensive versus less intensive glucose reduction did not show clear effects on the risk of kidney failure.<sup>113</sup> Subsequent studies have further shown clear differences between different classes of glucose-lowering therapies.

#### SGLT2 inhibitors

SGLT2 (SLC5A2) inhibitors were developed to reduce blood glucose in people with diabetes by blocking proximal tubular glucose reabsorption, thus inducing glycosuria (figure 2). Early studies in type 2 diabetes identified that these drugs significantly reduce proteinuria and have a clear beneficial effect on kidney haemodynamics. This is manifest clinically as an acute reduction in eGFR (approximately 3–5 mL/min per 1.73 m<sup>2</sup>) followed by stabilisation of kidney function compared with either placebo or sulfonylurea therapy, benefits that are observed on top of renin–angiotensin–aldosterone system blockade. (appendix p 8).<sup>114</sup> Several cardiovascular safety studies mandated by regulatory authorities have successively shown reductions in composite outcomes based on reductions in eGFR or



doubling of creatinine, kidney failure, and death due to kidney disease.<sup>115–117</sup> However, these trials were done in people at high risk of cardiovascular disease, less than 25% of whom had kidney disease.

The first primary kidney trial of SGLT2 inhibitors assessing efficacy on major clinical outcomes was the CREDENCE study.<sup>118</sup> A total of 4401 participants with albuminuric diabetic kidney disease (albuminuria 300–5000 mg/g, eGFR 30–90 mL/min per 1.73 m<sup>2</sup>) were randomly assigned to receive canagliflozin or placebo (appendix p 8). The trial was stopped early for efficacy after the primary outcome (doubling of serum creatinine, kidney failure, or death due to cardiovascular or kidney disease) was reduced by 30%, with similar reductions in a range of renal outcomes, including kidney failure and the need for dialysis or kidney transplantation.<sup>118</sup> Major cardiovascular events (myocardial infarction, stroke, or cardiovascular death) and hospitalisations for heart failure were also significantly reduced. On the basis of these findings, major treatment guidelines worldwide have now been updated to recommend SGLT2 inhibitors for people with diabetic kidney disease.

The kidney-protective effects of SGLT2 inhibition have also been observed in people with non-diabetic kidney disease. The DAPA-CKD trial<sup>119</sup> showed that dapagliflozin reduces the risk of sustained 50% decline in eGFR, kidney failure, or death due to cardiovascular or kidney disease by 44% in people with chronic kidney disease (eGFR 25–75 mL/min per 1.73 m<sup>2</sup>, urine albumin-to-creatinine ratio 200–5000 mg/g), with clear and independent benefits irrespective of diabetes status or aetiology of chronic kidney disease.<sup>120</sup> The trial also showed that dapagliflozin substantially reduced the risk of hospitalisation for heart failure or cardiovascular death and all-cause mortality, irrespective of diabetes status. On the basis of these findings, SGLT2 inhibitors are anticipated to be routinely offered to people with albuminuric chronic kidney disease, regardless of the presence of diabetes. A trial of empagliflozin in non-diabetic kidney disease that includes people with low or normal albuminuria is ongoing.<sup>121</sup> The mechanisms underpinning the cardiovascular and kidney benefits of SGLT2 inhibition are an area of active investigation, but are probably multifactorial, including reductions in intraglomerular pressure, favourable effects on the extracellular fluid compartment, and multiple direct effects on cellular and metabolic functions.<sup>122</sup>

#### DPP4 inhibitors

DPP4 inhibitors are widely used to improve glycaemic control in people with type 2 diabetes. The effects on hard kidney outcomes, such as progression of kidney disease and end-stage renal disease, were formally assessed in the CARMELINA trial,<sup>123</sup> in which almost 7000 participants with type 2 diabetes enriched for either or both cardiovascular disease and kidney disease (reduced glomerular filtration rate, increased albuminuria, or both) were

randomly assigned to linagliptin or placebo. The trial showed that linagliptin did not increase the risk of cardiovascular events, but also did not reduce the risk of a composite renal outcome of 40% eGFR decline, kidney failure, or renal death, despite over 600 kidney-related endpoints being observed in the trial. Therefore, the available data suggest that DPP4 inhibitors do not meaningfully reduce the risk of kidney disease progression in patients with type 2 diabetes.


#### GLP-1 receptor agonists

There have not yet been any outcome studies sufficiently powered to detect effects on clinically important kidney outcomes with the use of GLP-1 receptor agonists, another class of drugs developed in the past 10 years to improve glucose control in type 2 diabetes. GLP-1 receptor agonists have been shown to reduce the risk of cardiovascular events in meta-analyses of completed cardiovascular outcome trials.<sup>124</sup> Similar meta-analyses of these trials suggest that the risk of a composite kidney outcome (including progression of albuminuria, substantial losses of renal function [40% or 57% reductions in eGFR], renal failure, or kidney-related death) is significantly reduced by GLP-1 receptor agonists. However, this reduction appears to be primarily driven by effects on albuminuria; no other clear benefit was observed on kidney outcomes.<sup>124</sup> A dedicated kidney outcome study (FLOW, NCT03819153) is currently underway, specifically recruiting people with chronic kidney disease and comparing the effects of semaglutide versus placebo on the risk of major kidney and cardiovascular outcomes.

#### Examples of treatment of primary glomerulonephritides and cystic disorders

IgA nephropathy, the most common idiopathic glomerulonephritis worldwide, is currently treated by optimising blood pressure control with ACE inhibitors or angiotensin receptor blockers, along with lifestyle modifications, such as salt and protein restriction and weight loss.<sup>125</sup> The role of corticosteroids in managing IgA nephropathy is controversial because of the conflicting evidence from randomised trials that yielded both negative and positive data.<sup>126,127</sup> New therapeutic strategies for IgA nephropathy are an area of active investigation, and several agents are currently being tested, including combined angiotensin and endothelin receptor blockade and drugs targeting complement pathways.<sup>128,129</sup>

Primary membranous nephropathy is another globally prevalent glomerulonephritis. Although the discovery that it is an autoimmune condition has led to substantial changes in the diagnosis, treatment, and monitoring of this disease,<sup>130–132</sup> many of these patients will develop spontaneous remission; as with other causes of proteinuric chronic kidney disease, optimal supportive care should include the maximum tolerated ACE inhibitor or angiotensin receptor blockade therapy.<sup>133</sup> For patients with more severe proteinuria, at high risk of disease



**Figure 3: Association of eGFR and albuminuria with hazard ratio of cardiovascular events**

Association of eGFR (creatinine-based eGFR presented with cystatin C-based eGFR for comparison) and albuminuria with hazard ratio of cardiovascular events after adjustment for traditional cardiovascular risk factors. Data from the Chronic Kidney Disease Prognosis Consortium of 637 315 individuals from 24 cohorts (ie, general population, high risk of chronic kidney disease, and established chronic kidney disease), followed up for a mean of 8.9 years, during which time 10 605 cardiovascular deaths, 6283 coronary heart disease events, 180 stroke events, and 2066 heart failure events occurred. Cystatin C-based eGFR data from a meta-analysis of ten general population cohorts with 64 010 participants, of whom 3193 died from cardiovascular causes during follow-up. eGFR=estimated glomerular filtration rate. The blue shaded area around the line represents the 95% CI. Adapted from Matsushita and colleagues<sup>42</sup> by permission of Elsevier and from Shlipak and colleagues.<sup>143</sup>

progression, or both, other treatments are recommended, including alkylating agents such as cyclophosphamide,<sup>134</sup> calcineurin inhibitors,<sup>135,136</sup> rituximab,<sup>137</sup> or a combination of these agents.<sup>138</sup> For other primary glomerular diseases, there is a paucity of evidence from randomised controlled trials.<sup>139</sup> Autosomal dominant polycystic kidney disease is another non-glomerular disease common worldwide and for which tolvaptan, a vasopressin receptor antagonist, has shown an effect in slowing the rate of kidney growth and glomerular filtration rate decline<sup>140</sup> in early and late stages.<sup>141</sup>

These traditional and emerging pharmacotherapies strategies are summarised in table 1 and in figure 1 (see also appendix pp 3–4).

### Addressing cardiovascular risk during chronic kidney disease progression

Cardiovascular disease is a leading cause of death in patients with chronic kidney disease, and is therefore a major focus of preservative care in this population.<sup>142</sup> Lower eGFR and higher albuminuria are independently associated with a higher risk of cardiovascular events (in addition to the risk conferred by traditional risk factors such as blood pressure, high cholesterol, and lifestyle) (figure 3).<sup>42,143</sup>

Whereas the traditional shared risk factors such as overweight, hypertension, diabetes, dyslipidaemia, and smoking are associated with cardiovascular disease in patients with earlier chronic kidney disease stages, a host of non-traditional risk factors magnify the risk of cardiovascular disease, especially in patients with advanced chronic kidney disease. Some of these so-called kidney-specific factors include inflammation with or without protein-energy wasting, mineral and bone disorders, and endothelial dysfunction.<sup>144–151</sup>

The management of cardiovascular disease in chronic kidney disease is challenging (appendix p 9).<sup>152</sup> These challenges include interpretation of cardiac biomarkers that are used to diagnose myocardial infarction, the exclusion of individuals with more advanced chronic kidney disease from cardiovascular outcome trials, and the absence of a clear benefit of revascularisation for individuals with chronic kidney disease and stable coronary artery disease.<sup>153–156</sup> The mainstays of cardiovascular risk reduction in chronic kidney disease include lifestyle modification (table 2), blood pressure reduction with renin–angiotensin–aldosterone pathway modulators, lipid reduction with statins, and specific glucose-lowering drugs that have been shown to reduce cardiovascular outcomes in people with type 2 diabetes (appendix pp 3–4).<sup>124,157</sup> Evidence suggests that in people with non-dialysis-dependent chronic kidney disease, higher physical activity is associated with slower disease progression,<sup>158</sup> lower risk of cardiovascular events and mortality,<sup>159,160</sup> and some improvements in kidney function and blood pressure;<sup>161</sup> in patients receiving dialysis, improvements in physical function and health-related quality of life are also observed.<sup>162</sup> Smoking is associated with a greater risk of incident chronic kidney disease<sup>163</sup> and increased risk of all-cause mortality in people with chronic kidney disease.<sup>164</sup> Dietary sodium restriction improves albuminuria, hypertension, fluid status, and arterial stiffness in people with and without chronic kidney disease,<sup>165–167</sup> which is also more likely to harbour salt-sensitive hypertension.<sup>168</sup> Higher plant-based sources of protein and dietary fibres may have salutary effects,<sup>52</sup> whereas higher red meat intake may be deleterious to patients with chronic kidney disease.<sup>169</sup> Protein intake recommendations vary depending on the stage of the disease, acute kidney injury events, and need for dialysis.<sup>70</sup> Weight reduction improves blood pressure in chronic kidney disease,<sup>170,171</sup> notwithstanding the obesity paradox in patients requiring dialysis.<sup>59</sup>

However, there is insufficient evidence to support commencing lipid-lowering therapy for primary prevention of cardiovascular disease in most people with advanced chronic kidney disease requiring dialysis, especially in the absence of high concentrations of serum LDLs.<sup>172</sup> In people with type 2 diabetes, guidelines recommend that SGLT2 inhibitors and GLP-1 receptor agonists be prioritised in people with chronic kidney disease.<sup>173,174</sup> Although antiplatelets are often offered for secondary cardiovascular prevention, the relative benefits and harms for primary prevention are unknown,<sup>175</sup> as is the role of anticoagulation with warfarin versus direct acting oral anticoagulants in advanced chronic kidney disease.<sup>176</sup> In a randomised controlled trial of patients with a history of acute coronary syndrome in the previous 3 months, type 2 diabetes, and low concentrations of HDLs, apabetalone (a novel epigenetic modulator) added to standard therapy resulted in a 52% reduction in the risk of major adverse cardiovascular events among subgroups

	Kidney-preserving care level of relevance	Supportive care level of relevance	Comments
Slowing chronic kidney disease progression	Strongly relevant	Possibly relevant to moderately relevant	Important goal across both domains of care, but not the primary goal of supportive care
Preventing or delaying dialysis	Strongly relevant	Strongly relevant	Both domains aim to prevent or delay initiation of dialysis
Symptom management	Possibly relevant to moderately relevant	Strongly relevant	Addressing symptoms is one of the key priorities of supportive care, but should also have an important role in preservative management
Health-related quality of life and patient-reported outcomes	Moderately relevant to strongly relevant	Strongly relevant	Each domain aims to improve quality of life through different approaches: kidney-preserving care prioritises aggressive treatments to restore and protect health and quality of life, whereas supportive care places a greater focus on maintaining quality of life by minimising invasive interventions
Prioritising overall survival with life-prolonging and kidney-prolonging care	Strongly relevant	Not relevant to possibly relevant	Preventing complications, addressing comorbidities, and prolonging survival are among the highest priorities of kidney-preserving care, whereas alleviating symptoms is prioritised over prolonging survival with supportive care
Cardiovascular health	Strongly relevant	Not relevant to possibly relevant	Reducing cardiovascular risk is a fundamental aspect of kidney-preserving care, and less of a priority with supportive care
Minimising risk of acute kidney injury and infection	Strongly relevant	Possibly relevant to moderately relevant	Important aspect of both domains
Diet and lifestyle modifications	Strongly relevant	Possibly relevant	Dietary strategies (eg, plant-dominant, low-protein diet) are an important component of kidney-preserving care, whereas dietary approaches can be more flexible with supportive care, focusing on quality of life and comfort care
Chronic kidney disease pharmacotherapy	Strongly relevant	Possibly relevant	Targeted and non-targeted pharmacotherapies aimed at preserving kidney function are less of a priority with supportive care, in which pharmacotherapy is mostly for symptom management
Treating uraemia and complications of chronic kidney disease	Strongly relevant	Possibly relevant to moderately relevant	Priorities in the management of chronic kidney disease-related complications are primarily to avoid unpleasant symptoms with supportive care
Preserving residual kidney function after transition to dialysis and less frequent dialysis	Strongly relevant	Possibly relevant	Kidney-preserving care places a greater emphasis on protecting residual kidney function (eg, incremental transition to dialysis), whereas palliative dialysis can be infrequent as well as include decremental dialysis regimens
Shared decision making and advance care planning	Moderately relevant	Strongly relevant	Fundamental concept in the care of all people with kidney disease and that is often more clearly highlighted under supportive care, including palliative care and hospice medicine

See also figure 1 for schematic representations of the domains within the chronic kidney disease management chart.

**Table 3: Similarities and distinctions between kidney-preserving management and supportive and palliative care**

of participants with pre-existing, stage 3 chronic kidney disease.<sup>177,178</sup> Patient-reported outcomes, including health-related quality of life, should be considered a key outcome for a holistic assessment of interventions in all cardiovascular outcome trials involving patients with chronic kidney disease.<sup>19,179</sup>

### Acute kidney injury

Patients with chronic kidney disease, especially in more advanced stages (eGFR of less than 30 mL/min per 1.73 m<sup>2</sup>) often do not exhibit linear progression of disease, which might be related to superimposed episodes of acute kidney injury or other factors.<sup>180</sup> Some (but not all) studies suggest that each acute kidney injury event might accelerate progression of chronic kidney disease.<sup>181,182</sup> Therefore, preventing acute kidney injury is an important component of the management of chronic kidney disease. This prevention involves avoiding acute kidney injury-associated drug combinations (eg, ACE inhibitors or angiotensin receptor blockers in conjunction with loop diuretics and non-steroidal anti-inflammatory drugs)<sup>183</sup> and preventing infections that can precipitate hypotension or septic shock necessitating the use of potentially nephrotoxic antimicrobials. Other contributors to acute kidney injury include cardiovascular events, particularly

decompensated heart failure leading to venous congestion and impaired kidney blood flow, or coronary artery bypass and other major surgeries with possible intraoperative hypotensive episodes.<sup>184,185</sup>

### Role of supportive care and of palliative and hospice medicine

People with advanced chronic kidney disease, particularly those with kidney failure, often have a high symptom burden that substantially affects their health-related quality of life.<sup>186</sup> Although there is little evidence from randomised controlled trials, observational studies suggest that chronic dialysis might not be associated with improved survival in some patients older than 80 years with a high burden of comorbidities, and that at least some of these patients might regret their decision to commence dialysis in view of treatment-related complications, high symptom burden, and poor quality of life.<sup>187,188</sup> These factors have led to an increased recognition of the importance of supportive care of kidney failure (figure 1).

There are similarities and distinctions between kidney-preserving management and supportive care, including palliative care and hospice medicine (table 3). Both strategies are expected to minimise the risk of adverse events or complications, such as acute kidney injury, as chronic kidney disease progresses, although

kidney-preserving management is more strongly focused on kidney function longevity. Whereas palliative care strategies are often considered for those with advanced age (ie, older than 80 years) or more severe comorbidities, such as terminal cancer,<sup>21–24,26–33</sup> kidney-preserving management is a kidney-sustaining and life-sustaining strategy for all individuals and at any stage of chronic kidney disease.

The goal of supportive care is to improve symptoms and quality of life through a multidisciplinary approach that incorporates shared decision making, detailed communication with patients and their care partners, advanced care planning, and psychological and social support.<sup>189,190</sup> Standardised tools to identify those who might benefit the most from supportive care are not widely validated, and thus treatment decisions, such as fluid management, must be individualised—including for those who decide to reduce dialysis dose and frequency to a minimum (the so-called decremental or palliative dialysis), or to withdraw completely from renal replacement therapy (figure 1).<sup>191</sup> Novel prediction tools have been developed to identify individuals who would have the highest mortality in the first year after transitioning to dialysis and who might therefore benefit from palliative care,<sup>45</sup> but these tools need to be more widely validated. The decision must be according to the free choice of the patient, without pressure by family members and care partners or health-care professionals, and it should not be influenced by rationing dialysis care such as during COVID-19 pandemic surges or other resource constraints.<sup>17</sup> Importantly, preservation of kidney function and supportive care are entirely complementary and should be considered as parts of the full spectrum of conservative management of chronic kidney disease, depending on the severity of the disease and on the goals of the individual (figure 1; table 3).

### Infection control and management of chronic kidney disease in the COVID-19 pandemic

Evidence suggests that uraemia is associated with worse immune response, as exemplified by the diminished antibody response to hepatitis B vaccination or by the higher risk of infections as kidney function worsens, not otherwise explained by concurrent comorbidities, and by the historical observation that autoimmune diseases such as systemic lupus erythematosus are less aggressive once patients have uraemia.<sup>192</sup> Similarly, some infections, such as hepatitis C, might cause chronic kidney disease if untreated,<sup>193,194</sup> or can lead to faster disease progression and greater mortality in the case of pre-existing chronic kidney disease.<sup>195</sup> Conversely, treatment of hepatitis C might possibly have a salutary effect in preserving kidney function.<sup>196</sup> Patients with chronic kidney disease have a two times higher risk of death after respiratory infections;<sup>197</sup> therefore, influenza and pneumococcal vaccinations might be an indirect way to prevent acute kidney injury and avoid chronic kidney disease

progression, although this approach has not been assessed in clinical trials.<sup>198</sup>

During the COVID-19 pandemic, data suggest a two times increase in mortality from COVID-19 in the presence of chronic kidney disease.<sup>199</sup> Despite screening and isolation of affected patients, outbreaks cannot always be prevented because some infected individuals can have long incubation periods or be asymptomatic carriers.<sup>199</sup> To maintain adequate staffing and to protect patients, outpatient chronic kidney disease care has undergone a radical transformation with the use of telemedicine and remote care to guide dietary and therapeutic decisions.<sup>200</sup> Advance care planning and ensuring completion of renal replacement therapy plans are of even higher importance because COVID-19 can cause acute kidney injury through septic shock, cytokine release, or direct renal tropism of the virus.<sup>201,202</sup> Ongoing and future trials are expected to examine whether ACE receptor modulators or other modulators of kidney function can avert COVID-19 involvement in acute kidney injury and chronic kidney disease progression.<sup>201–203</sup>

### Global and regional disparities in preserving kidney care

Preserving kidney function in people with chronic kidney disease is confounded by regional and global health inequalities, disparities in health-care models, pharmaceutical industry policies, and geopolitical and fiscal complexities.<sup>204</sup> Income disparities, poverty, and social disadvantages have a dominant effect on the risk of incident chronic kidney disease and its progression.<sup>205,206</sup> Many people at risk of chronic kidney disease in low-income and middle-income countries are not provided with appropriate infrastructure for screening and identification of chronic kidney disease. The awareness of chronic kidney disease is relatively poor, and opportunities for updating knowledge and education on preserving kidney health are scarce.<sup>207</sup>

In low-income and middle-income countries, diet and lifestyle modification might offer an inexpensive approach for primary, secondary, and tertiary prevention of chronic kidney disease, notwithstanding the little scientific evidence to justify population-based programmes for lower sodium intake, promotion of plant-based diets, and adequate hydration. Whereas pharmacotherapy and dialysis treatment are almost universally accessible in high-income countries, many people in low-income and middle-income countries are unable to access these options.<sup>208</sup> The out-of-pocket expenditure for chronic kidney disease preserving pharmacotherapy is high relative to income; for instance, in India, the high costs of SGLT2 inhibitors or renal replacement therapy are more likely to push people into the poverty range than communicable diseases.<sup>209</sup>

In some high-income nations, such as the USA, there are racial disparities in chronic kidney disease care.<sup>210</sup>

Black Americans are far more likely to develop chronic kidney disease and exhibit faster disease progression than White Americans.<sup>211</sup> This disparity might be related to socioeconomic factors such as access to care, exposing many Black Americans to higher chronic kidney disease risks; furthermore the higher prevalence of a pathogenic *APOL1* allele in Black Americans might partly explain the high burden of chronic kidney disease in this population.<sup>212</sup> The current creatinine-based eGFR equations have a race index for Black Americans, which inflates the eGFR value by as high as 16% compared with non-Black Americans with the same serum creatinine concentration.<sup>213</sup> In an effort to reduce racial disparities in chronic kidney disease care, race indices have been suggested to be removed from eGFR equations to enable a more commensurate approach to the burden of chronic kidney disease.<sup>214</sup> Race-free measurements, such as serum cystatin C-derived eGFR equations, could be used instead, especially given their more linear associations with clinical outcomes (figure 3).<sup>143,214</sup>

### Conclusion remarks and future steps

The primary goal of the conservative management of chronic kidney disease through kidney-preserving care is to slow chronic kidney disease progression to prolong dialysis-free time while striving to achieve the greatest quality of life and survival. These strategies also include effective treatment of renal and non-renal comorbidities and their associated symptoms.<sup>18</sup> There are similarities and distinctions between kidney-preserving management and supportive care in chronic kidney disease. Dietary interventions, such as low-salt and plant-dominant, low-protein diets, and pharmacotherapies, such as renin-angiotensin-aldosterone pathway modulators, SGLT2 inhibitors, and newer non-steroidal mineralocorticoid receptor antagonists should be used to the best possible extent. Adequately powered, randomised trials to evaluate the effects of new pharmacotherapies to slow the progression of chronic kidney disease and prevent its complications, especially in early-stage disease, should be prioritised. When dialysis therapy is necessary and kidney transplantation cannot be offered, a gradual and incremental transition to dialysis might preserve residual kidney function for longer. Some patients might benefit from palliative dialysis for ultrafiltration to manage fluid overload and its associated symptoms.<sup>215</sup> Future studies of strategies to replace or complement dialysis should include, but not be limited to, intestinal dialysis,<sup>216,217</sup> diaphoresis therapy for control of uraemia, and management of fluid and electrolytes,<sup>218,219</sup> on the basis of the results from studies in relevant animal models.<sup>220</sup> Future studies are needed to refine current interventions and to examine novel models and strategies to prolong kidney and patient survival without dialysis, if possible, and to help patients to live longer and well with chronic kidney disease.<sup>19</sup>

### Contributors

All authors contributed equally to the conception, interpretation of the relevant literature, writing, and critical revision of this Seminar.

### Declaration of interests

KK-Z reports commercial honoraria and support from Abbott, AbbVie, Alexion, Amgen, AstraZeneca, Aveo, Chugai, DaVita, Fresenius, Genentech, Haymarket Media, Hospira, Kabi, Keryx, Novartis, Pfizer, Relypsa, Resverlogix, Sandoz, Sanofi, Shire, Vifor, UpToDate, and ZS Pharma. DN is part of the steering group of GlaxoSmithKline-funded studies investigating aspects of kidney disease in sub-Saharan Africa. BLN reports travel support from Janssen and consultancy fees (trial steering committee member) from Bayer paid to institution. VP has served on steering committees, advisory boards, or given scientific presentations supported by AbbVie, Amgen, Astellas, AstraZeneca, Bayer, Baxter, Bristol Myers Squibb, Boehringer Ingelheim, Chinook, Dimerix, Durect, Eli Lilly, Gilead Sciences, GlaxoSmithKline, Janssen, Merck Sharp & Dohme, Metavant, Mitsubishi Tanabe, Mundipharma, Novartis, Novo Nordisk, Pfizer, Pharmalink, Relypsa, Retrophin, Sanofi, Servier, Vifor, Vitae, UpToDate, and Tricida. THJ declares no competing interests.

### Acknowledgments

THJ is supported by the Singapore National Medical Research Council. This work was supported by research grant K24-DK091419 from the National Institutes of Health and by philanthropic grants from Harold Simmons and Joseph Lee.

### References

- 1 Webster AC, Nagler EV, Morton RL, Masson P. Chronic kidney disease. *Lancet* 2017; **389**: 1238–52.
- 2 Bikbov B, Purcell CA, Levey AS, et al. Global, regional, and national burden of chronic kidney disease, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2020; **395**: 709–33.
- 3 Xie Y, Bowe B, Mokdad AH, et al. Analysis of the Global Burden of Disease study highlights the global, regional, and national trends of chronic kidney disease epidemiology from 1990 to 2016. *Kidney Int* 2018; **94**: 567–81.
- 4 Foreman KJ, Marquez N, Dolgert A, et al. Forecasting life expectancy, years of life lost, and all-cause and cause-specific mortality for 250 causes of death: reference and alternative scenarios for 2016–40 for 195 countries and territories. *Lancet* 2018; **392**: 2052–90.
- 5 Ku E, Lee BJ, Wei J, Weir MR. Hypertension in CKD: core curriculum 2019. *Am J Kidney Dis* 2019; **74**: 120–31.
- 6 Pearce N, Caplin B. Let's take the heat out of the CKDu debate: more evidence is needed. *Occup Environ Med* 2019; **76**: 357–59.
- 7 Bowe B, Artimovich E, Xie Y, Yan Y, Cai M, Al-Aly Z. The global and national burden of chronic kidney disease attributable to ambient fine particulate matter air pollution: a modelling study. *BMJ Glob Health* 2020; **5**: e002063.
- 8 Cockwell P, Fisher LA. The global burden of chronic kidney disease. *Lancet* 2020; **395**: 662–64.
- 9 Silverwood RJ, Pierce M, Thomas C, et al. Association between younger age when first overweight and increased risk for CKD. *J Am Soc Nephrol* 2013; **24**: 813–21.
- 10 Zarantonello D, Rhee CM, Kalantar-Zadeh K, Brunori G. Novel conservative management of chronic kidney disease via dialysis-free interventions. *Curr Opin Nephrol Hypertens* 2021; **30**: 97–107.
- 11 Cano N, Fernandez JP, Lacombe P, et al. Statistical selection of nutritional parameters in hemodialyzed patients. *Kidney Int Suppl* 1987; **22**: S178–80.
- 12 Song MK. Quality of life of patients with advanced chronic kidney disease receiving conservative care without dialysis. *Semin Dial* 2016; **29**: 165–69.
- 13 Bello AK, Levin A, Lunney M, et al. Status of care for end stage kidney disease in countries and regions worldwide: international cross sectional survey. *BMJ* 2019; **367**: l5873.
- 14 Kurella Tamura M, Desai M, Kappahh KI, Thomas IC, Asch SM, Chertow GM. Dialysis versus medical management at different ages and levels of kidney function in veterans with advanced CKD. *J Am Soc Nephrol* 2018; **29**: 2169–77.

- 15 Kalantar-Zadeh K, Kovesdy CP, Streja E, et al. Transition of care from pre-dialysis prelude to renal replacement therapy: the blueprints of emerging research in advanced chronic kidney disease. *Nephrol Dial Transplant* 2017; **32** (suppl 2): ii91–98.
- 16 O'Connor NR, Kumar P. Conservative management of end-stage renal disease without dialysis: a systematic review. *J Palliat Med* 2012; **15**: 228–35.
- 17 Kalantar-Zadeh K, Wightman A, Liao S. Ensuring choice for people with kidney failure—dialysis, supportive care, and hope. *N Engl J Med* 2020; **383**: 99–101.
- 18 Rhee CM, Nguyen DV, Nyamathi A, Kalantar-Zadeh K. Conservative vs preservative management of chronic kidney disease: similarities and distinctions. *Curr Opin Nephrol Hypertens* 2020; **29**: 92–102.
- 19 Kalantar-Zadeh K, Kam-Tao Li P, Tantisattamo E, et al. Living well with kidney disease by patient and care-partner empowerment: kidney health for everyone everywhere. *Kidney Int* 2021; **99**: 278–84.
- 20 Teehan BP, Schleifer CR, Brown JM, Sigler MH, Raimondo J. Urea kinetic analysis and clinical outcome on CAPD. A five year longitudinal study. *Adv Perit Dial* 1990; **6**: 181–85.
- 21 Brown MA, Collett GK, Josland EA, Foote C, Li Q, Brennan FP. CKD in elderly patients managed without dialysis: survival, symptoms, and quality of life. *Clin J Am Soc Nephrol* 2015; **10**: 260–68.
- 22 Carson RC, Juszczak M, Davenport A, Burns A. Is maximum conservative management an equivalent treatment option to dialysis for elderly patients with significant comorbid disease? *Clin J Am Soc Nephrol* 2009; **4**: 1611–19.
- 23 Chandna SM, Da Silva-Gane M, Marshall C, Warwicker P, Greenwood RN, Farrington K. Survival of elderly patients with stage 5 CKD: comparison of conservative management and renal replacement therapy. *Nephrol Dial Transplant* 2011; **26**: 1608–14.
- 24 Da Silva-Gane M, Wellsted D, Greenshields H, Norton S, Chandna SM, Farrington K. Quality of life and survival in patients with advanced kidney failure managed conservatively or by dialysis. *Clin J Am Soc Nephrol* 2012; **7**: 2002–09.
- 25 Hemmelgarn BR, James MT, Manns BJ, et al. Rates of treated and untreated kidney failure in older vs younger adults. *JAMA* 2012; **307**: 2507–15.
- 26 Joly D, Anglicheau D, Alberti C, et al. Octogenarians reaching end-stage renal disease: cohort study of decision-making and clinical outcomes. *J Am Soc Nephrol* 2003; **14**: 1012–21.
- 27 Kamar FB, Tam-Tham H, Thomas C. A description of advanced chronic kidney disease patients in a major urban center receiving conservative care. *Can J Kidney Health Dis* 2017; **4**: 2054358117718538.
- 28 Murtagh FE, Marsh JE, Donohoe P, Ekbal NJ, Sheerin NS, Harris FE. Dialysis or not? A comparative survival study of patients over 75 years with chronic kidney disease stage 5. *Nephrol Dial Transplant* 2007; **22**: 1955–62.
- 29 Reindl-Schwaighofer R, Kainz A, Kammer M, Dumfarth A, Oberbauer R. Survival analysis of conservative vs dialysis treatment of elderly patients with CKD stage 5. *PLoS One* 2017; **12**: e0181345.
- 30 Seow YY, Cheung YB, Qu LM, Yee AC. Trajectory of quality of life for poor prognosis stage 5D chronic kidney disease with and without dialysis. *Am J Nephrol* 2013; **37**: 231–38.
- 31 Shum CK, Tam KF, Chak WL, Chan TC, Mak YF, Chau KF. Outcomes in older adults with stage 5 chronic kidney disease: comparison of peritoneal dialysis and conservative management. *J Gerontol A Biol Sci Med Sci* 2014; **69**: 308–14.
- 32 Teruel JL, Burguera Vion V, Gomis Couto A, et al. Choosing conservative therapy in chronic kidney disease. *Nefrologia* 2015; **35**: 273–79.
- 33 Verberne WR, Geers AB, Jellema WT, Vincent HH, van Delden JJ, Bos WJ. Comparative survival among older adults with advanced kidney disease managed conservatively versus with dialysis. *Clin J Am Soc Nephrol* 2016; **11**: 633–40.
- 34 Li PK, Garcia-Garcia G, Lui SF, et al. Kidney health for everyone everywhere—from prevention to detection and equitable access to care. *Kidney Int* 2020; **97**: 226–32.
- 35 Stengel B, Tarver-Carr ME, Powe NR, Eberhardt MS, Brancati FL. Lifestyle factors, obesity and the risk of chronic kidney disease. *Epidemiology* 2003; **14**: 479–87.
- 36 Komenda P, Ferguson TW, Macdonald K, et al. Cost-effectiveness of primary screening for CKD: a systematic review. *Am J Kidney Dis* 2014; **63**: 789–97.
- 37 Gansevoort RT, Correa-Rotter R, Hemmelgarn BR, et al. Chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. *Lancet* 2013; **382**: 339–52.
- 38 Astor BC, Matsushita K, Gansevoort RT, et al. Lower estimated glomerular filtration rate and higher albuminuria are associated with mortality and end-stage renal disease. A collaborative meta-analysis of kidney disease population cohorts. *Kidney Int* 2011; **79**: 1331–40.
- 39 Matsushita K, van der Velde M, Astor BC, et al. Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality in general population cohorts: a collaborative meta-analysis. *Lancet* 2010; **375**: 2073–81.
- 40 Gansevoort RT, Matsushita K, van der Velde M, et al. Lower estimated GFR and higher albuminuria are associated with adverse kidney outcomes. A collaborative meta-analysis of general and high-risk population cohorts. *Kidney Int* 2011; **80**: 93–104.
- 41 van der Velde M, Matsushita K, Coresh J, et al. Lower estimated glomerular filtration rate and higher albuminuria are associated with all-cause and cardiovascular mortality. A collaborative meta-analysis of high-risk population cohorts. *Kidney Int* 2011; **79**: 1341–52.
- 42 Matsushita K, Coresh J, Sang Y, et al. Estimated glomerular filtration rate and albuminuria for prediction of cardiovascular outcomes: a collaborative meta-analysis of individual participant data. *Lancet Diabetes Endocrinol* 2015; **3**: 514–25.
- 43 Levin A, Stevens PE, Bilous RW, et al. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int Suppl* 2013; **3**: 1–150.
- 44 Tangri N, Stevens LA, Griffith J, et al. A predictive model for progression of chronic kidney disease to kidney failure. *JAMA* 2011; **305**: 1553–59.
- 45 Obi Y, Nguyen DV, Zhou H, et al. Development and validation of prediction scores for early mortality at transition to dialysis. *Mayo Clin Proc* 2018; **93**: 1224–35.
- 46 Obi Y, Rhee CM, Mathew AT, et al. Residual kidney function decline and mortality in incident hemodialysis patients. *J Am Soc Nephrol* 2016; **27**: 3758–68.
- 47 Kovesdy CP, Furth SL, Zoccali C, World Kidney Day Steering Committee. Obesity and kidney disease: hidden consequences of the epidemic. *Can J Kidney Health Dis* 2017; **4**: 2054358117698669.
- 48 Chang AR, Grams ME, Ballew SH, et al. Adiposity and risk of decline in glomerular filtration rate: meta-analysis of individual participant data in a global consortium. *BMJ* 2019; **364**: k5301.
- 49 Tonneijck L, Muskiet MHA, Smits MM, et al. Glomerular hyperfiltration in diabetes: mechanisms, clinical significance, and treatment. *J Am Soc Nephrol* 2017; **28**: 1023–39.
- 50 Herrington WG, Smith M, Bankhead C, et al. Body-mass index and risk of advanced chronic kidney disease: prospective analyses from a primary care cohort of 1.4 million adults in England. *PLoS One* 2017; **12**: e0173515.
- 51 Look AHEAD Research Group. Effect of a long-term behavioural weight loss intervention on nephropathy in overweight or obese adults with type 2 diabetes: a secondary analysis of the Look AHEAD randomised clinical trial. *Lancet Diabetes Endocrinol* 2014; **2**: 801–09.
- 52 Kalantar-Zadeh K, Joshi S, Schlueter R, et al. Plant-dominant low-protein diet for conservative management of chronic kidney disease. *Nutrients* 2020; **12**: E1931.
- 53 Joshi S, McMacken M, Kalantar-Zadeh K. Plant-based diets for kidney disease: a guide for clinicians. *Am J Kidney Dis* 2021; **77**: 287–96.
- 54 Scirica BM, Bohula EA, Dwyer JP, et al. Lorcaserin and renal outcomes in obese and overweight patients in the CAMELLIA-TIMI 61 trial. *Circulation* 2019; **139**: 366–75.
- 55 Chang AR, Grams ME, Navaneethan SD. Bariatric surgery and kidney-related outcomes. *Kidney Int Rep* 2017; **2**: 261–70.
- 56 Chang AR, Chen Y, Still C, et al. Bariatric surgery is associated with improvement in kidney outcomes. *Kidney Int* 2016; **90**: 164–71.
- 57 Cohen RV, Pereira TV, Aboud CM, et al. Effect of gastric bypass vs best medical treatment on early-stage chronic kidney disease in patients with type 2 diabetes and obesity: a randomized clinical trial. *JAMA Surg* 2020; **155**: e200420.

- 58 Naderi N, Kleine CE, Park C, et al. Obesity paradox in advanced kidney disease: from bedside to the bench. *Prog Cardiovasc Dis* 2018; **61**: 168–81.
- 59 Kalantar-Zadeh K, Rhee CM, Chou J, et al. The obesity paradox in kidney disease: how to reconcile it with obesity management. *Kidney Int Rep* 2017; **2**: 271–81.
- 60 Rhee CM, Ahmadi SF, Kalantar-Zadeh K. The dual roles of obesity in chronic kidney disease: a review of the current literature. *Curr Opin Nephrol Hypertens* 2016; **25**: 208–16.
- 61 Tovar-Palacio C, Tovar AR, Torres N, et al. Proinflammatory gene expression and renal lipogenesis are modulated by dietary protein content in obese Zucker fa/fa rats. *Am J Physiol Renal Physiol* 2011; **300**: F263–71.
- 62 Kalantar-Zadeh K, Fouque D. Nutritional management of chronic kidney disease. *N Engl J Med* 2017; **377**: 1765–76.
- 63 Koppe L, Fouque D. The role for protein restriction in addition to renin-angiotensin-aldosterone system inhibitors in the management of CKD. *Am J Kidney Dis* 2019; **73**: 248–57.
- 64 Levey AS, Greene T, Beck GJ, et al. Dietary protein restriction and the progression of chronic renal disease: what have all of the results of the MDRD study shown? Modification of Diet in Renal Disease Study group. *J Am Soc Nephrol* 1999; **10**: 2426–39.
- 65 Peterson JC, Adler S, Burkart JM, et al. Blood pressure control, proteinuria, and the progression of renal disease. The Modification of Diet in Renal Disease Study. *Ann Intern Med* 1995; **123**: 754–62.
- 66 Rhee CM, Ahmadi SF, Kovesdy CP, Kalantar-Zadeh K. Low-protein diet for conservative management of chronic kidney disease: a systematic review and meta-analysis of controlled trials. *J Cachexia Sarcopenia Muscle* 2018; **9**: 235–45.
- 67 Fouque D, Laville M. Low protein diets for chronic kidney disease in non diabetic adults. *Cochrane Database Syst Rev* 2009; **3**: CD001892.
- 68 Kasiske BL, Lakatua JD, Ma JZ, Louis TA. A meta-analysis of the effects of dietary protein restriction on the rate of decline in renal function. *Am J Kidney Dis* 1998; **31**: 954–61.
- 69 Chewcharat A, Takkavatakar N, Wongrattanagorn S, et al. The effects of restricted protein diet supplemented with ketoanalogue on renal function, blood pressure, nutritional status, and chronic kidney disease-mineral and bone disorder in chronic kidney disease patients: a systematic review and meta-analysis. *J Ren Nutr* 2020; **30**: 189–99.
- 70 Kovesdy CP, Kopple JD, Kalantar-Zadeh K. Management of protein-energy wasting in non-dialysis-dependent chronic kidney disease: reconciling low protein intake with nutritional therapy. *Am J Clin Nutr* 2013; **97**: 1163–77.
- 71 Moore LW, Byham-Gray LD, Scott Parrott J, et al. The mean dietary protein intake at different stages of chronic kidney disease is higher than current guidelines. *Kidney Int* 2013; **83**: 724–32.
- 72 Kalantar-Zadeh K, Moore LW. Does kidney longevity mean healthy vegan food and less meat or is any low-protein diet good enough? *J Ren Nutr* 2019; **29**: 79–81.
- 73 Kim H, Caulfield LE, Garcia-Larsen V, et al. Plant-based diets and incident CKD and kidney function. *Clin J Am Soc Nephrol* 2019; **14**: 682–91.
- 74 Ko GJ, Rhee CM, Kalantar-Zadeh K, Joshi S. The effects of high-protein diets on kidney health and longevity. *J Am Soc Nephrol* 2020; **31**: 1667–79.
- 75 Joshi S, Shah S, Kalantar-Zadeh K. Adequacy of plant-based proteins in chronic kidney disease. *J Ren Nutr* 2019; **29**: 112–17.
- 76 Mafra D, Borges NA, Lindholm B, Shiels PG, Evenepoel P, Stenvinkel P. Food as medicine: targeting the uraemic phenotype in chronic kidney disease. *Nat Rev Nephrol* 2021; **17**: 153–71.
- 77 Koppe L, Mafra D, Fouque D. Probiotics and chronic kidney disease. *Kidney Int* 2015; **88**: 958–66.
- 78 Chauveau P, Combe C, Fouque D, Aparicio M. Vegetarianism: advantages and drawbacks in patients with chronic kidney diseases. *J Ren Nutr* 2013; **23**: 399–405.
- 79 Lau WL, Savoj J, Nakata MB, Vaziri ND. Altered microbiome in chronic kidney disease: systemic effects of gut-derived uremic toxins. *Clin Sci (Lond)* 2018; **132**: 509–22.
- 80 Wu IW, Hsu KH, Lee CC, et al. p-Cresyl sulphate and indoxyl sulphate predict progression of chronic kidney disease. *Nephrol Dial Transplant* 2011; **26**: 938–47.
- 81 Koppe L, Pillon NJ, Vella RE, et al. p-Cresyl sulfate promotes insulin resistance associated with CKD. *J Am Soc Nephrol* 2013; **24**: 88–99.
- 82 Wu IW, Hsu KH, Hsu HJ, et al. Serum free p-cresyl sulfate levels predict cardiovascular and all-cause mortality in elderly hemodialysis patients—a prospective cohort study. *Nephrol Dial Transplant* 2012; **27**: 1169–75.
- 83 Carrero JJ, González-Ortiz A, Avesani CM, et al. Plant-based diets to manage the risks and complications of chronic kidney disease. *Nat Rev Nephrol* 2020; **16**: 525–42.
- 84 Hung S-C, Kuo K-L, Peng C-H, et al. Volume overload correlates with cardiovascular risk factors in patients with chronic kidney disease. *Kidney Int* 2014; **85**: 703–09.
- 85 Tsai Y-C, Tsai J-C, Chen S-C, et al. Association of fluid overload with kidney disease progression in advanced CKD: a prospective cohort study. *Am J Kidney Dis* 2014; **63**: 68–75.
- 86 Hung SC, Kuo KL, Peng CH, Wu CH, Wang YC, Tarng DC. Association of fluid retention with anemia and clinical outcomes among patients with chronic kidney disease. *J Am Heart Assoc* 2015; **4**: e001480.
- 87 Kimura G, Dohi Y, Fukuda M. Salt sensitivity and circadian rhythm of blood pressure: the keys to connect CKD with cardiovascular events. *Hypertens Res* 2010; **33**: 515–20.
- 88 Bovée DM, Visser WJ, Middel I, et al. A randomized trial of distal diuretics versus dietary sodium restriction for hypertension in chronic kidney disease. *J Am Soc Nephrol* 2020; **31**: 650–62.
- 89 Agarwal R, Cramer AE, Balmes-Fenwick M, Sinha AD, Ouyang F, Tu W. Design and baseline characteristics of the Chlorothalidone in Chronic Kidney Disease (CLICK) trial. *Am J Nephrol* 2020; **51**: 542–52.
- 90 Kovesdy CP, Matsushita K, Sang Y, et al. Serum potassium and adverse outcomes across the range of kidney function: a CKD Prognosis Consortium meta-analysis. *Eur Heart J* 2018; **39**: 1535–42.
- 91 Clegg DJ, Hill Gallant KM. Plant-based diets in CKD. *Clin J Am Soc Nephrol* 2019; **14**: 141–43.
- 92 Sarwar CMS, Bhagat AA, Anker SD, Butler J. Role of hyperkalemia in heart failure and the therapeutic use of potassium binders. *Handb Exp Pharmacol* 2017; **243**: 537–60.
- 93 Georgianos PI, Agarwal R. Revisiting RAAS blockade in CKD with newer potassium-binding drugs. *Kidney Int* 2018; **93**: 325–34.
- 94 Kraut JA, Kurtz I. Metabolic acidosis of CKD: diagnosis, clinical characteristics, and treatment. *Am J Kidney Dis* 2005; **45**: 978–93.
- 95 Navaneethan SD, Shao J, Buysse J, Bushinsky DA. Effects of treatment of metabolic acidosis in CKD: a systematic review and meta-analysis. *Clin J Am Soc Nephrol* 2019; **14**: 1011–20.
- 96 Wesson DE, Mathur V, Tangri N, et al. Veverimer versus placebo in patients with metabolic acidosis associated with chronic kidney disease: a multicentre, randomised, double-blind, controlled, phase 3 trial. *Lancet* 2019; **393**: 1417–27.
- 97 Brenner BM, Cooper ME, de Zeeuw D, et al. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001; **345**: 861–69.
- 98 Lewis EJ, Hunsicker LG, Clarke WR, et al. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001; **345**: 851–60.
- 99 Fried LF, Emanuele N, Zhang JH, et al. Combined angiotensin inhibition for the treatment of diabetic nephropathy. *N Engl J Med* 2013; **369**: 1892–903.
- 100 Parving HH, Brenner BM, McMurray JJ, et al. Cardiorenal end points in a trial of aliskiren for type 2 diabetes. *N Engl J Med* 2012; **367**: 2204–13.
- 101 Yusuf S, Teo KK, Pogue J, et al. Telmisartan, ramipril, or both in patients at high risk for vascular events. *N Engl J Med* 2008; **358**: 1547–59.
- 102 Bakris GL, Agarwal R, Anker SD, et al. Effect of finerenone on chronic kidney disease outcomes in type 2 diabetes. *N Engl J Med* 2020; **383**: 2219–29.
- 103 Bombardieri AS, Kshirsagar AV, Amamoo MA, Klemmer PJ. Change in proteinuria after adding aldosterone blockers to ACE inhibitors or angiotensin receptor blockers in CKD: a systematic review. *Am J Kidney Dis* 2008; **51**: 199–211.
- 104 Jafar TH, Schmid CH, Landa M, et al. Angiotensin-converting enzyme inhibitors and progression of nondiabetic renal disease. A meta-analysis of patient-level data. *Ann Intern Med* 2001; **135**: 73–87.

- 105 Xie X, Atkins E, Lv J, et al. Effects of intensive blood pressure lowering on cardiovascular and renal outcomes: updated systematic review and meta-analysis. *Lancet* 2016; **387**: 435–43.
- 106 Jafar TH, Stark PC, Schmid CH, et al. Progression of chronic kidney disease: the role of blood pressure control, proteinuria, and angiotensin-converting enzyme inhibition: a patient-level meta-analysis. *Ann Intern Med* 2003; **139**: 244–52.
- 107 Obi Y, Kalantar-Zadeh K, Shintani A, Kovesdy CP, Hamano T. Estimated glomerular filtration rate and the risk-benefit profile of intensive blood pressure control amongst nondiabetic patients: a post hoc analysis of a randomized clinical trial. *J Intern Med* 2018; **283**: 314–27.
- 108 Sim JJ, Shi J, Kovesdy CP, Kalantar-Zadeh K, Jacobsen SJ. Impact of achieved blood pressures on mortality risk and end-stage renal disease among a large, diverse hypertension population. *J Am Coll Cardiol* 2014; **64**: 588–97.
- 109 Kovesdy CP, Lu JL, Molnar MZ, et al. Observational modeling of strict vs conventional blood pressure control in patients with chronic kidney disease. *JAMA Intern Med* 2014; **174**: 1442–49.
- 110 Malhotra R, Craven T, Ambrosius WT, et al. Effects of intensive blood pressure lowering on kidney tubule injury in CKD: a longitudinal subgroup analysis in SPRINT. *Am J Kidney Dis* 2019; **73**: 21–30.
- 111 Alicic RZ, Rooney MT, Tuttle KR. Diabetic kidney disease: challenges, progress, and possibilities. *Clin J Am Soc Nephrol* 2017; **12**: 2032–45.
- 112 Perkovic V, Heerspink HL, Chalmers J, et al. Intensive glucose control improves kidney outcomes in patients with type 2 diabetes. *Kidney Int* 2013; **83**: 517–23.
- 113 Coca SG, Ismail-Beigi F, Haq N, Krumholz HM, Parikh CR. Role of intensive glucose control in development of renal end points in type 2 diabetes mellitus: systematic review and meta-analysis intensive glucose control in type 2 diabetes. *Arch Intern Med* 2012; **172**: 761–69.
- 114 Heerspink HJL, Desai M, Jardine M, Balis D, Meininger G, Perkovic V. Canagliflozin slows progression of renal function decline independently of glycemic effects. *J Am Soc Nephrol* 2017; **28**: 368–75.
- 115 Wanner C, Inzucchi SE, Lachin JM, et al. Empagliflozin and progression of kidney disease in type 2 diabetes. *N Engl J Med* 2016; **375**: 323–34.
- 116 Perkovic V, de Zeeuw D, Mahaffey KW, et al. Canagliflozin and renal outcomes in type 2 diabetes: results from the CANVAS Program randomised clinical trials. *Lancet Diabetes Endocrinol* 2018; **6**: 691–704.
- 117 Mosenzon O, Wiviott SD, Cahn A, et al. Effects of dapagliflozin on development and progression of kidney disease in patients with type 2 diabetes: an analysis from the DECLARE-TIMI 58 randomised trial. *Lancet Diabetes Endocrinol* 2019; **7**: 606–17.
- 118 Perkovic V, Jardine MJ, Neal B, et al. Canagliflozin and renal outcomes in type 2 diabetes and nephropathy. *N Engl J Med* 2019; **380**: 2295–306.
- 119 Heerspink HJL, Stefánsson BV, Chertow GM, et al. Rationale and protocol of the Dapagliflozin And Prevention of Adverse outcomes in Chronic Kidney Disease (DAPA-CKD) randomized controlled trial. *Nephrol Dial Transplant* 2020; **35**: 274–82.
- 120 Wheeler DC, Stefánsson BV, Jongs N, et al. Effects of dapagliflozin on major adverse kidney and cardiovascular events in patients with diabetic and non-diabetic chronic kidney disease: a prespecified analysis from the DAPA-CKD trial. *Lancet Diabetes Endocrinol* 2021; **9**: 22–31.
- 121 Herrington WG, Preiss D, Haynes R, et al. The potential for improving cardio-renal outcomes by sodium-glucose co-transporter-2 inhibition in people with chronic kidney disease: a rationale for the EMPA-KIDNEY study. *Clin Kidney J* 2018; **11**: 749–61.
- 122 Verma S, McMurray JVV. SGLT2 inhibitors and mechanisms of cardiovascular benefit: a state-of-the-art review. *Diabetologia* 2018; **61**: 2108–17.
- 123 Rosenstock J, Perkovic V, Johansen OE, et al. Effect of linagliptin vs placebo on major cardiovascular events in adults with type 2 diabetes and high cardiovascular and renal risk: the CARMELINA randomized clinical trial. *JAMA* 2019; **321**: 69–79.
- 124 Kristensen SL, Rørth R, Jhund PS, et al. Cardiovascular, mortality, and kidney outcomes with GLP-1 receptor agonists in patients with type 2 diabetes: a systematic review and meta-analysis of cardiovascular outcome trials. *Lancet Diabetes Endocrinol* 2019; **7**: 776–85.
- 125 Floege J, Barbour SJ, Catran DC, et al. Management and treatment of glomerular diseases (part 1): conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Controversies Conference. *Kidney Int* 2019; **95**: 268–80.
- 126 Rauen T, Eitner F, Fitzner C, et al. Intensive supportive care plus immunosuppression in IgA nephropathy. *N Engl J Med* 2015; **373**: 2225–36.
- 127 Lv J, Zhang H, Wong MG, et al. Effect of oral methylprednisolone on clinical outcomes in patients with IgA nephropathy: the TESTING randomized clinical trial. *JAMA* 2017; **318**: 432–42.
- 128 Barratt J, Rovin B, Diva U, Mercer A, Komers R. Implementing the kidney health initiative surrogate efficacy endpoint in patients with IgA nephropathy (the PROTECT Trial). *Kidney Int Rep* 2019; **4**: 1633–37.
- 129 Zipfel PF, Wiech T, Rudnick R, Afonso S, Person F, Skerka C. Complement inhibitors in clinical trials for glomerular diseases. *Front Immunol* 2019; **10**: 2166.
- 130 Beck LH Jr, Bonegio RGB, Lambeau G, et al. M-type phospholipase A2 receptor as target antigen in idiopathic membranous nephropathy. *N Engl J Med* 2009; **361**: 11–21.
- 131 Beck LH Jr, Fervenza FC, Beck DM, et al. Rituximab-induced depletion of anti-PLA2R autoantibodies predicts response in membranous nephropathy. *J Am Soc Nephrol* 2011; **22**: 1543–50.
- 132 Hofstra JM, Beck LH Jr, Beck DM, Wetzels JF, Salant DJ. Anti-phospholipase A<sub>2</sub> receptor antibodies correlate with clinical status in idiopathic membranous nephropathy. *Clin J Am Soc Nephrol* 2011; **6**: 1286–91.
- 133 Polanco N, Gutiérrez E, Covarsí A, et al. Spontaneous remission of nephrotic syndrome in idiopathic membranous nephropathy. *J Am Soc Nephrol* 2010; **21**: 697–704.
- 134 van den Brand JAJG, Ruggenti P, Chianca A, et al. Safety of rituximab compared with steroids and cyclophosphamide for idiopathic membranous nephropathy. *J Am Soc Nephrol* 2017; **28**: 2729–37.
- 135 Praga M, Barrio V, Juárez GF, Luño J. Tacrolimus monotherapy in membranous nephropathy: a randomized controlled trial. *Kidney Int* 2007; **71**: 924–30.
- 136 Catran DC, Appel GB, Hebert LA, et al. Cyclosporine in patients with steroid-resistant membranous nephropathy: a randomized trial. *Kidney Int* 2001; **59**: 1484–90.
- 137 Fervenza FC, Appel GB, Barbour SJ, et al. Rituximab or cyclosporine in the treatment of membranous nephropathy. *N Engl J Med* 2019; **381**: 36–46.
- 138 Trivin-Avillach C, Beck LH Jr. Management of membranous nephropathy after MENTOR. *Clin J Am Soc Nephrol* 2020; **15**: 415–17.
- 139 Levey AS, Gansevoort RT, Coresh J, et al. Change in albuminuria and GFR as end points for clinical trials in early stages of CKD: a scientific workshop sponsored by the National Kidney Foundation in collaboration with the US Food and Drug Administration and European Medicines Agency. *Am J Kidney Dis* 2020; **75**: 84–104.
- 140 Torres VE, Chapman AB, Devuyst O, et al. Tolvaptan in patients with autosomal dominant polycystic kidney disease. *N Engl J Med* 2012; **367**: 2407–18.
- 141 Torres VE, Chapman AB, Devuyst O, et al. Tolvaptan in later-stage autosomal dominant polycystic kidney disease. *N Engl J Med* 2017; **377**: 1930–42.
- 142 Thomas B, Matsushita K, Abate KH, et al. Global cardiovascular and renal outcomes of reduced GFR. *J Am Soc Nephrol* 2017; **28**: 2167–79.
- 143 Shlipak MG, Matsushita K, Ärnlöv J, et al. Cystatin C versus creatinine in determining risk based on kidney function. *N Engl J Med* 2013; **369**: 932–43.
- 144 Gansevoort RT, Correa-Rotter R, Hemmelgarn BR, et al. Chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. *Lancet* 2013; **382**: 339–52.
- 145 Watson CL, Gold MR. Effect of intracellular and extracellular acidosis on sodium current in ventricular myocytes. *Am J Physiol* 1995; **268**: H1749–56.
- 146 Silswal N, Touchberry CD, Daniel DR, et al. FGF23 directly impairs endothelium-dependent vasorelaxation by increasing superoxide levels and reducing nitric oxide bioavailability. *Am J Physiol Endocrinol Metab* 2014; **307**: E426–36.



- 147 Yilmaz MI, Sonmez A, Saglam M, et al. FGF-23 and vascular dysfunction in patients with stage 3 and 4 chronic kidney disease. *Kidney Int* 2010; **78**: 679–85.
- 148 Lv J, Chen J, Wang M, Yan F. Klotho alleviates indoxyl sulfate-induced heart failure and kidney damage by promoting M2 macrophage polarization. *Aging (Albany NY)* 2020; **12**: 9139–50.
- 149 Cozzolino M, Dusso AS, Slatopolsky E. Role of calcium-phosphate product and bone-associated proteins on vascular calcification in renal failure. *J Am Soc Nephrol* 2001; **12**: 2511–16.
- 150 Rubinger D, Friedlaender MM, Silver J, Kopolovic Y, Czaczkes WJ, Popovtzer MM. Progressive vascular calcification with necrosis of extremities in hemodialysis patients: a possible role of iron overload. *Am J Kidney Dis* 1986; **7**: 125–29.
- 151 Lioufas N, Toussaint ND, Pedagogos E, et al. Can we IMPROVE cardiovascular outcomes through phosphate lowering in CKD? Rationale and protocol for the Impact of Phosphate Reduction On Vascular End-points in Chronic Kidney Disease (IMPROVE-CKD) study. *BMJ Open* 2019; **9**: e024382.
- 152 Sarnak MJ, Amann K, Bangalore S, et al. Chronic kidney disease and coronary artery disease: JACC state-of-the-art review. *J Am Coll Cardiol* 2019; **74**: 1823–38.
- 153 Bangalore S, Maron DJ, O'Brien SM, et al. Management of coronary disease in patients with advanced kidney disease. *N Engl J Med* 2020; **382**: 1608–18.
- 154 Wanner C, Amann K, Shoji T. The heart and vascular system in dialysis. *Lancet* 2016; **388**: 276–84.
- 155 Konstantinidis I, Nadkarni GN, Yacoub R, et al. Representation of patients with kidney disease in trials of cardiovascular interventions: an updated systematic review. *JAMA Intern Med* 2016; **176**: 121–24.
- 156 O'Lone E, Viececchi AK, Craig JC, et al. Cardiovascular outcomes reported in hemodialysis trials. *J Am Coll Cardiol* 2018; **71**: 2802–10.
- 157 Arnott C, Li Q, Kang A, et al. Sodium-glucose cotransporter 2 inhibition for the prevention of cardiovascular events in patients with type 2 diabetes mellitus: a systematic review and meta-analysis. *J Am Heart Assoc* 2020; **9**: e014908.
- 158 Robinson-Cohen C, Littman AJ, Duncan GE, et al. Physical activity and change in estimated GFR among persons with CKD. *J Am Soc Nephrol* 2014; **25**: 399–406.
- 159 MacKinnon HJ, Wilkinson TJ, Clarke AL, et al. The association of physical function and physical activity with all-cause mortality and adverse clinical outcomes in nondialysis chronic kidney disease: a systematic review. *Ther Adv Chronic Dis* 2018; **9**: 209–26.
- 160 Afsar B, Siroopol D, Aslan G, et al. The impact of exercise on physical function, cardiovascular outcomes and quality of life in chronic kidney disease patients: a systematic review. *Int Urol Nephrol* 2018; **50**: 885–904.
- 161 Zhang L, Wang Y, Xiong L, Luo Y, Huang Z, Yi B. Exercise therapy improves eGFR, and reduces blood pressure and BMI in non-dialysis CKD patients: evidence from a meta-analysis. *BMC Nephrol* 2019; **20**: 398.
- 162 Mallamaci F, Pisano A, Tripepi G. Physical activity in chronic kidney disease and the EXerCise Introduction To Enhance trial. *Nephrol Dial Transplant* 2020; **35** (suppl 2): ii18–22.
- 163 Xia J, Wang L, Ma Z, et al. Cigarette smoking and chronic kidney disease in the general population: a systematic review and meta-analysis of prospective cohort studies. *Nephrol Dial Transplant* 2017; **32**: 475–87.
- 164 Staplin N, Haynes R, Herrington WG, et al. Smoking and adverse outcomes in patients with CKD: the Study of Heart and Renal Protection (SHARP). *Am J Kidney Dis* 2016; **68**: 371–80.
- 165 Saran R, Padilla RL, Gillespie BW, et al. A randomized crossover trial of dietary sodium restriction in stage 3–4 CKD. *Clin J Am Soc Nephrol* 2017; **12**: 399–407.
- 166 Jardine MJ, Li N, Ninomiya T, et al. Dietary sodium reduction reduces albuminuria: a cluster randomized trial. *J Ren Nutr* 2019; **29**: 276–84.
- 167 McMahon EJ, Bauer JD, Hawley CM, et al. A randomized trial of dietary sodium restriction in CKD. *J Am Soc Nephrol* 2013; **24**: 2096–103.
- 168 Luzardo L, Noboa O, Boggia J. Mechanisms of salt-sensitive hypertension. *Curr Hypertens Rev* 2015; **11**: 14–21.
- 169 Koeth RA, Wang Z, Levison BS, et al. Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nat Med* 2013; **19**: 576–85.
- 170 Navaneethan SD, Yehner H, Moustarah F, Schreiber MJ, Schauer PR, Beddhu S. Weight loss interventions in chronic kidney disease: a systematic review and meta-analysis. *Clin J Am Soc Nephrol* 2009; **4**: 1565–74.
- 171 Lambert K, Beer J, Dumont R, et al. Weight management strategies for those with chronic kidney disease: a consensus report from the Asia Pacific Society of Nephrology and Australia and New Zealand Society of Nephrology 2016 renal dietitians meeting. *Nephrology (Carlton)* 2018; **23**: 912–20.
- 172 Herrington WG, Emberson J, Mihaylova B, et al. Impact of renal function on the effects of LDL cholesterol lowering with statin-based regimens: a meta-analysis of individual participant data from 28 randomised trials. *Lancet Diabetes Endocrinol* 2016; **4**: 829–39.
- 173 Rydén L, Grant PJ, Anker SD, et al. ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD: the Task Force on diabetes, pre-diabetes, and cardiovascular diseases of the European Society of Cardiology (ESC) and developed in collaboration with the European Association for the Study of Diabetes (EASD). *Eur Heart J* 2013; **34**: 3035–87.
- 174 Davies MJ, D'Alessio DA, Fradkin J, et al. Management of hyperglycaemia in type 2 diabetes, 2018. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia* 2018; **61**: 2461–98.
- 175 Palmer SC, Di Micco L, Razavian M, et al. Effects of antiplatelet therapy on mortality and cardiovascular and bleeding outcomes in persons with chronic kidney disease: a systematic review and meta-analysis. *Ann Intern Med* 2012; **156**: 445–59.
- 176 Ha JT, Neuen BL, Cheng LP, et al. Benefits and harms of oral anticoagulant therapy in chronic kidney disease: a systematic review and meta-analysis. *Ann Intern Med* 2019; **171**: 181–89.
- 177 Ray KK, Nicholls SJ, Buhr KA, et al. Effect of apabetalone added to standard therapy on major adverse cardiovascular events in patients with recent acute coronary syndrome and type 2 diabetes: a randomized clinical trial. *JAMA* 2020; **323**: 1565–73.
- 178 Kalantar-Zadeh K, Schwartz GG, Nicholls SJ, et al. Effect of apabetalone on major adverse cardiovascular events in patients with chronic kidney disease, type-2 diabetes mellitus and recent acute coronary syndrome: results from the BETonMACE randomized controlled trial. *Clin J Am Soc Nephrol* 2021; **16**: 705–16.
- 179 Ware JE Jr, Richardson MM, Meyer KB, Gandek B. Improving CKD-specific patient-reported measures of health-related quality of life. *J Am Soc Nephrol* 2019; **30**: 664–77.
- 180 Kovedy CP, Naseer A, Sumida K, et al. Abrupt decline in kidney function precipitating initiation of chronic renal replacement therapy. *Kidney Int Rep* 2017; **3**: 602–09.
- 181 Coca SG, Singanamala S, Parikh CR. Chronic kidney disease after acute kidney injury: a systematic review and meta-analysis. *Kidney Int* 2012; **81**: 442–48.
- 182 Rifkin DE, Coca SG, Kalantar-Zadeh K. Does AKI truly lead to CKD? *J Am Soc Nephrol* 2012; **23**: 979–84.
- 183 Lapi F, Azoulay L, Yin H, Nessim SJ, Suissa S. Concurrent use of diuretics, angiotensin converting enzyme inhibitors, and angiotensin receptor blockers with non-steroidal anti-inflammatory drugs and risk of acute kidney injury: nested case-control study. *BMJ* 2013; **346**: e8525.
- 184 Gaipov A, Molnar MZ, Potukuchi PK, et al. Acute kidney injury following coronary revascularization procedures in patients with advanced CKD. *Nephrol Dial Transplant* 2019; **34**: 1894–901.
- 185 Grams ME, Sang Y, Coresh J, et al. Acute kidney injury after major surgery: a retrospective analysis of veterans health administration data. *Am J Kidney Dis* 2016; **67**: 872–80.
- 186 Wyld M, Morton RL, Hayen A, Howard K, Webster AC. A systematic review and meta-analysis of utility-based quality of life in chronic kidney disease treatments. *PLoS Med* 2012; **9**: e1001307.
- 187 Murtagh FEM, Marsh JE, Donohoe P, Ekbal NJ, Sheerin NS, Harris FE. Dialysis or not? A comparative survival study of patients over 75 years with chronic kidney disease stage 5. *Nephrol Dial Transplant* 2007; **22**: 1955–62.
- 188 Saeed F, Ladwig SA, Epstein RM, Monk RD, Duberstein PR. Dialysis regret: prevalence and correlates. *Clin J Am Soc Nephrol* 2020; **15**: 957–63.

- 189 Murtagh FEM, Burns A, Moranne O, Morton RL, Naicker S. Supportive care: comprehensive conservative care in end-stage kidney disease. *Clin J Am Soc Nephrol* 2016; **11**: 1909–14.
- 190 Siriwardana AN, Hoffman AT, Brennan FP, Li K, Brown MA. Impact of renal supportive care on symptom burden in dialysis patients: a prospective observational cohort study. *J Pain Symptom Manage* 2020; **60**: 725–36.
- 191 Crail S, Walker R, Brown M. Renal supportive and palliative care: position statement. *Nephrology (Carlton)* 2013; **18**: 393–400.
- 192 Cheigh JS, Kim H, Stenzel KH, et al. Systemic lupus erythematosus in patients with end-stage renal disease: long-term follow-up on the prognosis of patients and the evolution of lupus activity. *Am J Kidney Dis* 1990; **16**: 189–95.
- 193 Saadi G, Kalantar-Zadeh K, Almasio P, et al. Hepatitis C virus infection and global kidney health: the consensus proceedings of the International Federation of Kidney Foundations. *Afr J Nephrol* 2020; **23**: 159–68.
- 194 Ladino M, Pedraza F, Roth D, Hepatitis C virus infection in chronic kidney disease. *J Am Soc Nephrol* 2016; **27**: 2238–46.
- 195 Molnar MZ, Alhourani HM, Wall BM, et al. Association of hepatitis C viral infection with incidence and progression of chronic kidney disease in a large cohort of US veterans. *Hepatology* 2015; **61**: 1495–502.
- 196 Roth D, Nelson DR, Bruchfeld A, et al. Grazoprevir plus elbasvir in treatment-naïve and treatment-experienced patients with hepatitis C virus genotype 1 infection and stage 4–5 chronic kidney disease (the C-SURFER study): a combination phase 3 study. *Lancet* 2015; **386**: 1537–45.
- 197 Su G, Iwagami M, Qin X, et al. Kidney disease and mortality in patients with respiratory tract infections: a systematic review and meta-analysis. *Clin Kidney J* 2020; **14**: 602–11.
- 198 McDonald HI, Thomas SL, Millett ERC, Quint J, Nitsch D. Do influenza and pneumococcal vaccines prevent community-acquired respiratory infections among older people with diabetes and does this vary by chronic kidney disease? A cohort study using electronic health records. *BMJ Open Diabetes Res Care* 2017; **5**: e000332.
- 199 Corbett RW, Blakey S, Nitsch D, et al. Epidemiology of COVID-19 in an urban dialysis center. *J Am Soc Nephrol* 2020; **31**: 1815–23.
- 200 Kalantar-Zadeh K, Moore LW. Renal telenutrition for kidney health: leveraging telehealth and telemedicine for nutritional assessment and dietary management of patients with kidney disorders. *J Ren Nutr* 2020; **30**: 471–74.
- 201 Sise ME, Baggett MV, Shepard JO, Stevens JS, Rhee EP. Case 17-2020: a 68-year-old man with Covid-19 and acute kidney injury. *N Engl J Med* 2020; **382**: 2147–56.
- 202 Battle D, Soler MJ, Sparks MA, et al. Acute kidney injury in COVID-19: emerging evidence of a distinct pathophysiology. *J Am Soc Nephrol* 2020; **31**: 1380–83.
- 203 Zamai L. The yin and yang of ACE/ACE2 pathways: the rationale for the use of renin-angiotensin system inhibitors in COVID-19 patients. *Cells* 2020; **9**: E1704.
- 204 Stanifer JW, Muiru A, Jafar TH, Patel UD. Chronic kidney disease in low- and middle-income countries. *Nephrol Dial Transplant* 2016; **31**: 868–74.
- 205 Chang TI, Lim H, Park CH, et al. Association between income disparities and risk of chronic kidney disease: a nationwide cohort study of seven million adults in Korea. *Mayo Clin Proc* 2020; **95**: 231–42.
- 206 Moosa MR, Norris KC. Sustainable social development: tackling poverty to achieve kidney health equity. *Nat Rev Nephrol* 2021; **17**: 3–4.
- 207 Ene-Iordache B, Perico N, Bikbov B, et al. Chronic kidney disease and cardiovascular risk in six regions of the world (ISN-KDDC): a cross-sectional study. *Lancet Glob Health* 2016; **4**: e307–19.
- 208 Liyanage T, Ninomiya T, Jha V, et al. Worldwide access to treatment for end-stage kidney disease: a systematic review. *Lancet* 2015; **385**: 1975–82.
- 209 Engelgau MM, Karan A, Mahal A. The economic impact of non-communicable diseases on households in India. *Global Health* 2012; **8**: 9.
- 210 Kovesdy CP, Norris KC, Boulware LE, et al. Association of race with mortality and cardiovascular events in a large cohort of US veterans. *Circulation* 2015; **132**: 1538–48.
- 211 Derose SF, Rutkowski MP, Crooks PW, et al. Racial differences in estimated GFR decline, ESRD, and mortality in an integrated health system. *Am J Kidney Dis* 2013; **62**: 236–44.
- 212 Norris KC, Williams SF, Rhee CM, et al. Hemodialysis disparities in African Americans: the deeply integrated concept of race in the social fabric of our society. *Semin Dial* 2017; **30**: 213–23.
- 213 Ahmed S, Nutt CT, Eneanya ND, et al. Examining the potential impact of race multiplier utilization in estimated glomerular filtration rate calculation on African-American care outcomes. *J Gen Intern Med* 2021; **36**: 464–71.
- 214 Norris KC, Eneanya ND, Boulware LE. Removal of race from estimates of kidney function: first, do no harm. *JAMA* 2021; **325**: 135–37.
- 215 Kazory A. Fluid overload as a major target in management of cardiorenal syndrome: Implications for the practice of peritoneal dialysis. *World J Nephrol* 2017; **6**: 168–75.
- 216 Kajbafzadeh AM, Sabetkish N, Sabetkish S. Establishment of colonic dialysis model in uremic rats by right nephrectomy and left partial nephrectomy. *J Pediatr Urol* 2018; **14**: 159.e1–8.
- 217 Dai S, Dai Y, Peng J, Xie X, Ning J. Simplified colonic dialysis with hemodialysis solutions delays the progression of chronic kidney disease. *QJM* 2019; **112**: 189–96.
- 218 Keller RW Jr, Kopple JD, Kalantar-Zadeh K. Perspiration interventions for conservative management of kidney disease and uremia. *Curr Opin Nephrol Hypertens* 2020; **29**: 57–63.
- 219 Hanafusa N, Lodebo BT, Shah A, Kopple JD. Is there a role for diaphoresis therapy for advanced chronic kidney disease patients? *J Ren Nutr* 2017; **27**: 295–302.
- 220 Keller RW, Bailey JL, Wang Y, Klein JD, Sands JM. Urea transporters and sweat response to uremia. *Physiol Rep* 2016; **4**: e12825.

© 2021 Elsevier Ltd. All rights reserved.