

UC Davis

UC Davis Previously Published Works

Title

Genotypic and tissue-specific variation of Populus nigra transcriptome profiles in response to drought

Permalink

<https://escholarship.org/uc/item/31g444fd>

Journal

Scientific Data, 9(1)

ISSN

2052-4463

Authors

Eckert, Christian
Wildhagen, Henning
Paulo, Maria João
et al.

Publication Date

2022

DOI

10.1038/s41597-022-01417-z

Peer reviewed



OPEN

DATA DESCRIPTOR

Genotypic and tissue-specific variation of *Populus nigra* transcriptome profiles in response to drought

Christian Eckert^{1,8}, Henning Wildhagen^{2,8}, Maria João Paulo³, Simone Scalabrin⁴, Johannes Ballauff¹, Sabine K. Schnabel³, Vera Vendramin⁴, Joost J. B. Keurentjes⁵, Marie-Béatrice Bogeat-Triboulot⁶, Gail Taylor⁷ & Andrea Polle¹

Climate change is one of the most important challenges for mankind in the far and near future. In this regard, sustainable production of woody crops on marginal land with low water availability is a major challenge to tackle. This dataset is part of an experiment, in which we exposed three genetically differentiated genotypes of *Populus nigra* originating from contrasting natural habitats to gradually increasing moderate drought. RNA sequencing was performed on fine roots, developing xylem and leaves of those three genotypes under control and moderate drought conditions in order to get a comprehensive dataset on the transcriptional changes at the whole plant level under water limiting conditions. This dataset has already provided insight in the transcriptional control of saccharification potential of the three *Populus* genotypes under drought conditions and we suggest that our data will be valuable for further in-depth analysis regarding candidate gene identification or, on a bigger scale, for meta-transcriptome analysis.

Background & Summary

Ongoing climate change, entailing more frequent and severe drought events, put biomass productivity at an increasing risk^{1–3}, especially on marginal land. Considering the increasing demand for both, food production and feedstock, energy crop production systems should preferentially utilize perennial crops grown on marginal sites⁴. Consequently, there is a need to develop new germplasms of perennial biomass crops characterized by high productivity and the ability to maintain productivity under water limited growth conditions⁵. To this end, it is pivotal to understand the underlying physiological mechanisms and molecular drivers of drought stress responses and growth adjustment of trees, especially mechanisms underlying adjustments of growth in response to water deprivation.

Because of its adaptation to a broad range of habitats, its fast growth and available molecular and genomic resources, the genus *Populus* is considered as a model system for woody biomass plants⁶. The publication of the full genome sequence of *Populus trichocarpa*⁷ fostered the application of systems biology approaches on a multitude of *Populus* species, including most widespread species like *P. tremuloides* and its European counterpart *P. tremula*. In Europe, *P. nigra*, European black poplar, is also widely distributed, and this is reflected in significant phenotypic variation in growth rates, tree architecture and leaf size⁸. *P. nigra* is a candidate for second generation biofuels that use lignocellulosic biomass^{9–11} as well as a raw material for pulp and paper production¹². As a pioneer species, *P. nigra* can also be used to preserve local biodiversity, as it is able to outcompete exotic poplar species. Therefore, it is used in restoration and protection of riparian forest sites¹³. In Eastern Europe, *P. nigra* is

¹Forest Botany and Tree Physiology, University of Goettingen, Büsgenweg 2, Göttingen, Germany. ²HAWK University of Applied Sciences and Arts, Faculty of Resource Management, Büsgenweg 1a, 37077, Göttingen, Germany.

³Biometris, Wageningen UR Wageningen Plant Research, Droevendaalsesteeg 1, Wageningen, The Netherlands.

⁴IGA Technology Services, via Jacopo Linussio 51, Udine, Italy. ⁵Laboratory of Genetics, Wageningen University & Research, Droevendaalsesteeg 1, Wageningen, The Netherlands. ⁶Université de Lorraine, AgroParisTech, INRAE, UMR Silva, Nancy, France. ⁷Department of Plant Sciences, University of California, One Shields Ave, Davis, CA, USA.

⁸These authors contributed equally: Christian Eckert, Henning Wildhagen. ✉e-mail: henning.wildhagen@hawk.de

planted for soil protection and used for restoration areas polluted by industrial usage¹⁴. As a species perceived as both, economically and ecologically important, *P. nigra* is targeted in studies aiming to elucidate the genetic basis of variation in adaptive traits^{8,15,16}.

In order to characterize traits that play a conserved role across populations adapted to contrasting natural habitats we included three *P. nigra* genotypes originating from areas with different water availability. The genotypes were previously shown to be genetically differentiated⁸. This genetic variance between the three subgroups originates from the last glacial period, where there were three refugia for *P. nigra*⁸ in Europe from which the species recolonized Europe. Among the three genotypes, the Spanish genotype grows on the driest land, while the Italian genotype derives from the most humid area^{8,17}.

Here, trees of each genotype were exposed to a gradually increasing, precisely controlled, moderate drought for five weeks before harvest. Transcriptome profiles of the developing xylem, fine roots, and fully-expanded young leaves were prepared in order to get a holistic insight into the transcriptional reprogramming in specific tissues and between different genotypes. Of these tissues roots sense drying soil first^{18,19}; the xylem is the water conducting system in plants²⁰; and leaves play the key role in transpiration and gas exchange²¹. Thus, this study covers all pivotal tissues involved in the water balance of plants.

Our data set opens the possibility for analyses in various directions. One could either look at conserved gene clusters across all three genotypes to identify gene families that are important for drought acclimation or other important traits. An example for this kind is the investigation of the transcriptional drought response in the developing xylem²², which constitutes a subset of the present data. Weighted gene correlation network analysis identified genes correlated significantly with the saccharification potential of the wood, which was enhanced under drought²². Interestingly, no genes involved in lignin biosynthesis were found to be correlated with drought, but polysaccharide biosynthesis genes were upregulated, underpinning the improved saccharification potential²². This study gave a first glimpse in the power of the data set generated in this study. In-depth analyses of the whole data set regarding the acclimation of specific tissues to drier conditions and the crosstalk between tissues has not been done yet. In addition, investigations of the intraspecific genetic variation can assist identifying genetic markers that can be used for predictive breeding. Previous studies have mostly concentrated on intraspecific variation in single tissues^{23,24}. Our dataset combines data on different tissues and genotypes. We therefore believe that this data set will be of keen interest for the scientific community and can add a piece to the puzzle to understand drought adaptation of trees.

Methods

Plant material. Three genotypes of *Populus nigra* L., originating from natural populations in France, Italy and Spain were studied. The three genotypes represent clones of individual trees sampled from the populations Drôme 6 (FR-6), La Zelata (IT1), and Ebro 2 (SP-2)⁸. Cuttings were planted in 10-liter plastic pots filled with a 1:1 (v/v) mixture of peat and sand fertilized with a slow-release fertilizer (4 g L⁻¹ of Nutricote T100, 13:13:13 NPK and micronutrients; FERTIL S.A.S, Boulogne Billancourt, France). Plants were grown in two chambers of a glasshouse located at Champenoux, France (48°45′09.3″N, 6°20′27.6″E), under natural light conditions. Growth conditions in the greenhouse were affected by weather conditions, but the temperature was adjusted to not exceed 28 °C, and daily maxima of irradiance ranged from 73–478 W m⁻². Plants were watered three times per day to field capacity on a custom-made automated watering system.

Drought experiment. After six weeks of growth, plants of each genotype were randomly assigned to either a control or a drought treatment. The plants were allocated to the two greenhouse chambers in balanced proportions according to genotype and treatment. Plants were exposed to drought by gradually decreasing the available soil water content. The regulation of soil water availability was based on the soil relative extractable water content (REW_{soil}), which is defined as: $REW_{soil} = (SWC - \text{water content at wilting point}) / (\text{water content at field capacity} - \text{water content at wilting point})$, with SWC: soil volumetric water content; water content at wilting point = 3%; water content at field capacity = 32%. Control plants were watered to 85% REW_{soil} three times per day for the whole 5-week period of the experiment. For drought-treated plants, a target level of 20% REW_{soil} was defined, which was reached two weeks after starting to gradually withhold water. Plants were watered at this target level of REW_{soil} for the following three weeks^{22,25} (Fig. 1a).

After five weeks of control or drought treatment, all plants were harvested destructively. We used four individual plants from each genotype and water level (3 genotypes × 4 plants × 2 water levels = 24 plants) for the harvest of 3 tissues (= 72 samples for further analysis). Different tissues per plant were harvested in parallel by several persons to achieve rapid sampling times. Leaf number 10 from the top of each plant was cut off, weighed and flash frozen in liquid nitrogen. The stem was cut at the base, the bark was removed from the lower part and the developing xylem, a soft tissue, was scratched from the surface and immediately frozen in liquid nitrogen. The root system with soil was carefully removed from the pot and briefly washed under tap water to clean fine roots, which were then cut off. The surface water from the collected fine roots was quickly dabbed off by tissue paper and then roots were frozen in liquid nitrogen. The samples were stored at -80 °C. For the extraction of RNA, each tissue was milled to a fine powder keeping the sample frozen by cooling the device under liquid nitrogen. Of each sample 150 mg of frozen tissue was weighed into the extraction medium for RNA. The procedure of RNA sampling to count table is depicted in Fig. 1b.

RNA extraction, library preparation and sequencing. Total RNA was extracted from homogenized samples of a young fully expanded leaf, fine roots, and developing xylem of four biological replicates per genotype and treatment using the CTAB protocol²⁶ with minor modifications described in²⁷. After checking quality and integrity (see Technical Validation), 2 µg of total RNA were used for library preparation using the ‘TruSeq mRNA Sample Prep kit v2’ (Illumina, San Diego, CA, USA), following the manufacturer’s instructions. Libraries were




Fig. 1 Experimental setup and data processing pipeline. (a) Water availability was reduced for two weeks to reach 20% soil extractable water (SEW) and was kept constant for 3 more weeks until harvest. (b) Flow chart of RNA sample processing from harvest to data analysis.

then processed with Illumina cBot for cluster generation on the flowcell, following the manufacturer's instructions and sequenced in 50 bp single-end mode at the 6-fold multiplex on the Illumina HiSeq2000 (Illumina, San Diego, CA, USA).

Data Records

Raw data of RNA-seq analysis are deposited at the NCBI short read archive under SRP numbers SRP095832²⁸ (dev. xylem samples) and SRP101711²⁹ (fine root and leaf samples). Each biological replicate refers to a single SRA Sample Accession (SRS accession, Online-only Table 1). Raw count data are available at figshare data




Fig. 2 Quality assessment of RNA sequencing data. **(a)** Mean quality phred scores of individual samples, **(b)** Per Sequence Quality Scores as counts per phred score **(c)** GC content of individual samples. FastQC results for single samples were summarized using MultiQC.

repository under <https://doi.org/10.6084/m9.figshare.17031842.v3>³⁰. Tables for differentially expressed genes for each tissue and genotype are available under <https://doi.org/10.6084/m9.figshare.19382603.v1>³¹.

Technical Validation

RNA integrity was determined using Agilent 2100 Bioanalyzer RNA Nano assay (Agilent technologies, Santa Clara, CA, USA). Average RIN values were 7.9 ± 0.5 for the developing xylem, 7.2 ± 0.4 for fine roots and 7.2 ± 0.3 for leaf samples. qRT-PCR validation of selected genes has been performed by Wildhagen and colleagues²².




Fig. 3 Initial analysis of RNAseq data from control- or drought-stress samples of fine roots, developing xylem and leaf samples of three genotypes of *P. nigra* originating from Spain, France and Italy. (a–d) Principal component analyses of VST normalized RNA-seq count data (a) PCA of all tissues and ecotypes; (b) PCA of leaf samples; (c) PCA of developing xylem samples; (d) PCA of fine root samples; (e) Venn diagram showing the distribution of DEGs under drought between the three tissues regardless of the genotype. (f–h) Venn diagrams showing the intersections of drought-regulated DEGs between the genotypes for fine roots (f), developing xylem (g) and leaf (h).

Quality assessment. FastQC Version 0.11.9 was applied to the clean reads to assess the quality of the sequence reads³². A summary of FastQC reports was generated using MultiQC³³. Figure 2 shows that the Mean Quality Scores and the Per Sequence Quality Scores of all sequencing results were of high quality indicated by the green color in the diagrams. Per Sequence GC content was consistent at 44–45%.

Data filtering and processing. Raw sequence reads were processed with the Python package Cutadapt v1.4.2³⁴ to remove residual adapter contamination. Reads were subsequently trimmed to remove low-quality reads (option `-trim_qual_left/right = 25`) and reads shorter than 40 nucleotides, using the PRINSEQ software v.lite-0.20.4³⁵. Trimmed reads were aligned to the *Populus trichocarpa* v2.1 transcriptome database⁷ using TopHat2 v2.0.10³⁶. A count table was generated using the Python package HTSeq v0.6.1³⁷. Python codes for all procedures can be found in document FigShare Python Codes³⁸. The raw count table is available under FigShare Raw Count Table³⁰.

Principal component analysis. As a first step of exploratory data analysis, a PCA across all samples of the RNA-seq data was performed. Normalization of counts was performed using the VST (variance stabilizing transformation) function of DESeq2 v1.32.0 package for R^{39,40}. The plot of the first two PCs revealed a strong clustering according to tissue type, suggesting strong variation of transcript abundance among tissues (Fig. 3a). Further PCA analyses of leaf (Fig. 3b), developing xylem (Fig. 3c) and fine roots (Fig. 3d) transcriptomes revealed separate clustering of the different genotypes. This result indicates putatively interesting differential gene expression patterns between the different origins (Fig. 3b–d). PCA analysis has been performed with R⁴⁰ using the packages DESeq2 v1.32.0³⁹. The R code for the PCA is available under Figshare R Code PCA⁴¹.

Initial analysis of differentially expressed genes. Analyses of the count data were done with the R package DESeq2, version 1.34.0³⁹. Normalisation of count tables was done based on the ‘median ratio method’⁴² implemented in the function ‘estimateSizeFactors’. We applied an unspecific filtering⁴³ to keep only those genes to which at least one read per million reads of library size aligned in at least four samples.

The analysis of differential expression was carried out by fitting two-factorial negative binomial generalized linear models (function ‘DESeq’) to the count data. To assess the significance of a ‘treatment’ main effect, a full model with ‘treatment’ and ‘genotype’ main effects was set up, against which a reduced model without ‘treatment’ main effect was tested with the function ‘nbinomLRT’. The sets of genes showing a significant drought main effect were used for a genotype specific analysis of drought effects. For this purpose the data set was split according to tissue and genotype, and for each combination of tissue and genotype, a full model with ‘treatment’ main effect was set up, against which a reduced model with intercept only was tested with the function ‘nbinomLRT’. The R code for the DEG analysis is available under Figshare R Code DEG analysis⁴⁴. Venn diagrams have been made using an online Venn diagram tool (<https://bioinformatics.psb.ugent.be/webtools/Venn/>).

Analysis of the drought main effect, i.e. without differentiation of the genotypes revealed that the majority of differentially expressed genes (DEGs) were tissue specific under drought (92.8% in fine roots, 62% in the developing xylem and 86.3% in the leaf, Fig. 3e). Interestingly, no shared gene was found to be differentially regulated in all three tissues. When these drought related DEGs shown in Fig. 3e were further analyzed on genotype level, it was found that the majority of drought-related DEGs in fine roots (Fig. 3f) and leaves (Fig. 3h) were genotype specific. However, in the developing xylem (Fig. 3g) there were no genotype specific DEGs found for the Spanish and French genotype. This initial analysis shows, that this dataset can be used to identify conserved drought acclimation processes that are shared by all three genotypes, as well as genotype specific drought responses of *P. nigra*.

Code availability

Figshare R Code DEG analysis: <https://doi.org/10.6084/m9.figshare.19382594.v144>.

Figshare Python Codes: <https://doi.org/10.6084/m9.figshare.17031869.v238>.

Figshare R Code for PCA analysis: <https://doi.org/10.6084/m9.figshare.17031884.v241>.

Received: 23 November 2021; Accepted: 23 May 2022;

Published online: 14 June 2022

References

- Allen, C. D. *et al.* A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* **259**, 660–684 (2010).
- Schaller, N. *et al.* Ensemble of european regional climate simulations for the winter of 2013 and 2014 from HadAM3P-RM3P. *Scientific Data* (2018).
- Klein, T. & Hartmann, H. Climate change drives tree mortality. *Science (New York, N.Y.)* **362** (2018).
- Karp, A. & Richter, G. M. Meeting the challenge of food and energy security. *J Exp Bot* **62**, 3263–3271 (2011).
- Polle, A., Chen, S. L., Eckert, C. & Harfouche, A. Engineering drought resistance in forest trees. *Front. Plant Sci.* (2019).
- Allwright, M. R. & Taylor, G. Molecular breeding for improved second generation bioenergy crops. *Trends in Plant Science* **21**, 43–54 (2016).
- Tuskan, G. A. *et al.* The genome of black cottonwood, *Populus trichocarpa* (Torr. & Gray). *Science* **313**, 1596–1604 (2006).
- DeWoody, J., Trewin, H. & Taylor, G. Genetic and morphological differentiation in *Populus nigra* L.: Isolation by colonization or isolation by adaptation? *Molecular Ecology* **24**, 2641–2655 (2015).
- Allwright, M. R. *et al.* Biomass traits and candidate genes for bioenergy revealed through association genetics in coppiced European *Populus nigra* (L.). *Biotechnology for Biofuels* **9**, 195 (2016).
- Gupta, A. *et al.* Bioethanol production from hemicellulose rich *Populus nigra* involving recombinant hemicellulases from *Clostridium thermocellum*. *Bioresour Technol* **165**, 205–213 (2014).
- Guerra, F. P. *et al.* Association genetics of chemical wood properties in black poplar (*Populus nigra*). *New Phytologist* **197**, 162–176 (2013).
- Wu, Q., Chen, H. L., Wang, B. B. & Cao, B. B. Analysis on fast-growing black poplar branch used as raw materials for APMP pulping and papermaking. *Applied Mechanics and Materials* **448–453**, 972–977 (2014).
- Vanden Broeck, A. *et al.* Reintroduced native *Populus nigra* in restored floodplain reduces spread of exotic poplar species. *Front. Plant Sci.* (2021).
- Vanden Broeck, A. EUFORGEN Technical Guidelines for genetic conservation and use for European black poplar (*Populus nigra*). (2003).

15. Rohde, A., Bastien, C. & Boerjan, W. Temperature signals contribute to the timing of photoperiodic growth cessation and bud set in poplar. *Tree Physiology* **31**, 472–482 (2011).
16. Fabbri, F. *et al.* Phenotypic plasticity, QTL mapping and genomic characterization of bud set in black poplar. *BMC Plant Biol* **12**, 1–16 (2012).
17. Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. & Jarvis, A. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**, 1965–1978 (2005).
18. Hamanishi, E. T. & Campbell, M. M. Genome-wide responses to drought in forest trees. *Forestry (Lond)* **84**, 273–283 (2011).
19. Brunner, I., Herzog, C., Dawes, M. A., Arend, M. & Sperisen, C. How tree roots respond to drought. *Front. Plant Sci.* **6** (2015).
20. Myburg, A. A., Lev-Yadun, S. & Sederoff, R. R. Xylem structure and function. In *Encyclopedia of Life Science* (American Cancer Society, 2013).
21. Stålfelt, M. G. The stomata as a hydrophotic regulator of the water deficit of the plant. *Physiologia Plantarum* **8**, 572–593 (1955).
22. Wildhagen, H. *et al.* Genes and gene clusters related to genotype and drought-induced variation in saccharification potential, lignin content and wood anatomical traits in *Populus nigra*. *Tree Physiol* **38**, 320–339 (2018).
23. Lucani, C. J., Brodrick, T. J., Jordan, G. & Mitchell, P. J. Intraspecific variation in drought susceptibility in *Eucalyptus globulus* is linked to differences in leaf vulnerability. *Functional Plant Biol.* **46**, 286–293 (2019).
24. Zhang, L., Liu, B., Zhang, J. & Hu, J. Insights of molecular mechanism of xylem development in five black poplar cultivars. *Front. Plant Sci.* **11** (2020).
25. Bogeat-Triboulot, M. B. *et al.* Additive effects of high growth rate and low transpiration rate drive differences in whole plant transpiration efficiency among black poplar genotypes. *Environmental and Experimental Botany* (2019).
26. Chang, S., Puryear, J. & Cairney, J. A simple and efficient method for isolating RNA from pine trees. *Plant Molecular Biology Reporter* **11**, 113–116 (1993).
27. Janz, D. *et al.* Salt stress induces the formation of a novel type of ‘pressure wood’ in two *Populus* species. *New Phytologist* **194**, 129–141 (2012).
28. NCBI Sequence Read Archive <https://identifiers.org/ncbi/insdc.sra:SRP095832> (2017).
29. NCBI Sequence Read Archive <https://identifiers.org/ncbi/insdc.sra:SRP101711> (2018).
30. Eckert, C., Wildhagen, H. & Polle, A. Raw Count Table. *figshare* <https://doi.org/10.6084/m9.figshare.17031842.v3> (2021).
31. Eckert, C., Wildhagen, H. & Polle, A. DEG tables. *figshare* <https://doi.org/10.6084/m9.figshare.19382603.v1> (2022).
32. Andrews, S. FastQC: A quality control tool for high throughput sequence data. (2010).
33. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).
34. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet journal* **17**, 10–12 (2011).
35. Schmieder, R. & Edwards, R. Quality control and preprocessing of metagenomic datasets. *Bioinformatics* **27**, 863–864 (2011).
36. Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biology* **14**, R36 (2013).
37. Anders, S., Pyl, P. T. & Huber, W. HTSeq - a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).
38. Eckert, C., Schnabel, S. K., Paulo, M. J., Wildhagen, H. & Polle, A. Fig Share Python Codes. *figshare* <https://doi.org/10.6084/m9.figshare.17031869.v2> (2022).
39. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, (2014).
40. R Core Team. R: A language and environment for statistical computing. R foundation for statistical computing. (<https://www.R-project.org/>, 2018).
41. Eckert, C., Ballauff, J. & Polle, A. R Code for PCA Analysis. *figshare* <https://doi.org/10.6084/m9.figshare.17031884.v2> (2022).
42. Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome Biol* **11**, 1–12 (2010).
43. Bourgon, R., Gentleman, R. & Huber, W. Independent filtering increases detection power for high-throughput experiments. *PNAS* **107**, 9546–9551 (2010).
44. Eckert, C., Wildhagen, H. & Polle, A. Figshare R Code DEG analysis. *figshare* <https://doi.org/10.6084/m9.figshare.19382594.v1> (2022).

Acknowledgements

We thank T. Klein (Laboratory for Radio-Isotopes, Göttingen, Germany) for RNA isolation. We acknowledge the providers of the original *P. nigra* genotypes ‘France 6J-29’ (INRA, Paris, France represented by G. Pilate) and ‘Spain RIN2-new’ (CITA, Zaragoza, Spain, represented by JV Lacasa Azlor) and C. Bastien (INRA, Orleans, France) for providing the stock cuttings. This work was conducted in the frame of the WATBIO project (Development of improved perennial biomass crops for water-stressed environments), which is a collaborative research project funded from the European Union’s Seventh Programme for research, technological development and demonstration under grant agreement No. 311929. UMR Silva was supported by the French National Research Agency through the Laboratory of Excellence ARBRE (ANR-12- LABXARBRE-01). We acknowledge support by HAWK for funding the article-processing charge.

Author contributions

C.E.: Manuscript writing, data analysis, revision of the manuscript. J.B.: Data analysis. A.P.: Revision of the Manuscript, experiment planning, study supervision. H.W.: Designed and performed the greenhouse experiment and contributed to data analysis, writing and revision of the manuscript. M.B.B.T.: supervised and performed the greenhouse experiment, revision of the manuscript. S.K.S.: data preparation & analysis, revision of the manuscript. J.K.: data preparation & analysis, revision of the manuscript. M.J.P.: data preparation & analysis. G.T.: experiment planning, revision of the manuscript. S.S.: RNA sequencing & revision of the manuscript. V.V.: RNA sequencing.

Funding

Open Access funding enabled and organized by Projekt DEAL.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to H.W.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022