Lawrence Berkeley National Laboratory

Recent Work

Title

SPECIFIC SEQUESTERING AGENTS FOR THE ACTINIDES. 7. SYNTHESIS AND STRUCTURAL CHEMISTRY OF TETRAKIS(THIOHYDROXAMATO)HAFNIUM(IV) IN $\text{Hf}(\text{CH}_3\text{C}_6\text{H}_4(\text{S})\text{N}(\text{O})\text{CH}_3)_4 \cdot \text{C}_2\text{H}_5\text{OH}$.

Permalink

<https://escholarship.org/uc/item/31h2q1x3>

Authors

Abu-Dari, Kamal
Raymond, K.N.

Publication Date

1981-07-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Molecular Research Division

RECEIVED
LAWRENCE
BERKELEY
LABORATORY

SEP 10 1981

LIBRARY AND
DOCUMENTS SECTION

Submitted as a Note to Inorganic Chemistry

SPECIFIC SEQUESTERING AGENTS FOR THE ACTINIDES.

7. SYNTHESIS AND STRUCTURAL CHEMISTRY OF

TETRAKIS(THIOHYDROXAMATO)HAFNIUM(IV) IN

$\text{Hf}(\text{CH}_3\text{C}_6\text{H}_4(\text{S})\text{N}(\text{O})\text{CH}_3)_4 \cdot \text{C}_2\text{H}_5\text{OH}$

Kamal Abu-Dari and Kenneth N. Raymond

July 1981

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782

LBL-13173
2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Specific Sequestering Agents for the Actinides. 7. Synthesis and
Structural Chemistry of Tetrakis(thiohydroxamato)hafnium(IV)

By

Kamal Abu-Dari and Kenneth N. Raymond*

Contribution from the Department of Chemistry (and Materials and Molecular
Research Division, Lawrence Berkeley Laboratory) University of California,
Berkeley, California 94720

This work was supported by the Director, Office of Energy Research,
Office of Basic Energy Sciences, Chemical Sciences Division of the
U.S. Department of Energy under Contract Number W-7405-ENG-48.

Introduction

A continuing project in this laboratory has been the design and synthesis of specific sequestering agents for the actinide(IV) ions. The degree of success so far achieved in this goal has been reviewed recently.¹⁻⁴ The approach used relies on the similar coordination chemistry of high-spin Fe^{3+} and Pu^{4+} ^{2,3,5} and the molecular architecture of the siderophores (ferric-ion specific chelating agents of microbial origin). While hydroxamate and catecholate chelating groups are the primary functional groups in siderophores, the thiohydroxamate chelating unit also occurs in at least one compound of natural origin.⁶

The tetrakis(catecholato)metallate(IV) complexes of Th(IV), U(IV) and Hf(IV) have all been prepared and structurally characterized.^{7,8} They are isostructural, with the D_{2d} coordination symmetry of the trigonal-faced dodecahedron. Both the hydroxamate complexes tetrakis(N-isopropyl-3,3-dimethylbutanohydroxamato)thorium(IV)¹ and tetrakis(N-phenylbenzohydroxamato)hafnium(IV)⁹ have this same D_{2d} geometry. In contrast, the more sterically hindered complex, tetrakis(N-isopropyl-2,3-dimethylpropanahydroxamato)thorium(IV), is distorted substantially toward a cubic coordination polyhedron.¹ No tetrakis(thiohydroxamate) complexes have been structurally characterized to date.

Determination of the coordination geometry of the unconstrained tetrakis(bidentate)metallate(IV) complexes are a fundamental component in the design of an octadentate chelating agent that incorporates four such functionalities to form an optimum metal coordination environment. While we have prepared thiohydroxamate complexes of Th(IV), no crystals suitable for structure analysis have been obtained. Instead, we report

the synthesis and structural characterization of the hafnium complex, the title compound.

Experimental

Synthesis of Compounds

Tetrakis(N-methyl-p-thiotolylhydroxamato)hafnium(IV). Ethanol.

An aqueous solution of hafnium tetrachloride (1.0 g, 3.12 mmol) in 10 mL water was added to a stirred aqueous solution of sodium N-methyl-p-thiotolylhydroxamate (2.53 g, 12.5 mmol); the mixture was stirred for five minutes and the white precipitate was filtered, washed with water, and dried in vacuum over P_2O_5 . The compound was recrystallized from acetone-ethanol solution by slow evaporation. The crystals lose solvent easily in air. A sample for analysis was dried under high vacuum. The complexes are very hygroscopic and were partially hydrated just prior to analysis.

Anal. Calc. for $Hf(C_9H_{10}NOS)_4$: C, 48.13; H, 4.49; N, 6.23; S, 14.27.

Calc. for $Hf(C_9H_{10}NOS)_4 \cdot H_2O$: C, 47.13; H, 4.61; N, 6.10; S, 14.00.

Found: C, 47.43; H, 4.74; N, 6.12; S, 13.05%.

Thorium complexes with N-methylthiobenzohydroxamate and N-methyl-p-thiotolylhydroxamate were prepared by similar methods and were recrystallized from chloroform-isopropyl alcohol by slow evaporation. Again, the dried compounds are very hygroscopic. Analysis for tetrakis(N-methyl-p-thiotolylhydroxamato)thorium(IV) is:

Anal. Calc. for $Th(C_9H_{10}NOS)_4 \cdot H_2O$: C, 44.53; H, 4.36; N, 5.77.

Found: C, 44.25; H, 4.26; N, 5.66%.

Structure Determination

The crystal diffraction symmetry and lack of systematic absences imply the triclinic space groups $P1$ or $P\bar{1}$. The latter is confirmed by the final structure. The crystal density was determined by the flotation technique in bromoform/heptane solution. Crystal and unit cell data are given in Table I. Intensity data were collected on an Enraf-Nonius CAD-4 automated diffractometer controlled by a PDP-8/E computer, using monochromatic MoK_α radiation.^{10,11} The 8231 data were reduced to F^2 and $\sigma(F^2)$ as previously described¹¹⁻¹³ and averaged to give 7507 reflections with $F^2 > 3\sigma(F^2)$.

The structure was solved by direct phasing (MULTAN), Fourier and least-squares techniques. The positions of the phenyl hydrogen atoms were calculated and found to match those found in the difference Fourier. The phenyl, and most methyl and some ethanol, hydrogen atoms were introduced as fixed atoms in the final refinement. Full-matrix least-squares refinement with 470 variables and using the 7507 reflections with $F^2 > 3\sigma(F^2)$, led to convergence with $R = 2.8\%$ and $R_w = 4.4\%$.¹⁴

Description of the Structure and Discussion

While thorium(IV) complexes with N-unsubstituted thiohydroxamic acids have been obtained, they are unstable toward hydrolysis. Green tetrakis(thiohydroxamato)uranium(IV) complexes have also been obtained under oxygen free conditions [the metal undergoes oxidation to uranium(VI) in the presence of oxygen]. In contrast, the colorless hafnium complex reported here is quite stable and its structure consists of tetrakis(N-methyl-p-thiotolylhydroxamato)hafnium(IV) and ethanol molecules of

crystallization (Figure 1). The structure of the complex and the atom numbering scheme used within it is shown in Figure 2. Relevant bond lengths and angles of the thiohydroxamato molecules are given in Table V. The thiohydroxamato rings are planar and bend away from the hafnium atom such that the Hf-to-plane distances are 0.62, 0.11, 0.77 and 0.34 Å for chelate rings 1, 2, 3 and 4, respectively.

Bond distances in the thiohydroxamato ring compare well with the corresponding bond distances in other thiohydroxamato complexes.¹⁵ The N-O and N-C bond distances, 1.355(4) and 1.303(7) Å, in tetrakis(N-methyl-p-thiotolylhydroxamato)hafnium(IV) are slightly shorter than the corresponding bond distances, 1.37(2) and 1.32(2) Å, found in the tetrakis(N-phenylbenzohydroxamato)hafnium(IV) structure.⁹ This reflects the increase in contribution of structure Ib to the bonding in thiohydroxamato complexes.^{13,15-17}

Detailed analyses of eight-coordinate polyhedra have been carried out by several workers.¹⁸⁻²¹ The general conclusion is that the trigonal-faced dodecahedron (D_{2d}) and the square antiprism (D_{4d}) lie very close in energy, with the bicapped trigonal prism (C_{2v}) at slightly higher energy,²¹ (Figure 3). Both the bicapped trigonal prism and the square

antiprism can be generated from the intermediate dodecahedron by changing the length of one or two b edges.²¹ A set of shape parameters have been proposed to describe eight-coordinate geometries, where δ is the dihedral angle between faces of the polyhedron defined by the ligand atoms, and ϕ measures the nonplanarity of the trapezoidal-type atoms BABA in the dodecahedron (Figure 3). The ϕ and the δ values for the compound tetrakis(N-methyl-p-thiotolylhydroxamato)hafnium(IV) are given in Table VI, along with the corresponding calculated values for the ideal dodecahedron, square antiprism and bicapped trigonal prism. The average value for ϕ (15°) is very close to that for C_{2v} geometry (14.1°). The δ values also support the conclusion that the solid state thiohydroxamato hafnium complex has a geometry close to a bicapped trigonal prism. The largest deviations from C_{2v} geometry are for t_1 edges and for the diagonal edges of the tetragonal face. These deviations are ascribed to the difference in the size of the sulfur and oxygen atoms.

In conclusion, the thorium and hafnium complexes of thiohydroxamic acids have been prepared and characterized. The structure of the compound tetrakis(N-methyl-p-thiotolylhydroxamato)hafnium(IV) displays a bicapped trigonal prismatic coordination polyhedron, a polyhedron not observed for the catecholato and hydroxamato complexes studied to date.

Acknowledgment

We thank the U.S.A.I.D. (K.A.D.) and N.I.H. for support. This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division of the U.S. Department of Energy under Contract Number W-7405-ENG-48.

Supplementary Material

Table II, III and IV and a listing of structure factor amplitudes.

Ordering information is given on current masthead page.

References

1. Previous paper in this series: Smith, W. L.; Raymond, K. N.,
J. Am. Chem. Soc. 1981, 103, 3341. A preliminary account of this work appeared at the 1980 Fall Meeting of the American Chemical Society, Abstract 300.
2. Smith, W. L.; Raymond, K. N. Structure and Bonding, Vol. 43, 159-186, 1981.
3. Raymond, K. N.; Smith, W. L.; Weitzl, F. L.; Durbin, P. W.; Jones, E. S.; Abu-Dari, K.; Sofen, S. R.; Cooper, S. R. In "Lanthanides and Actinides"; Edelstein, N. M., Ed.; ACS Symposium Series, American Chemical Society: Washington, D.C., 1980, 143-172; Weitzl, F. L.; Raymond, K. N.; Smith, W. L.; Howard, T. R.
J. Am. Chem. Soc. 1978, 100, 1170-2.
4. New York Times, September 13, 1980, p 8.
5. Weitzl, F. L.; Raymond, K. N. 1980 Fall Meeting of the American Chemical Society, Abstract 109.
6. Raymond, K. N.; Carrano, C. J. Acc. Chem. Res. 1979, 12, 183-190.
7. Sofen, S. R.; Abu-Dari, K.; Freyberg, D. P.; Raymond, K. N.
J. Am. Chem. Soc. 1978, 100, 7882-7887
8. Sofen, S. R.; Cooper, S. R.; Raymond, K. N. Inorg. Chem. 1979, 18, 1611-16.
9. Teanqui, D.; Laugier, J.; Boyer, P.; Vullket, P. Acta Crystallog. 1978, B34, 767-73.
10. The Enraf-Nonius Fortran IV system, which operates the CAD-4 diffractometer, and locally written or modified programs for the CDC-7600 have been described in Refs. 12 and 13.

11. The intensities of a unique set of reflections ($\pm h, \pm k, \pm l$) with $2^\circ < 2\theta < 53^\circ$ were measured, using a θ - 2θ scan technique [the duplicates were measured for $(0, \pm k, \pm l)$ as a check]. The instrument parameters used in data collection were those described in Ref. 13. A p factor of 0.03 was used in data reduction.
12. Baker, E. C.; Brown, L. D.; Raymond, K. N. Inorg. Chem. 1975, 14, 1376-79.
13. Abu-Dari, K.; Ekstrand, J. D.; Freyberg, D. P.; Raymond, K. N. Inorg. Chem. 1979, 18, 108-12.
14. Definitions of the indicators are: $R = (\sum ||F_o| - |F_c||) / \sum |F_o|$;
 $R_w = [\sum w(|F_o| - |F_c|)^2 / \sum w F_o^2]^{1/2}$, the error in an observation of unit weight is: $[(\sum w(|F_o| - |F_c|)^2) / (N_o - N_v)]^{1/2}$ where N_o is the number of observations and N_v is the number of variables. The final error in an observation of unit weight is 1.63. The largest peaks in the final difference Fourier were $< 0.35 \text{ e}^-/\text{\AA}^3$. Table II gives the positions and thermal parameters of the non-hydrogen atoms. Table III lists the rms amplitudes of vibration of these atoms. Table IV gives the positional and thermal parameters assigned to the hydrogen atoms. These tables are all available as supplementary material; see the paragraph at the end of the paper.
15. Freyberg, D. P.; Abu-Dari, K.; Raymond, K. N. Inorg. Chem. 1979, 18, 3037-43.
16. Abu-Dari, K.; Freyberg, D. P.; Raymond, K. N. Inorg. Chem. 1978, 18, 2427-31.
17. Abu-Dari, K.; Raymond K. N. Inorg. Chem. 1980, 20, 2034-40.
18. Blight, D. G.; Kepert, D. L. Inorg. Chem. 1972, 11, 1556-63.

19. Muettterties, E. L.; Wright, C. M. Q. Rev. Chem. Soc. 1967, 21, 109-53.
20. Porai-Koshits, M. A.; Aslanov, L. A. Zh. Strukt. Khim. 1972, 13, 266-76.
21. Muettterties, E. L.; Guggenberger, L. J. J. Am. Chem. Soc. 1974, 96, 1748-56.

Table I

Summary of Crystal Data for $\text{Hf}(\text{C}_9\text{H}_{10}\text{NOS})_4 \cdot \text{C}_2\text{H}_5\text{OH}$

Formula weight	945.5
Space group	$P\bar{1}$
Cell constants ^a	
<u>a</u>	10.501(1) Å
<u>b</u>	10.958(1) Å
<u>c</u>	18.684(2) Å
α	97.388(8)°
β	93.136(8)°
γ	98.205(9)°
Cell volume	2014.4(4)
Formula units/cell, Z	2
Calculated density	1.49 g cm ⁻³
Observed density	1.50(1) g cm ⁻³
Crystal dimensions	0.35 x 0.30 x 0.18 mm
Absorption coefficient, $\mu \text{ MoK}_{\alpha}$	28.5 cm ⁻¹

^a Ambient temperature of 23°C; MoK_{α_1} radiation, $\lambda = 0.70926$ Å.

Table V

Bond distances and angles for Hf(N-methyl-p-thiotolylhydroxamato)hafnium(IV)^a

	Ring 1	Ring 2	Ring 3	Ring 4
Hf-S	2.688(1)	2.673(1)	2.699(1)	2.653(1)
Hf-O	2.175(3)	2.139(3)	2.144(3)	2.141(3)
C-S	1.709(4)	1.694(4)	1.701(4)	1.695(4)
N-O	1.365(4)	1.352(4)	1.347(5)	1.357(4)
C-N	1.284(5)	1.310(5)	1.301(5)	1.317(5)
C-C(ph)	1.484(5)	1.485(5)	1.476(6)	1.478(6)
CH ₃ -N	1.434(6)	1.465(5)	1.461(5)	1.459(5)
O-Hf-S	70.11	71.67(7)	70.28(8)	72.35(7)
S-C-N	118.8(3)	119.7(3)	118.5(3)	119.3(3)
S-C-C(ph)	120.2(3)	120.0(3)	119.3(3)	119.6(3)
N-C-C(ph)	121.1(3)	120.3(3)	122.2(3)	121.1(4)
O-N-C(CH ₃)	111.1(4)	112.0(3)	111.8(3)	111.3(3)
O-N-C	120.7(3)	120.4(3)	120.8(3)	120.8(3)

^aBond (Å) or angle (deg).

Table VI
Shape parameters for $\text{Hf}(\text{N-methyl-p-thiotolylhydroxamate})_4$

1. Dihedral angles, δ (degrees)

Edge ^b	Porai-Koshits/Aslanov notation for C_{2v}	Observed angle	Calculated angle for ideal		
			D_{2d}	D_{4d}	C_{2v}
$S_2(S_1S_3)S_4$	P_1	58.3	51.3	52.4	60.8
$O_1(O_2O_4)O_3$	P_1	58.7	51.3	52.4	60.8
$S_1(S_2S_3)O_3$	P_1	62.6	62.5	52.4	60.8
$O_2(O_1O_4)S_4$	P_1	45.6	62.5	52.4	60.8
$S_2(O_3S_3)S_4$	P_2	66.7	51.3	76.2	60.8
$O_1(O_4S_4)O_3$	P_2	75.7	51.3	76.2	60.8
$O_3(S_3S_4)S_1$	P_2	71.0	62.5	76.2	60.8
$O_2(O_3O_4)S_4$	P_2	62.6	62.5	76.2	60.8
$O_1(O_2S_2)O_3$	t_1	82.1	51.3	76.2	90.0
$S_2(O_1S_1)S_4$	t_1	74.5	51.3	76.2	90.0
$O_2(O_3S_2)S_3$	t_2	47.5	29.5	52.4	48.2
$O_4(O_1S_4)S_1$	t_2	45.3	29.5	52.4	48.2
$O_4(O_2O_3)S_2$	t_2	39.5	62.5	52.4	48.2
$O_1(S_1S_4)S_3$	t_2	58.2	62.5	52.4	48.2
$O_1(S_1S_2)S_3$	h_1	71.4	62.5	76.2	70.9
$O_4(O_1O_2)S_2$	h_1	65.4	62.5	76.2	70.9
$O_4(O_3S_4)S_3$	h_2	21.1	29.5	0.0	21.8
$O_2(O_1S_2)S_1$		9.8	29.5	0.0	0.0

2. Twist angle in the BAAB trapezoid, ϕ

Observed	Calculated for		
	D_{2d}	D_{4d}	C_{2v}
15	0.0	24.5	14.1

^aSee Figure 3.

^bIn defining the dihedral angle, atoms in parenthesis represent the edge, the other two atoms form the two intersecting triangles with the edge.

Figure Captions

Figure 1. Packing diagram for tetrakis(N-methyl-p-thiotolylhydroxamato)-hafnium(IV)·C₂H₅OH.

Figure 2. A perspective view and atom numbering scheme for tetrakis(N-methyl-p-thiotolylhydroxamato)hafnium(IV).

Figure 3. The coordination polyhedron of Hf(N-methyl-p-thiotolylhydroxamate)₄ compared to the ideal dodecahedron, bicapped trigonal prism and square antiprism.

XBL 799-11359

XBL 799-11358

D_{2d}

D_{4d}

C_{2v}

$Hf(THIO)_4$

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720