

Lawrence Berkeley National Laboratory

Recent Work

Title

THE EFFECTS OF 8-METHYL LIPOIC ACID ON THE EVOLUTION OF OXYGEN AND REDUCTION OF CARBON DIOXIDE DURING PHOTOSYNTHESIS

Permalink

<https://escholarship.org/uc/item/31s8r3cw>

Authors

Bassham, J.A.
Egster, Horst
Edmonston, Frances
et al.

Publication Date

1963-08-01

UCRL-10972

c.2

University of California

Ernest O. Lawrence
Radiation Laboratory

THE EFFECTS OF 8-METHYL LIPOIC ACID ON
THE EVOLUTION OF OXYGEN AND REDUCTION
OF CARBON DIOXIDE DURING PHOTOSYNTHESIS

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.

[REDACTED]

[REDACTED]

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

To be published in Biochem and Biophys.

UCRL-10972

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

THE EFFECTS OF 8-METHYL LIPOIC ACID ON THE EVOLUTION OF OXYGEN
AND REDUCTION OF CARBON DIOXIDE DURING PHOTOSYNTHESIS

J. A. Bassham, Horst Ege, Frances Edmonston, and Martha Kirk

August 1963

UCRL 10972

THE EFFECTS OF 8-METHYL LIPOIC ACID ON THE EVOLUTION OF OXYGEN AND REDUCTION
OF CARBON DIOXIDE DURING PHOTOSYNTHESIS*

J.A. Bassham, Horst Egster†, Frances Edmonston and Martha Kirk
Bio-Organic Chemistry Group, Lawrence Radiation Laboratory, University of
California, Berkeley 4, California

Kinetic studies of transient changes in the pattern of ^{14}C -labeled compounds formed during photosynthesis with $^{14}\text{CO}_2$ have given much information about the operation of the carbon reduction cycle *in vivo*. Such changes accompanying the change from light to dark (Calvin and Massini, 1952; Bassham *et al.*, 1956) and the sudden change from high levels of CO_2 pressure to very low levels of CO_2 pressure (Wilson and Calvin, 1955) provided an important part of the evidence on which the carbon reduction cycle of photosynthesis was formulated (Bassham *et al.*, 1954). Introduction of a chemical inhibitor capable of interfering with electron transport from the photochemical stage of photosynthesis to the carbon reduction pathway could be a tool for further studies of this type.

It has been suggested (Bassham, 1963) that disulfide compounds such as lipoic acid or some enzyme bound disulfides might function in the electron transport from substances at the potential of reduced PPNR to the site of carbon reduction. We therefore chose an analog of lipoic acid in the hope of interfering with such a function. Also, we hoped to obtain information about a possible reductive carboxylation reaction *in vivo* in the light.

Chlorella pyrenoidosa, grown in continuous culture (Bassham and Calvin, 1957) was harvested and resuspended in a mineral nutrient solution described elsewhere (Bassham and Kirk, 1963). 80 ml of a 2% (wet packed volume / suspension volume) suspension of algae were placed in the steady state apparatus. The work described in this paper was sponsored by the U.S. Atomic Energy Commission.

*Fellow of the German Federal Ministry of Scientific Research.

(Bassham and Kirk, 1963). However, the pH control (automatic addition of 0.1 N NH₄ OH) was set at 5.0. Air, containing 2% CO₂, was passed through the vessel which was illuminated from one side by an incandescent lamp, est. 33,000 lux incident, and from the other side by a bank of alternate white and blue fluorescent lamps, est. 8000 lux incident. After one hour of photosynthesis, the system was closed and ¹⁴CO₂ was introduced. The rates of oxygen evolution and carbon dioxide and ¹⁴CO₂ uptake were continuously observed (Bassham and Kirk, 1960; Smith *et al.*, 1961).

Previous studies (Bassham and Kirk, 1960) showed that phosphoglyceric acid (PGA) and the sugar phosphates of the carbon reduction cycle become "saturated" with radicarbon after some 5 min of photosynthesis in the presence of ¹⁴CO₂ of constant specific radioactivity. After 6 min, 6 samples of the algae suspension were taken at 30 sec intervals into weighed tubes containing methanol to give a final concentration of 80% methanol, to stop the enzymic reactions. 10 mg of 8-methyl lipoic acid in 200 μ l of ethanol was then injected into the algae suspension. Control experiments showed that 200 μ l of ethanol alone cause no measurable change in photosynthesis rate or labeling pattern.

Samples of the algae suspension were taken at times shown in the Figs. All algal samples were subsequently extracted, concentrated and analyzed by two-dimensional paper chromatography and radicautography as described previously (Bassham and Calvin, 1957). The amount of ¹⁴C in each compound was determined by means of the automatic spot counter (Noses and Lomborg, 1961, 1962).

The course and degree of inhibition with several levels of 8-methyl lipoic acid and with lipoic acid are indicated in Table I.

Figs. 1 and 2 show the amount of ¹⁴C-labeled compounds obtained from the samples taken during the course of experiment 3 (see Table I). At in-

Table I. INHIBITION OF PHOTOSYNTHESIS IN Chlorella pyrenoidosa BY 8-METHYL LIPOIC ACID AND BY LIPOIC ACID.

Exp. Conditions	Time (min)	micromoles/min/cm ³ algae (wet pack)	
		O ₂ evolution	CO ₂ uptake
1 initial after 1 mg 8-Me lipoic acid	before	10.9	8.2
	2-10	7.8	5.5
2 initial after 5 mg 8-Me lipoic acid	before	8.8	7.6
	2-8	1.3	1.3
	8-12	recovering	
	12-20	7.0	4.9
3 initial after 10 mg 8-Me lipoic acid	before	12.9	10.6
	0-20	0	0
	20-26	recovering	
	26-40	4.2	2.9
4 initial after 10 mg lipoic acid	before	10.2	7.6
	3-10	0.2	0.3
	14-20	5.4	4.3
	26-40	8.7	7.0

previous studies with the steady state apparatus (Bassham and Kirk, 1960; Smith *et al.*, 1961) the specific activity of the ¹⁴CO₂ has been maintained at a constant measured value during the course of the experiment. The amount of ¹⁴C found experimentally in each compound has been converted to the micromoles of carbon in the compound in the active reservoir by dividing the amount of ¹⁴C by the ratio ¹⁴C/(¹²C + ¹⁴C) in the CO₂ administered. This result has been further converted to give the micromoles of carbon in the compound per cm³ of wet packed algae.

Fig. 1 shows a sudden, large drop in the level of PGA and a sudden rise in the levels of fructose-1,6-diphosphate and sedoheptulose-1,7-diphosphate upon introduction of the inhibitor. The rate of decrease of the PGA pool during the first 15 sec with the inhibitor may be calculated as $(2.4 - 0.7)/0.25 = 6.8$ micromoles of ¹⁴C/min/cm³ of algae. This rate may be compared with the rate of uptake of carbon dioxide from Table I, which was 10.6 micromoles per min. The rate of increase in sucrose, which is accelerated once the fructose diphosphate has increased, occurs in the

Figs. 1 and 2. Effects of 8-methyl lipoic acid on labeling of compounds with ^{14}C during photoynthesis with $^{14}\text{CO}_2$.

M.U.B. 1982


Fig. 1.

- ▲ Sucrose
- Fructose-1,6-diphosphate
- Glucose-1,6-diphosphate
- Sedoheptulose-1,7-diphosphate
- 3-Phosphoglyceric acid

M.U. 31459

Fig. 2.

- △ Glucose-6-phosphate
- Sedoheptulose-7-phosphate
- Fructose-6-phosphate
- Ribulose-1,5-diphosphate






Fig. 2

-5-

absence of any appreciable increase in uridine diphosphoglucose (not shown but included in a similar to the "glucose diphosphates").

Changes in the levels of some of the sugar monophosphates and of ribulose-1,5-diphosphate are shown in Fig. 2. Labeled fructose and sedoheptulose monophosphates decrease rapidly at first, then undergo a small positive transient, before continuing to decline gradually. Glucose-6-phosphate undergoes a gradual but large decrease in radioactivity during the time that sucrose increases. Considering the great change in phosphoglyceric acid and the complete inhibition of photosynthesis, it is surprising that ribulose diphosphate changes in concentration by only a small amount during the same period.

In terms of the photosynthetic carbon reduction cycle as usually expressed (Fassham *et al.*, 1954) the more immediate effects of the addition of inhibitor seem to suggest either a stimulation in the reduction of PGA (which is difficult to rationalize with the apparent inhibition of CO_2 uptake and O_2 evolution) or perhaps more plausibly, an inhibition of carboxylation reactions leading to the formation of PGA.

In the latter case, since ribulose-1,5-diphosphate neither accumulates nor disappears it is necessary to postulate that its formation from ribulose-5-phosphate is also inhibited. Inhibition of this reaction (presumably catalyzed by a kinase) may be somehow related to inhibition of the conversion of fructose and sedoheptulose diphosphates to their respective monophosphates (presumably through the action of phosphatases).

Other explanations of the inhibitor effects (such as inhibition of an *in vivo* reductive carboxylation of hexose and heptose diphosphates) are possible, but their serious consideration must await further experimental evidence. The immediacy and magnitude of effects caused by 8-methyl

lipoic acid and lipoic acids, indicate important and primary roles for disulfide groups in the primary reactions for carbon reduction during photosynthesis.

REFERENCES

- Bassham, J.A., Adv. in Enzymol. XV, 39 (1963).
Bassham, J.A., Benson, A.A., Kay, L.D., Barris, A.Z., Wilson, A.T., and Calvin, M., J. Am. Chem. Soc. 76, 1760 (1954).
Bassham, J.A. and Calvin, M., "The Path of Carbon in Photosynthesis", Prentice-Hall, Englewood Cliffs, N.J., 1957.
Bassham, J.A. and Kirk, M., Biochim. Biophys. Acta 43, 447 (1960).
Bassham, J.A. and Kirk, M., "Microalgae and Photosynthetic Bacteria", p. 493, 1963. (Special issue, Plant and Cell Physiol., Japan.)
Bassham, J.A., Shibata, K., Steenberg, K., Bourdon, J., and Calvin, M., J. Am. Chem. Soc. 78, 4129 (1956).
Calvin, M. and Bassham, J.A., "The Photosynthesis of Carbon Compounds", W.A. Benjamin, Inc., New York, 1962.
Calvin, M. and Massini, P., Experientia VIII, 445 (1952).
Noses, V. and Lonberg-Holm, K.K., Anal. Biochem. 2, 11 (1963).
Smith, D.C., Bassham, J.A. and Kirk, M., Biochim. Biophys. Acta 48, 299 (1961).
Wilson, A.T. and Calvin, M., J. Am. Chem. Soc. 77, 5948 (1955).

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

