Lawrence Berkeley National Laboratory

Recent Work

Title

Mesons Produced by Neutrons from the Cyclotron

Permalink

<https://escholarship.org/uc/item/31t7m4hr>

Authors

Smith, F.M.
Gardner, Eugene
Bradner, Hugh

Publication Date

1949-11-23

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNCLASSIFIED

UCRL-527

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

MESONS PRODUCED BY NEUTRONS FROM THE CYCLOTRON

F. M. Smith, Eugene Gardner, and Hugh Bradner

November 23, 1949

Berkeley, California

MESONS PRODUCED BY NEUTRONS FROM THE CYCLOTRON

F. M. Smith, Eugene Gardner, and Hugh Bradner

Radiation Laboratory, Department of Physics
University of California
Berkeley, California

In photographic plates exposed to high energy neutrons from the 184-inch Berkeley cyclotron, we have observed stars in which one of the outgoing particles is a meson. This type of star is well known from cosmic ray studies¹, and the only new feature is that the stars we are

¹ Lattes, Muirhead, Occhialini, and Powell, *Nature* 159, 694 (1947)

studying were initiated by particles from the cyclotron. The arrangement of apparatus is shown in Fig. 1. As indicated in the figure, the neutrons used for bombarding the plates are produced in the cyclotron when a beam of 345 Mev protons strikes an internal target. The plates are placed in a position similar to that used in cloud chamber studies² of mesons

² W. Hartsough, E. Hayward, and W. M. Powell, *Phys. Rev.* 75, 905 (1949)

produced by neutrons from the cyclotron.

An example of a photographic emulsion star in which one of the outgoing particles is a meson is shown at point A of Fig. 2. The star at A consists of two tracks, a proton (upper track) and a π^- -meson (lower track). From the grain density changes along the tracks it is clear that both proton and meson were moving from left to right. The meson came to rest in the emulsion at point B and initiated a one-prong star; the one prong is the straight heavy track which begins at B. This event was found in a preliminary survey of meson production by neutrons from the cyclotron.

-2-

The survey indicated that the neutron beam offered excellent opportunities for meson work, and systematic studies are being planned. Inasmuch as it may be some time before any results are available from these studies, we shall give additional details of the preliminary survey. In this survey we scanned hastily an area of 8 sq. cm. of one plate which was exposed as shown in Fig. 1. The plate used was an Ilford C.2 unbacked emulsion with a thickness before development of 300 microns. The exposure time was about 5 minutes at a distance of about 60 feet from the target. We found a total of 5 stars in which one of the outgoing particles was a meson. In addition, there were 4 meson tracks which could not be followed back to the points at which the mesons were formed. In the same plate area we found 5×10^4 stars at which no meson tracks were observed. We hesitate to use these numbers to estimate the cross section for the production of mesons by neutrons because of the uncertainty in the meson count. In this survey we made no attempt to find all of the mesons or to estimate the fraction of the total that might have been missed. Even with a more careful search we would not expect to see the higher energy mesons with this plate, since the C.2 emulsion does not record mesons of energy greater than about 10 Mev.

We are indebted to Professor Ernest O. Lawrence for his continued interest in this work. We also wish to thank Professor R. L. Thornton for many helpful discussions. The first work on this problem was done by Professor C. M. G. Lattes, who has kindly given us the benefit of his experience. The photomicrograph shown in Fig. 2 was prepared by Mr. A. J. Oliver. This work was done under the auspices of the Atomic Energy Commission.

Figure Captions

Fig. 1 - Sketch showing position of photographic plate for exposure to neutrons from the cyclotron. (Not to scale.)

Fig. 2 - Photographic emulsion star in which one of the outgoing particles is a π^- -meson. Meson is formed at A and comes to rest at B.

FIG. 1

SKETCH SHOWING POSITION OF PHOTOGRAPHIC PLATE FOR
EXPOSURE TO NEUTRONS FROM THE CYCLOTRON. (NOT TO SCALE.)

FIG. 2
PHOTOGRAPHIC EMULSION STAR IN WHICH ONE OF THE OUTGOING PARTICLES IS A π^- -MESON. MESON IS FORMED AT A AND COMES TO REST AT B.