UC San Diego

UC San Diego Previously Published Works

Title

Kisspeptin neurones do not directly signal to RFRP-3 neurones but RFRP-3 may directly modulate a subset of hypothalamic kisspeptin cells in mice.

Permalink

<https://escholarship.org/uc/item/335186bp>

Journal

Journal of neuroendocrinology, 25(10)

ISSN

0953-8194

Authors

Poling, MC
Quennell, JH
Anderson, GM
et al.

Publication Date

2013-10-01

DOI

10.1111/jne.12084

Peer reviewed

ORIGINAL ARTICLE

Kisspeptin Neurones do not Directly Signal to RFRP-3 Neurones but RFRP-3 may Directly Modulate a Subset of Hypothalamic Kisspeptin Cells in Mice

M. C. Poling*, J. H. Quennell†, G. M. Anderson‡ and A. S. Kauffman*

*Department of Reproductive Medicine, University of California San Diego, La Jolla, CA, USA.

†Department of Anatomy, School of Medical Sciences, University of Otago, Dunedin, New Zealand.

‡Centre for Neuroendocrinology, University of Otago, Dunedin, New Zealand.

Journal of Neuroendocrinology

The neuropeptides kisspeptin (encoded by *Kiss1*) and RFamide-related peptide-3 (also known as GnIH; encoded by *Rfrp*) are potent stimulators and inhibitors, respectively, of reproduction. Whether kisspeptin or RFRP-3 might act directly on each other's neuronal populations to indirectly modulate reproductive status is unknown. To examine possible interconnectivity of the kisspeptin and RFRP-3 systems, we performed double-label *in situ* hybridisation (ISH) for the RFRP-3 receptors, *Gpr147* and *Gpr74*, in hypothalamic *Kiss1* neurones of adult male and female mice, as well as double-label ISH for the kisspeptin receptor, *Kiss1r*, in *Rfrp*-expressing neurones of the hypothalamic dorsal-medial nucleus (DMN). Only a very small proportion (5–10%) of *Kiss1* neurones of the anteroventral periventricular region expressed *Gpr147* or *Gpr74* in either sex, whereas higher co-expression (approximately 25%) existed in *Kiss1* neurones in the arcuate nucleus. Thus, RFRP-3 could signal to a small, primarily arcuate, subset of *Kiss1* neurones, a conclusion supported by the finding of approximately 35% of arcuate kisspeptin cells receiving RFRP-3-immunoreactive fibre contacts. By contrast to the former situation, no *Rfrp* neurones co-expressed *Kiss1r* in either sex, and *Tacr3*, the receptor for neurokinin B (NKB; a neuropeptide co-expressed with arcuate kisspeptin neurones) was found in <10% of *Rfrp* neurones. Moreover, kisspeptin-immunoreactive fibres did not readily appose RFRP-3 cells in either sex, further excluding the likelihood that kisspeptin neurones directly communicate to RFRP-3 neurones. Lastly, despite abundant NKB in the DMN region where RFRP-3 soma reside, NKB was not co-expressed in the majority of *Rfrp* neurones. Our results suggest that RFRP-3 may modulate a small proportion of kisspeptin-producing neurones in mice, particularly in the arcuate nucleus, whereas kisspeptin neurones are unlikely to have any direct reciprocal actions on RFRP-3 neurones.

Key words: RFRP-3, GnIH, *Gpr147*, *Gpr74*, *Gpr54*, kisspeptin, *Kiss1*, *Kiss1r*, *Tacr3*, *Tac2*, neurokinin B, reproduction, hypothalamus

Correspondence to:
Dr Alexander S. Kauffman,
Department of Reproductive Medicine,
Leichtag Building, Room 3A-15,
University of California, San Diego,
9500 Gilman Drive, #0674, La Jolla,
CA 92093, USA (e-mail:
akauffman@ucsd.edu).

doi: 10.1111/jne.12084

Neuropeptides of the arginine-phenylalanine-amide (RFamide) family have been demonstrated to have potent modulatory effects on a variety of physiological functions, including reproduction (1). Two members of this family, kisspeptin (encoded by *Kiss1*) and RFamide-related peptide 3 (RFRP-3; encoded by *Rfrp*), have been shown to regulate mammalian reproductive function through central mechanisms but have opposing effects on the reproductive axis in mice, with kisspeptin stimulating and RFRP-3 inhibiting reproduction, respectively.

The kisspeptin system, which includes kisspeptin and its receptor, *Kiss1r* (formerly known as *Gpr54*), is considered stimulatory and essential for reproductive function. Human patients or rodents lacking functional *Kiss1* or *Kiss1r* genes suffer from impaired puberty and hypogonadotrophic hypogonadism, presenting with low levels of gonadotrophins and sex steroids, underdeveloped gonads, impaired sexual development, and infertility (2–5). Exogenous kisspeptin administration potently stimulates the secretion of luteinising hormone (LH) and follicle-stimulating hormone (5–9), working

centrally through a gonadotrophin-releasing hormone (GnRH)-dependent mechanism (10,11). Kisspeptin can directly activate GnRH neurones, as determined via c-fos induction (a marker of neuronal activation) in GnRH cells (6,11) and stimulation of electrical firing of GnRH neurones in brain explants (12,13). Anatomical support for a direct kisspeptin effect on GnRH cells includes the presence of kisspeptin neuronal fibres appositions on GnRH neurones (14–16) and high *Kiss1r* expression in the majority of GnRH neurones (5,11,12). Within the rodent brain, kisspeptin/*Kiss1* mRNA somata are found in two primary populations: the rostral hypothalamic continuum of the anteroventral periventricular nucleus and neighbouring rostral periventricular nucleus (AVPV/PeN), as well as the arcuate nucleus (ARC) (10,14). In the ARC, kisspeptin neurones highly co-express both neurokinin B (NKB; encoded by the *Tac2* gene) (17) and dynorphin, giving rise to the terminology KNDy neurones, although the exact roles of these co-transmitters are still being elucidated.

By contrast to kisspeptin, RFRP-3 has potent inhibitory actions on both GnRH neuronal activity and LH secretion in most rodent species (18–20). RFRP-3 is produced from a precursor peptide encoded by the *Rfrp* gene (21) and is the mammalian orthologue of avian gonadotrophin-inhibiting hormone (GnIH) (22,23). Through immunohistochemical assessment, RFRP-3-immunoreactive (IR) cells are found exclusively in the region of the dorsal-medial nucleus of the hypothalamus (DMN) of rodents (23,24), mirroring the selective expression of *Rfrp* mRNA in this region, as determined by *in situ* hybridisation (ISH) (25,26). In rodents, some GnRH neurones are contacted by RFRP-3 axonal fibres (23,27,28) and a subset of GnRH neurones express Gpr147, a high affinity receptor for RFRP-3 (26,28). In addition, RFRP-3 can bind to a second G-protein coupled receptor, Gpr74, with lower affinity (21,25), although this receptor is not expressed in GnRH neurones (26), and its relevance for the reproductive actions of RFRP-3 is currently unknown.

Although both kisspeptin and RFRP-3 appear to modulate the reproductive axis in part by direct effects on GnRH, it is possible that these two neuropeptides may also influence reproductive status via indirect pathways. To this end, it is currently unclear if there is modulatory cross-talk between these two neuropeptide populations. In addition to projecting to some GnRH cells, RFRP-3-IR fibres also project to a variety of brain regions that do not have GnRH neurones, including the AVPV, lateral hypothalamic area, paraventricular nucleus and ARC (23,24,27–29), and appositions of RFRP-3 fibres on some kisspeptin cells in the AVPV/PeN have been observed in female mice (28). Moreover, the RFRP-3 receptors, Gpr147 and Gpr74, are also expressed in several hypothalamic non-GnRH regions, including the periventricular nucleus, paraventricular nucleus and ARC (26,28,30,31). Additionally, RFRP-3 has been functionally shown to inhibit the electrical firing of some ARC kisspeptin neurones (32), suggesting that RFRP-3 may be able to directly regulate this kisspeptin population. However, whether ARC kisspeptin neurones actually express RFRP-3 receptors in animals of either sex has not been addressed. Similarly, the possibility of kisspeptin neurones regulating RFRP-3 neurones, either through kisspeptin itself or one of its co-transmitters, such as NKB, has not yet been

explored. Indeed, kisspeptin fibres have been observed in the DMN, and some *Kiss1r* expression has also been reported in this area (33), as has *Tacr3* (the receptor for NKB, a co-transmitter of ARC kisspeptin neurones) (34). Thus, there may be unilateral or bilateral communication between the RFRP-3 and kisspeptin populations to fine-tune each other's actions on the reproductive axis, although this has not yet been thoroughly examined.

To begin to address the possible anatomical interconnectivity of the kisspeptin and RFRP-3 systems, we used double-label ISH and immunocytochemistry to determine: (i) if one or both of the RFRP-3 receptors are expressed in *Kiss1* cells of either the AVPV/PeN or the ARC of males and females; (ii) if the kisspeptin or NKB receptors are co-expressed with *Rfrp* neurones in the DMN; (iii) if kisspeptin axonal fibres are found apposing RFRP-3 cells in the DMN; and (iv) if *Rfrp* neurones co-express *Tac2* (the gene encoding NKB), which is also known to be highly expressed in the DMN.

Materials and methods

Animals, gonadectomies and tissue collection

Adult C57BL6 mice of both sexes were housed under a 12 : 12 light/dark cycle (lights off 18.00 h) with food and water available *ad lib*. For some experiments, mice were anaesthetised and bilaterally gonadectomised (GDX) 1 week before sacrifice, as described previously (35,36). For *in situ* hybridisation studies, GDX mice or gonadal-intact mice (females in di-oestrus, as determined by vaginal smears) were anaesthetised with isoflurane and sacrificed by rapid decapitation. Brains were collected, frozen on dry ice and stored at -80°C . Five coronal series of 20- μm brain sections were cut on a cryostat, thaw-mounted onto Superfrost Plus slides (Menzel-Gläser, Braunschweig, Germany), and stored at -80°C until use in *in situ* hybridisation. For immunohistochemistry experiments, gonadal-intact male and GDX male and female mice were perfused with 4% paraformaldehyde in 0.1 M phosphate buffer for brain collection. Coronal (30- μm thick) sections throughout the caudal hypothalamus, containing the entire ARC and DMN, were cut from each brain on a sliding microtome with a freezing stage.

For all experiments, each group consisted of four to six animals. All experiments were conducted in accordance with the NIH Animal Care and Use Guidelines and with approval of the Animal Care and Use Committee of the University of California, San Diego, and the University of Otago Animal Ethics Committee.

Double-label ISH

The following cRNA ISH riboprobes have been previously described and validated: *Rfrp*, *Kiss1r*, *Gpr74*, *Gpr147* (26); *Kiss1* (10); *Tac2* (17,37); *Gnrh* (38). *Tacr3* was cloned from adult mouse hypothalamic cDNA into pBluescript II SK(–) transcription plasmid (Stratagene, La Jolla, CA, USA) as described previously (26) and corresponds to bases 286–691 of the mouse *Tacr3* sequence (NM_021382).

Double-label ISH assays were performed as described previously (17,39). For double-label assays studying ARC KNDy cells, we used *Tac2* as a designator for KNDy neurones because pilot studies indicated that *Tac2* expression per cell was stronger than *Kiss1*, allowing for better detection with the fluorescent digoxigenin (DIG) probe. Briefly, slide-mounted brain sections encompassing the hypothalamus were fixed in 4% paraformaldehyde, pre-treated with acetic anhydride, rinsed in $2 \times \text{SSC}$ (sodium citrate, sodium chloride), delipidated in chloroform, dehydrated in ethanols and air-dried.

Radiolabelled (^{33}P) antisense *Gpr147*, *Gpr74*, *Kiss1r*, *Tac2* or *Tacr3* (0.05 pmol/ml) and DIG-labelled *Rfrp*, *Kiss1*, *Tac2* or *Gnrh* riboprobes (digoxigenin labelling kit, dilution 1 : 500;) were combined with tRNA, denatured by boiling, and dissolved together in hybridisation buffer. The probe mix was then applied to slides (100 μl /slide), and slides were hybridised at 55 °C overnight. Slides were cover-slipped and placed in a 55 °C humidity chamber overnight. The slides were then washed in 4 $\times$ SSC and placed into RNase A treatment for 30 min at 37 °C, and then in RNase buffer without RNase at 37 °C for 30 min. After washing in 2 $\times$ SSC at room temperature, slides were washed in 0.1 $\times$ SSC at 62° C for 1 h. Slides were then incubated in 2 $\times$ SSC with 0.05% Triton X-100 containing 3% sheep serum (NSS) for 75 min at room temperature and then incubated overnight at room temperature with anti-DIG antibody conjugated to alkaline phosphatase [(Roche) diluted 1 : 500 in Buffer 1 containing 1% NSS and 0.3% Triton X-100]. The next day, slides were washed with Buffer 1 and incubated with Vector Red alkaline phosphatase substrate (Vector Laboratories, Burlingame, CA, USA) for 1 h at room temperature. The slides were then air-dried, dipped in NTB emulsion (Kodak, Rochester, NY, USA), stored at 4 °C, and developed and cover-slipped 9–11 days later.

In situ hybridisation slides were analysed with an automated image processing system (Dr Don Clifton, University of Washington, Seattle, WA, USA) by an individual who was unaware of the treatment group of each slide (40). DIG-containing cells (*Kiss1*, *Tac2*, *Rfrp* or *Gnrh* cells) were identified by fluorescence microscopy and the grain-counting software was used to quantify silver grains (representing *Gpr147*, *Gpr74*, *Kiss1r*, *Tacr3* or *Tac2* mRNA) overlying each cell. Signal-to-background ratios for individual cells were calculated, and a cell was considered double-labelled if its ratio was > 3.

Immunohistochemistry for kisspeptin and RFRP-3 in the DMN and ARC

For dual-label immunohistochemistry of kisspeptin and RFRP-3, all steps were separated by four 10-min washes in 50 mM Tris-buffered saline containing 0.5% Triton X-100 (TBS-TX). After blocking in TBS-TX containing 1% bovine serum albumin and 1% normal donkey serum, sections were incubated overnight at 4 °C in sheep anti-mouse kisspeptin-52 (AC053; kindly provided by Dr Alain Caraty, National Institute for Agronomic Research, France; dilution 1 : 2000) and rabbit anti-sparrow GnIH (PAC 123/124; kindly provided by Dr George Bentley, University of California, Berkeley; dilution 1 : 5000) in blocking solution. Sections were then incubated for 2 h at room temperature in biotinylated donkey anti-sheep (dilution 1 : 500; Jackson ImmunoResearch, Bar Harbor, ME, USA) and Alexa Fluor 488 donkey anti-rabbit (dilution 1 : 500; Molecular Probes, Carlsbad, CA, USA). Subsequently, sections were incubated for 1 h in Alexa Fluor 568-streptavidin (dilution 1 : 500; Molecular Probes). Staining was observed with an LSM 710 confocal microscope (Carl Zeiss, Oberkochen, Germany) using a $\times$ 63 objective lens and laser excitation lines and filters for 488 or 543 nm. Stacks of images, collected at intervals of 600 nm, were analysed offline using IMAGEJ (National Institutes of Health, Bethesda, MD, USA). In the DMN, twenty RFRP-3 cell bodies were visualised per mouse and all kisspeptin-IR contacts recorded. In the ARC, 29–50 kisspeptin cell bodies were visualised per mouse and all RFRP-3 contacts were recorded. Contacts were defined as no black pixel between the fibre and the soma. The omission of any of the primary antibodies resulted in complete absence of staining.

Statistical analysis

All data are expressed as the mean $\pm$ SEM for each group. In all experiments, differences were analysed by Student's t-test or by two-way ANOVA, followed by post-hoc comparisons for individual sex/treatment groups via Fisher's (protected) least significant difference. $P < 0.05$ was considered

statistically different. All analyses were performed using Statview, version 5.0.1 (SAS Institute, Cary, NC, USA).

Results

Experiment 1: Only a small proportion of AVPV Kiss1 neurones express *Gpr147* or *Gpr74*

We previously reported that 12–15% of *Kiss1* neurones in the AVPV/PeN of female mice express *Gpr147*, both in ovary-intact (di-oestrous) and oestradiol-treated conditions (28). However, it is unknown whether a similar proportion of AVPV/PeN *Kiss1* neurones in male mice express *Gpr147*. Additionally, the co-expression of *Gpr74* with *Kiss1* in the AVPV/PeN in either sex has not previously been determined. In the present experiment, we used double-label ISH to determine the co-expression of *Gpr147* or *Gpr74* mRNA in *Kiss1* neurones in the AVPV/PeN of gonadally-intact female (di-oestrous) and male mice. Such co-expression was not also examined in GDX mice because *Kiss1* expression is almost undetectable in the AVPV/PeN in the GDX state (35). As expected, there was a pronounced sex difference in the number of detectable AVPV/PeN *Kiss1* neurones, with females having several fold more *Kiss1* neurones than males (data not shown) (41). In terms of RFRP-3 receptors, we found an overall low abundance of both *Gpr147* and *Gpr74* expression in the AVPV/PeN region, unlike other regions, such as the paraventricular nucleus and thalamus where *Gpr147* and *Gpr74* mRNAs, respectively, were more highly expressed. In agreement with a previous study (28), quantitatively, only 12% of AVPV/PeN *Kiss1* neurones expressed *Gpr147* in females, and a similar proportion was observed in males (Fig. 1). An even smaller proportion (5–6%) of AVPV/PeN *Kiss1* neurones co-expressed *Gpr74* in either sex (Fig. 1). There were no statistical differences between the sexes for co-expression of either RFRP-3 receptor with *Kiss1* in the AVPV/PeN. The relative amount of *Gpr147* or *Gpr74* mRNA per *Kiss1* cell, reflected by the number of silver grains in each *Kiss1* cell, also did not differ between sexes (not shown).

Experiment 2: A moderate proportion of KNDy neurones in the ARC express *Gpr147* or *Gpr74* and receive contacts from RFRP-3 fibres

The ARC *Kiss1* population highly co-expresses *Tac2* (which encodes NKB) and is referred to as the KNDy neurone population. Here, we determined whether either *Gpr147* or *Gpr74* is co-expressed in ARC KNDy neurones of adult male and female mice. We found that *Gpr147* mRNA was moderately expressed in approximately 25% of KNDy neurones in gonadally-intact female (di-oestrous) and male mice (Fig. 2A–C). Similar levels of *Gpr147* co-expression were found in GDX mice of both sexes (Fig. 2). Similar to *Gpr147*, 23–25% of ARC KNDy neurones in gonadally-intact male and female mice co-expressed *Gpr74* mRNA, with slightly lower but not significantly different co-expression in GDX mice (Fig. 2D–E). For both assays, GDX mice had considerably more KNDy neurones than gonadally-intact mice ($P < 0.05$; data not shown), as expected as a result of the known stimulatory effects of GDX on KNDy cells in rodents (42,43). There were no statistical differences between sexes with

Fig. 1. Expression of *Gpr147* and *Gpr74* in anteroventral periventricular nucleus and neighbouring rostral periventricular nucleus (AVPV/PeN) *Kiss1* neurones by double-label *in situ* hybridisation. (A) Representative photomicrographs of double-label *in situ* hybridisation of *Kiss1* (red fluorescence) and *Gpr147* (silver grains) in an intact male. 3V, third ventricle. (B) *Kiss1* neurones co-expressing *Gpr147* (green arrowhead) and *Kiss1* neurones with no co-expression of *Gpr147* (blue arrows). (C) Quantification of the percentage co-expression of *Gpr147* in AVPV/PeN *Kiss1* neurones between gonadally-intact females (F) and males (M). There were no significant differences in co-expression between the groups. (D) Representative photomicrographs of double-label *in situ* hybridisation of *Kiss1* (red fluorescence) and *Gpr74* (silver grains) in a di-oestrous female. (E) *Kiss1* neurones co-expressing *Gpr74* (green arrowhead) and *Kiss1* neurones with no co-expression of *Gpr147* (blue arrows). (F) Quantification of the percentage co-expression of *Gpr147* in *Kiss1* neurones between gonadally-intact females (F) and males (M). There were no statistical differences between the sexes.

respect to the degree of *Gpr147* or *Gpr74* co-expression in ARC KNDy neurones and no group differences were observed in the relative amount of receptor mRNA per KNDy neurone (silver grains per double-labelled cell; not shown). The co-expression for the two RFRP-3 receptors within KNDy neurones was evenly dispersed throughout the ARC and not noticeably different between in any anatomical subregion within the KNDy neurone population.

To further examine possible RFRP-3 neurone to kisspeptin neurone interactions, we next used double-label IHC to assess potential RFRP-3 fibre contacts on kisspeptin neurones in the ARC of female mice. For this analysis, GDX females were used because this gonadal state allows for the identification of kisspeptin cell bodies in the ARC, unlike gonadally-intact mice in which the dense kisspeptin fibre network obscures cell bodies. Supporting our receptor co-expression data above, RFRP-3 fibres were observed to appose a moderate proportion of ARC kisspeptin cells. Quantification determined that approximately 35% of kisspeptin soma in the ARC received RFRP-3 fibre contacts (Fig. 3).

Experiment 3: RFRP-3 neurones do not highly express *Kiss1r* or *Tacr3* or receive axonal contacts from kisspeptin neurones

Experiments 1 and 2 indicated that a small population of kisspeptin neurones in the AVPV/PeN, and more so in the ARC, could be

responsive to RFRP-3 signalling. Next, we determined whether the reciprocal relationship might also exist: could kisspeptin neurones also act on *Rfrp* neurones? To address this, we measured the degree of *Kiss1r* co-expression in *Rfrp* neurones and also determined the degree to which kisspeptin fibres contact RFRP-3 neurones in the DMN. Double-label ISH for *Kiss1r* in *Rfrp* neurones in adult male and female mice revealed that essentially all *Rfrp* neurones (> 99%) lacked *Kiss1r* (Fig. 4A). Indeed, *Kiss1r* was surprisingly absent from the DMN, despite previous ISH data demonstrating the high expression of *Kiss1r* in this nucleus (33). As a positive control, pronounced *Kiss1r* mRNA expression was observed in the habenula, a known region of *Kiss1r* expression (not shown). To ensure that the lack of *Kiss1r* in *Rfrp* neurones was not a result of technical reasons, a second set of slides from the rostral hypothalamus of adult females was concurrently assayed for *Kiss1r* expression in *Gnrh* neurones (Fig. 4C,D) along with *Kiss1r* in *Rfrp* neurones. Although > 85% of GnRH neurones expressed *Kiss1r*, no *Rfrp* neurones expressed *Kiss1r* (Fig. 4C,D), consistent with the previous assay.

In the ARC, kisspeptin neurones co-express NKB, which could potentially be used by ARC KNDy neurones to communicate with RFRP-3 neurones via *Tacr3* signalling. Indeed, *Tacr3* (the NKB receptor) is highly expressed in the DMN region, along with NKB fibres (34,44), although it is unknown whether this specifically includes

Fig. 2. Expression of *Gpr147* and *Gpr74* in arcuate nucleus (ARC) KNDy neurones by double-label *in situ* hybridisation. (A) Representative photomicrographs of double-label *in situ* hybridisation of *Tac2* (red fluorescence) and *Gpr147* (silver grains) in the ARC of a di-oestrous female. (B) ARC *Tac2* neurones co-expressing *Gpr147* (green arrowhead) and *Tac2* neurones with no co-expression of *Gpr147* (blue arrows). (C) Quantification of the percentage co-expression of *Gpr147* in *Tac2* neurones between gonadally-intact females (F) and males (M) and gonadectomised (GDX) M and F. There were no significant differences in co-expression between any of the groups. (D) Representative photomicrographs of double-label *in situ* hybridisation of *Tac2* (red fluorescence) and *Gpr74* (silver grains) in the ARC of a di-oestrous female. (E) *Tac2* neurones co-expressing *Gpr74* (green arrowhead) and a *Gpr74* neurone that is not expressing *Tac2* (yellow arrow). (F) Quantification of the percentage co-expression of *Gpr74* in *Tac2* neurones between gonadally-intact females (F) and males (M) and gonadectomised (GDX) M and F. These experimental groups were not statistically different.

RFRP-3 neurones. Using double-label ISH for *Tacr3* and *Rfrp*, we observed robust staining for both mRNAs in the DMN of mice of both sexes (Fig. 4e,f). However, quantitative analysis determined that *Tacr3* mRNA was absent in most *Rfrp* neurones. Less than

10% of *Rfrp* neurones expressed *Tacr3* in gonadally-intact males and females (Fig. 4g), and similar low co-expression levels were quantified in GDX mice of both sexes, with *Tacr3* being detected in only approximately 8% of *Rfrp* neurones (Fig. 4g). There were no

Fig. 3. Representative photomicrograph of RFRP-3 immunoreactive fibre (green fluorescence) in apposition with an arcuate nucleus (ARC) kisspeptin neurone (red fluorescence) in a female mouse. Immunohistochemical analysis revealed RFRP-3 contacts with approximately 35% of kisspeptin cell bodies in the ARC of gonadectomised (GDX) female mice. The figure is a collapsed stack of several confocal optical sections. A green triangle denotes example RFRP-3 fibre immunoreactivity. A red triangle denotes an ARC kisspeptin cell not receiving RFRP-3 input. Yellow triangle denotes RFRP-3 fibre contacting an ARC kisspeptin cell.

statistical differences between sexes or gonadal state in the proportion of cells expressing *Tacr3* or the relative *Tacr3* mRNA level per *Rfrp* cell.

In a complementary experiment, we used double-label IHC to assess potential kisspeptin fibre contacts, which could arise from either the AVPV/PeN kisspeptin population and/or ARC kisspeptin/NKB (KNDy) cells, on RFRP-3 neurones in the DMN. Matching the receptor co-expression data, almost no RFRP-3 cells were observed with contacts from kisspeptin fibres (Fig. 5A). Quantification of the staining revealed that, on average, just 4% of RFRP-3-immunoreactive cells received apparent contacts from kisspeptin-containing fibres in gonadally-intact adult male mice (Fig. 5B). Similar results were observed in GDX mice, with only approximately 3% and 7% of RFRP-3 cells receiving kisspeptin fibre appositions in males and females, respectively (Fig. 5B). There were no statistical differences in the degree of kisspeptin-RFRP-3 contacts between intact and GDX mice or between sexes.

Experiment 4: Is NKB a co-neuropeptide with RFRP-3?

Tac2 mRNA, which codes for NKB, is known to be highly expressed in the DMN (45), although its co-expression in RFRP-3 neurones is unknown. We used double-label ISH to determine whether *Rfrp* neurones are the same DMN cells (or an overlapping population) as those expressing *Tac2*. However, despite strong expression of both genes in the DMN region, we found that *Rfrp* and *Tac2* neurones in the DMN are mostly distinct populations, with relatively low levels of co-expression (Fig. 6A,B). Quantitatively, approximately 12% of *Rfrp* neurones co-express *Tac2* in adult mice of both sexes (Fig. 6C) under both gonadal-intact and GDX conditions. There were no sta-

tistical differences in the proportion of *Rfrp* neurones co-expressing *Tac2* between sexes or treatment group (Fig. 6C) and there were no differences in the grains per cell representing relative *Tac2* mRNA levels in the double-labelled cells (not shown). We did not attempt to quantify the degree of reciprocal co-localisation of DMN *Tac2* neurones expressing *Rfrp*. However, in general, we consistently noted significantly more total *Tac2* neurones than *Rfrp* neurones in the DMN region, indicating that the proportion of *Tac2* cells co-expressing *Rfrp* would be notably lower than the 12% of *Rfrp* neurones found to co-express *Tac2*.

Discussion

Despite the potent and reciprocal activities of kisspeptin and RFRP-3 on the reproductive axis, the comprehensive interconnectivity of these two neuropeptide systems has not been thoroughly investigated. In the present study, we determine whether the receptors for RFRP-3, Gpr147 and Gpr74 were expressed in either population of hypothalamic kisspeptin neurones and whether the kisspeptin receptor, or that for NKB, was expressed in RFRP-3 cells. We found that the majority of AVPV/PeN *Kiss1* neurones do not express either of the receptors known to mediate the actions of RFRP-3, whereas a moderate percentage of kisspeptin cells in the ARC do co-express RFRP-3 receptors. Conversely, *Kiss1r* (kisspeptin receptor) was absent in almost all *Rfrp* neurones and almost no RFRP-3 neurones receive appositions from kisspeptin axonal fibres. Moreover, *Tacr3*, the receptor for ARC kisspeptin's co-transmitter NKB, was not highly expressed in most RFRP-3 cells. Overall, our anatomical data suggest that the kisspeptin and RFRP-3 neuronal systems likely act independently on the GnRH-pituitary axis and may only have notable communication with each other at the level of RFRP-3 signalling to ARC kisspeptin cells.

The various mechanisms by which RFRP-3 neurones might regulate the reproductive axis are not fully known. A good part of the reproductive modulation of RFRP-3 appears to take place through the inhibition of GnRH release, and antagonising the GnRH receptor abolishes the stimulatory effect of an RFRP-3 antagonist, RF9, on LH secretion (28). These data suggest that RFRP-3 provides an inhibitory tone upstream of GnRH signalling because blockade of RFRP-3 signalling is only effective at stimulating LH when GnRH signalling pathways are functional. However, using several different techniques, we previously found only low to moderate co-expression of the RFRP-3 receptors, Gpr147 and Gpr74, in GnRH neurones, with a majority of GnRH cells not expressing either receptor (26,28), matching the finding that only a subset of GnRH neurones changing their firing rate after RFRP-3 treatment (19). The ability of RFRP-3 to inhibit GnRH and LH despite a majority of GnRH cells not expressing RFRP-3 receptors could indicate that some RFRP-3-mediated inhibition on GnRH may occur indirectly. Although the possibility of RFRP-3 acting on the pituitary has been hypothesised, RFRP-3 neurones of rodents are not hypophysiotrophic because they are unable take up peripherally administered retrograde tracers (46). These data exclude the possibility of RFRP-3 acting directly on the pituitary of rodents, which differs from the ovine model, where RFRP-3 can be measured in portal blood (47).

Fig. 4. Expression of *Kiss1r* in *Rfrp* and gonadotrophin-releasing hormone (GnRH) neurons and expression of *Tacr3* in *Rfrp* neurons by double-label *in situ* hybridisation. (A) Representative photomicrographs of double-label *in situ* hybridisation of *Rfrp* (red fluorescence) and *Kiss1r* (silver grains) in the dorsal-medial nucleus (DMN) of a di-oestrous female. *Rfrp* neurons lacking *Kiss1r* are marked with blue arrows. (B) Quantification of the percentage co-expression of *Kiss1r* in *Rfrp* neurons between gonadally-intact females and males. There were no significant differences in co-expression between sexes. (C) Representative photomicrographs of double-label *in situ* hybridisation of *Kiss1r* (silver grains) and GnRH (red fluorescence) in the preoptic area (POA) of an intact male. Green arrowheads identify double-labeled cells. (D) Quantification of the percentage co-expression of *Kiss1r* in *Rfrp* or GnRH neurons in di-oestrous females and intact males (percentages averaged across all animals). (E) Representative photomicrographs of double-label *in situ* hybridisation of *Rfrp* (red fluorescence) and *Tacr3* (silver grains) in the DMN of an intact male mouse. (F) *Rfrp* neurons co-expressing *Tacr3* (green arrowhead) and a cell expressing *Tacr3* without *Rfrp* (yellow arrow). (G) Quantification of the percentage co-expression of *Tacr3* in *Rfrp* neurons between gonadally-intact females (F) and males (M) and gonadectomised (GDX) M and F. There were no significant differences in co-expression between any of the groups.

Fig. 5. Immunohistochemistry for kisspeptin fibres and RFRP-3 cell bodies in the mouse dorsal-medial nucleus (DMN). (A) Representative images of RFRP-3 cell bodies and fibres (green fluorescence) and kisspeptin fibres (red fluorescence) in the DMN of a gonadally-intact male (left) and gonadectomised (GDX) female (right). (B) Quantification of the percentage of RFRP-3 neurones receiving contacts from kisspeptin fibres in the DMN of intact and GDX male and female mice. Almost no RFRP-3 cells were contacted and there was no statistical difference between gonadal states. IR, immunoreactive.

Given that only a subset of GnRH neurones expresses RFRP-3 receptors, we speculated that RFRP-3 may also regulate the GnRH axis through an intermediate neuropeptide population(s), such as kisspeptin neurones. Kisspeptin is a potent stimulator of GnRH release (10), although the 'upstream' circuitry that regulates the synthesis and secretion of kisspeptin is poorly understood. Thus, we hypothesised that RFRP-3 may be an upstream factor that negatively modulates kisspeptin neurones, thereby reducing GnRH activation. This possibility was supported by data indicating that RFRP-3 fibres appose some AVPV/PeN kisspeptin neurones in female mice (28) and are also present in the ARC where kisspeptin neurones also reside (23,24,29). However, based on our present findings, it appears that a large majority of kisspeptin neurones, in both the AVPV/PeN and the ARC, are lacking receptors for RFRP-3. This was especially apparent in the AVPV/PeN, suggesting that kisspeptin neurones in that nucleus are unlikely to be significantly regulated by direct RFRP-3 signalling. In the ARC, however, a moderate proportion (approximately 25%) of kisspeptin cells co-expressed Gpr147 or Gpr74, and almost 35% of ARC kisspeptin neurones receive RFRP-3 fibre contacts, indicating that there could

be some functional regulation of kisspeptin neurones by RFRP-3 in this specific brain region. Even so, it is not clear what the functional significance of such communication would be given the lack of RFRP-3 receptors and fibre contacts in such a large proportion of these ARC KNDy cells. Because we could not perform triple labelling experiments, we do not know whether the same ARC kisspeptin cells that express Gpr147 also express Gpr74, or whether different kisspeptin cells express each of the two RFRP-3 receptors. If the latter scenario is correct, then not only would a larger proportion of ARC kisspeptin cells than what we observed (approximately 25%) actually be responsive to RFRP-3 signals, but also the differing affinities of RFRP-3 for these receptor subtypes might enable graded or differing responses of different kisspeptin cells to the same RFRP-3 stimulus. Although the maximum percentage of ARC kisspeptin neurones directly modulated by RFRP-3 signalling is likely capped at approximately one-third, as a result of the proportion of KNDy neurones with RFRP-3 fibre appositions and RFRP-3 receptors, it remains possible that such RFRP-3 signalling may still affect the entire KNDy population indirectly via the reciprocally interconnected nature of the KNDy neurone network. Of note, we

Fig. 6. Expression *Tac2* in *Rfrp* neurones in the dorsal-medial nucleus (DMN) by double-label *in situ* hybridisation. (A) Representative photomicrographs of double-label *in situ* hybridisation of *Rfrp* (red fluorescence) and *Tac2* (silver grains) in the DMN of a di-oestrous female. (B) Higher magnification of *Rfrp* neurones co-expressing *Tac2* (green arrowhead) and a *Tac2* neurone that is not expressing *Rfrp* (yellow arrow). (C) Quantification of the percentage co-expression of *Tac2* in *Rfrp* neurones between gonadally-intact females (F) and males (M) and gonadectomised (GDX) M and F. These experimental groups were not statistically different.

used *Tac2* expression to represent KNDy neurones because, in our hands, almost all *Tac2* neurones (> 95%) co-express *Kiss1* in the ARC of mice (A. S. Kauffman, unpublished observations). Thus, *Tac2* expression in the mouse ARC faithfully reflects *Kiss1* expressing neurones.

One interesting possible role for RFRP-3-kisspeptin interactions that has been hypothesised is the regulation of the preovulatory GnRH/LH surge, an event driven by kisspeptin and suppressed by RFRP-3 (18,48). RFRP-3 neuronal activity declines at the time of the LH surge (29), as does the hypothalamic concentration of RFRP-3 peptide (M. Z. Rizwan and G. M. Anderson, unpublished data). It is conceivable that this decline reduces inhibitory RFRP-3 tone on kisspeptin neurones, allowing the increased kisspeptin drive to trigger the GnRH/LH surge (15). Such speculation would be consistent with previous reports of RFRP-3 causing the suppression of cellular activity, as well as a reduced kisspeptin neuronal firing rate in the AVPV (18,19), a key brain region implicated in the LH surge event. However, this is less consistent with our present finding of minimal RFRP-3 receptors in AVPV kisspeptin neurones. Indeed, most RFRP-3 receptors in kisspeptin neurones were located in the ARC rather than the AVPV, and the former brain region is not implicated in the LH surge in rodents. Thus, our findings suggest that any effects of RFRP-3 on AVPV kisspeptin neurones to govern the LH surge would likely be indirect on those neurones.

Initial ISH studies targeting *Kiss1r* suggested that it was highly expressed in the DMN (33), and kisspeptin fibres have been observed in the DMN (49), supporting the possibility that kisspeptin signalling may interface with RFRP-3 neurones. Our present results, however, strongly exclude the possibility of kisspeptin acting on RFRP-3 neurones through *Kiss1r* because no *Rfrp* neurones expressed the mRNA for this receptor. By contrast to the initial study (33), we find no evidence of significant *Kiss1r* mRNA in the DMN area, at least under the conditions examined. This was not a result of the poor sensitivity of our *Kiss1r* ISH because we observed high *Kiss1r* expression in GnRH neurones and other brain regions, as expected. To complement these receptor expression data, we also determined whether kisspeptin-containing fibres apposed RFRP-3 cells. In agreement with the ISH data, these immunohistochemistry results also excluded the likelihood of kisspeptin neurones targeting RFRP-3 neurones because almost all RFRP-3 neurones were devoid of kisspeptin fibre appositions. Importantly, these fibre apposition data also indicate that, as a result of the lack of physical connectivity, it is highly unlikely that kisspeptin neurones utilise other co-neuropeptides, such as NKB or dynorphin, to act directly on *Rfrp* neurones. This was supported by our finding that *Tacr3*, the receptor for NKB (a co-transmitter with kisspeptin from ARC cells) was absent in the vast majority of *Rfrp* neurones, despite robust *Tacr3* expression elsewhere nearby in the DMN. Any NKB interaction on

the small subset of RFRP-3 neurones expressing *Tacr3* would likely arise from non-ARC NKB neurones because almost no kisspeptin fibres appose RFRP-3 neurones (ARC KNDy neurone fibres would contain kisspeptin as well as NKB).

Most hypothalamic neuropeptide populations tend to produce more than one neuropeptide or neurotransmitter. Yet, the potential co-transmitters that may also be expressed in and released by RFRP-3 neurones are unknown. In the present study, we therefore also examined whether NKB was co-expressed with RFRP-3 because many neurones in the DMN highly express *Tac2*. However, despite a little degree of overlap, we found that *Tac2* was not expressed in the large majority of *Rfrp* neurones. These data suggest that the RFRP-3 and NKB neuronal populations residing in the DMN are, for the most part, distinct and separate neuropeptide populations. Thus, it presently remains unknown whether RFRP-3 neurones also secrete additional co-transmitters or not.

In summary, the data reported in the present study exclude the likelihood of RFRP-3 acting on any sizable proportion of kisspeptin neurones in the AVPV/PeN in either sex, although they suggest that RFRP-3 may potentially provide some direct regulation to a moderate subset of ARC KNDy cells. Additionally, the possibility of a reciprocal action of direct kisspeptin or NKB on RFRP-3 neurones was strongly excluded because no *Rfrp* neurones express *Kiss1r* and almost no RFRP-3-IR neurones receive kisspeptin appositions or express *Tacr3*. Finally, the majority of *Rfrp* neurones lack *Tac2*, indicating that NKB is not a co-expressed with RFRP-3 and that these are likely two separate neuronal populations in the DMN. These data demonstrate that kisspeptin is acting independently of, and in parallel with, the RFRP-3 system to govern the reproductive axis, whereas some of the effects of RFRP-3 on reproduction may potentially be derived via actions on a subset of ARC kisspeptin cells. Whether RFRP-3 also acts elsewhere in the brain to indirectly modulate reproduction remains unexplored, although this could possibly include other regions such as the paraventricular nucleus, lateral hypothalamus, thalamus and amygdala where RFRP-3 fibres have been reported (24) and where *Gpr147* or *Gpr74* mRNAs are notably expressed (50).

Acknowledgements

This research was supported by NSF grant IOS-1025893 and NIH grant R01 HD065856. Additional support was provided by the Eunice Kennedy Shriver NICHD/NIH (SCCPIR) grant U54-HD012303 (University of California San Diego). The authors thank Sheila J. Semaan and Kristen P. Tolson for their technical assistance and comments on the manuscript.

Received 25 April 2013,
revised 24 July 2013,
accepted 3 August 2013

References

- 1 Tsutsui K, Bentley GE, Kriegsfeld LJ, Osugi T, Seong JY, Vaudry H. Discovery and evolutionary history of gonadotrophin-inhibitory hormone and kisspeptin: new key neuropeptides controlling reproduction. *J Neuroendocrinol* 2010; **22**: 716–727.
- 2 Funes S, Hedrick JA, Vassileva G, Markowitz L, Abbondanzo S, Golovko A, Yang S, Monsma FJ, Gustafson EL. The KiSS-1 receptor GPR54 is essential for the development of the murine reproductive system. *Biochem Biophys Res Commun* 2003; **312**: 1357–1363.
- 3 Seminara SB, Messenger S, Chatzidaki EE, Thresher RR, Acierno JS Jr, Shagoury JK, Bo-Abbas Y, Kuohung W, Schwinof KM, Hendrick AG, Zahn D, Dixon J, Kaiser UB, Slaugenhaupt SA, Gusella JF, O'Rahilly S, Carlton MB, Crowley WF Jr, Aparicio SA, Colledge WH. The GPR54 gene as a regulator of puberty. *N Engl J Med* 2003; **349**: 1614–1627.
- 4 de Roux N, Genin E, Carel JC, Matsuda F, Chaussain JL, Milgrom E. Hypogonadotropic hypogonadism due to loss of function of the KiSS1-derived peptide receptor GPR54. *Proc Natl Acad Sci USA* 2003; **100**: 10972–10976.
- 5 Messenger S, Chatzidaki EE, Ma D, Hendrick AG, Zahn D, Dixon J, Thresher RR, Malinge I, Lomet D, Carlton MB, Colledge WH, Caraty A, Aparicio SA. Kisspeptin directly stimulates gonadotropin-releasing hormone release via G protein-coupled receptor 54. *Proc Natl Acad Sci USA* 2005; **102**: 1761–1766.
- 6 Kauffman AS, Park JH, McPhie-Lalmansingh AA, Gottsch ML, Bodo C, Hohmann JG, Pavlova MN, Rohde AD, Clifton DK, Steiner RA, Rissman EF. The kisspeptin receptor GPR54 is required for sexual differentiation of the brain and behavior. *J Neurosci* 2007; **27**: 8826–8835.
- 7 Kinoshita M, Tsukamura H, Adachi S, Matsui H, Uenoyama Y, Iwata K, Yamada S, Inoue K, Ohtaki T, Matsumoto H, Maeda K. Involvement of central metastin in the regulation of preovulatory luteinizing hormone surge and estrous cyclicity in female rats. *Endocrinology* 2005; **146**: 4431–4436.
- 8 Navarro VM, Castellano JM, Fernandez-Fernandez R, Tovar S, Roa J, Mayen A, Barreiro ML, Casanueva FF, Aguilar E, Dieguez C, Pinilla L, Tena-Sempere M. Effects of KiSS-1 peptide, the natural ligand of GPR54, on follicle-stimulating hormone secretion in the rat. *Endocrinology* 2005; **146**: 1689–1697.
- 9 Navarro VM, Castellano JM, Fernandez-Fernandez R, Tovar S, Roa J, Mayen A, Nogueiras R, Vazquez MJ, Barreiro ML, Magni P, Aguilar E, Dieguez C, Pinilla L, Tena-Sempere M. Characterization of the potent luteinizing hormone-releasing activity of KiSS-1 peptide, the natural ligand of GPR54. *Endocrinology* 2005; **146**: 156–163.
- 10 Gottsch ML, Cunningham MJ, Smith JT, Popa SM, Acohido BV, Crowley WF, Seminara S, Clifton DK, Steiner RA. A role for kisspeptins in the regulation of gonadotropin secretion in the mouse. *Endocrinology* 2004; **145**: 4073–4077.
- 11 Irwig MS, Fraley GS, Smith JT, Acohido BV, Popa SM, Cunningham MJ, Gottsch ML, Clifton DK, Steiner RA. Kisspeptin activation of gonadotropin-releasing hormone neurons and regulation of KiSS-1 mRNA in the male rat. *Neuroendocrinology* 2004; **80**: 264–272.
- 12 Han SK, Gottsch ML, Lee KJ, Popa SM, Smith JT, Jakawich SK, Clifton DK, Steiner RA, Herbison AE. Activation of gonadotropin-releasing hormone neurons by kisspeptin as a neuroendocrine switch for the onset of puberty. *J Neurosci* 2005; **25**: 11349–11356.
- 13 Pielecka-Fortuna J, Chu Z, Moenter SM. Kisspeptin acts directly and indirectly to increase gonadotropin-releasing hormone neuron activity and its effects are modulated by estradiol. *Endocrinology* 2008; **149**: 1979–1986.
- 14 Clarkson J, Herbison AE. Postnatal development of kisspeptin neurons in mouse hypothalamus: sexual dimorphism and projections to gonadotropin-releasing hormone neurons. *Endocrinology* 2006; **147**: 5817–5825.
- 15 Yeo SH, Herbison AE. Projections of arcuate nucleus and rostral periventricular kisspeptin neurons in the adult female mouse brain. *Endocrinology* 2011; **152**: 2387–2399.
- 16 Lehman MN, Merkley CM, Coolen LM, Goodman RL. Anatomy of the kisspeptin neural network in mammals. *Brain Res* 2010; **1364**: 90–102.

- 17 Navarro VM, Gottsch ML, Chavkin C, Okamura H, Clifton DK, Steiner RA. Regulation of GnRH secretion by kisspeptin/dynorphin/neurokinin B neurons in the arcuate nucleus of the mouse. *J Neurosci* 2009; **29**: 11859–11866.
- 18 Anderson GM, Relf HL, Rizwan MZ, Evans JJ. Central and peripheral effects of RFamide-related peptide-3 on luteinizing hormone and prolactin secretion in rats. *Endocrinology* 2009; **150**: 1834–1840.
- 19 Ducret E, Anderson GM, Herbison AE. RFamide-related peptide-3, a mammalian gonadotropin-inhibitory hormone ortholog, regulates gonadotropin-releasing hormone neuron firing in the mouse. *Endocrinology* 2009; **150**: 2799–2804.
- 20 Wu M, Dumalska I, Morozova E, van den Pol AN, Alreja M. Gonadotropin inhibitory hormone inhibits basal forebrain vGluT2-gonadotropin-releasing hormone neurons via a direct postsynaptic mechanism. *J Physiol* 2009; **587**: 1401–1411.
- 21 Hinuma S, Shintani Y, Fukusumi S, Iijima N, Matsumoto Y, Hosoya M, Fujii R, Watanabe T, Kikuchi K, Terao Y, Yano T, Yamamoto T, Kawamata Y, Habata Y, Asada M, Kitada C, Kurokawa T, Onda H, Nishimura O, Tanaka M, Ibata Y, Fujino M. New neuropeptides containing carboxy-terminal RFamide and their receptor in mammals. *Nat Cell Biol* 2000; **2**: 703–708.
- 22 Tsutsui K, Saigoh E, Ukena K, Teranishi H, Fujisawa Y, Kikuchi M, Ishii S, Sharp PJ. A novel avian hypothalamic peptide inhibiting gonadotropin release. *Biochem Biophys Res Commun* 2000; **275**: 661–667.
- 23 Kriegsfeld LJ, Mei DF, Bentley GE, Ubuka T, Mason AO, Inoue K, Ukena K, Tsutsui K, Silver R. Identification and characterization of a gonadotropin-inhibitory system in the brains of mammals. *Proc Natl Acad Sci USA* 2006; **103**: 2410–2415.
- 24 Yano T, Iijima N, Kahirah K, Hinuma S, Tanaka M, Ibata Y. Localization and neuronal response of RFamide related peptides in the rat central nervous system. *Brain Res* 2003; **982**: 156–167.
- 25 Liu Q, Guan XM, Martin WJ, McDonald TP, Clements MK, Jiang Q, Zeng Z, Jacobson M, Williams DL Jr, Yu H, Bomford D, Figueroa D, Mallee J, Wang R, Evans J, Gould R, Austin CP. Identification and characterization of novel mammalian neuropeptide FF-like peptides that attenuate morphine-induced antinociception. *J Biol Chem* 2001; **276**: 36961–36969.
- 26 Poling MC, Kim J, Dhamija S, Kauffman AS. Development, sex steroid regulation, and phenotypic characterization of RFamide-related peptide (Rfrp) gene expression and RFamide receptors in the mouse hypothalamus. *Endocrinology* 2012; **153**: 1827–1840.
- 27 Johnson MA, Tsutsui K, Fraley GS. Rat RFamide-related peptide-3 stimulates GH secretion, inhibits LH secretion, and has variable effects on sex behavior in the adult male rat. *Horm Behav* 2007; **51**: 171–180.
- 28 Rizwan MZ, Poling MC, Corr M, Cornes PA, Augustine RA, Quennell JH, Kauffman AS, Anderson GM. RFamide-related peptide-3 receptor gene expression in GnRH and kisspeptin neurons and GnRH-dependent mechanism of action. *Endocrinology* 2012; **153**: 3770–3779.
- 29 Gibson EM, Humber SA, Jain S, Williams WP III, Zhao S, Bentley GE, Tsutsui K, Kriegsfeld LJ. Alterations in RFamide-related peptide expression are coordinated with the preovulatory luteinizing hormone surge. *Endocrinology* 2008; **149**: 4958–4969.
- 30 Quennell JH, Rizwan MZ, Relf HL, Anderson GM. Developmental and steroidogenic effects on the gene expression of RFamide related peptides and their receptor in the rat brain and pituitary gland. *J Neuroendocrinol* 2010; **22**: 309–316.
- 31 Son YL, Ubuka T, Millar RP, Kanasaki H, Tsutsui K. Gonadotropin-inhibitory hormone inhibits GnRH-induced gonadotropin subunit gene transcriptions by inhibiting AC/cAMP/PKA-dependent ERK pathway in LbetaT2 cells. *Endocrinology* 2012; **153**: 2332–2343.
- 32 Fu LY, van den Pol AN. Kisspeptin directly excites anorexigenic proopiomelanocortin neurons but inhibits orexigenic neuropeptide Y cells by an indirect synaptic mechanism. *J Neurosci* 2010; **30**: 10205–10219.
- 33 Lee DK, Nguyen T, O'Neill GP, Cheng R, Liu Y, Howard AD, Coulombe N, Tan CP, Tang-Nguyen AT, George SR, O'Dowd BF. Discovery of a receptor related to the galanin receptors. *FEBS Lett* 1999; **446**: 103–107.
- 34 Sunkin SM, Ng L, Lau C, Dolbeare T, Gilbert TL, Thompson CL, Hawrylycz M, Dang C. Allen Brain Atlas: an integrated spatio-temporal portal for exploring the central nervous system. *Nucleic Acids Res* 2013; In Press.
- 35 Smith JT, Cunningham MJ, Rissman EF, Clifton DK, Steiner RA. Regulation of Kiss1 gene expression in the brain of the female mouse. *Endocrinology* 2005; **146**: 3686–3692.
- 36 Smith JT, Dungan HM, Stoll EA, Gottsch ML, Braun RE, Eacker SM, Clifton DK, Steiner RA. Differential regulation of Kiss-1 mRNA expression by sex steroids in the brain of the male mouse. *Endocrinology* 2005; **146**: 2976–2984.
- 37 Kauffman AS, Navarro VM, Kim J, Clifton D, Steiner RA. Sex differences in the regulation of Kiss1/NKB neurons in juvenile mice: implications for the timing of puberty. *Am J Physiol Endocrinol Metab* 2009; **297**: E1212–E1221.
- 38 Finn PD, Steiner RA, Clifton DK. Temporal patterns of gonadotropin-releasing hormone (GnRH), c-fos, and galanin gene expression in GnRH neurons relative to the luteinizing hormone surge in the rat. *J Neurosci* 1998; **18**: 713–719.
- 39 Poling MC, Kauffman AS. Sexually dimorphic testosterone secretion in prenatal and neonatal mice is independent of kisspeptin-kiss1r and GnRH signaling. *Endocrinology* 2012; **153**: 782–793.
- 40 Chowen JA, Clifton DK. Semiquantitative analysis of cellular somatostatin mRNA levels by *in situ* hybridization histochemistry. *Methods Neurosci* 1991; **5**: 137–158.
- 41 Kauffman AS, Gottsch ML, Roa J, Byquist AC, Crown A, Clifton DK, Hoffman GE, Steiner RA, Tena-Sempere M. Sexual differentiation of Kiss1 gene expression in the brain of the rat. *Endocrinology* 2007; **148**: 1774–1783.
- 42 Rance NE, Bruce TR. Neurokinin B gene expression is increased in the arcuate nucleus of ovariectomized rats. *Neuroendocrinology* 1994; **60**: 337–345.
- 43 Navarro VM, Castellano JM, McConkey SM, Pineda R, Ruiz-Pino F, Pinilla L, Clifton DK, Tena-Sempere M, Steiner RA. Interactions between kisspeptin and neurokinin B in the control of GnRH secretion in the female rat. *Am J Physiol Endocrinol Metab* 2011; **300**: E202–E210.
- 44 Krajewski SJ, Burke MC, Anderson MJ, McMullen NT, Rance NE. Forebrain projections of arcuate neurokinin B neurons demonstrated by anterograde tract-tracing and monosodium glutamate lesions in the rat. *Neuroscience* 2010; **166**: 680–697.
- 45 Marksteiner J, Sperk G, Krause JE. Distribution of neurons expressing neurokinin B in the rat brain: immunohistochemistry and *in situ* hybridization. *J Comp Neurol* 1992; **317**: 341–356.
- 46 Rizwan MZ, Porteous R, Herbison AE, Anderson GM. Cells expressing RFamide-related peptide-1/3, the mammalian gonadotropin-inhibitory hormone orthologs, are not hypophysiotropic neuroendocrine neurons in the rat. *Endocrinology* 2009; **150**: 1413–1420.
- 47 Smith JT, Young IR, Veldhuis JD, Clarke IJ. Gonadotropin-inhibitory hormone (GnIH) secretion into the ovine hypophyseal portal system. *Endocrinology* 2012; **153**: 3368–3375.
- 48 Khan AR, Kauffman AS. The role of kisspeptin and RFRP-3 neurons in the circadian-timed preovulatory luteinizing hormone surge. *J Neuroendocrinol* 2012; **24**: 131–143.
- 49 Clarkson J, d'Anglemont de Tassigny X, Colledge WH, Caraty A, Herbison AE. Distribution of kisspeptin neurones in the adult female mouse brain. *J Neuroendocrinol* 2009; **21**: 673–682.
- 50 Gouarderes C, Puget A, Zajac JM. Detailed distribution of neuropeptide FF receptors (NPFF1 and NPFF2) in the rat, mouse, octodon, rabbit, guinea pig, and marmoset monkey brains: a comparative autoradiographic study. *Synapse* 2004; **51**: 249–269.