

UC Davis

UC Davis Previously Published Works

Title

Binding of a fluorescence reporter and a ligand to an odorant-binding protein of the yellow fever mosquito, *Aedes aegypti*

Permalink

<https://escholarship.org/uc/item/33q037v9>

Authors

Leal, Gabriel M

Leal, Walter S

Publication Date

2014

DOI

10.12688/f1000research.5879.2

Peer reviewed



RESEARCH NOTE

REVISED

Binding of a fluorescence reporter and a ligand to an odorant-binding protein of the yellow fever mosquito, *Aedes aegypti* [v2; ref status: indexed, <http://f1000r.es/4yp>]

Gabriel M. Leal^{1,2}, Walter S. Leal¹¹Department of Molecular and Cellular Biology, University of California, Davis, Davis, CA, 95616, USA²Davis Senior High School, Davis, CA, 95616, USA

v2 First published: 12 Dec 2014, 3:305 (doi: [10.12688/f1000research.5879.1](https://doi.org/10.12688/f1000research.5879.1))
 Latest published: 09 Jan 2015, 3:305 (doi: [10.12688/f1000research.5879.2](https://doi.org/10.12688/f1000research.5879.2))

Abstract

Odorant-binding proteins (OBPs), also named pheromone-binding proteins when the odorant is a pheromone, are essential for insect olfaction. They solubilize odorants that reach the port of entry of the olfactory system, the pore tubules in antennae and other olfactory appendages. Then, OBPs transport these hydrophobic compounds through an aqueous sensillar lymph to receptors embedded on dendritic membranes of olfactory receptor neurons. Structures of OBPs from mosquito species have shed new light on the mechanism of transport, although there is considerable debate on how they deliver odorant to receptors. An OBP from the southern house mosquito, *Culex quinquefasciatus*, binds the hydrophobic moiety of a mosquito oviposition pheromone (MOP) on the edge of its binding cavity. Likewise, it has been demonstrated that the orthologous protein from the malaria mosquito binds the insect repellent DEET on a similar edge of its binding pocket. A high school research project was aimed at testing whether the orthologous protein from the yellow fever mosquito, AegOBP1, binds DEET and other insect repellents, and MOP was used as a positive control. Binding assays using the fluorescence reporter N-phenyl-1-naphthylamine (NPN) were inconclusive. However, titration of NPN fluorescence emission in AegOBP1 solution with MOP led to unexpected and intriguing results. Quenching was observed in the initial phase of titration, but addition of higher doses of MOP led to a stepwise increase in fluorescence emission coupled with a blue shift, which can be explained at least in part by formation of MOP micelles to house stray NPN molecules.

Open Peer Review

Referee Status:

Invited Referees

1 2

REVISED

version 2

published
09 Jan 2015

report



version 1

published
12 Dec 2014

report



report

- 1 **Kostas Iatrou**, National Center for Scientific Research "Demokritos" Greece,
Spyros Zographos, National Hellenic Research Foundation Greece
Spyros Zographos, National Hellenic Research Foundation Greece
- 2 **Paolo Pelosi**, University of Pisa Italy

Discuss this article

Comments (0)

Corresponding author: Walter S. Leal (wsleal@ucdavis.edu)

How to cite this article: Leal GM and Leal WS. **Binding of a fluorescence reporter and a ligand to an odorant-binding protein of the yellow fever mosquito, *Aedes aegypti*** [v2; ref status: indexed, <http://f1000r.es/4yp>] *F1000Research* 2015, 3:305 (doi: [10.12688/f1000research.5879.2](https://doi.org/10.12688/f1000research.5879.2))

Copyright: © 2015 Leal GM and Leal WS. This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Grant information: This work was supported in part from unrestricted gifts from Fuji Flavor Co. and Bedoukian Research Inc.

Competing interests: No competing interests were disclosed.

First published: 12 Dec 2014, 3:305 (doi: [10.12688/f1000research.5879.1](https://doi.org/10.12688/f1000research.5879.1))

First indexed: 22 Jan 2015, 3:305 (doi: [10.12688/f1000research.5879.2](https://doi.org/10.12688/f1000research.5879.2))

REVISED Amendments from Version 1

In the revised version of this article we included: (1) citation of a paper by Sun *et al.* (2012) which showed a similar increase in fluorescence during titration with high doses of (E)- β -farnesene, (2) new data regarding titration of NPN fluorescence with MOP in ammonium acetate buffer, pH 5.5 (see Figure 5C), and (3) a short statement how low affinity of mosquito OBPs for DEET could explain, at least in part, the need for applications of high doses of DEET.

See referee reports

Introduction

Over the past decade progress towards our understanding of the molecular basis of mosquito olfaction has been remarkable. It was not until the sunset of last century that odorant receptor (OR) genes have been identified in the genome of the fruit fly, *Drosophila melanogaster*¹⁻³ and thereafter in mosquitoes and various insect species (see review⁴), and less than a decade since the unique topology of ORs, with an intracellular N-terminus and an extracellular C-terminus⁵, has been elucidated. Although previously known from moth species⁶, it was about a decade ago that the first odorant-binding proteins (OBPs) from mosquitoes have been isolated and identified⁷. By now the complete repertoire of olfactory genes, including *OBP*, *OR* and *ionotropic receptor (IR)* genes, have been identified in the three major mosquito species: the yellow fever mosquito, *Aedes aegypti*⁸, the malaria mosquito, *Anopheles gambiae*⁹, and the southern house mosquito, *Culex quinquefasciatus*¹⁰. There is growing evidence in the literature that OBPs and ORs play a crucial role in the sensitivity and selectivity of the insect's olfactory system⁴. Mosquito ORs have been deorphanized and demonstrated to be essential for the reception of physiologically and behaviorally relevant odorants^{9,11}, including oviposition attractants¹²⁻¹⁴, insect repellents¹⁵ and a signature compound (sulcatone) for human host preference¹⁶. Elucidation of the three-dimensional (3D) structures of mosquito OBPs¹⁷⁻²¹ along with knockdown experiments^{22,23} and binding assays²⁴⁻²⁷ strongly suggest that these olfactory proteins are involved in the transport of odorant from the ports of entry of olfactory sensilla (the pore tubules) to ORs housed on dendritic membranes of olfactory receptor neurons.

There are typically two binding assays to “de-orphanize” OBPs, i.e., to measure their binding affinities and specificity towards physiologically and behaviorally relevant odorants (ligands). They are the cold binding assay²⁸ so named because – as opposed to its predecessors – it does not require radioactive ligands and a fluorescence reporter assay^{29,30}. The former is based on separation of bound and unbound OBPs, followed by extraction of bound ligands and their quantification by gas chromatography. In the latter a test OBP is bound to a fluorescence reporter, *N*-phenyl-1-naphthylamine (NPN, Figure 1), and subsequently increasing amounts of a test ligand are added. Decreasing NPN fluorescence emission is inferred as NPN displacement, i.e., the test ligand is assumed to compete for the binding site initially occupied by NPN. The fluorescence reporter assay is such a facile method that we envisioned it could be used even in a high school research project.





Figure 1. Structures of a fluorescence reporter and a mosquito oviposition pheromone. *N*-phenyl-1-naphthylamine (NPN) is widely used in binding assays with insect OBPs. (5*R*,6*S*)-6-acetoxy-5-hexadecanolide (MOP) is an attractant first isolated from eggs of *Cx. quinquefasciatus*³⁷, but it is known to bind not only to CquiOBP1, but also to its orthologous proteins, i.e., AaegOBP1 and AgamOBP1¹⁹.

The 3D structures of the malaria mosquito OBP, AgamOBP1²¹ bound to polyethylene glycol (PEG) and AgamOBP1 complex with DEET¹⁸, suggested that AgamOBP1 could be a DEET carrier. For this high school project we asked the question whether DEET and other insect repellents (picaridin, IR3535, and PMD) would bind to AaegOBP1³¹ (also named AaegOBP39^{32,33}), an orthologue of AgamOBP1 from the yellow fever mosquito with similar 3D structure²⁰. In the course of this investigation, we found evidence suggesting that AaegOBP1 might bind simultaneously the fluorescence reporter and an odorant.

Materials and methods

Protein preparations

AaegOBP1 (AY189223)³¹ was expressed in LB medium with transformed BL21(DE3) cell (Agilent Technologies, Santa Clara, CA) according to a protocol for periplasmic expression of insect OBPs³⁴. Proteins were extracted with 10 mM Tris-HCl, pH 8 by three cycles of freeze and thaw³⁵. After centrifuging at 16,000×g to remove debris, AaegOBP1 was isolated from the supernatant and purified by a series of ion-exchange and gel filtration chromatographic steps, as previously described²⁰. The purest fractions were combined and desalted, according to a previous protocol²⁰. Then, AaegOBP1 was delipidated following an earlier protocol³⁶ with small modifications. In short, hydroxyalkoxypropyl-dextran Type VI resin (H2658, Sigma, St. Louis, MI) (1g) was suspended in HPLC grade methanol (20 ml), transferred to a glass column (i.d., 8.5 mm) with a stopper, washed with 60 ml of methanol and then washed and finally equilibrated with 50 mM citric acid buffer, pH 4.5. AaegOBP1

(ca. 2 mg per batch) in 50 mM citric acid buffer, pH 4.5 was mixed with the equilibrated resin in a 15 ml Falcon tube, and incubated at room temperature in a high speed rotating extractor (Taitec, Tokyo, Japan) at 50 rpm. The mixture was then transferred to a glass column and AegOBP1 was eluted with citric acid buffer and analyzed by SDS-gel electrophoresis. The purest fractions were desalted on four 5-ml HiTrap desalting columns (GE Healthcare Life Sciences) in tandem by using water as mobile phase. Protein concentration was measured by the Quick Start Bradford Protein Assay (Bio-Rad, Hercules, CA).

Fluorescence assays

Fluorescence measurements were done on a RF-5301 spectrofluorophotometer (Shimadzu, Kyoto, Japan) equipped with a magnetic stir bar. Samples in a 2-ml cell were excited at 337 nm, with the emission spectra recorded from 350 to 500 nm. Both emission and excitation slit were set a 5 nm. Data were recorded in high sensitivity, with automatic response time, fast scan speed, and sample pitch of 1 nm. AegOBP1 samples (10 µg/ml; ca. 0.7 µM, unless otherwise specified) were prepared in 100 mM ammonium acetate buffers. NPN titration were performed with acetate buffers pH 5.5 or pH 7. The other experiments, unless otherwise indicated, were done with acetate buffer pH 7. The fluorescence reporter and ligands were added by 0.5 or 1 µl aliquots of 1, 5, or 10 mM solutions in methanol. For displacement assays, 1 µl of 10 mM NPN (unless otherwise specified) was added, the solution was stirred in the cell for at least 10 min, stirring was ceased and spectra recorded. Then one aliquot of the test ligand was added, mixed for 2 min, and then the spectra were recorded. For NPN titration, the protein sample was stirred for 2 min, spectra recorded, 0.5 or 1 µl of 1 mM NPN solution was added and stirred for 2 min before recording. To avoid possible interferences, the light path was open only during recording and stirring was ceased at least 10 s before spectra were acquired.

Data were analyzed with GraphPad Prism 6 (La Jolla, CA). For clarity, traces were reconstructed with GraphPad by transferring recorded data without normalization. To draw Figure 4, data were normalized (fluorescence recorded with AegOBP1 and NPN, 100%) and for each concentration of the ligand mean ± SEM from three experiments were calculated in an Excel datasheet and transferred into Prism. Dissociation constants for NPN were determined by nonlinear regression curve fitting, one site and specific binding. MOP dissociation constant was calculated by measuring its competition for NPN binding. Thus, data were analyzed by nonlinear regression curve fitting (one site fits K_i), using the concentration of NPN (typically 5000 nM as HotNM) and K_d for NPN in NM (HotKdNM).

Chemicals

NPN and DEET (*N,N*-diethyl-3-methylbenzamide) were acquired from Sigma-Aldrich. MOP and PMD (*p*-mentan-3,8-diol) were gifts from Bedoukian Research, Inc. Picaridin (butan-2-yl 2-(2-hydroxyethyl)piperidine-1-carboxylate) and IR3535 (ethyl 3-[acetyl(butyl)amino]propanoate) were gifts from Dr. Kamal Chauhan (USA, ARS, Beltsville).

Results and discussion

Dataset 1. Update 1. Fluorescence reporter assay data with assessing binding of insect repellents to the yellow fever mosquito (*Culex quinquefasciatus*) odorant binding protein AegOBP1

<http://dx.doi.org/10.5256/f1000research.5879.d41724>

Fluorescence reporter was *N*-phenyl-1-naphthylamine (NPN). Insect repellents used were DEET, PMD, Picaridin and IR3535. Mosquito oviposition pheromone was used as a positive control. Please see ReadMe file for details regarding each file. Please see the associated article for methods. The raw data for Figure 5C has been added in this version.

Binding assays with insect repellents

In preparation for binding assays of AegOBP1 with insect repellents, we first measured the dissociation constant, K_d , for NPN: $3.31 \pm 0.48 \mu\text{M}$ ($n = 3$). Subsequently, we measured fluorescence quenching by adding aliquots of insect repellents to a protein solution pre-equilibrated with 5 µM of NPN. To minimize solvent effect and reduce experimental error, we added 0.5 µl of 5 mM solutions of test ligands using a 2 µl pipette. As a positive control, we used a racemic solution of the mosquito oviposition pheromone (*5R,6S*)-6-acetoxy-5-hexadecanolide (MOP)³⁷ (Figure 1), which has been previously demonstrated with the cold binding assay to bind to AegOBP1 with apparently high affinity¹⁹. Titration with DEET showed minor reduction in fluorescence intensity (Figure 2) thus suggesting weak binding. By contrast, addition of 1.25 µM MOP led to almost one-third




Figure 2. NPN fluorescence emission spectra. NPN bound to AegOBP1 was excited at 337 nm and its emission spectra (black trace) was recorded. Then, increasing doses of DEET were added and finally one aliquot of MOP was added.

reduction in fluorescence intensity. Titration with other commercially available insect repellents, namely, picaridin, IR3535, and PMD gave similar results as DEET. Although our results suggest that all four repellents bound to AaegOBP1, it seems their affinities were too low to accurately measure dissociation constants. To complete the project and allow the high school investigator to measure at least one dissociation constant, we titrated MOP and this experiment led to unexpected and interesting results.

Evidence for micelle formation

Addition of MOP to solutions of AaegOBP1 pre-incubated with NPN caused a stepwise decrease in fluorescence intensity (2.5 μM to 10–12.5 μM doses), but rather than saturation further addition of MOP led to fluorescence increase and a blue shift. The senior investigator assumed it was an experimental error and repeated the

experiments (Figure 3). Quenching was observed when MOP was added up to 10–12.5 μM , but fluorescence increased thereafter and the maxima excitation wavelength shifted: AaegOBP1-NPN only, max 445 nm; AaegOBP1-NPN plus 2.5 μM MOP, 449 nm; AaegOBP1-NPN plus 20 μM MOP, 433 nm. Similar increase in fluorescence has been previously observed with high concentrations of (*E*)- β -farnesene when titrating NPN fluorescence in the presence of an aphid OBP. Although unlikely, we tested in our case whether this unexpected fluorescence emission could be generated by MOP itself when bound to AaegOBP1³⁸. The fluorescence emission levels generated even with AaegOBP1 plus 20 μM MOP (highest dose and no NPN) were indeed too low (Figure 3) to explain the overall increase in fluorescence. We repeated these experiments and observed a clear U-shape curve with a minimum at 10–12.5 μM (Figure 4). We measured the dissociation constant for MOP ($2.64 \pm 0.16 \mu\text{M}$, $n = 3$)




Figure 3. Binding of MOP to AaegOBP1. Following addition of NPN, fluorescence emission spectra were recorded with increasing doses of MOP. Note the decrease in fluorescence intensity (quenching) as the doses increases up to 10 μM and an increase in fluorescence and blue shift at higher doses. In a separate experiment, included in the lower part of the figure for comparison, fluorescence emission spectra were recorded with AaegOBP1 alone and after addition of MOP, but in the absence of NPN.




Figure 4. Effect of MOP on fluorescence emission of NPN bound to AaegOBP1. Emission maxima were normalized to display mean \pm SEM from three experiments. MOP dissociation constant was calculated for the decreasing phase (0–12.5 μM). Note the increase in fluorescence emission thereafter.

by considering only the first phase of the curve, i.e., by using the data generated by quenching or NPN replacement. Although the above experiments were conducted with reasonable low concentrations of ligands as compared to typical experiments^{29,30}, we next examined the possibility of micelle formation with higher doses of MOP. We repeated titration of MOP using the same doses of the ligand, but reducing the concentrations of protein (0.35 μM) and fluorescence reporter (NPN, 2.5 μM) (Figure 5). When added to ammonium acetate buffer at pH 7 (Figure 5B) or AegOBP1 in the same buffer (Figure 5A), NPN fluoresced with emission maxima at 469 and 446 nm, respectively. Addition of MOP (2.5–10 μM) led

to quenching of NPN in protein solution, but no significant change of NPN fluorescence in buffer solution. Addition of higher doses of MOP to a buffer solution, however, suggested the formation of micelles given the increase in fluorescence and blue shift observed at 12.5 and 15 μM of MOP at pH 7 (Figure 5B) and at 15 and 17.5 μM at pH 5.5 (Figure 5C), although we do not know the critical micelle concentration for MOP. The increase in fluorescence and blue shift were more pronounced in the presence of protein (Figure 5A). It is, therefore, possible that the increase in fluorescence is a combination of micelle formation and other factor(s), which cannot be dissected by these experiments.




Figure 5. Titration of NPN fluorescence emission with MOP. (A) NPN (2.5 μM) was added to a solution of AegOBP1 (0.35 μM) in ammonium acetate buffer, pH 7. NPN (2.5 μM) was added to ammonium acetate buffer, (B) pH 7 or (C) pH 5.5.

Lastly, we compared the fluorescence emission spectra obtained by titrating AaegOBP1 solutions at low and high pH values (Figure 6). Interestingly, NPN showed a higher affinity for AaegOBP1 at pH 5.5 than at pH 7. Additionally, the emission spectra at low pH were blue shifted relative to pH 7 thus suggesting that at low pH NPN is accommodated in a more hydrophobic environment. It has been previously demonstrated that AaegOBP1 undergoes a pH-dependent conformational change. Although AaegOBP1 does not bind MOP at low pH, it has higher affinity for the fluorescence reporter: $K_d = 1.07 \pm 0.15 \mu\text{M}$, pH 5.5; $K_d = 3.31 \pm 0.48 \mu\text{M}$, pH 7. Lack of binding to odorants at low pH has been observed with the *Culex*

orthologous protein, CquiOBP1²⁴ and other OBPs, but insect fatty carriers bind ligands at low and high pH values³⁹.

Conclusion

A clear mechanistic explanation for the findings reported here must await further structural experimental data, particularly elucidation of crystal structures of AaegOBP1 bound to MOP and NPN separately as well as simultaneously. There are currently five structures of mosquito OBP1s deposited in Protein Data Bank (PDB), namely, AgamOBP1-PEG (PDB entry, 2ERB)²¹ (Figure 7A,B), AaegOBP1-PEG (3K1E)²⁰, CquiOBP1-MOP (NMR, 2L2C; crystal,




Figure 6. NPN fluorescence emission spectra obtained by titration at two pH values. Emission spectra at pH 5.5 (top traces) were considerably blue shifted relative to pH 7 (lower traces). Fluorescence intensity was also relatively higher at lower pH.




Figure 7. Structures of AaegOBP1 and CquiOBP1 bound to PEG and MOP, respectively. (A and C) Hydrophobicity surfaces of AaegOBP1 and CquiOBP1. (B and D) Ribbon displays of the same structures. A potential secondary binding site for MOP is highlighted with circles. It is occupied by PEG in AaegOBP1 but “empty” in CquiOBP1. The central cavity is highlighted in (D) with a dashed circle and shows that only the polar head (lactone moiety) of MOP is housed in the core of the protein. Figure prepared with UCSF Chimera software.

30GN)¹⁹ (Figure 7C,D), AgamOBP1-DEET (3N7H)¹⁸, AgamOBP1-sulcatone (4FQT)¹⁷. Unfortunately, the only OBP-NPN complex (3SOB)⁴⁰ deposited in PDB is for an OBP from the European honey bee, AmelOBP14, which differs from classical OBPs for having two, instead of three, disulfide bridges. Here, NPN is bound in the central cavity of the protein. In CquiOBP1, MOP (Figure 1) has its long lipid tail bound to a hydrophobic tunnel formed between helices 4 and 5 (Figure 7D) and only its lactone/ acetyl ester polar moiety is accommodated in part of the central cavity (Figure 7D, dashed circle). It is, therefore, feasible that MOP and NPN were bound simultaneously, and given the vicinity between the two ligands MOP could cause quenching of NPN fluorescence. It has been shown that in AgamOBP1 DEET is localized at the edge of the binding pocket in the equivalent hydrophobic tunnel that accommodates the lipid tail of MOP in CquiOBP1 (Figure 7D). Providing that NPN would bind in the central cavity, as in AmelOBP14, the distance between DEET and NPN would prevent quenching and, therefore, the “lack of binding” suggested by DEET titration (Figure 2) might be interpreted with caution. If indeed mosquito OBPs have low affinity for DEET, it may explain, at least in part, the need to apply high doses of insect repellents. The unusual increase in fluorescence observed here might be explained at least in part by micelle formation. Unbound NPN, either displaced from AaegOBP1 or remaining in solution, could be housed in MOP-derived micelles and in this hydrophobic environment a blue shift and fluorescence increase are expected. It is also conceivable that at higher doses of MOP a second molecule of this ligand binds to AaegOBP1. There is another hydrophobic moiety bordered by helices $\alpha 1$ and $\alpha 4$ and occupied by PEG in the “apo-AgamOBP1”, which could possibly accommodate another ligand (Figure 7, highlighted with circles). If so, NPN could be accommodated in a more hydrophobic environment thus causing a blue shift and additional increase in fluorescence. This change in NPN environment could be triggered by a conformational change. Of notice, NPN fluorescence emission was blue shifted at acidic pH (5.5) compared to neutral pH (7) (Figure 6). Thus in the acidic conformation of AaegOBP1 NPN was more protected from the solvent, i.e., it is likely to be localized in a more hydrophobic environment. Previously, we have observed binding of two ligands to an insect OBP. The pheromone-binding protein from the

silkmoth moth, *Bombyx mori*, has been crystallized with two molecules of the bell pepper odorant, 2-isobutyl-3-methoxy-pyrazine⁴¹. Likewise, fatty acid binding proteins have been demonstrated to bind two molecules of the same ligand, oleic acid⁴². Recently, it has been suggested that DEET and NPN might bind simultaneously to AgamOBP1¹⁷, but experimental evidence showing increase in NPN fluorescence and blue shift data was missing. The hypotheses put forward here on the basis of our findings must await experimental evidence, in particular X-ray crystallography studies. Studies to test these hypotheses may lead to more effective fluorescence reporters and a better understanding of OBP odorant binding.

Data availability

F1000Research: Dataset 1. Update 1. Fluorescence reporter assay data with assessing binding of insect repellents to the yellow fever mosquito (*Culex quinquefasciatus*) odorant binding protein AaegOBP1, [10.5256/f1000research.5879.d41724](https://doi.org/10.5256/f1000research.5879.d41724)⁴³

Author contributions

WSL designed the experiments. GML and WSL carried out the research. WSL analyzed the data and wrote the manuscript. All authors revised the manuscript and agreed to its final content.

Competing interests

No competing interests were disclosed.

Grant information

This work was supported in part from unrestricted gifts from Fuji Flavor Co. and Bedoukian Research Inc.

Acknowledgements

We thank Dr. Robert Bedoukian (Bedoukian Research Inc.) for MOP and PMD samples, and Dr. Kamal Chauhan (USDA, ARS) for picaridin and IR3535 samples. We are indebted to Mr. Garrison Buss (UC Davis), Drs. Yuko Ishida (Toyama University, Japan), Julien Pelletier (Keele University, UK), Pingxi Xu (UC Davis) and David Wilson (UC Davis) for comments and suggestion to improve an earlier version of the manuscript.

References

- Clyne PJ, Warr CG, Freeman MR, *et al.*: **A novel family of divergent seven-transmembrane proteins: candidate odorant receptors in *Drosophila***. *Neuron*. 1999; **22**(2): 327–38.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gao Q, Chess A: **Identification of candidate *Drosophila* olfactory receptors from genomic DNA sequence**. *Genomics*. 1999; **60**(1): 31–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Vosshall LB, Amrein H, Morozov PS, *et al.*: **A spatial map of olfactory receptor expression in the *Drosophila* antenna**. *Cell*. 1999; **96**(5): 725–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Leal WS: **Odorant reception in insects: roles of receptors, binding proteins, and degrading enzymes**. *Annu Rev Entomol*. 2013; **58**: 373–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Benton R, Sachse S, Michnick SW, *et al.*: **Atypical membrane topology and heteromeric function of *Drosophila* odorant receptors in vivo**. *PLoS Biol*. 2006; **4**(2): e20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vogt RG, Riddiford LM: **Pheromone binding and inactivation by moth antennae**. *Nature*. 1981; **293**(5828): 161–3
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ishida Y, Cornet AJ, Leal WS: **Identification and cloning of a female antenna-specific odorant-binding protein in the mosquito *Culex quinquefasciatus***. *J Chem Ecol*. 2002; **28**(4): 867–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bohbot J, Pitts RJ, Kwon HW, *et al.*: **Molecular characterization of the *Aedes aegypti* odorant receptor gene family**. *Insect Mol Biol*. 2007; **16**(5): 525–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Carey AF, Wang G, Su CY, *et al.*: **Odorant reception in the malaria mosquito *Anopheles gambiae***. *Nature*. 2010; **464**(7285): 66–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Leal WS, Choo YM, Xu P, *et al.*: **Differential expression of olfactory genes in the southern house mosquito and insights into unique odorant receptor gene isoforms**. *Proc Natl Acad Sci U S A*. 2013; **110**(46): 18704–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wang G, Carey AF, Carlson JR, *et al.*: **Molecular basis of odor coding in the malaria vector mosquito *Anopheles gambiae***. *Proc Natl Acad Sci U S A*. 2010;

- 107(9): 4418–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Hughes DT, Pelletier J, Luetje CW, *et al.*: **Odorant receptor from the southern house mosquito narrowly tuned to the oviposition attractant skatole.** *J Chem Ecol.* 2010; **36**(8): 797–800.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 13. Pelletier J, Hughes DT, Luetje CW, *et al.*: **An odorant receptor from the southern house mosquito *Culex pipiens quinquefasciatus* sensitive to oviposition attractants.** *PLoS One.* 2010; **5**(4): e10090.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 14. Zhu F, Xu P, Barbosa RM, *et al.*: **RNAi-based demonstration of direct link between specific odorant receptors and mosquito oviposition behavior.** *Insect Biochem Mol Biol.* 2013; **43**(10): 916–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 15. Xu P, Choo YM, De La Rosa A, *et al.*: **Mosquito odorant receptor for DEET and methyl jasmonate.** *Proc Natl Acad Sci U S A.* 2014; **111**(46): 16592–16597.
[PubMed Abstract](#) | [Publisher Full Text](#)
 16. McBride CS, Baier F, Omondi AB, *et al.*: **Evolution of mosquito preference for humans linked to an odorant receptor.** *Nature.* 2014; **515**(7526): 222–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 17. Murphy EJ, Booth JC, Davrazou F, *et al.*: **Interactions of *Anopheles gambiae* odorant-binding proteins with a human-derived repellent: implications for the mode of action of n,n-diethyl-3-methylbenzamide (DEET).** *J Biol Chem.* 2013; **288**(6): 4475–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 18. Tsitsanou KE, Thireou T, Drakou CE, *et al.*: ***Anopheles gambiae* odorant binding protein crystal complex with the synthetic repellent DEET: implications for structure-based design of novel mosquito repellents.** *Cell Mol Life Sci.* 2012; **69**(2): 283–97.
[PubMed Abstract](#) | [Publisher Full Text](#)
 19. Mao Y, Xu X, Xu W, *et al.*: **Crystal and solution structures of an odorant-binding protein from the southern house mosquito complexed with an oviposition pheromone.** *Proc Natl Acad Sci U S A.* 2010; **107**(44): 19102–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 20. Leite NR, Krogh R, Xu W, *et al.*: **Structure of an odorant-binding protein from the mosquito *Aedes aegypti* suggests a binding pocket covered by a pH-sensitive “Lid”.** *PLoS One.* 2009; **4**(11): e8006.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 21. Wogulis M, Morgan T, Ishida Y, *et al.*: **The crystal structure of an odorant binding protein from *Anopheles gambiae*: evidence for a common ligand release mechanism.** *Biochem Biophys Res Commun.* 2006; **339**(1): 157–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
 22. Biessmann H, Andronopoulou E, Biessmann MR, *et al.*: **The *Anopheles gambiae* odorant binding protein 1 (AgamOBP1) mediates indole recognition in the antennae of female mosquitoes.** *PLoS One.* 2010; **5**(3): e9471.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 23. Pelletier J, Guidolin A, Syed Z, *et al.*: **Knockdown of a mosquito odorant-binding protein involved in the sensitive detection of oviposition attractants.** *J Chem Ecol.* 2010; **36**(3): 245–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 24. Leal WS, Barbosa RM, Xu W, *et al.*: **Reverse and conventional chemical ecology approaches for the development of oviposition attractants for *Culex* mosquitoes.** *PLoS One.* 2008; **3**(8): e3045.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 25. Lagarde A, Spinelli S, Qiao H, *et al.*: **Crystal structure of a novel type of odorant-binding protein from *Anopheles gambiae*, belonging to the C-plus class.** *Biochem J.* 2011; **437**(3): 423–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
 26. Qiao H, He X, Schymura D, *et al.*: **Cooperative interactions between odorant-binding proteins of *Anopheles gambiae*.** *Cell Mol Life Sci.* 2011; **68**(10): 1799–813.
[PubMed Abstract](#) | [Publisher Full Text](#)
 27. Iovinella I, Bozza F, Caputo B, *et al.*: **Ligand-binding study of *Anopheles gambiae* chemosensory proteins.** *Chem Senses.* 2013; **38**(5): 409–19.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. Leal WS, Chen AM, Ishida Y, *et al.*: **Kinetics and molecular properties of pheromone binding and release.** *Proc Natl Acad Sci U S A.* 2005; **102**(15): 5386–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. Ban L, Zhang L, Yan Y, *et al.*: **Binding properties of a locust’s chemosensory protein.** *Biochem Biophys Res Commun.* 2002; **293**(1): 50–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 30. Ban L, Scaloni A, Brandazza A, *et al.*: **Chemosensory proteins of *Locusta migratoria*.** *Insect Mol Biol.* 2003; **12**(2): 125–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Ishida Y, Chen AM, Tsuruda JM, *et al.*: **Intriguing olfactory proteins from the yellow fever mosquito, *Aedes aegypti*.** *Naturwissenschaften.* 2004; **91**(9): 426–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. Zhou JJ, He XL, Pickett JA, *et al.*: **Identification of odorant-binding proteins of the yellow fever mosquito *Aedes aegypti*: genome annotation and comparative analyses.** *Insect Mol Biol.* 2008; **17**(2): 147–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
 33. Zhou JJ, He XL, Pickett JA, *et al.*: **Corrigendum.** *Insect Mol Biol.* 2008; **17**(4): 445.
[Publisher Full Text](#)
 34. Wojtasek H, Leal WS: **Conformational change in the pheromone-binding protein form *Bombyx mori* induced by pH and by interaction with membrane.** *J Biol Chem.* 1999; **274**(43): 30950–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 35. Leal WS: **Duality monomer-dimer of the pheromone-binding protein from *Bombyx mori*.** *Biochem Biophys Res Commun.* 2000; **268**(2): 521–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 36. Lautenschlager C, Leal WS, Clardy J: **Coil-to-helix transition and ligand release of *Bombyx mori* pheromone-binding protein.** *Biochem Biophys Res Commun.* 2005; **335**(4): 1044–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. Laurence BB, Pickett JA: **erythro-6-Acetoxy-5-hexadecanolide, the major component of a mosquito oviposition attractant pheromone.** *J Chem Soc Chem Commun.* 1982; **1982**(1): 59–60.
[Publisher Full Text](#)
 38. Sun YF, De Biasio F, Qiao HL, *et al.*: **Two odorant-binding proteins mediate the behavioural response of aphids to the alarm pheromone (E)- β -farnesene and structural analogues.** *PLoS One.* 2012; **7**(3): e32759.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Ishida Y, Ishibashi J, Leal WS: **Fatty acid solubilizer from the oral disk of the blowfly.** *PLoS One.* 2013; **8**(1): e51779.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 40. Spinelli S, Lagarde A, Iovinella I, *et al.*: **Crystal structure of *Apis mellifera* OBP14, a C-minus odorant-binding protein, and its complexes with odorant molecules.** *Insect Biochem Mol Biol.* 2012; **42**(1): 41–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Lautenschlager C, Leal WS, Clardy J: ***Bombyx mori* pheromone-binding protein binding nonpheromone ligands: implications for pheromone recognition.** *Structure.* 2007; **15**(9): 1148–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 42. Cai J, Lucke C, Chen Z, *et al.*: **Solution structure and backbone dynamics of human liver fatty acid binding protein: fatty acid binding revisited.** *Biophys J.* 2012; **102**(11): 2585–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 43. Leal GM, Leal SW: **Dataset 1 in ‘Binding of a fluorescence reporter and a ligand to an odorant-binding protein of the yellow fever mosquito, *Aedes aegypti*’.** *F1000Research.* 2014.
[Data Source](#)

Open Peer Review

Current Referee Status:



Version 2

Referee Report 22 January 2015

doi:[10.5256/f1000research.6433.r7334](https://doi.org/10.5256/f1000research.6433.r7334)



Kostas Iatrou¹, Spyros Zographos²

¹ Institute of Biosciences and Applications, National Center for Scientific Research “Demokritos”, Athens, Greece

² Institute of Biology, Medicinal Chemistry & Biotechnology, National Hellenic Research Foundation, Athens, Greece

We are happy to see that our comments have been considered to be useful to the authors and appreciate the additional experimentation that was carried out in response to one of our comments. We do realize, of course, that molecular modeling by itself would not provide definitive answers to the working hypotheses put forward by the authors in relation to the binding pockets of AaegOBP1. Moreover, given that modeling is not an area of in-depth expertise for the specific group and that solid answers may eventually only come through X-ray crystallography, we feel that the authors have exhausted their capability to interpret fully their carefully documented observations. We would therefore encourage them to pursue a future investigation on this very interesting matter in collaboration with another relevant group. At the same time, we approve the indexing of the paper in its revised form.

We have read this submission. We believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Competing Interests: No competing interests were disclosed.

Version 1

Referee Report 02 January 2015

doi:[10.5256/f1000research.6281.r7176](https://doi.org/10.5256/f1000research.6281.r7176)



Paolo Pelosi

Department of Chemistry and Agricultural Biotechnology, University of Pisa, Pisa, Italy

The phenomenon described in this paper is well known and documented in many papers. However, it has never been directly examined and explained in detailed. Therefore, it is nice and useful to have a focused study to describe and dissect such apparently anomalous behaviour once and for all.

I fully agree with the Authors that the formation of micelles is the most likely explanation. We have come

across this same phenomenon several times and I have always discussed this fact with my students hypothesising the formation of micelles as the most likely reason behind this. A brief explanation of some anomalous binding curves can also be found in some of our published papers, most recently in [Sun et al. \(2012\)](#).

When a ligand capable of forming micelles also has affinity for the protein, we observe a decrease of fluorescence, followed by an increase when titrating the protein (the U curve observed in this paper). When the ligand has poor affinity for the protein, we only observe a constant increase in fluorescence. Sometimes we have also recorded a complex behaviour: the intensity of fluorescence experiences an increase at low concentration values of the ligand, then drops when more ligand is added. In this case, the phenomenon could be explained by assuming that the ligand enters the binding pocket without displacing the fluorescent probe. As the Authors point out such facts can occur and have been documented with OBPs and CSPs. The increase of fluorescence in such case would be the result of the increased hydrophobicity of the binding pocket due to the presence of a ligand, usually a highly hydrophobic molecule, as in the case of many pheromones of Lepidoptera and Diptera. As the concentration of the ligand increases, competition with bound 1-NPN can take over producing a decrease in fluorescence.

This study could be complemented (but not necessarily) by monitoring the intrinsic fluorescence of the tryptophan, which appropriately is located inside the binding pocket of the protein. Particularly in the case of DEET, which is an aromatic compound, if the molecule binds to OBP without being able to displace 1-NPN, we should observe a strong quenching of the tryptophan fluorescence.

Overall the paper is well written and the observed phenomenon clearly described and explained.

I have only one minor concern:

The Authors report the emission spectrum of 1-NPN alone (in Tris buffer at pH 7.4) with a maximum around 470 nm and in the presence of protein at about 440. In my experience, I found that 1-NPN in buffer produces a peak with a maximum at 480, which is shifted in the presence of a binding protein to values generally between 400 and 410 nm, in some cases even below 400. This has been observed with a large number of OBPs, including some of mosquitoes, although not with the specific OBP used in this study. I suggest that the Authors double check these data, also in relationship to the instrument calibration.

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Competing Interests: No competing interests were disclosed.

Author Response 02 Jan 2015

Walter Leal, UC Davis, USA

We appreciate your time and effort to evaluate our article, and are delighted to hear that it meets your approval. We will certainly cite [Sun et al. \(2012\)](#) in the revised version of the article. The apparent formation of (E)- β -farnesene micelle, as shown in [your Figure 5](#), skipped our attention, perhaps because this phenomenon has never been examined and explained in detail. We will certainly give the appropriate credit in the revised version. Regarding your minor concern, we have verified wavelength accuracy per vendor's instruction manual. We are, therefore, confident that the

data set reported is accurate. It is worth pointing out, however, that apparent discrepancies may be explained at least in part by the proteins studied. For AaegOBP1, the NPN peaks at pH values 7 and 5 are quite different (our Figure 6), but in the presence of CquiOBP1 (Figure 5 in [Leal *et al.*, 2008](#)) or AfunOBP1 (Figure 9, [Xu *et al.*, 2010](#)) the wavelength for the peaks at high and low pH are nearly the same, i.e., ca. 400 nm. All these studies were performed in our laboratory with the same instrument. Again, thank you very much for your suggestions. Sincerely, WSL & GML

Competing Interests: No competing interests were disclosed.

Referee Report 29 December 2014

doi:10.5256/f1000research.6281.r7040



Kostas Iatrou¹, Spyros Zographos²

¹ Institute of Biosciences and Applications, National Center for Scientific Research "Demokritos", Athens, Greece

² Institute of Biology, Medicinal Chemistry & Biotechnology, National Hellenic Research Foundation, Athens, Greece

This manuscript describes interesting results obtained in the course of what of an investigation initially designed as a high school project undertaken in order to deduce whether AaegOBP1, an odorant binding protein of the yellow fever mosquito *Aedes aegypti*, binds DEET and/or other known mosquito repellents such as icaridin, PMD and IR3535. The relevant experiments consisted of classical binding competition assays by the tested repellents against an AaegOBP1 pre-bound fluorescent reporter molecule, NPN, causing reductions in NPN-emitted fluorescent quenching with the latter serving as measure of mosquito repellent binding to AaegOBP1 resulting in displacement of the pre-bound NPN.

While the experiments suggested that the specific OBP may only bind the tested repellents with limited affinity relative to NPN, they also produced results that could not have been predicted *a priori*. The first concerned an unexpected property of a mosquito (*Culex quiquefasciatus*) oviposition pheromone (MOP) that was used as positive control for binding to AaegOBP1. Thus, while titration AaegOBP1/NPN complexes by increasing quantities of MOP produced the anticipated reduction in NPN fluorescence, titrations with higher MOP doses led to gradual increases of fluorescence emitted by NPN accompanied by a wavelength shift toward the blue region of the spectrum. To explain this finding as well as the parallel observation that the same phenomenon also occurs at the same MOP concentrations in the absence of AaegOBP1, the authors have postulated the formation of MOP micelles forming a highly hydrophobic environment to which displaced and free NPN may bind.

The second intriguing finding has been that at a low pH of 5.5 at which AaegOBP1 is unable to bind MOP, this protein binds NPN with higher affinity relative to a neutral pH, causing higher emitted fluorescence with a concomitant blue-shift in the emission wavelength suggestive of the formation of a higher hydrophobicity environment to which NPN is bound. Based on these findings as well as the crystal structures of CquiOBP1 and *Anopheles gambiae* AgamOBP1, both AaegOBP1 orthologs, in complex with MOP and DEET, respectively, as well as the complex of the honey bee AmelOBP14 with NPN, the authors postulate the possibility that NPN and MOP could bind simultaneously to AaegOBP1 at a neutral pH. In turn, this possibility suggests that caution should also be exercised for the postulated conclusion regarding the low affinity binding of DEET to AaegOBP1, because DEET binding to a separate pocket

might not necessarily result in displacement of NPN.

Suggestions:

- For the first set of observations related to the postulated micelle formation by MOP at concentrations of 12.5 μM or higher, the hypothesized explanation is quite reasonable. A dynamic light scattering experiment using MOP in buffer alone could further strengthen the postulated hypothesis. Moreover, a NPN titration experiment similar to that shown in Fig. 5B but at a pH 5.5, which should result in protonation e.g. of the acetoxy-group of MOP, could reveal whether an increase in micelle size occurs or not. This latter experiment could also provide additional suggestive evidence for the postulated creation of a more hydrophobic environment for NPN binding in AaegOBP1 at the acidic pH.
- For the structural considerations presented in the conclusions, as the authors indicate, co-crystallization of AaegOBP1 with NPN, MOP or both, will be required in order for conclusive interpretations to be drawn. Nevertheless, it is not clear to us why in a case of simultaneous binding of NPN and MOP (AaegOBP1-MOP-NPN complex), NPN should move to a different binding pocket of higher hydrophobicity producing higher fluorescence emission and a blue shift, only at higher MOP concentrations and not at lower ones. If, on the other hand, the requirement for higher MOP concentrations is interpreted as indicative of the formation of AaegOBP1-MOP-NPN-MOP complexes, a docking model should indicate whether enough space exists in the L-shaped tunnel of the AaegOBP1 monomer for simultaneous binding of 3 molecules.
- For the low apparent affinity of AaegOBP1 for DEET, it is indeed possible that the binding of DEET and NPN to AaegOBP1 are not mutually exclusive, hence the low reduction in emitted NPN fluorescence in the presence of increasing concentrations of DEET. A docking model should indicate whether the possibility of nearby binding sites or even overlapping ones for NPN and DEET is predicted, which would lead to fluorescence quenching rather than reduction due to NPN displacement.
- Finally, the authors should provide a concluding statement as to whether and how these interesting findings relate to the contributions of OBPs in the mosquito's olfactory function under normal conditions.

We have read this submission. We believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Competing Interests: No competing interests were disclosed.

Author Response 01 Jan 2015

Walter Leal, UC Davis, USA

First of all, we would like to thank you for the time and effort to process and evaluate our article. We were delighted to read the laudatory comments in your report. We have considered carefully your suggestions (thank you very much), and performed an additional experiment, which will be incorporated in Figure 5C. Specifically, we performed NPN titration with MOP in ammonium acetate buffer, pH 5.5. Although we agree that the suggested docking experiments (bullets 2 and 3) might add to the discussion, ultimately the structural hypotheses raised in the article shall be supported or refuted by X-ray crystallography-based structural evidence. While none of the authors is well versed in molecular modeling, we have the expertise and collaboration in place to rigorously test the structural hypotheses. In collaboration with our UCD colleague, Dr. David Wilson, Gabriel

has been able to crystallize other OBPs, and we are confident that he/we will succeed in crystallizing AaegOBP1 complexes and address these questions. Obviously, time is uncertain in crystallography and the scope of the new work seems to belong to future publication(s). In the revised version of our *F1000Research* article, we will add comments regarding contributions of OBPs to mosquito olfaction, as suggested. Additionally, we will add results shown further evidence of micelle formation even at low pH, except that the effect was manifested at slightly higher concentrations of MOP. In sum, figures B and C are almost identical, but the blue shift and fluorescence increase were clearly observed starting at 15 and 17.5 μM at pH 5.5 (revised Fig. 5C) as compared to 12.5 and 15 μM (Fig. 5B). We do hope that now the article meets your approval. Sincerely, WSL & GML

Competing Interests: No competing interests were disclosed.
