

UC Riverside

UC Riverside Previously Published Works

Title

Symbiotic nitrogen fixation by rhizobia—the roots of a success story

Permalink

<https://escholarship.org/uc/item/33n0s9h9>

Authors

Masson-Boivin, Catherine
Sachs, Joel L

Publication Date

2018-08-01

DOI

10.1016/j.pbi.2017.12.001

Peer reviewed



Symbiotic nitrogen fixation by rhizobia — the roots of a success story

Catherine Masson-Boivin¹ and Joel L Sachs²

By evolving the dual capacity of intracellular survival and symbiotic nitrogen fixation in legumes, rhizobia have achieved an ecological and evolutionary success that has reshaped our biosphere. Despite complex challenges, including a dual lifestyle of intracellular infection separated by a free-living phase in soil, rhizobial symbiosis has spread horizontally to hundreds of bacterial species and geographically throughout the globe. This symbiosis has also persisted and been reshaped through millions of years of history. Here, we summarize recent advances in our understanding of the molecular mechanisms, ecological settings, and evolutionary pathways that are collectively responsible for this symbiotic success story. We offer predictions of how this symbiosis can evolve under new influences and for the benefit of a burgeoning human population.

Addresses

¹ LIPM, Université de Toulouse, INRA, CNRS, Castanet-Tolosan cedex, France

² Department of Evolution Ecology and Organismal Biology, University of California, Riverside, CA, USA

Corresponding author: Masson-Boivin, Catherine (catherine.masson@inra.fr)

Current Opinion in Plant Biology 2018, 44:7–15

This review comes from a themed issue on **Biotic interactions**

Edited by **Sebastian Schornack** and **Caroline Gutjahr**

<https://doi.org/10.1016/j.pbi.2017.12.001>

1369-5266/© 2017 Published by Elsevier Ltd.

Introduction

Biological nitrogen fixation, the capacity to convert atmospheric dinitrogen (N₂) to a reduced form, is an ancient innovation exclusively achieved by Bacteria and Archaea and is one of the most significant ecological services that microbes offer to eukaryotes. A variety of organisms, including animals, plants, fungi and protists, form symbiotic interactions with N₂-fixing bacteria, but these associations are generally loose and with little evidence of evolutionary modification in either partner [1]. Yet a few Angiosperm lineages have evolved complex intracellular and highly efficient symbioses with N₂-fixing bacteria. These symbioses are manifested by the formation of

specialized root (or occasionally stem) organs called nodules that are massively colonized by bacterial partners and act as miniature N₂-fixing factories for the plant. Nodulation in plants first evolved around 100 MYA leading to nodulation capacity in 70% of legume species, and in several lineages of plants, mainly so-called actinorhizal plants, distributed across three Angiosperm orders [2,3]. In parallel, the ability to fix nitrogen with legumes has spread to hundreds of species in alpha-Proteobacteria and beta-Proteobacteria [4], referred to as rhizobia, while the ability to nodulate actinorhizal plants has been restricted to the *Frankia* genus in Actinobacteria. Phylogenomic approaches are making tremendous progress in identifying the genetic innovations that allowed plants to establish a nitrogen fixation symbiosis with bacteria [5]. In this review, we explore the drivers of evolutionary success of rhizobia, scrutinizing the origin and evolution of key symbiotic innovations, and the effect of host and environment in shaping rhizobial adaptation. Finally, we consider future directions to select and design efficient, host-adapted, nitrogen fixers for productive and sustainable agriculture.

Molecular keys

Molecular mechanisms underlying the rhizobium-legume symbiosis can be divided into functions that instigate two main stages: (i) nodule formation and invasion, which are distinct yet genetically tightly coordinated processes, and (ii) symbiotic nitrogen fixation that most often occurs within nodule cells.

All rhizobia are equipped with the MoFe containing variant of the nitrogenase enzyme system [6], which is the only biological system able to fix nitrogen. Rhizobia have solved the paradox between their strict aerobic status and the extreme sensitivity of nitrogenase to oxygen by inducing new organs, the nodules, which exhibit an extremely low free oxygen atmosphere and synthesize a form of plant hemoglobin — leghemoglobin — that facilitates O₂ diffusion to bacteroids.

The last decade has changed the paradigm that the capacity to instigate nodule formation is solely determined by a canonical set of rhizobial nodulation (*nod*) genes. Nodulation can be achieved by two additional strategies, the type three secretion system (T3SS) and a genetically uncharacterized mechanism (Box 1). These mechanisms may correspond to three different ‘strategies’ to activate a host genetic program governing the development of nodules. This program involves a

Box 1 Nodulation strategies and their origins.

Three nodulation strategies have been identified in rhizobia to date: the Nod, T3SS and non-Nod/non-T3SS strategies, whose characteristics are detailed in Figure 1. *Frankia* could possibly use a fourth mechanism to nodulate actinorhizal plants since their symbiotic signal appears chemically different from lipo-chitooligosaccharides [68].

In the Nod strategy (reviewed in [7,8,69]), strain-specific lipo-chitooligosaccharides called Nod factors (NFs) are produced under the control of nodulation genes (*nod/nol/noe* genes collectively referred to as *nod* genes). Their core structure is determined by the common *nodABC* genes. NFs are perceived by plant receptors (LysM-receptor like kinases) that activate the CSSP. In the T3SS strategy [70**,71*], T3SS effectors, whose nature and function are under investigation, activate CSSP components by bypassing NF recognition [70**]. The mechanism of the third nodulation strategy is still unknown, but it involves neither *nod* nor T3SS functions. No non-nodulating bacterial mutants could be obtained [72*,73], suggesting that essential genes are required for the induction of nodule formation. Nod-independent and T3SS-independent nodulation occurs via CSSP activation, since most CSSP components have been identified in *Aeschynomene evenia* [74] -that is only nodulated via this strategy-, some of which have been shown to be involved in symbiosis in this legume [75].

nod genes have likely been recruited from non-symbiotic soil bacteria since homologs are widely present in eubacteria. *nodA*, the acyl transferase long thought to be unique to rhizobia, has now been found in the Firmicutes (e.g. *Paenibacillus* ABC68285.1) and Actinobacteria (e.g. SBW22889.1 and *Rhodococcus* ADW94999.1) phyla. T3SS are widely present in rhizobia [71*,76,77], where they are generally controlled by the *nodD* regulator, suggesting that these systems have been co-opted by Nod-rhizobia for symbiotic purposes during evolution [78]. It would be interesting to determine whether nodulation-inducing T3SS are always associated to a Nod strategy and are under *nodD* control. This, together with the fact that T3SS-based nodulation does not lead to a mutualistic symbiosis so far, would suggest that T3SS effectors recruited to improve nodulation may sometimes be efficient enough to supersede NFs. Phylogenetic analyses revealed a single clade of non-Nod/non-T3SS *Bradyrhizobium* and a single emergence of the capacity to interact with such *Bradyrhizobium* in *Aeschynomene* [79]. This suggests that non-Nod/non-T3SS nodulation is a derived and more recent character than the *nod*-mediated nodulation [80].

common symbiotic signaling pathway (CSSP) highly conserved in plants able to form N₂-fixing or arbuscular mycorrhizal (AM) associations [7]. The Nod strategy has been extensively studied and reviewed [8] whereas our understanding of the two other mechanisms is still in its infancy. Each strategy appears to correspond to different degrees of adaptation and efficiency of nitrogen fixation (Box 1).

In addition to the basic genetic toolset needed to instigate nodule formation is a range of bacterial functions that promote infection and intracellular accommodation and whose requirement and/or genetic basis depend on the rhizobial species. In many rhizobia infection requires the production of surface polysaccharides [9,10] that are recognized by plant receptors [11]. In legumes producing nodule specific cysteine-rich (NCR) peptides [12],

peptide transporters (BacA or BclA) are required for the differentiation of internalized bacteria into functional bacteroids [13,14]. Other bacterial functions can also improve or impede the infection process and thus modulate host range: examples are type-3 and type-4 secretion systems (T3SS, T4SS) that translocate effectors in plant cells and that trigger or suppress plant defense reaction [15,16] and the HrrP peptidase which cleaves host-encoded signaling NCR peptides [17].


Evolutionary trends

The existence of three genetically distinct nodulation strategies suggests that the nodulation trait in rhizobia has evolved multiple times. How these strategies emerged and in which order is a challenging question (see Box 1). AM fungi produce signaling Myc Factors (structurally similar to NFs [18]), and both of these lipooligosaccharides (LCOs) can induce CSSP components in plants [19]. However, *nod* homologous genes have not been reported in the only complete AM fungus genome available so far [20], thus indicating that LCO synthesis can be mediated by genes unrelated to *nod* genes. Instead, *nod* genes might have originated from non-symbiotic bacteria (see Box 1) that had co-opted a plant program initially devoted to establishing symbioses with AM fungi [19]. NFs suppress MAMP-triggered immunity in non-legumes [21**] and are perceived by receptor-like kinases (LysM RLK) involved in immune signaling [19]. These findings together with the fact that T3SS deliver effector proteins directly into the plant cell to induce effector-triggered immunity argue for an initial pathogenic stage in the rhizobium-legume evolution. The primary role of T3SS effectors and LCOs in suppressing immunity might have further evolved to support nodulation.

Among the mechanisms used to induce nodule formation, the Nod strategy has been most successful as it is used by most of the rhizobia investigated so far. One hypothesis to explain this success is that *nod* and *nif* genes are tightly linked on mobile genetic elements (MGE), either large plasmids or genomic islands, with a few exceptions [22]. Clustering on MGEs can allow the rapid and joint dissemination of key symbiotic traits [23,24*]. Transfer of MGEs is predicted to be most frequent during the free-living rhizospheric stage where bacteria live in diverse biofilm consortia [25], and where transfer can be enhanced by plant flavonoids that also induce the expression of *nod* genes [26], establishing a possible link between symbiosis functioning and evolution. Rhizobia are closely related to non-symbiotic taxa, and ancestral state reconstruction suggests that nodulation and nitrogen fixation functions have been successfully transferred to both saprophytic and parasitic bacteria [27].

The first step towards symbiosis is transfer of a limited set of key genes likely coupled with the genetic rewiring of the recipient genome (Figure 2a), as can occur in

Figure 1



Nodulation strategies in legume symbionts. *, Possesses the Nod and T3SS strategies. **, Whether the use of a T3SS is the unique nodulation strategy in this strain is not known. ***, Possesses the Nod and non-Nod/non-T3SS strategies. A, *Aeschynomene*. IT, infection thread.

pathogens [28]. Indeed most of the genes involved in nodulation competitiveness or later stages, including surface polysaccharides crucial for infection, do not belong to the symbiotic MGE [29] and must be recruited from the genomic background. A step by step evolution, including MGE transfer followed by genomic remodeling under plant selection pressure, is supported by the existence of clear phenotypic shifts driven by adaptive mutations in experimental evolution studies [30[•]] (Figure 2b). Post-HGT evolution of emerging rhizobia may have been accelerated by error-prone DNA polymerases that are present on half of the rhizobial symbiotic plasmids [31[•]] and facilitated by IS transposition [32].


Host controls

To maximize their benefits from symbiosis legumes have evolved a suite of 'host control' traits that select for compatible rhizobia and defend against strains that provide little or no benefit. Nodule formation and infection are determined via multiple systems of specificity

(Figure 3). Host control over nodulation can be subtle, such as when legumes form fewer nodules with rhizobial strains that tend to provide less benefit to the host [33,34] or nodulation can be blocked completely, such as when the *Rj4* allele in soybeans triggers gene-for-gene resistance against toxin-producing *Bradyrhizobium* that reduce host growth [35]. Yet, multiple experiments suggest that legumes cannot specifically detect the potential for a rhizobial symbiont to fix N_2 prior to nodule organogenesis. Hosts that are co-inoculated with effective rhizobia and isogenic strains lacking N_2 -fixation function are often nodulated with equal frequency by both [36,37]. This is a common pattern in symbioses: hosts cannot reliably discriminate between beneficial and harmful symbionts during initial colonization, because there is no reason for symbionts to signal honestly.

After nodule organogenesis is underway, legumes can respond to nodules that house ineffective rhizobia (strains that do not fix nitrogen) and reduce bacterial

Figure 2



Evolution of Nod-rhizobia. **(a)** Model for the evolution of Nod-rhizobia. *nod/nif*-containing MGEs are horizontally transferred to diverse soil bacteria, conferring symbiotic potential that depends on the recipient genome. The resulting chimeric bacteria subsequently evolved by genome remodeling under legume selection pressure – which can take indefinite time – allowing the recruitment of local regulatory and metabolic functions to better thrive in the new endosymbiotic environment (Figure 3). Adapted from [81]. **(b)** Experimental evolution of symbiotic *Ralstonia*. The evolution of a Nod-rhizobium is being tested in the laboratory. A rhizobial (*Cupriavidus taiwanensis*) symbiotic plasmid carrying *nod* and *nif* genes was introduced into the strictly extracellular plant pathogen *Ralstonia solanacearum*, allowing the resulting transconjugant to only induce root hair curling, which is the first symbiotic step. The chimeric *Ralstonia* was further evolved under selection pressure of *Mimosa pudica*, the natural host of *C. taiwanensis* [82]. Nodulation and infection, were acquired and improved step by step in parallel lineages. Mutualism was not achieved after two years as bacteria did not support plant growth. Clear phenotypic shifts mainly resulted from the neutralization of a bacterial function (T3SS, mutation in *hrcV*) that impairs the symbiotic process or the rewiring of the virulence regulatory pathway of *R. solanacearum* (mutations targeting *hrpG*, *prhI*, *vsrA*, *efpR* or *Rsc0965* genes) [30,64,65]. This experiment illustrates how the rewiring of a genetic network regulating virulence allows a radically different type of symbiotic interaction and contributes to ecological transitions.


within-nodule growth rates, a trait termed ‘sanctions’ [34,38–42]. But there is little mechanistic understanding of how legumes detect and sanction ineffective strains. Models predict that legumes target ineffective rhizobia at the level of the whole nodule, hence that sanctions are modulated dependent upon the total amount of fixed nitrogen provided by individual nodules whether they are clonal or contain mixed strain infections [43,44]. But growing evidence suggests that hosts can sanction ineffective rhizobia even when individual nodules are co-infected by a mix of effective and ineffective strains [34,40]. Furthermore, recent experimental data show that legumes can sanction ineffective strains in a cell-autonomous fashion; wherein the host plant induces programmed cell death of cells that house ineffective rhizobia [45,46]. These results bring us closer to a mechanistic understanding of how legumes detect and defend against ineffective rhizobial genotypes. Experiments and mathematical modelling showed that rare

nitrogen-fixing symbionts can invade a population dominated by non-fixing bacteria with a probability that depends on ecological factors [46]. Hence, host sanctions may have played a major role in the spread of the mutualistic trait during evolution. The imperfect match between the nodulation and nitrogen fixation capacities together with the loose selection process, that is, non-fixers do not become extinct within nodules [46], helps maintain genetically diverse rhizobial communities in the soil and shape the ecology and evolution of rhizobia. The ecological correlates of rhizobial fitness, including both effective and symbiotically ineffective strains, is the topic of the next section.

Ecological correlates and drivers

Rhizobia are often strictly defined in functional terms of nodulation and nitrogen fixation capacity, but this definition becomes blurred when considering the evolutionary flexibility that can occur between the multiple lifestyles

Figure 3



Multistep molecular surveillance mechanism in the rhizobium-legume symbiosis. Successful mutualism relies on the mutual recognition of a series of molecular signals between the symbiotic partners, allowing or impairing bacterial proliferation at each stage (nodulation, infection, intracellular survival) and determining host specificity. Plant receptors (LysM domain containing receptor-like kinases such as NFR or SymRK) recognize species-specific lipo-chitooligosaccharides (NFs) synthesized by Nod-rhizobia in response to plant compounds (flavonoids). These receptors trigger the plant genetic program leading to nodule organogenesis and infection. This plant genetic program can also be activated by bacterial T3SS effectors. Microbe-associated molecular patterns (MAMPs) or other bacterial T3/4SS effectors can induce plant immunity mediated by intracellular or plasma membrane localized receptors. Exopolysaccharides (EPS) recognized by plant (EPR3) facilitate infection [11]. NCRs, which are extremely abundant defensin-like peptides in the nodules of some legumes (Inverted Repeat-Lacking Clade and *Aeschynomene* legumes), can have a positive or negative impact on the intracellular stage. Some NCRs control the differentiation of intracellular bacteria into enlarged polyploid bacteroids [83,84], a step required for N_2 -fixation but which can be impaired by bacterial peptidases (HrrPs) [17]. The bactericidal activity of other peptides (NFS1 NFS2) affects the intracellular survival of specific bacterial genotypes [85,86]. The bacterial BacA/BclA transporter promotes the import of NCR peptides [13,87]. Molecules identified in different rhizobium-legume systems are represented here. Adapted from [70*].

of rhizobial taxa [47,48]. For instance, classically defined rhizobia are mutualists that provide fixed nitrogen to hosts (i.e. Nod^+/Fix^+), but close relatives can be ineffective strains that form nodules but fail to fix nitrogen (i.e. Nod^+/Fix^-), or non-nodulating strains that do not have the genetic machinery to form nodules (i.e. Nod^-/Fix^- ; [49,50]), but nonetheless thrive in the rhizosphere and can gain access to nodules by coinfecting with nodulating strains [54,55]. Legume hosts play a major role in promoting Nod^+/Fix^+ strains given that the presence of legume hosts increases the abundance of nodule forming rhizobia in soils [51] and selects for rhizobia with genetically compatible *nod* loci [24*,52,53]. But the relative role of selection by hosts remains poorly understood. As described above, legumes can favor Fix^+ rhizobia over Fix^- under lab conditions [34,38–42,46*]. But non-fixing rhizobia nonetheless persist in a diversity of settings [54], and multiple models could explain their persistence. Nod^+/Fix^- can be favored because of their ability to exploit hosts [55], because some hosts have lost the

capacity to discriminate cheaters [56], or because of extrinsic factors, such as enriched soil nitrogen that can erase fitness effects of Fix^+ versus Fix^- strains [57]. Non-nodulating strains are the least understood among the three lifestyles described above. In some cases, these strains are deeply diverged from genotypes that nodulate legumes [47] and thus might have little relevance for the legume-rhizobium interaction. In other cases these strains are closely related to nodulating genotypes and are active competitors for persistence in the rhizosphere and within host nodules [58]. Moreover, recurrent evolutionary transitions from Nod^+ to Nod^- status suggests that selection might favor a Nod^- strategy under some conditions [50]. Consistent with this latter hypothesis, some Nod^- strains can outnumber related Nod^+ rhizobia in the host rhizosphere and can dominate large geographic areas [59]. Moreover, some work suggests that cosmopolitan rhizobial genotypes reach high frequency because they can catabolize diverse carbon sources in the rhizosphere [60]. To fully understand the drivers of rhizobial fitness we

must better understand these alternative strategies, the evolutionary pathways between them, and the genomic and ecological forces that drive these shifts [61].

New human influences

Legume crops contribute >30% of human nutritional nitrogen and represent a significant potential for sustainable crop productivity without the economic or environmental costs of chemical fertilization [62]. One common method has been to inoculate legume crops with ‘elite’ rhizobial strains; genotypes that fix substantial amounts of nitrogen for compatible hosts under defined greenhouse conditions. However, under agricultural conditions, legumes often do not recover the elite strains, and instead become infected with the indigenous rhizobia that fix little or no N for the host (i.e. the longstanding ‘rhizobial competition problem’; [63]).

Several approaches could further improve rhizobia as successful inocula under field conditions. One technique is to experimentally select for desired rhizobial traits such as the ability to efficiently form nodules and fix nitrogen. Experimental evolution has been applied under lab conditions converting non-rhizobia into strains that instigate nodule formation and infection on novel hosts [64,65] or resulting in an improved rhizobial capacity to provide benefits to hosts [33]. A more holistic approach attempts to select on whole consortia of beneficial soil microbes using a process described as ‘microbiome engineering’, which leverages host control traits described above to select microbial consortia that optimize plant yield (i.e. rapid growth, seed production, etc.; [66]. A final approach has been to avoid using rhizobia altogether and instead to engineer nitrogenase genes into the genomes of key crop plants, especially cereals. However, nitrogen fixation requires multiple functional modules and research has yet to accomplish this in any plant species [67].

Conclusion

Decades of research have clarified the main (Nod) strategy used by rhizobia to induce nitrogen-fixing root nodules and the reasons for its evolutionary success. Future challenges will be to evaluate the prevalence of the alternative nodulation strategies, and to understand their molecular mechanisms and evolution. A better understanding of how natural symbioses have established and evolved, as well as developing innovative tools for optimizing these associations and engineering new symbioses, should assist in engineering plant-associated microbiota for microbe-assisted crop improvement and developing nitrogen-fixing associations with cereals that would revolutionize world agriculture.

Acknowledgements

Catherine Masson-Boivin was supported by funds from the French National Research Agency (ANR-16-CE20-0011-01), and the French Laboratory of Excellence project ‘TULIP’ (ANR-10-LABX-41). Joel Sachs was supported

by grants from the US National Science Foundation, including DEB 1150278 and 1738009.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Kneip C, Lockhart P, Voss C, Maier UG: **Nitrogen fixation in eukaryotes — new models for symbiosis**. *BMC Evol Biol* 2007, **7**.
2. Werner GDA, Cornwell WK, Sprent JI, Kattge J, Kiers ET: **A single evolutionary innovation drives the deep evolution of symbiotic N₂-fixation in angiosperms**. *Nat Commun* 2014, **5**.
3. Van Velzen R, Holmer R, Bu F, Rutten L, van Zeijl A, Liu W, Santuari L, Cao Q, Sharma T, Shen D *et al.*: **Parallel loss of symbiosis genes in relatives of nitrogen-fixing non-legume Parasponia**. *bioRxiv* 2017 <http://dx.doi.org/10.1101/169706>.
4. Remigi P, Zhu J, Young JPW, Masson-Boivin C: **Symbiosis within symbiosis: evolving nitrogen-fixing legume symbionts**. *Trends Microbiol* 2016, **24**:63–75.54.
5. Delaux P-M, Radhakrishnan G, Oldroyd G: **Tracing the evolutionary path to nitrogen-fixing crops**. *Curr Opin Plant Biol* 2015, **26**:95–99.
6. Raymond J, Siefert JL, Staples CR, Blankenship RE: **The natural history of nitrogen fixation**. *Mol Biol Evol* 2004, **21**:541–554.
7. Oldroyd GED: **Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants**. *Nat Rev Microbiol* 2013, **11**:252–263.
8. Perret X, Staehelin C, Broughton WJ: **Molecular basis of symbiotic promiscuity**. *Microbiol Mol Biol Rev* 2000, **64**:180–201.
9. Jones KM, Kobayashi H, Davies BW, Taga ME, Walker GC: **How rhizobial symbionts invade plants: the Sinorhizobium–Medicago model**. *Nat Rev Microbiol* 2007, **5**:619–633.
10. Mathis R, Van Gijsegem F, De Rycke R, D’Haeze W, Van Maelsaeke E, Anthonio E, Van Montagu M, Holsters M, Vereecke D: **Lipopolysaccharides as a communication signal for progression of legume endosymbiosis**. *Proc Natl Acad Sci U S A* 2005, **102**:2655–2660.
11. Kawaharada Y, Kelly S, Nielsen MW, Hjuler CT, Gysel K, Muszynski A, Carlson RW, Thygesen MB, Sandal N, Asmussen MH *et al.*: **Receptor-mediated exopolysaccharide perception controls bacterial infection**. *Nature* 2015, **523**:308–312.
12. Mergaert P, Uchiumi T, Alunni B, Evanno G, Cheron A, Catrice O, Mausset A, Barloy-Hubler F, Galibert F, Kondoroski A *et al.*: **Eukaryotic control on bacterial cell cycle and differentiation in the rhizobium-legume symbiosis**. *Proc Natl Acad Sci U S A* 2006, **103**:5230–5235.
13. Guefrachi I, Pierre O, Timchenko T, Alunni B, Barriere Q, Czernic P, Villaecija-Aguilar JA, Verly C, Bourge M, Fardoux J *et al.*: **Bradyrhizobium BclA is a peptide transporter required for bacterial differentiation in symbiosis with Aeschynomene legumes**. *Mol Plant Microbe Interact* 2015, **28**:1155–1166.
14. Haag AF, Baloban M, Sani M, Kerscher B, Pierre O, Farkas A, Longhi R, Boncompagni E, Herouart D, Dall’Angelo S *et al.*: **Protection of Sinorhizobium against host cysteine-rich antimicrobial peptides is critical for symbiosis**. *PLoS Biol* 2011, **9**.
15. Hubber A, Vergunst AC, Sullivan JT, Hooykaas PJJ, Ronson CW: **Symbiotic phenotypes and translocated effector proteins of the Mesorhizobium loti strain R7A VirB/D4 type IV secretion system**. *Mol Microbiol* 2004, **54**:561–574.
16. Bartsev AV, Deakin WJ, Boukli NM, McAlvin CB, Stacey G, Malnoe P, Broughton WJ, Staehelin C: **NopL, an effector protein of Rhizobium sp. NGR234, thwarts activation of plant defense reactions**. *Plant Physiol* 2004, **134**:871–879.

17. Price PA, Tanner HR, Dillon BA, Shabab M, Walker GC, Griffiths JS: **Rhizobial peptidase HrrP cleaves host-encoded signaling peptides and mediates symbiotic compatibility**. *Proc Natl Acad Sci U S A* 2015, **112**:15244-15249.
18. Mailet F, Poinot V, Andre O, Puech-Pages V, Haouy A, Gueunier M, Cromer L, Giraudet D, Formey D, Niebel A *et al.*: **Fungal lipochitooligosaccharide symbiotic signals in arbuscular mycorrhiza**. *Nature* 2011, **469**:58-U1501.
19. Zhang XW, Dong WT, Sun JH, Feng F, Deng YW, He ZH, Oldroyd GED, Wang ET: **The receptor kinase CERK1 has dual functions in symbiosis and immunity signalling**. *Plant J* 2015, **81**:258-267.
20. Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, Charron P, Duensing N, Frey NFD, Gianinazzi-Pearson V *et al.*: **Genome of an arbuscular mycorrhizal fungus provides insight into the oldest plant symbiosis**. *Proc Natl Acad Sci U S A* 2013, **110**:20117-20122.
21. Liang Y, Cao YR, Tanaka K, Thibivilliers S, Wan JR, Choi J, Kang CH, Qiu J, Stacey G: **Nonlegumes respond to rhizobial Nod factors by suppressing the innate immune response**. *Science* 2013, **341**:1384-1387.
- Shows that NFs can suppress MAMP-associated immunity in legumes.
22. Lee KB, De Backer P, Aono T, Liu CT, Suzuki S, Suzuki T, Kaneko T, Yamada M, Tabata S, Kupfer DM *et al.*: **The genome of the versatile nitrogen fixer *Azorhizobium caulinodans* ORS571**. *BMC Genomics* 2008, **9**.
23. Nandasena KG, O'Hara GW, Tiwari RP, Sezmis E, Howieson JG: **In situ lateral transfer of symbiosis islands results in rapid evolution of diverse competitive strains of mesorhizobia suboptimal in symbiotic nitrogen fixation on the pasture legume *Biserrula pelecinus* L.** *Environ Microbiol* 2007, **9**:2496-2511.
24. Sullivan JT, Patrick HN, Lowther WL, Scott DB, Ronson CW: **Nodulating strains of *Rhizobium loti* arise through chromosomal symbiotic gene transfer in the environment**. *Proc Natl Acad Sci U S A* 1995, **92**:8985-8989.
- First demonstration of transfer of MGE to diverse recipient genomes.
25. Edwards J, Johnson C, Santos-Medellin C, Lurie E, Podishetty NK, Bhatnagar S, Eisen JA, Sundareshan V: **Structure, variation, and assembly of the root-associated microbiomes of rice**. *Proc Natl Acad Sci U S A* 2015, **112**:E911-E920.
26. Ling J, Wang H, Wu P, Li T, Tang Y, Naseer N, Zheng HM, Masson-Boivin C, Zhong ZT, Zhu J: **Plant nodulation inducers enhance horizontal gene transfer of *Azorhizobium caulinodans* symbiosis island**. *Proc Natl Acad Sci U S A* 2016, **113**:13875-13880.
27. Sachs JL, Skophammer RG, Regus JU: **Evolutionary transitions in bacterial symbiosis**. *Proc Natl Acad Sci U S A* 2011, **108**:10800-10807.
28. Shapiro BJ, Levade I, Kovacicova G, Taylor RK, Almagro-Moreno S: **Origins of pandemic *Vibrio cholerae* from environmental gene pools**. *Nat Microbiol* 2017, **2**.
29. Pobigaylo N, Szymczak S, Nattkemper TW, Becker A: **Identification of genes relevant to symbiosis and competitiveness in *Sinorhizobium meliloti* using signature-tagged mutants**. *Mol Plant Microbe Interact* 2008, **21**:219-231.
30. Capela D, Marchetti M, Clerissi C, Perrier A, Guetta D, Gris C, Valls M, Jauneau A, Cruveiller S, Rocha EPC *et al.*: **Recruitment of a lineage-specific virulence regulatory pathway promotes intracellular infection by a plant pathogen experimentally evolved into a legume symbiont**. *Mol Biol Evol* 2017, **34**:2503-2521.
- Shows that the rewiring of a genetic network regulating virulence allows a radically different type of symbiotic interaction during experimental evolution of a plant pathogen into legume symbionts.
31. Remigi P, Capela D, Clerissi C, Tasse L, Torchet R, Bouchez O, Batut J, Cruveiller S, Rocha EP, Masson-Boivin C: **Transient hypermutagenesis accelerates the evolution of legume endosymbionts following horizontal gene transfer**. *PLoS Biol* 2014, **12**:e1001942.
- Experimentally shows that the co-transfer of *nod* genes with error-prone polymerases accelerates the evolution of rhizobia.
32. Zhao R, Liu LX, Zhang YZ, Jiao J, Cui WJ, Zhang B, Wang XL, Li ML, Chen Y, Xiong *et al.*: **Adaptive evolution of rhizobial symbiotic compatibility mediated by co-evolved insertion sequences**. *ISME J* 2017 <http://dx.doi.org/10.1038/ismej.2017.136>.
33. Heath KD, Tiffin P: **Stabilizing mechanisms in a legume-rhizobium mutualism**. *Evolution* 2009, **63**:652-662.
34. Sachs JL, Russell JE, Lii YE, Black KC, Lopez G, Patil AS: **Host control over infection and proliferation of a cheater symbiont**. *J Evol Biol* 2010, **23**:1919-1927.
35. Tang F, Yang SM, Liu JG, Zhu HY: **Rj4, a gene controlling nodulation specificity in soybeans, encodes a thaumatin-like protein but not the one previously reported**. *Plant Physiol* 2016, **170**:26-32.
36. Hahn M, Studer D: **Competitiveness of a *nif⁻* *Bradyrhizobium japonicum* mutant against the wild-type strain**. *FEMS Microbiol Lett* 1986:143-148.
37. Amarger N: **Competition for nodule formation between effective and ineffective strains of *Rhizobium meliloti***. *Soil Biol Biochem* 1981, **13**:475-480.
38. Kiers ET, Rousseau RA, West SA, Denison RF: **Host sanctions and the legume-rhizobium mutualism**. *Nature* 2003, **425**:78-81.
39. Oono R, Anderson CG, Denison RF: **Failure to fix nitrogen by non-reproductive symbiotic rhizobia triggers host sanctions that reduce fitness of their reproductive clonemates**. *Proc R Soc B Biol Sci* 2011, **278**:2698-2703.
40. Regus JU, Gano KA, Hollowell AC, Sachs JL: **Efficiency of partner choice and sanctions in *Lotus* is not altered by nitrogen fertilization**. *Proc R Soc B Biol Sci* 2014, **281**.
41. Simms EL, Taylor DL, Povich J, Shefferson RP, Sachs JL, Urbina M, Tausczik Y: **An empirical test of partner choice mechanisms in a wild legume-rhizobium interaction**. *Proc R Soc B Biol Sci* 2006, **273**:77-81.
42. Regus JU, Gano KA, Hollowell AC, Sofish V, Sachs JL: ***Lotus* hosts delimit the mutualism-parasitism continuum of *Bradyrhizobium***. *J Evol Biol* 2015, **28**:447-456.
43. West SA, Kiers ET, Pen I, Denison RF: **Sanctions and mutualism stability: when should less beneficial mutualists be tolerated?** *J Evol Biol* 2002, **15**:830-837.
44. West SA, Kiers ET, Simms EL, Denison RF: **Sanctions and mutualism stability: why do rhizobia fix nitrogen?** *Proc R Soc B Biol Sci* 2002, **269**:685-694.
45. Regus JU, Quides KJ, O'Neill MR, Suzuki R, Savory EA, Chang JH, Sachs J: **Cell autonomous sanctions in legumes target ineffective rhizobia in nodules with mixed infections**. *Am J Bot* 2017, **104**:1299-1312.
- Discovery of cell-autonomy of legume sanctions in determinate nodules.
46. Daubech B, Remigi P, Doin de Moura G, Marchetti M, Pouzet C, Auriac MC, Gokhale CS, Masson-Boivin C, Capela D: **Spatio-temporal control of mutualism in legumes helps spread symbiotic nitrogen fixation**. *Elife* 2017 <http://dx.doi.org/10.7554/eLife.28683>.
- Evidence of cell-autonomy of legume sanctions in indeterminate nodules. Shows that rare nitrogen fixing variants invade a non-fixing population in appropriate conditions.
47. VanInsberghe D, Maas KR, Cardenas E, Strachan CR, Hallam SJ, Mohn WW: **Non-symbiotic *Bradyrhizobium* ecotypes dominate North American forest soils**. *ISME J* 2015, **9**:2435-2441.
48. Parker MA: **The spread of *Bradyrhizobium* lineages across host legume clades: from *Abarema* to *Zygia***. *Microbial Ecol* 2015, **69**:630-640.
49. Denison RF, Kiers ET: **Lifestyle alternatives for rhizobia: mutualism, parasitism, and forgoing symbiosis**. *FEMS Microbiol Lett* 2004, **237**:187-193.
50. Sachs JL, Russell JE, Hollowell AC: **Evolutionary instability of symbiotic function in *Bradyrhizobium japonicum***. *PLoS One* 2011, **6**:10.

51. Thies JE, Wooster PL, Singleton PW: **Enrichment of *Bradyrhizobium* spp. populations in soil due to cropping of the homologous host legume.** *Soil Biol Biochem* 1995, **27**: 633-636.
52. Nandasena KG, O'Hara GW, Tiwari RP, Howieson JG: **Rapid in situ evolution of nodulating strains for *Biserrula pelecinus* L. through lateral transfer of a symbiosis island from the original mesorhizobial inoculant.** *Appl Environ Microbiol* 2006, **72**: 7365-7367.
53. Parker MA: **Legumes select symbiosis island sequence variants in *Bradyrhizobium*.** *Mol Ecol* 2012, **21**:1769-1778.
54. Sachs JL, Simms EL: **The origins of uncooperative rhizobia.** *Oikos* 2008, **117**:961-966.
55. Porter SS, Simms EL: **Selection for cheating across disparate environments in the legume-rhizobium mutualism.** *Ecol Lett* 2014, **17**:1121-1129.
56. Kiers ET, Hutton MG, Denison RF: **Human selection and the relaxation of legume defences against ineffective rhizobia.** *Proc R Soc B Biol Sci* 2007, **274**:3119-3126.
57. Weese DJ, Heath KD, Dentinger BTM, Lau JA: **Long-term nitrogen addition causes the evolution of less-cooperative mutualists.** *Evolution* 2015, **69**:631-642.
58. Gano-Cohen KA, Stokes PJ, Blanton MA, Wendlandt CE, Hollowell AC, Regus JU, Kim D, Patel S, Pahua VJ, Sachs JL: **Nonnodulating *Bradyrhizobium* spp. modulate the benefits of legume-rhizobium mutualism.** *Appl Environ Microbiol* 2016, **82**:5259-5268.
59. Hollowell AC, Regus JU, Gano KA, Bantay R, Centeno D, Pham J, Lyu J, Moore D, Bernardo A, Lopez G *et al.*: **Epidemic spread of symbiotic and non-symbiotic *Bradyrhizobium* genotypes across California.** *Microbial Ecol* 2016, **71**:700-710.
60. Hollowell AC, Regus JU, Turissini D, Gano-Cohen KA, Bantay R, Bernardo A, Moore D, Pham J, Sachs JL: **Metapopulation dominance and genomic-island acquisition of *Bradyrhizobium* with superior catabolic capabilities.** *Proc R Soc B Biol Sci* 2016, **283**.
61. Sachs JL, Skophammer RG, Bansal N, Stajich JE: **Evolutionary origins and diversification of proteobacterial mutualists.** *Proc R Soc B Biol Sci* 2014, **281**:9.
62. Ferguson BJ, Indrasumunar A, Hayashi S, Lin MH, Lin YH, Reid DE, Gresshoff PM: **Molecular analysis of legume nodule development and autoregulation.** *J Integr Plant Biol* 2010, **52**: 61-76.
63. Yates RJ, Howieson JG, Reeve WG, O'Hara GW: **A re-appraisal of the biology and terminology describing rhizobial strain success in nodule occupancy of legumes in agriculture.** *Plant Soil* 2011, **348**:255-267.
64. Marchetti M, Capela D, Glew M, Cruveiller S, Chane-Woon-Ming B, Gris C, Timmers T, Poinso V, Gilbert LB, Heeb P *et al.*: **Experimental evolution of a plant pathogen into a legume symbiont.** *PLoS Biol* 2010, **8**.
65. Guan SH, Gris C, Cruveiller S, Pouzet C, Tasse L, Leru A, Maillard A, Medigue C, Batut J, Masson-Boivin C *et al.*: **Experimental evolution of nodule intracellular infection in legume symbionts.** *ISME J* 2013, **7**:1367-1377.
66. Mueller UG, Sachs JL: **Engineering microbiomes to improve plant and animal health.** *Trends Microbiol* 2015, **23**:606-617.
67. Yang JG, Xie XQ, Yang MX, Dixon R, Wang YP: **Modular electron-transport chains from eukaryotic organelles function to support nitrogenase activity.** *Proc Natl Acad Sci U S A* 2017, **114**:E2460-E2465.
- Role of some NCR peptides in selecting specific bacterial genotypes.
68. Chabaud M, Gherbi H, Piroilles E, Vaissayre V, Fournier J, Moukouanga D, Franche C, Bogusz D, Tisa LS, Barker DG *et al.*: **Chitinase-resistant hydrophilic symbiotic factors secreted by Frankia activate both Ca²⁺ spiking and NIN gene expression in the actinorhizal plant *Casuarina glauca*.** *New Phytol* 2016, **209**:86-93.
69. D'Haese W, Holsters M: **Nod factor structures, responses, and perception during initiation of nodule development.** *Glycobiology* 2002, **12**:79R-105R.
70. Okazaki S, Kaneko T, Sato S, Saeki K: **Hijacking of leguminous nodulation signaling by the rhizobial type III secretion system.** *Proc Natl Acad Sci U S A* 2013, **110**:17131-17136.
- Shows that *Bradyrhizobium* T3SS can activate nodulation in legumes.
71. Okazaki S, Tittabutr P, Teulet A, Thouin J, Fardoux J, Chaintreuil C, Gully D, Arrighi JF, Furuta N, Miwa H *et al.*: **Rhizobium-legume symbiosis in the absence of Nod factors: two possible scenarios with or without the T3SS.** *ISME J* 2016, **10**:64-74.
- Evidence of two distinct Nod-alternative strategies, the T3SS and non-Nod/non T3SS.
72. Giraud E, Moulin L, Vallenet D, Barbe V, Cytryn E, Avarre JC, Jaubert M, Simon D, Cartieaux F, Prin Y *et al.*: **Legumes symbioses: absence of Nod genes in photosynthetic bradyrhizobia.** *Science* 2007, **316**:1307-1312.
- First evidence of an alternative strategy to the Nod strategy.
73. Bonaldi K, Gourion B, Fardoux J, Hannibal L, Cartieaux F, Boursot M, Vallenet D, Chaintreuil C, Prin Y, Nouwen N *et al.*: **Large-scale transposon mutagenesis of photosynthetic *Bradyrhizobium* sp. strain ORS278 reveals new genetic loci putatively important for nod-independent symbiosis with *Aeschynomene indica*.** *Mol Plant Microbe Interact* 2010, **23**: 760-770.
74. Chaintreuil C, Rivallan R, Bertoli DJ, Klopp C, Gouzy J, Courtois B, Leleux P, Martin G, Rami JF, Gully D *et al.*: **A gene-based map of the Nod factor-independent *Aeschynomene evenia* sheds new light on the evolution of nodulation and legume genomes.** *DNA Res* 2016, **23**:365-376.
75. Fabre S, Gully D, Poitout A, Patrel D, Arrighi JF, Giraud E, Czernic P, Cartieaux F: **Nod factor-independent nodulation in *Aeschynomene evenia* required the common plant-microbe symbiotic toolkit.** *Plant Physiol* 2015, **169**:2654-2664.
76. Sugawara M, Epstein B, Badgley BD, Unno T, Xu L, Reese J, Gyaneshwar P, Denny R, Mudge J, Bharti AK *et al.*: **Comparative genomics of the core and accessory genomes of 48 *Sinorhizobium* strains comprising five genospecies.** *Genome Biol* 2013, **14**.
77. Mazurier S, Lemunier M, Hartmann A, Siblot S, Lemanceau P: **Conservation of type III secretion system genes in *Bradyrhizobium* isolated from soybean.** *FEMS Microbiol Lett* 2006, **259**:317-325.
78. Deakin WJ, Broughton WJ: **Symbiotic use of pathogenic strategies: rhizobial protein secretion systems.** *Nat Rev Microbiol* 2009, **7**:312-320.
79. Chaintreuil C, Arrighi JF, Giraud E, Miche L, Moulin L, Dreyfus B, Munive-Hernandez JA, Villegas-Hernandez MD, Bena G: **Evolution of symbiosis in the legume genus *Aeschynomene*.** *New Phytol* 2013, **200**:1247-1259.
80. Okubo T, Fukushima S, Minamisawa K: **Evolution of *Bradyrhizobium-Aeschynomene* mutualism: living testimony of the ancient world or highly evolved state?** *Plant Cell Physiol* 2012, **53**:2000-2007.
81. Masson-Boivin C, Giraud E, Perret X, Batut J: **Establishing nitrogen-fixing symbiosis with legumes: how many rhizobium recipes?** *Trends Microbiol* 2009, **17**:458-466.
82. Marchetti M, Jauneau A, Capela D, Remigi P, Gris C, Batut J, Masson-Boivin C: **Shaping bacterial symbiosis with legumes by experimental evolution.** *Mol Plant Microbe Interact* 2014, **27**: 956-964.
83. Czernic P, Gully D, Cartieaux F, Moulin L, Guefrachi I, Patrel D, Pierre O, Fardoux J, Chaintreuil C, Nguyen P *et al.*: **Convergent evolution of endosymbiont differentiation in dalbergioid and inverted repeat-lacking clade legumes mediated by nodule-specific cysteine-rich peptides.** *Plant Physiol* 2015, **169**: 1254-1265.
84. Van de Velde W, Zehirov G, Szatmari A, Debreczeny M, Ishihara H, Kevei Z, Farkas A, Mikulass K, Nagy A, Tiricz H *et al.*: **Plant peptides govern terminal differentiation of bacteria in symbiosis.** *Science* 2010, **327**:1122-1126.

85. Wang Q, Yang SM, Liu JG, Terecskei K, Abraham E, Gombor A, Domonkos A, Szucs A, Kormoczi P, Wang T *et al.*: **Host-secreted antimicrobial peptide enforces symbiotic selectivity in *Medicago truncatula***. *Proc Natl Acad Sci U S A* 2017, **114**: 6854-6859.

Role of some NCR peptides in selecting specific bacterial genotypes.

86. Yang SM, Wang Q, Fedorova E, Liu JG, Qin QL, Zheng QL, Price PA, Pan HR, Wang D, Griffiths JS *et al.*: **Microsymbiont**

discrimination mediated by a host-secreted peptide in *Medicago truncatula*. *Proc Natl Acad Sci U S A* 2017, **114**: 6848-6853.

Role of some NCR peptides in selecting specific bacterial genotypes.

87. Marlow VL, Haag AF, Kobayashi H, Fletcher V, Scocchi M, Walker GC, Ferguson GP: **Essential role for the BacA protein in the uptake of a truncated eukaryotic peptide in *Sinorhizobium meliloti***. *J Bacteriol* 2009, **191**:1519-1527.