

UC Irvine

UC Irvine Previously Published Works

Title

Tinnitus and hyperacusis: central noise, gain and variance

Permalink

<https://escholarship.org/uc/item/3558t02x>

Author

Zeng, Fan-Gang

Publication Date

2020-12-01

DOI

10.1016/j.cophys.2020.10.009

Peer reviewed



Published in final edited form as:

Curr Opin Physiol. 2020 December ; 18: 123–129. doi:10.1016/j.cophys.2020.10.009.

Tinnitus and hyperacusis: Central noise, gain and variance

Fan-Gang Zeng

Departments of Anatomy and Neurobiology, Biomedical Engineering, Cognitive Sciences, and Otolaryngology – Head and Neck Surgery, Center for Hearing Research, University of California Irvine

Abstract

Tinnitus is a phantom auditory sensation in the absence of external sounds, while hyperacusis is an atypical sensitivity to external sounds that leads them to be perceived as abnormally loud or even painful. Both conditions may reflect the brain's over-compensation for reduced input from the ear. The present work differentiates between two compensation models: The additive central noise compensates for hearing loss and is likely to generate tinnitus, whereas the multiplicative central gain compensates for hidden hearing loss and is likely to generate hyperacusis. Importantly, both models predict increased variance in central representations of sounds, especially a nonlinear increase in variance by the central gain. The increased central variance limits the amount of central compensation and reduces temporal synchrony, which can explain the insufficient central gain reported in the literature. Future studies need to collect trial-by-trial firing variance data so that the present variance-based model can be falsified.

Keywords

Central Variance; Central Noise; Central Gain; Temporal Coding; Dynamic Range Normalization

Introduction

Tinnitus is a phantom perception of a sound in the absence of external sounds, whereas hyperacusis is an excessive response to external sounds [1]. Tinnitus affects 15% of the general population and is typically perceived as a ringing sensation in the ear or head. Individuals with hyperacusis cannot tolerate conventional loud sounds or even report painful responses to these sounds. At first glance, tinnitus and hyperacusis are opposites, but they are intertwined clinically. Nearly half of tinnitus sufferers report some degree of hyperacusis, while most hyperacusis sufferers have tinnitus [2].

Contact information: 110 Medical Science E, University of California Irvine, Irvine, CA 92697-5320, USA, Phone: (949) 824-1539, fzenng@uci.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of interest statement

The author owns stock in Axonics, DiaNavi, Nurotron, Syntiant, Velox and Xense.

Hearing loss is a common and main risk factor for both conditions [3, 4]. Recent advances have broadened the definition of hearing loss, which traditionally was defined by elevated audiometric thresholds between 125 and 8000 Hz. A key advance is the discovery of “hidden” hearing loss, in which there is no elevation of audio metric thresholds but there is reduced auditory nerve activity due to cochlear synaptopathy and secondary nerve degeneration [5]. Although some individuals with tinnitus do not have any hearing loss, they may suffer from hidden hearing loss [6, 7]. Indeed, studies on tinnitus without hearing loss often found evidence for the presence of hidden hearing loss and its involvement in hyperacusis in humans [7-10] or animals [11-13]. The current view is that both tinnitus and hyperacusis are consequences of the central auditory system’s compensatory responses to the reduced peripheral input due to either overt or hidden hearing loss.

At present, the exact central compensatory mechanisms are not clear. Hearing loss causes massive changes in the central auditory system, from to notopic reorganization and hyperactivity to altered network connectivity within and beyond the auditory system [14-18]. Some of these central changes cause tinnitus and hyperacusis but others may not. For example, excessive to notopic reorganization was thought of as a main mechanism for tinnitus [19], but a recent brain imaging study found the opposite; only those hearing-impaired individuals who have incomplete tonotopic reorganization develop tinnitus [20]. In contrast, central hyperactivity has been consistently observed from the cochlear nucleus to the auditory cortex, evidenced by increased spontaneous firing rate or restored driven activity [21-26]. Because hearing loss actually reduces both spontaneous and driven activity in the auditory nerve fibers [27, 28], such central hyperactivity is considered to be the result of a compensatory response to hearing loss. Tinnitus and hyperacusis are likely a consequence of this central compensation or over-compensation [7, 29, 30]. In addition to the bottom-up compensatory mechanism, several top-down mechanisms such as noise cancellation, predictive errors or attention may influence the generation and behavior of tinnitus and hyperacusis [31-34].

The present work builds upon the previous suggestions that tinnitus is caused by increased central noise and hyperacusis by increased central gain [30]. Here I propose a third mechanism, namely central variance, which may affect the relative contributions of central noise and central gain to tinnitus and hyperacusis.

First consider what happens when the total central activity is the sum of compensatory additive central noise (N) and driven activity (D). Then the total mean activity is the sum of the two individual means:

$$\text{Mean}(N + D) = \text{Mean}(N) + \text{Mean}(D) \quad (1)$$

while the total variance is the sum of the two variances [p140 in 35]:

$$\text{Var}(N + D) = \text{Var}(N) + \text{Var}(D) \quad (2)$$

Alternatively, applying a gain (g) to the driving activity results in a proportional increase in the total mean activity:

$$\text{Mean}(gD) = g\text{Mean}(D); \quad (3)$$

however, the total variance increases nonlinearly by the square of the gain [p106 in 35]:

$$\text{Var}(gD) = g^2\text{Var}(D) \quad (4)$$

Thus, increasing central noise and increasing central gain lead to different predictions for how central variance changes. The next section considers how these different effects on central variance may help explain tinnitus and hyperacusis.

Tinnitus: Increased central noise and variance

Figure 1(a) shows schematically the effect of hearing loss on auditory nerve activity as a function of sound level (left panel) and time (right panel). Compared with the normal mean activity (blue lines), hearing loss elevates the hearing threshold (double-angled green line below the x-axis, left panel) and reduces the overall auditory nerve mean activity (solid green lines in both panels) [28]. To a first-order approximation [36], the firing variance is proportional to the mean firing rate (dashed lines, left panel; or amplitude fluctuations, right panel).

Figure 1(b) shows the effects of adding a central noise on loudness represented at a higher level of auditory processing, both as a function of sound level (left panel) and time (right panel). The additive central noise (represented by two upward angled black lines, left panel) restores normal loudness growth at high sound levels (solid orange sloping line, left panel) but raises the baseline loudness at and below threshold (solid orange horizontal line above “Tinnitus”, both panels). Similarly, the central noise increases the baseline variance (dashed orange horizontal line, left panel, and amplitude fluctuations above “Tinnitus”, right panel) but not the slope of the normal variance function (dashed blue line).

These schematical effects are consistent with behavioral observations that individuals with hearing loss have a normal slope of the loudness function but abnormally large loudness at the threshold [37]. In other words, the central noise compensates for hearing loss by restoring normal loudness growth above threshold, but generates tinnitus at or below the hearing threshold.

Hyperacusis: Increased central gain and variance

Figure 2(a) shows a schematic similar to Figure 1(a), except for hidden hearing loss. Unlike in traditional hearing loss, hidden hearing loss does not elevate the hearing threshold, but instead produces a shallower-than-normal auditory nerve rate-level function (solid purple line=mean; dashed purple line=variance, left panel) and reduced activity (purple line, right panel).

Figure 2(b) shows the effects of increasing central gain on loudness as a function of sound level (left panel) and time (right panel). The multiplicative central gain (a factor of two in this case) restores mean normal loudness growth at all sound levels (solid diagonal red

line=solid blue line, left panel) but increases the amount of the central variance by a factor of four. As a result, the quadrupled variance produces hyperacusis at high sound levels (double-headed red line, left panel; and solid red lines outside the box reflecting amplitude fluctuations beyond normal loudness range, right panel).

Central variance limits the amount of compensation

This analysis of a central noise model assumes that internal noise and driven activities are summed up in the central auditory system [38]. If the internal and driven activities are not additive but instead multiplicative [26], then the resulting variance can be 10-100 times larger than the individual variance, depending on the squared terms of both individual means and variances [39]. Similarly, if the actual central gain compensation involves a cascade of amplifiers [13, 40], then the central variance can grow geometrically. In reality, an individual may have mixed hearing loss, requiring both forms of compensation, which further exacerbate the central variance problem.

An inevitable consequence of the large central variance is a severe limitation on the amount of central compensation that can be accommodated. To minimize the likelihood of tinnitus and hyperacusis, neither the central noise nor the central gain would be expected to reach the level that restores the normal mean value. This lower-than-expected compensation limit is more pronounced for increases in central gain than for additive noise because the variance increases much more when the compensation is multiplicative rather than additive. Thus, it is not too surprising that recent studies found insufficient central gain because they all used the normal mean response as the criterion [41-44].

Central variance reduces temporal synchrony

Considering how increased variance influences central representations reconciles other conflicting results in the literature. For example, early studies found a reduced auditory brainstem response for wave I but a normal or even increased wave V response in human tinnitus subjects with normal audiograms [7, 45]. Recent studies could not replicate these findings in a large cohort of similar human tinnitus subjects [44, 46] or in subjects with a history of noise exposure [47]. Instead, several studies found delayed auditory brainstem responses [48, 49].

The auditory brainstem response is determined by the total synchronized activity across all responding neurons [50, see their Fig. 13]. It is possible that early studies used subjects with a relatively small central gain while recent studies used subjects with a larger central gain. The linearly increasing gain cannot make up for the dys-synchrony caused by the nonlinearly increasing variance, such that a synchronized amplitude measure, like wave V, underestimates the true central gain. In this case, more dys-synchrony actually indicates larger gain. Indeed, there is evidence showing that temporal measures are more sensitive than amplitude measures in detecting hidden hearing loss in humans [51, 52]. Of course, single-unit recordings would not have this gain vs. variance dilemma, but at present no studies have reported trial-to-trial firing rate variance from single units.

Dynamic range normalization

The central variance problem can be reduced by dynamic range normalization [53], which is widely used in the nervous system to adapt to external signals [54] and internal noise [55] in order to preserve stimulus sensitivity and prevent response saturation [56-59]. The neural system also needs to maintain the same average energy, e.g., the same average firing rate [60]. Activation of the fast-spiking auditory pathway [18] may be such a normalization mechanism. Failure to normalize the increased gain and variance may lead to not only tinnitus and hyperacusis [20, 31, 32] but also excitation spread towards other brain areas controlling pain, fear or other emotional responses [61-63].

Conclusion

There are two potential mechanisms allowing the central system to compensate for reduced peripheral input: Increasing central noise or increasing central gain. An inevitable consequence of both mechanisms is an increase in central variance, which can help explain both tinnitus and hyperacusis. An additive central noise, which compensates for elevated hearing thresholds like those seen in traditional hearing loss, is more likely to induce tinnitus than hyperacusis. In contrast, an increase in multiplicative central gain, which compensates for hidden hearing loss, is more likely to induce hyperacusis. The effects of compensatory mechanisms on central variance have been ignored in previous studies, but may account for the conflicting results in the generation and behavior of tinnitus and hyperacusis.

Acknowledgements

The work was partially supported by the National Institutes of Health (5R01 DC015587). I thank Barbara Shinn-Cunningham, Paul Fuchs, Nick Lesica, Michelle Kaplowicz, Phillip Tran and Ye Yang for commenting on the manuscript.

References


- [1]. Ke J, Du Y, Tyler RS, Perreau A, Mancini PC, Complaints of People with Hyperacusis, *Journal of the American Academy of Audiology* (2020).
- [2]. Anari M, Axelsson A, Eliasson A, Magnusson L, Hypersensitivity to sound--questionnaire data, audiometry and classification, *Scandinavian audiology* 28(4) (1999) 219–30. [PubMed: 10572967]
- [3]. Baguley D, McFerran D, Hall D, Tinnitus, *Lancet* 382(9904) (2013) 1600–7. [PubMed: 23827090]
- [4]. Hertzano R, Lipford EL, Depireux D, Noise: Acoustic Trauma to the Inner Ear, *Otolaryngologic clinics of North America* 53(4) (2020) 531–542. [PubMed: 32362563]
- [5]. Kujawa SG, Liberman MC, Adding insult to injury: cochlear nerve degeneration after "temporary" noise-induced hearing loss, *J Neurosci* 29(45) (2009) 14077–85. [PubMed: 19906956]
- [6]. Hazell JW, Jastreboff PJ, Tinnitus. I: Auditory mechanisms: a model for tinnitus and hearing impairment, *J. Otolaryngol* 19(1) (1990) 1–5.
- [7]. Schaeffe R, McAlpine D, Tinnitus with a normal audiogram: physiological evidence for hidden hearing loss and computational model, *J. Neurosci* 31(38) (2011) 13452–7. [PubMed: 21940438]
- [8]. Gu JW, Halpin CF, Nam EC, Levine RA, Melcher JR, Tinnitus, diminished sound-level tolerance, and elevated auditory activity in humans with clinically normal hearing sensitivity, *Journal of neurophysiology* 104(6) (2010) 3361–70. [PubMed: 20881196]

- [9]. Fournier P, Hebert S, Gap detection deficits in humans with tinnitus as assessed with the acoustic startle paradigm: does tinnitus fill in the gap?, *Hearing research* 295 (2013) 16–23. [PubMed: 22688322]
- [10]. Hebert S, Fournier P, Norena A, The auditory sensitivity is increased in tinnitus ears, *J Neurosci* 33(6) (2013) 2356–64. [PubMed: 23392665]
- [11]. Shaheen LA, Liberman MC, Cochlear Synaptopathy Changes Sound-Evoked Activity Without Changing Spontaneous Discharge in the Mouse Inferior Colliculus, *Front Syst Neurosci* 12 (2018) 59. [PubMed: 30559652]
- [12]. Thomas ME, Guercio GD, Drudik KM, de Villers-Sidani E, Evidence of Hyperacusis in Adult Rats Following Non-traumatic Sound Exposure, *Frontiers in systems neuroscience* 13 (2019) 55. [PubMed: 31708754]
- [13]. Wong E, Radziwon K, Chen GD, Liu X, Manno FA, Manno SH, Auerbach B, Wu EX, Salvi R, Lau C, Functional magnetic resonance imaging of enhanced central auditory gain and electrophysiological correlates in a behavioral model of hyperacusis, *Hearing research* 389 (2020) 107908. [PubMed: 32062293]
- [14]. Roberts LE, Eggermont JJ, Caspary DM, Shore SE, Melcher JR, Kaltenbach JA, Ringing Ears: The Neuroscience of Tinnitus, *J. Neurosci* 30(45) (2010) 14972–14979. [PubMed: 21068300]
- [15]. Shore SE, Roberts LE, Langguth B, Maladaptive plasticity in tinnitus--triggers, mechanisms and treatment, *Nat Rev Neurol* 12(3) (2016) 150–60. [PubMed: 26868680]
- [16]. Husain FT, Neural networks of tinnitus in humans: Elucidating severity and habituation, *Hearing research* 334 (2016) 37–48. [PubMed: 26415997]
- [17]. Chen YC, Li X, Liu L, Wang J, Lu CQ, Yang M, Jiao Y, Zang FC, Radziwon K, Chen GD, Sun W, Krishnan Muthaiah VP, Salvi R, Teng GJ, Tinnitus and hyperacusis involve hyperactivity and enhanced connectivity in auditory-limbic-arousal-cerebellar network, *Elife* 4 (2015) e06576. [PubMed: 25962854]
- [18]. Knipper M, van Dijk P, Schulze H, Mazurek B, Krauss P, Scheper V, Warnecke A, Schlee W, Schwabe K, Singer W, Braun C, Delano PH, Fallgatter AJ, Ehlis AC, Searchfield GD, Munk MHJ, Baguley DM, Ruttiger L, The Neural Bases of Tinnitus: Lessons from Deafness and Cochlear Implants, *J Neurosci* 40(38) (2020) 7190–7202. [PubMed: 32938634]
- [19]. Muhlnickel W, Elbert T, Taub E, Flor H, Reorganization of auditory cortex in tinnitus, *Proceedings of the National Academy of Sciences of the United States of America* 95(17) (1998) 10340–3. [PubMed: 9707649]
- [20]. Koops EA, Renken RJ, Lanting CP, van Dijk P, Cortical Tonotopic Map Changes in Humans Are Larger in Hearing Loss Than in Additional Tinnitus, *J Neurosci* 40(16) (2020) 3178–3185. [PubMed: 32193229]
- [21]. Kaltenbach JA, Tinnitus: Models and mechanisms, *Hear. Res* 276(1-2) (2011) 52–60. [PubMed: 21146597]
- [22]. Norena AJ, Eggermont JJ, Changes in spontaneous neural activity immediately after an acoustic trauma: implications for neural correlates of tinnitus, *Hearing research* 183(1-2) (2003) 137–53. [PubMed: 13679145]
- [23]. Chambers AR, Resnik J, Yuan Y, Whitton JP, Edge AS, Liberman MC, Polley DB, Central Gain Restores Auditory Processing following Near-Complete Cochlear Denervation, *Neuron* 89(4) (2016) 867–79. [PubMed: 26833137]
- [24]. Kaltenbach JA, Afman CE, Hyperactivity in the dorsal cochlear nucleus after intense sound exposure and its resemblance to tone-evoked activity: a physiological model for tinnitus, *Hearing research* 140(1-2) (2000) 165–72. [PubMed: 10675644]
- [25]. Shore SE, Wu C, Mechanisms of Noise-Induced Tinnitus: Insights from Cellular Studies, *Neuron* 103(1) (2019) 8–20. [PubMed: 31271756]
- [26]. Eggermont JJ, Animal models of spontaneous activity in the healthy and impaired auditory system, *Front Neural Circuits* 9 (2015) 19. [PubMed: 25983679]
- [27]. Liberman MC, Dodds LW, Single-neuron labeling and chronic cochlear pathology. II. Stereocilia damage and alterations of spontaneous discharge rates, *Hear. Res* 16(1) (1984) 43–53. [PubMed: 6511672]

- [28]. Heinz MG, Issa JB, Young ED, Auditory-nerve rate responses are inconsistent with common hypotheses for the neural correlates of loudness recruitment, *J Assoc Res Otolaryngol* 6(2) (2005) 91–105. [PubMed: 15952047]
- [29]. Norena AJ, An integrative model of tinnitus based on a central gain controlling neural sensitivity, *Neurosci. Biobehav. Rev* 35(5) (2011) 1089–109. [PubMed: 21094182]
- [30]. Zeng FG, An active loudness model suggesting tinnitus as increased central noise and hyperacusis as increased nonlinear gain, *Hearing research* 295 (2013) 172–9. [PubMed: 22641191]
- [31]. Sedley W, Tinnitus: Does Gain Explain?, *Neuroscience* 407 (2019) 213–228. [PubMed: 30690137]
- [32]. Vanneste S, Alsaman O, De Ridder D, Top-down and Bottom-up Regulated Auditory Phantom Perception, *J Neurosci* 39(2) (2019) 364–378. [PubMed: 30389837]
- [33]. Rauschecker JP, Leaver AM, Muhlau M, Tuning out the noise: limbic-auditory interactions in tinnitus, *Neuron* 66(6) (2010) 819–26. [PubMed: 20620868]
- [34]. Zeng FG, Richardson M, Turner K, Tinnitus Does Not Interfere with Auditory and Speech Perception, *J Neurosci* 40(31) (2020) 6007–6017. [PubMed: 32554549]
- [35]. Dekking M, A modern introduction to probability and statistics : understanding why and how, Springer, London, 2005.
- [36]. Johnson DH, Swami A, The transmission of signals by auditory-nerve fiber discharge patterns, *J. Acoust. Soc. Am* 74(2) (1983) 493–501. [PubMed: 6311884]
- [37]. Buus S, Florentine M, Growth of loudness in listeners with cochlear hearing losses: recruitment reconsidered, *J Assoc Res Otolaryngol* 3(2) (2002) 120–39. [PubMed: 12162363]
- [38]. Bender DA, Ni R, Barbour DL, Spontaneous activity is correlated with coding density in primary auditory cortex, *Journal of neurophysiology* 116(6) (2016) 2789–2798. [PubMed: 27707812]
- [39]. Ware R, Lad F, Approximating the distribution for sums of products of normal variables, Research-Paper 2003-15 Department of Mathematics and Statistics, University of Canterbury, University of Canterbury, New Zealand, 2003.
- [40]. Jiang C, Luo B, Manohar S, Chen GD, Salvi R, Plastic changes along auditory pathway during salicylate-induced ototoxicity: Hyperactivity and CF shifts, *Hearing research* 347 (2017) 28–40. [PubMed: 27989950]
- [41]. Radziwon K, Auerbach BD, Ding D, Liu X, Chen GD, Salvi R, Noise-Induced loudness recruitment and hyperacusis: Insufficient central gain in auditory cortex and amygdala, *Neuroscience* 422 (2019) 212–227. [PubMed: 31669363]
- [42]. Mohrle D, Hofmeier B, Amend M, Wolpert S, Ni K, Bing D, Klose U, Pichler B, Knipper M, Ruttiger L, Enhanced Central Neural Gain Compensates Acoustic Trauma-induced Cochlear Impairment, but Unlikely Correlates with Tinnitus and Hyperacusis, *Neuroscience* 407 (2019) 146–169. [PubMed: 30599268]
- [43]. Brotherton H, Turtle C, Plack CJ, Munro KJ, Schaette R, Earplug-induced changes in acoustic reflex thresholds suggest that increased subcortical neural gain may be necessary but not sufficient for the occurrence of tinnitus, *Neuroscience* 407 (2019) 192–199. [PubMed: 30890481]
- [44]. Hofmeier B, Wolpert S, Aldamer ES, Walter M, Thiericke J, Braun C, Zelle D, Ruttiger L, Klose U, Knipper M, Reduced sound-evoked and resting-state BOLD fMRI connectivity in tinnitus, *Neuroimage Clin* 20 (2018) 637–649. [PubMed: 30202725]
- [45]. Gu JW, Herrmann BS, Levine RA, Melcher JR, Brainstem auditory evoked potentials suggest a role for the ventral cochlear nucleus in tinnitus, *J Assoc Res Otolaryngol* 13(6) (2012) 819–33. [PubMed: 22869301]
- [46]. Shim HJ, An YH, Kim DH, Yoon JE, Yoon JH, Comparisons of auditory brainstem response and sound level tolerance in tinnitus ears and non-tinnitus ears in unilateral tinnitus patients with normal audiograms, *PloS one* 12(12) (2017) e0189157. [PubMed: 29253030]
- [47]. Prendergast G, Guest H, Munro KJ, Kluk K, Leger A, Hall DA, Heinz MG, Plack CJ, Effects of noise exposure on young adults with normal audiograms I: Electrophysiology, *Hearing research* 344 (2017) 68–81. [PubMed: 27816499]

- [48]. Mehraei G, Hickox AE, Bharadwaj HM, Goldberg H, Verhulst S, Liberman MC, Shinn-Cunningham BG, Auditory Brainstem Response Latency in Noise as a Marker of Cochlear Synaptopathy, *J Neurosci* 36(13) (2016) 3755–64. [PubMed: 27030760]
- [49]. Duron J, Monconduit L, Avan P, Auditory Brainstem Changes in Timing may Underlie Hyperacusis in a Salicylate-induced Acute Rat Model, *Neuroscience* 426 (2020) 129–140. [PubMed: 31846750]
- [50]. Starr A, Michalewski HJ, Zeng FG, Fujikawa-Brooks S, Linthicum F, Kim CS, Winnier D, Keats B, Pathology and physiology of auditory neuropathy with a novel mutation in the MPZ gene (Tyr145->Ser), *Brain* 126(Pt 7) (2003) 1604–19. [PubMed: 12805115]
- [51]. Paul BT, Bruce IC, Roberts LE, Evidence that hidden hearing loss underlies amplitude modulation encoding deficits in individuals with and without tinnitus, *Hear. Res* 344 (2017) 170–182. [PubMed: 27888040]
- [52]. Shaheen LA, Valero MD, Liberman MC, Towards a Diagnosis of Cochlear Neuropathy with Envelope Following Responses, *Journal of the Association for Research in Otolaryngology : JARO* 16(6) (2015) 727–45. [PubMed: 26323349]
- [53]. Carandini M, Heeger DJ, Normalization as a canonical neural computation, *Nat Rev Neurosci* 13(1) (2012) 51–62.
- [54]. Schwartz O, Simoncelli EP, Natural signal statistics and sensory gain control, *Nat Neurosci* 4(8) (2001) 819–25. [PubMed: 11477428]
- [55]. Dunn FA, Rieke F, The impact of photoreceptor noise on retinal gain controls, *Curr Opin Neurobiol* 16(4) (2006) 363–70. [PubMed: 16837189]
- [56]. Rabinowitz NC, Willmore BD, Schnupp JW, King AJ, Contrast gain control in auditory cortex, *Neuron* 70(6) (2011) 1178–91. [PubMed: 21689603]
- [57]. Dean I, Harper NS, McAlpine D, Neural population coding of sound level adapts to stimulus statistics, *Nat Neurosci* 8(12) (2005) 1684–9. [PubMed: 16286934]
- [58]. Zilany MS, Carney LH, Power-law dynamics in an auditory-nerve model can account for neural adaptation to sound-level statistics, *J Neurosci* 30(31) (2010) 10380–90. [PubMed: 20685981]
- [59]. Herrmann B, Maess B, Johnsrude IS, Aging Affects Adaptation to Sound-Level Statistics in Human Auditory Cortex, *J Neurosci* 38(8) (2018) 1989–1999. [PubMed: 29358362]
- [60]. Bakay WMH, Anderson LA, Garcia-Lazaro JA, McAlpine D, Schaette R, Hidden hearing loss selectively impairs neural adaptation to loud sound environments, *Nat Commun* 9(1) (2018) 4298.
- [61]. Zhang L, Wu C, Martel DT, West M, Sutton MA, Shore SE, Remodeling of cholinergic input to the hippocampus after noise exposure and tinnitus induction in Guinea pigs, *Hippocampus* 29(8) (2019) 669–682. [PubMed: 30471164]
- [62]. Asokan MM, Williamson RS, Hancock KE, Polley DB, Sensory overamplification in layer 5 auditory corticofugal projection neurons following cochlear nerve synaptic damage, *Nat Commun* 9(1) (2018) 2468. [PubMed: 29941910]
- [63]. Zhang GW, Sun WJ, Zingg B, Shen L, He J, Xiong Y, Tao HW, Zhang LI, A Non-canonical Reticular-Limbic Central Auditory Pathway via Medial Septum Contributes to Fear Conditioning, *Neuron* 97(2) (2018) 406–417 e4. [PubMed: 29290554]

(a) Hearing loss reduces the overall auditory nerve activity



(b) Additive central noise and variance induce tinnitus






Figure 1. Hearing loss, central noise and central variance.

(a) Schematic effects of hearing loss on auditory nerve activity as a function of sound level (left panel) and time (right panel). Hearing loss elevates the threshold (double-headed line below the x-axis). The normal activity is displayed as blue lines, whereas the activity with hearing loss is displayed as green lines. **(b)** Effects of adding central noise on loudness as a function of sound level (left panel) and time (right panel). The additive central noise (upward arrowed lines, left panel) restores normal loudness growth at high sound levels (solid sloping orange line=solid blue line, left panel) but increases loudness at the threshold or below (solid horizontal orange line, left panel). Both loudness mean and variance (dashed orange line, left panel) are the sum of their corresponding auditory nerve (green lines in Fig. 1(a)) and central noise values. The central noise generates tinnitus at or below the hearing threshold (labeled “Tinnitus” in both panels).

(a) Hidden hearing loss reduces the auditory nerve activity slope**(b) Nonlinear central gain and variance induce hyperacusis****Figure 2. Hidden hearing loss, central gain and central variance.**

(a) Schematic effects of hidden hearing loss on auditory nerve activity as a function of sound level (left panel) and time (right panel). Similar to Figure 1(a), except for the purple lines which represent the hidden hearing loss. **(b)** Effects of central gain on loudness as a function of sound level (left panel) and time (right panel). The multiplicative central gain (a factor of two in this case; “x2 gain” as shown by the right curved arrow line, left panel) restores mean normal loudness growth at all sound levels (solid diagonal red line=solid blue line, left panel) but increases the amount of the central variance by a factor of four (“x4 variance” as shown by the left curved arrow line, left panel). The nonlinearly increased central variance generates hyperacusis at high sound levels (labeled “Hyperacusis” in both panels).