

UC Irvine

Journal of Education and Teaching in Emergency Medicine

Title

Irreducible Traumatic Posterior Shoulder Dislocation

Permalink

<https://escholarship.org/uc/item/3599f93b>

Journal

Journal of Education and Teaching in Emergency Medicine, 2(1)

Authors

Collier, Blake
Trigger, Christopher

Publication Date

2017

DOI

10.5070/M521033770

Copyright Information

Copyright 2017 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Irreducible Traumatic Posterior Shoulder Dislocation




Blake Collier, DO* and Christopher Trigger, MD*

*Lakeland Health, Department of Emergency Medicine, St Joseph, MI

Correspondence should be addressed to Blake Collier at blcollier@lakelandhealth.org

Submitted: October 4, 2016; Accepted: November 4, 2016; Electronically Published: January 28, 2017; <https://doi.org/10.21980/J8V884>

Copyright: © 2017 Collier, et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>



History of present illness: A 22-year-old male presented to the emergency department complaining of right shoulder pain after a motocross accident. He was traveling at approximately 10 miles per hour around a turn when he lost control and was thrown over the handlebars, landing directly on his right shoulder. On arrival, he was holding his arm in adduction and internal rotation. An area of swelling was noted over his anterior shoulder. He was unable to abduct his shoulder. No humeral gapping was noted. He had normal neurovascular status distal to the injury.

Significant findings: Radiographs demonstrated posterior displacement of the humeral head on the “Y” view (see white arrow) and widening of the glenohumeral joint space on anterior-posterior view (see red arrow). The findings were consistent with posterior dislocation and a Hill-Sachs type deformity. Sedation was performed and reduction was attempted using external rotation, traction counter-traction. An immediate “pop” was felt during the procedure. Post-procedure radiographs revealed a persistent posterior subluxation with interlocking at posterior glenoid. Computed tomography (CT) revealed posterior dislocation with acute depressed impaction deformity medial to the biceps groove with the humeral head perched on the posterior glenoid, interlocked at reverse Hill-Sachs deformity (see blue arrow).

Discussion: Posterior shoulder dislocations are rare and represent only 2% of all shoulder dislocations. Posterior shoulder dislocations are missed on initial diagnosis in more than 60% of cases.¹ Posterior shoulder dislocations result from axial loading of the adducted and internally rotated shoulder, violent muscle contractions (resulting from seizures or electrocution), a direct posterior force applied to the anterior shoulder.¹ Physical findings include decreased anterior prominence of the humeral head, increased palpable posterior prominence of the humeral head below the acromion, increased palpable prominence of the coracoid, marked limitation of abduction, and complete absence of external rotation with a fixed internal rotation deformity.² Lesions commonly associated with traumatic posterior subluxation/dislocation are the reverse Hill-Sachs,³ a posterior labral detachment, glenohumeral ligament lesions,⁴ rotator cuff tears or posterior bony fractures.¹ In order to make an accurate diagnosis it is important to obtain adequate x-ray imaging, including a “Y” view.² Anteroposterior x-rays may show widening of the glenohumeral joint resembling a “light bulb” shape of the humeral head. However, definitive diagnosis is made by the “Y” view which shows the humeral head displaced posteriorly and no longer covering the glenoid fossa.⁶ Irreducible acute posterior dislocation of the shoulder is extremely rare⁵ and only one other case has been reported in the literature.⁷

Topics: Posterior shoulder, reverse Hill-Sachs, irreducible, ortho, orthopedics, shoulder dislocation.

References:

1. Rouleau DM, Hebert-Davies J. Incidence of associated injury in posterior shoulder dislocation: systematic review of the literature. *J Orthop Trauma*. 2012;26(4):246-251. doi: 10.1097/BOT.0b013e3182243909
2. Li HH, Liu SQ, Peng H. A neglected posterior fracture dislocation of the shoulder with ipsilateral distal humeral shaft fracture. *J Emerg Med*. 2011;41(2):154-156. doi: 10.1016/j.jemermed.2008.01.031
3. McLaughlin HL. Posterior dislocation of the shoulder. *J Bone Joint Surg Am*. 1952;24-A(3):584-590.

4. Ames JB, Millett PJ. Combined posterior osseous Bankart lesion and posterior humeral avulsion of the glenohumeral ligaments: a case report and pathoanatomic subtyping of “floating” posterior inferior glenohumeral ligament lesions. *J Bone Joint Surg Am.* 2011;93:e118(1)–(4). doi: 10.2106/JBJS.K.00010
5. Allard JC, Bancroft J. Irreducible posterior dislocation of the shoulder: MR and CT findings. *J Comput Assist Tomogr.* 1991;15(4):694-696. doi: 10.17795/soj-4023
6. Mimura T, Mori K, Matsusue Y, Tanaka N, Nishi Y, Kobayashi M. Closed reduction for traumatic posterior dislocation of the shoulder using the 'lever principle': two case reports and a review of the literature. *J Orthop Surg (Hong Kong).* 2006;14(3):336-339. doi: 10.1177/230949900601400321
7. Elmali N, Taşdemir Z, Sağlam F, Gülabi D, Baysal Ö. One-stage surgical treatment of neglected simultaneous bilateral locked posterior dislocation of shoulder: a case report and literature review. *Eklem Hastalik Cerrahisi.* 2015;26(3):175-180. doi: 10.5606/ehc.2015.36