

Deforestation decreases spatial turnover and alters the network interactions in soil bacterial communities

Author links open overlay panel [JingTian^{ab}](#) [NianpengHe^{ab}](#) [WeidongKong^{cb}](#) [YeDeng^{db}](#) [KaiFeng^{bd}](#) [Sophie](#)

[M.Green^e](#) [XiaoboWang^f](#) [JizhongZhou^{gh}](#) [YakovKuzyakovⁱ](#) [GuiruiYu^{ab}](#)

Show more

<https://doi.org/10.1016/j.soilbio.2018.05.007> [Get rights and content](#)

Highlights

- Deforestation decreased bacterial spatial turnover rate at large geographic scale.
- Soil pH and SOM availability strongly affected bacterial spatial turnover rate.
- Bacterial networks were more intricate in primary than in secondary forests.
- Bacterial taxa tended to co-occur in secondary forest network.
- The major connectors in the secondary forest network were *Proteobacteria*.

Abstract

Despite important progress in understanding the influence of [deforestation](#) on the bacterial α diversity and community structure at local scales, little is known about deforestation impacts in terms of spatial turnover and soil [bacterial community network interactions](#), especially at regional or global scales. To address this research gap, we examined the bacterial spatial turnover rate and the species networks in paired primary and [secondary forest](#) soils along a 3700-km north-south transect in eastern China using high-throughput [16S rRNA](#) gene [sequencing](#). The spatial turnover rate of bacterial communities was higher in [primary forests](#) than in secondary, suggesting deforestation increased biotic homogenization at a large geographic scale. [Multiple regression](#) on matrices analysis revealed that both geographic distance and soil properties (especially soil pH and organic matter availability) strongly affected bacterial spatial turnover. Through the [phylogenetic](#) molecular [ecological network](#) approach, we demonstrate that the bacterial network of primary forests was more intricate than in secondary forests.

This suggests that microbial species have greater niche-sharing and more interactions in primary forests as compared to secondary forests. On the other hand, the bacterial network in secondary forests was more modular, and the taxa tended to co-occur, with positive correlations accounting for 82% of all potential interactions. In conclusion, our findings demonstrate that anthropogenic deforestation has clear effects on bacterial spatial turnover and network interactions, with potential for serious consequences such as microbial diversity loss in primary forests.

- **Previous article in issue**
- **Next article in issue**

Keywords

Deforestation

Bacterial community structure

Spatial turnover rate

Microbial network interactions

Distance-decay relationship

1. Introduction

More and more [primary forests](#) are being cleared or strongly disturbed globally by human activities to make free areas for agriculture, wood production, human habitation and industry ([Gómez-Acata et al., 2016](#)). With increasing intensity of anthropogenic perturbations, more attention is being placed on [secondary forests](#), since they may act as buffer zones and serve as a habitat for forest plants, animals and microorganisms displaced from destroyed primary forests ([Brearley et al., 2004](#)). [Soil microbial communities](#) are engineers of important biogeochemical processes and play a critical role in regulating the functions and stability of an ecosystem ([Naeem and Li, 1997](#); [Fuhrman, 2009](#)). Many studies showed that [deforestation](#) changed soil bacterial α diversity, composition and community structure on local scales ([Jesus et al., 2009](#); [Bastida et al., 2015](#); [Gómez-Acata et al., 2016](#); [Wood et al., 2017](#)). However, little is known about the effects of deforestation on spatial turnover and species interactions of [bacterial community](#) at a large spatial scale, e.g. regional, continental or global. Distance-decay relationships describe the decrease in community similarity with increasing geographic distance, with the slope reflecting the spatial species turnover rate ([Nekola and White, 1999](#); [Soininen et al., 2007](#); [Hanson et al., 2012](#)). This trend has been reported for microorganisms across a broad range of habitats and spatial scales ([Horner-Devine et al., 2004](#); [Martiny et al., 2006](#); [Morlon et al., 2008](#); [Bell,](#)

[2010](#); [Wang et al., 2017](#); [Zhou et al., 2008](#)). Distance-decay relationships can be explained by two mechanisms: (1) similarity decays with distance due to environmental differences, which is attributable to niche-based community processes, with species differing in terms of their ability to adapt to environmental conditions ([Nekola and White, 1999](#); [Soininen et al., 2007](#)); (2) community similarity declines with distance due to an organisms' limited dispersal, even if the environment is completely homogeneous ([Soininen et al., 2007](#)). There are evidences suggesting that both mechanisms are important for shaping the distance-decay relationship ([Fierer and Jackson, 2006](#); [Dini-Andreote et al., 2015](#); [Ge et al., 2008](#); [Chu et al., 2010](#); [Xiong et al., 2012](#); [Hendershot et al., 2017](#)). However, most of these studies investigating spatial [microbial community](#) turnover rate have examined natural habitats such as those under natural forests, [grasslands](#) or marshes. Little is known about how microbial spatial turnover is affected by anthropogenic activities (e.g., long-term fertilization and elevated CO₂ levels) ([Liang et al., 2015](#); [Deng et al., 2016](#)) and only one study has focused on deforestation to date ([Rodrigues et al., 2013](#)). [Rodrigues et al. \(2013\)](#) found that conversion of Amazon [tropical rainforest](#) to cropland reduced the microbial turnover rate along a 10-km transect. However, it remains unclear whether deforestation influences microbial turnover rate across large spatial scales in different ecosystems and habitats. Clarifying the mechanisms that generate and maintain patterns of diversity is critical for predicting [ecosystem responses](#) to anthropogenic driven-changes.

Microorganisms coexist as complex arrays in many environments, and clarifying their interactions can provide insight into microbial diversity and function ([Hallam and McCutcheon, 2015](#); [Shi et al., 2016](#)). [Network analysis](#) has proven a powerful way to study the complex community organization and member interactions in different ecological systems ([Zhou et al., 2011](#)). These interactions can be positive (e.g., mutualism) or negative (e.g., competition), and maybe depicted by a network model ([Faust and Raes, 2012](#); [Zhou et al., 2011](#); [Deng et al., 2012](#)), in which each node represents a species and the edge linking two nodes represent the relationship between the two species ([Zhou et al., 2011](#); [Deng et al., 2012](#)). Network analysis can also identify keystone taxa that are critical for maintaining community structure and function ([Power et al., 1996](#); [Zhou et al., 2011](#); [Deng et al., 2012](#)). Recent studies have investigated microbial interactions in natural habitats or in response to human activities (e.g. elevated CO₂, [oil pollution](#), and land use change) ([Zhou et al., 2011](#); [Liang et al., 2015](#); [Deng et al., 2016](#); [Ma et al., 2016](#)), but the effects of deforestation on these interactions across large scales is unknown.

Deforestation is a very common [land use change](#) practice, which has far reaching environmental implications. With particular relevance to microbial habitats, deforestation leads to a broad number of environment modifications such as surface water losses and consequently more frequent soil drought ([Bagley et al., 2014](#)), decreased litter input and altered [biochemical composition](#) ([Zou et al., 1995](#)) and changed soil physicochemical characteristics ([Gómez-Acata et al., 2016](#)). Thus, we hypothesize that 1) deforestation may decrease the habitat differences among secondary forests, and consequently decrease the spatial microbial turnover rate at a large scale. In addition, the abundance and composition of key bacterial taxa have been shown to be sensitive to deforestation ([Bastida et al., 2015](#); [Navarrete et al., 2015](#); [Gómez-Acata et al., 2016](#)). Therefore, we also hypothesize that 2) deforestation may alter microbial interspecies interactions; bacterial taxa tend to present negative co-occurrence patterns as disturbance can promote microbial competition ([Violle et al., 2010](#); [Liang et al., 2015](#)). To test these hypotheses, soil was sampled in paired primary and nearby disturbed secondary forests in nine geographic regions along a 3700-km transect to investigate whether deforestation influences microbial spatial turnover and interaction patterns.

2. Materials and methods

2.1. Study sites and field sampling

The study was conducted at nine sites across a 3700-km north-south transect of eastern China (108.9° E, 18.7° N to 123.0° E, 51.8° N; [Supplementary Fig. S1](#)). The sampling sites ranged from cold temperate [coniferous forest](#) to [tropical rainforest](#) across a northern [latitudinal gradient](#) from 51° to 18°. These sites ranged in terms of mean annual temperature from -3.67 °C to 23.2 °C, and annual precipitation from 473 to 2266 mm. Soil physicochemical characteristic differences can be found in [Supplementary Table S1](#).

Within each [primary forest](#) site, four representative plots (30 m × 40 m) were established. Sampled primary forests were deemed well-protected national [nature reserves](#) that avoid any influence of human activities. Comparable four plots were established in nearby [secondary forests](#) which were severely disturbed by anthropogenic activities in the history. Soil was sampled from each plot by collecting 20 randomly selected cores (0–10 cm deep) that were thereafter, well mixed and homogenized. In total, 72 soil samples were collected in this study. Plant roots and leaves were carefully removed and discarded from the soil samples. A portion of each soil sample was transferred to a 50-ml [centrifuge](#) tube that was placed in an ice-box and

transferred to the laboratory. The soil tubes were stored at -80°C for [DNA](#) extraction. The remaining soil was used for physicochemical analyses.

2.2. Soil physicochemical analyses

Soil [organic carbon](#) (SOC) and total nitrogen (TN) contents were determined with a Vario EL III Elemental Analyzer (Elementar, Langenselbold, Germany). Soil total phosphorous (TP) content was determined using the [ammonium molybdate](#) method after $\text{H}_2\text{SO}_4\text{-H}_2\text{O}_2\text{-HF}$ digestion. Available phosphorus (AP) content was also determined using the ammonium molybdate method (no prior digestion). [Particulate organic carbon](#) (POC) content was determined according to [Cambardella and Elliott \(1992\)](#). The concentration of [dissolved organic carbon](#) (DOC) was determined according to [Jones and Willett \(2006\)](#). [Humus](#) composition was analyzed as reported in an earlier study ([Kumada, 1988](#)) with some modifications ([Zhang et al., 2011](#)). Soil pH was measured using a pH meter (1:2.5 w/v). Mean values for soil [physicochemical properties](#) in each forest sample are presented in [Supplementary Table S2](#).

The [soil organic matter](#) (SOM) [mineralization](#) rate was used to represent SOM availability ([Fierer and Jackson, 2006, 2007](#)) and was estimated by measuring the rates of CO_2 production over a 14-day incubation at 25°C after adjusting soil samples to 60% [water-holding capacity](#).

2.3. 16S rRNA gene amplicon sequencing and processing

DNA was extracted from 0.5 g of well-mixed soil using the PowerSoil kit (MoBio Laboratories, Carlsbad, CA, USA) according to the manufacturer's instructions. The V3-V4 [hypervariable regions](#) of bacterial [16S rRNA](#) genes were amplified using the [primers](#) 338F 5'-barcode-ACTCCTACGGGAGGCAGCAG-3' and 806R 5'-GGACTACHVGGGTWTCTAAT-3'. [PCR reactions](#) were performed in triplicate with a 20 μL mixture containing 4 μL of $5 \times$ FastPfu Buffer, 2 μL of 2.5 mM dNTPs, 0.8 μL of each primer (5 μM), 0.4 μL of FastPfu [Polymerase](#), and 10 ng of template DNA. The following thermal program was used for amplification: 95°C for 3 min, followed by 27 cycles at 95°C for 30 s, 55°C for 30 s, 72°C for 45 s and a final extension at 72°C for 10 min. PCR [amplicons](#) were extracted from 2% [agarose gels](#) and purified using an AxyPrep DNA [Gel Extraction](#) Kit (Axygen Biosciences, Union City, CA, USA) according to the manufacturer's instructions and quantified using QuantiFluor™ -ST (Promega, USA). The purified amplicons from all samples were pooled in equimolar concentrations. [Sequencing](#) was conducted on an Illumina MiSeq platform at Majorbio BioPharm Technology Co., Ltd. (Shanghai, China).

Raw sequences >200 bp with an average quality score >20 and without ambiguous base calls were quality processed using the Quantitative Insights into [Microbial Ecology](#) (QIIME) pipeline (version 1.17). [Operational taxonomic units](#) (OTUs) were clustered with a 97% similarity cutoff using UPARSE ([Edgar et al., 2011](#)) (version 7.1 <http://drive5.com/uparse/>). The taxonomic assignment was performed using the Ribosomal Database Project (RDP) classifier (<http://rdp.cme.msu.edu/>). To correct for sampling effort (number of analyzed sequences per sample), we used a randomly selected subset of 18460 sequences per sample for subsequent analysis.

2.4. Network construction and analysis

Networks were, separately, constructed for primary and secondary forests based on 16S rRNA gene sequence data. The framework for network construction can be divided into four key steps: (1) [metagenomic](#) sequence collection, (2) data [standardization](#), (3) pair-wise similarity estimation, and (4) adjacent matrix determination according to a random matrix theory (RMT)-based approach ([Zhou et al., 2011](#)).

Global network properties such as average degree (connectivity), average path length, and average clustering coefficient were characterized for the primary and secondary forest networks. The definitions and calculations of these indices have been previously described ([Zhou et al., 2011](#); [Deng et al., 2012](#)). The network module is a group of highly interconnected nodes (operational taxonomic units (OTUs) in this study) with few connections outside the group ([Zhou et al., 2011](#)). Modularity describes the extent, to which nodes attain more links within their own modules than expected for random linkages. Modules were detected by the greedy modularity optimization method ([Deng et al., 2012](#)). The connectivity of each node was determined based on its within-module connectivity (Z_i) and among-module connectivity (P_i) ([Guimera and Amaral, 2005](#)). Node topologies were organized into four categories: (1) module hubs (highly connected nodes within modules, $Z_i > 2.5$); (2) network hubs (highly connected nodes within the entire network, $Z_i > 2.5$ and $P_i > 0.62$); (3) connectors (nodes connecting modules, $P_i > 0.62$), and (4) peripherals (interconnected nodes in modules with few outside connections, $Z_i < 2.5$ and $P_i < 0.62$) ([Olesen et al., 2007](#); [Zhou et al., 2011](#); [Deng et al., 2012](#)). Topological network features in the primary and secondary forests were compared using a Student's t-test. The above analyses were carried out using the Molecular [Ecological Network](#) Analyses pipeline (<http://ieg2.ou.edu/MENA/>), with the networks being prepared as graphs using Cytoscape v.2.6.2 software ([Cline et al., 2007](#)).

2.5. Statistical analysis

A geographic distance matrix was calculated using latitudinal and longitudinal coordinates of each sampling site using a *gdist* function (R package *lmap*). Matrices of pairwise taxonomic distances between [bacterial communities](#) were analyzed using the Bray-Curtis method. The microbial spatial turnover rate were calculated by determining the slope of the distance-decay relationship ([Rodrigues et al., 2013](#); [Deng et al., 2016](#)), which was plotted as a logarithmic similarity against a logarithmic distance with the slope obtained by linear regression. To assess the significance of the distance-decay relationship curve, we examined whether the slopes were significantly less than zero with 1000 permutations ([Martiny et al., 2011](#)). The significance of slopes between primary and secondary forests was also evaluated by permutation.

[Multiple regression](#) on matrices (MRM) was used to assess linkages between the distance-decay relationship, and measured environmental variables and geographic distance ([Legendre et al., 1994](#)). Partial [regression coefficients](#) of an MRM model provided a measure of the rate of change in [microbial community](#) similarity for variables of interest when other variables were constant ([Martiny et al., 2011](#); [Deng et al., 2016](#)). R package *Ecodist* was used for MRM calculations ([Goslee and Urban, 2007](#)).

2.6. Data accessibility

The raw sequence data from this study were deposited in the SRA at the NCBI database with the assigned study SRP136586 and Biosamples SAMN08802961-SAMN08803032.

3. Results


3.1. Distribution of taxa and phylotypes

Of the 72 soil samples, a total of 1,356,031 and 1,334,898 qualified sequences were obtained for the primary and [secondary forests](#), respectively. When all samples were compared, at an equivalent [sequencing](#) depth of 18,460 per sample, soils from the [primary forests](#) showed higher OTU richness (1497) than those from the secondary forests (1360) at the 97% similarity level. The predominant phyla were *Acidobacteria*, *Actinobacteria*, Proteobacteria, [Chloroflexi](#) and Verrucomicrobia (relative abundance >3%) for both primary and secondary forests, which accounted for more than 84% of bacterial sequences ([Table S3](#)). In addition, the relative abundances at different taxonomical level were counted ([Table S3](#)). For example, the *Acidobacteria* and *alphaproteobacteria* are most dominant [phylotype](#) at the

class level, and the *Rhizobiales* is the most dominant phylotype at the order level (Table S3).

3.2. Bacterial community spatial turnover rate and driving factors in primary and secondary forests

There were significant distance-decay relationships for bacterial communities in both types of forests based on the OTU level ($p < 0.001$; Fig. 1). The slope for primary forests ($r = 0.075$) was steeper than that for secondary forests ($r = 0.068$; $p < 0.001$, permutation test). At the class level, we also observed steeper slopes for primary forests ($r = 0.047$) than that for secondary forests ($r = 0.042$) (Fig. S2; $p < 0.001$, permutation test). These results suggest that spatial turnover rate of soil bacterial community is higher in primary as compared to secondary forests. The average community β diversity was lower in secondary forests than in primary forests ($p < 0.05$; Fig. S3).



1. [Download high-res image \(373KB\)](#)
2. [Download full-size image](#)

Fig. 1. Distance-decay relationships of bacterial communities in primary and secondary forest soils.

For the geographic distances of any two samples among the nine forests, the slope of primary forests was 0.075, and the slope of secondary forests was 0.068; the permutation test indicated that two slopes were significantly different ($p < 0.001$).

MRM was used to assess the relative contribution of geographic distance and soil properties to the observed β diversity (Table 1). A large and significant proportion of the variability in bacterial communities were explained by the MRM model (63.9% and 80.3% for primary and secondary forests, respectively; $p < 0.001$). Soil pH had the greatest influence, with partial regression coefficients of $b = 0.452$ for primary and for 1.05 secondary forests, followed by SOC_{\min} ($b = 0.128$ – 0.139) (Table 1). Available P and geographic distance contributed minimally, but showed significant partial regression coefficients ($b = 0.024$ – 0.037) in both primary and secondary forests. In addition, total P content ($b = 0.131$), DOC ($b = 0.125$), and POC ($b = -0.168$) showed strong effects on β diversity in primary forests. Humic acid carbon (HUC) showed the significant influence on soil bacterial β diversity only in secondary forests.

Table 1. Contribution of environmental factors to the correlations by MRM on matrices analysis.

Factors	Primary forests ($R^2 = 0.639$, $p = 0.01$)		Secondary forests ($R^2 = 0.803$, $p = 0.01$)	
	Coefficient	p	Coefficient	p
distance	0.031	0.01^a	0.029	0.01
pH	0.452	0.01	1.05	0.01
SOC^b	-0.005	0.87	0.029	0.02
TN	-0.0017	0.98	-0.052	0.01
$\text{SOC}:\text{TN}$	0.0008	1.00	0.012	0.27
SOC_{\min}	0.139	0.01	0.128	0.01
TP	0.131	0.01	0.039	0.01
AP	0.037	0.01	0.024	0.03
DOC	0.125	0.01	0.021	0.16
POC	-0.168	0.01	-0.119	0.01
HUC	-0.023	0.53	0.124	0.01

a

Bold indicates significant effects $p < 0.01$.

b

SOC, soil organic carbon; TN, total nitrogen; SOC_{\min} , potential SOM mineralization rate; TP, total phosphorous; AP: available phosphorous; DOC, dissolved organic carbon; POC, particulate organic carbon; HUC, humic acid carbon.

3.3. Distinct bacterial networks in primary and secondary forests

The network in primary forests was larger and more complex than that in secondary forests. The primary forest contained ~ 2 times more nodes and a greater number of

links among nodes with a higher average connectivity ([Table 2](#), [Fig. 2](#)). The average geodesic distance was lower, whereas the average clustering coefficient and modularity were higher in secondary as compared to primary forest networks ([Table 2](#); $p < 0.05$). This indicates that secondary forest networks are more modular, with closer and better-connected nodes.

Table 2. Topological properties of the empirical [ecological networks](#) (MENs) of [microbial communities](#) in primary and [secondary forests](#) and their associate random MENs.

Community	Empirical networks								Random
	Similarity threshold (St)	Network size (n) ^a	Links	Avg. Connectivity (avgK)	Avg.path length (GD)	Avg.clustering coefficient (avgCC)	Modularity (No.of modules)	Avg.path length (GD)	
Primary forest	0.81	142	458	6.45	3.19 ^b	0.333 ^c	0.433 ^d	2.929 ± 0.055	0.116
Secondary forest	0.81	78	161	4.13	2.89 ^b	0.344 ^c	0.508 ^d	3.016 ± 0.082	0.093

a

Number of nodes in a network.

b

Significant difference ($p < 0.001$) in average path length between primary and secondary forests based on the students t-test with standard deviations derived from corresponding random networks.


c

Significant difference ($p < 0.001$) in average clustering coefficient between primary and secondary forests based on the students t-test with standard deviations derived from corresponding random networks.

d


Significant difference ($p < 0.001$) in modularity between primary and secondary forests based on the students t-test with standard deviations derived from corresponding random networks.

Primary forest



14 module, 142 nodes, 458 links (55% positive)

Secondary forest



14 modurle, 78 nodes, 161 links (82% positive)


1. [Download high-res image \(621KB\)](#)
2. [Download full-size image](#)

Fig. 2. [Network interactions](#) in primary and [secondary forest](#) soils.

Colors of nodes indicate different major phyla. A red link indicates negative interaction between two individual nodes, whereas a blue link indicates positive interaction. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

A total of 458 links were identified in primary forests, including 255 (55%) positive and 203 (45%) negative interactions ([Fig. 2](#)). In contrast, bacterial taxa tended to co-occur in secondary forest networks, with positive correlations accounting for 82% of potential interactions observed ([Fig. 2](#)). We assessed possible topological roles of taxa in the

networks based on Z_i and P_i values. The majority of nodes in the two networks were peripheral, and most of their links were inside the modules (Fig. 3). Only one node, belonging to phylum *Acidobacteria*, was identified as a module hub in the primary forest network (Fig. 3). Four nodes were classified as connectors in the primary forest network: one each belonging to *Acidobacteria* and *Actinobacteria* and two belonging to *Proteobacteria* (Fig. 3). Four of five identified connectors in the secondary forest network were *Proteobacteria* (primarily *Alphaproteobacteria*) and one was *Acidobacteria* (Fig. 3).



1. [Download high-res image \(209KB\)](#)
2. [Download full-size image](#)

Fig. 3. Z-P plot showing the classification of nodes to identify putative keystone species in primary and secondary forests.

Each symbol represents an OTU. Modules hubs have $Z_i > 2.5$, whereas connectors have $P_i > 0.62$. One module hub was classified as *Acidobacteria* in the primary forest network. Four connectors in the primary forest network: one each belonging to *Acidobacteria* and *Actinobacteria* and two belonging to *Proteobacteria*. Four connectors in the secondary forest network were *Proteobacteria* (primarily *Alphaproteobacteria*) and one connector was *Acidobacteria*.

4. Discussion

4.1. Effects of deforestation on bacterial community spatial turnover rate

Understanding the patterns of microbial diversity and the mechanisms that maintain them are critical for predicting the response of an ecosystem to environmental changes.

Our hypothesis that [bacterial community](#) similarity decreased with increasing distance in both primary and [secondary forests](#) ([Fig. 1](#)) was verified. Furthermore, the bacterial spatial turnover rate was reduced by [deforestation](#) ([Fig. 1](#)), resulting in more similar bacterial communities over space. Changes in β diversity can be caused by environmental variations and efficient dispersal ([Martiny et al., 2006](#); [Ramette and Tiedje, 2007](#); [Ge et al., 2008](#); [Hendershot et al., 2017](#)). Indeed, we found that a large and significant proportion of the variability in β diversity is explained by environmental variation and geographic distance ($R^2 = 0.64$ and 0.80 ; [Table 2](#)). However, based on the MRM model, [environmental factors](#) had strong effects on bacterial community similarity with high partial [regression coefficients](#) ([Table 2](#)). This suggests that environmental selection plays an important role in shaping the similarity of bacterial communities spatially after deforestation. Among environmental factors, soil pH was the most important factor with a partial regression coefficient of $b = 0.45\text{--}1.05$, followed by SOC_{\min} ($b = 0.128\text{--}0.139$) ([Table 1](#)). Consequently, changes in soil pH and SOM availability (as determined by SOM mineralization) caused by deforestation are major contributors to the decrease in β diversity. This is in agreement with previous studies demonstrating the importance of soil pH in influencing bacterial diversity on a large spatial scale ([Fierer and Jackson, 2006](#); [Liu et al., 2014](#); [Chu et al., 2010](#); [Griffiths et al., 2011](#); [Rousk et al., 2010](#)) and showing that SOC availability combined with taxa traits (e.g., abundance and distribution of dominant taxonomic groups) determines geographic distribution patterns ([Fierer et al., 2007](#); [Tian et al., 2018](#)).

An increase in spatial similarity of bacterial communities following deforestation indicates spatial biotic homogenization, which offers a simple prediction of the human impact on global [biodiversity](#) ([Olden and Poff, 2003](#)). Biotic homogenization describes the gradual transition to nonnative-dominated communities which results in increased spatial and temporal similarity in the taxonomic characteristics of once-disparate biota ([Olden and Poff, 2003](#)). Biotic homogenization of macro-organisms (e.g., plants and animals) is a common outcome of ecosystem conversion ([McKinney and Lockwood, 1999](#); [Smart et al., 2006](#)), but is seldom investigated in microorganisms ([Rodrigues et al., 2013](#)). Biotic homogenization of microorganisms can occur through loss of taxa with restricted geographic ranges (endemic taxa) or invasion of taxa with broad spatial ranges ([Olden and Poff, 2003](#)). There was no evidence for invasion by broadly distributed microbial taxa in this study, but we found that the deforestation reduced endemic taxa ([Supplementary Fig. S4](#)). This study purely focuses on the effects of deforestation on soil bacterial communities. However, it should be noted that soil fungi are also critical components of [microbial communities](#) and play crucial [functional](#)

roles in forest soils ([Peay et al., 2008](#); [He et al., 2017](#)). In contrast to bacterial communities, the fungal communities are more strongly affected by the vegetation ([Peay et al., 2013](#)). Therefore, it is expected that the effect of soil variables on fungal communities is less than that on bacterial composition. Nevertheless, further studies are necessary to investigate whether deforestation reduces fungal diversity and spatial turnover across a large scale. Biotic homogenization is now regarded as one of the most prominent forms of global biotic impoverishment. The long-term impact of diversity loss resulting from deforestation is unknown; additional studies are necessary to clarify the mechanisms underlying current trends to guide conservation efforts.

4.2. Effects of deforestation on bacterial network interactions

Understanding interactions between microbial taxon can reveal the structure of complex microbial communities across spatial gradients ([Barberán et al., 2012](#)). The RMT-based approach (which provides objectively defined thresholds for network construction) was used to model [phylogenetic](#) molecular [ecological networks](#) ([Fig. 2](#)) ([Zhou et al., 2011](#); [Deng et al., 2012](#)). Soil bacterial communities in [primary forests](#) formed larger (i.e., more nodes and links) and more complex (i.e., higher average K value) networks than those in secondary forests ([Table 2](#) and [Fig. 2](#)). Greater network complexity can stabilize communities with mixed interaction types ([Mougi and Kondoh, 2012](#); [Liang et al., 2015](#)); on the other hand, deforestation can reduce the stability of microbial communities.

Microbial networks in secondary forests were more modular than those in primary forests ([Fig. 2](#)). Modularity measures the connectivity between nodes within their own modules that would not occur by chance ([Zhou et al., 2011](#); [Deng et al., 2012](#)). In [ecology](#), a module is a group of species that interact strongly among themselves but little with species in other modules (known as compartmentalization) ([Olesen et al., 2007](#)). The modularity increased in secondary forests; meanwhile, the constituent nodes were more highly connected ([Table 2](#) and [Fig. 2](#)). This indicates that functional association and/or phylogenetic clustering of closely related species was higher in deforested areas. Interestingly, we found that covariation within modules was predominantly positive (82%) in secondary forests, suggesting the occurrence of extensive [mutualism](#) or [commensalism](#) ([Faust and Raes, 2012](#)). This disagrees with studies reporting that increasing disturbance promotes [interspecies competition](#) ([Violle et al., 2010](#); [Liang et al., 2015](#)), although this has been contradicted by others ([Grime, 1979](#); [Huston, 1979](#)). This result is also not consistent with our second hypothesis. Secondary forests have a higher SOC availability than primary forests (student's t-

test = 0.04, [Table S2](#)), presumably, a resource-rich environment reduced competitive pressures after deforestation ([Dini-Andreote et al., 2015](#)). This is supported by our finding that the main keystone nodes (OTUs) of secondary forests networks were *Proteobacteria* ([Fig. 3](#)). According to the growth processes of a network, keystone nodes are recognized as initiating components in networks ([Barabási, 2009](#); [Ma et al., 2016](#)). *Proteobacteria* are mainly copiotrophic; they are the initial metabolizers of labile carbon, and are therefore more abundant in soils with higher organic matter availability ([Fierer et al., 2007](#)). However, the extent of positive interactions among natural populations of bacteria is still debated ([Morris et al., 2012](#); [Ren et al., 2015](#)).

5. Conclusions

In conclusion, [deforestation](#) decreased the [bacterial community](#) spatial turnover rate, indicating biotic homogenization. We identified distinct topological features of networks in soils of primary and [secondary forests](#). The microbial network was larger and more complex in the soils of primary as compared to secondary forests, indicating greater potential for microbial interactions and niche-sharing. Taxonomic groups in secondary forests tended to be modular, and species were more closely connected than those in [primary forest](#) networks. [Keystone species](#) shifted to predominantly *Proteobacteria* after deforestation. Soil pH and SOM availability were the most important factors for the observed microbial β diversity and [networks interactions](#). Taken together, these findings provide a better understanding of the relationship between soil bacterial diversity and [forest ecosystems](#), as well as a framework for predicting continental-scale microbial responses to anthropogenic activities. Future studies are necessary to investigate the influence of deforestation on temporal dynamics of bacterial communities and their networks across large scales. Additionally, studies of the long-term influence of the loss of diversity after deforestation on forest ecosystem function are required.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (grant nos. [41571130041](#) and [31770560](#)); National Key R&D Program of China (grant no. [2017YFA0604803](#)), Youth Innovation Research Team Project (grant no. [LENOM2016Q0004](#)), and the National Environmental Research Council/Newton

Foundation of the UK (grant No. [NE/N007603/1](#)). We thank Prof. Evgenia Blagodatskaya for her useful suggestions on the earlier version of this manuscript. We also thank the anonymous reviewers and Editor for their helpful comments, which have greatly improved the manuscript.

Appendix A. Supplementary data

The following is the supplementary data related to this article:

[Download Word document \(471KB\)](#)[Help with docx files](#)

Supplementary material.

References

[Bagley et al., 2014](#)

J.E. Bagley, A.R. Desai, K.J. Harding, P.K. Snyder, J.A. Foley **Drought and deforestation: has land cover change influenced recent precipitation extremes in the Amazon?**

Journal of Climate, 27 (2014), pp. 345-361

[CrossRefView Record in Scopus](#)

[Barabási, 2009](#)

A.L. Barabási **Scale-free networks: a decade and beyond**

Science, 325 (2009), pp. 412-413

[CrossRefView Record in Scopus](#)

[Barberán et al., 2012](#)

A. Barberán, S.T. Bates, E.O. Casamayor, N. Fierer **Using network analysis to explore co-occurrence patterns in soil microbial communities**

The ISME Journal, 6 (2012), pp. 343-351

[CrossRefView Record in Scopus](#)

[Bastida et al., 2015](#)

F. Bastida, C. García, M. von Bergen, J.L. Moreno, H.H. Richnow, N. Jehmlich **Deforestation fosters bacterial diversity and the cyanobacterial community responsible for carbon fixation processes under semiarid climate: a metaproteomics study**

Applied Soil Ecology, 93 (2015), pp. 65-67

[ArticleDownload PDFView Record in Scopus](#)

[Bell, 2010](#)

T. Bell **Experimental tests of the bacterial distance-decay relationship**

The ISME Journal, 4 (2010), pp. 1357-1365

[CrossRefView Record in Scopus](#)

[Brearley et al., 2004](#)

F.Q. Brearley, S. Prajadinata, P.S. Kidd, J. Proctor, Suriantata **Structure and floristics of an old secondary rain forest in Central Kalimantan, Indonesia, and a comparison with adjacent primary forest**

Forest Ecology and Management, 195 (2004), pp. 385-397

[ArticleDownload PDFView Record in Scopus](#)

[Cambardella and Elliott, 1992](#)

C.A. Cambardella, E.T. Elliott **Particulate soil organic matter changes across a grassland cultivation sequence**

Soil Science Society of America Journal, 56 (1992), pp. 777-783

[CrossRefView Record in Scopus](#)

[Chu et al., 2010](#)

H. Chu, N. Fierer, C.L. Lauber, J.G. Caporaso, R. Knight, P. Grogan **Soil bacterial diversity in the Arctic is not fundamentally different from that found in other biomes**

Environmental Microbiology, 12 (2010), pp. 2998-3006

[CrossRefView Record in Scopus](#)

[Cline et al., 2007](#)

M.S. Cline, M. Smoot, E. Cerami, A. Kuchinsky, N. Landys, C. Workman, R. Christmas, I. Avila-Campilo, M. Creech, B. Gross, K. Hanspers, R. Isserlin, R. Kelley, S. Killcoyne, S. Lotia, S. Maere, J. Morris, K. Ono, V. Pavlovic, A.R. Pico, A. Vailaya, P.L. Wang, A. Adler, B.R. Conklin, L. Hood, M. Kuiper, C. Sander, I. Schmulevich, B. Schwikowski, G.J. Warner, T. Ideker, G.D. Bader **Integration of biological networks and gene expression data using Cytoscape**

Nature Protocols, 2 (2007), pp. 2366-2382

[CrossRefView Record in Scopus](#)

[Deng et al., 2012](#)

Y. Deng, Y.H. Jiang, Y.F. Yang, Z.L. He, F. Luo, J.Z. Zhou **Molecular ecological network analyses**

BMC Bioinformatics, 13 (2012)

[Deng et al., 2016](#)

Y. Deng, P. Zhang, Y.J. Qin, Q.C. Tu, Y.F. Yang, Z.L. He, C.W. Schadt, J.Z. Zhou **Network succession reveals the importance of competition in response to emulsified vegetable oil amendment for uranium bioremediation**

Environmental Microbiology, 18 (2016), pp. 205-218

[CrossRefView Record in Scopus](#)

[Dini-Andreote](#)

[et al., 2015](#)

F. Dini-Andreote, J.C. Stegen, J.D. van Elsas, J.F. Salles **Disentangling mechanisms that mediate the balance between stochastic and deterministic processes in microbial succession**

Proceedings of the National Academy of Sciences of the United States of America, 112 (11) (2015), pp. 1326-1332

[Edgar
et al.,
2011](#)

R.C. Edgar, B.J. Haas, J.C. Clemente, C. Quince, R. Knight **UCHIME improves sensitivity and speed of chimera detection**

Bioinformatics, 27 (2011), pp. 2194-2200

[CrossRefView Record in Scopus](#)

[F
a
u
s
t
-
a
n
d
-
R
a
e
s
.
-
2
0
1
2](#)

K. Faust, J. Raes **Microbial interactions: from networks to models**

Nature Reviews Microbiology, 10 (2012), pp. 538-550

[CrossRefView Record in Scopus](#)

[Fierer
and
Jackson
, 2006](#)

N. Fierer, R.B. Jackson **The diversity and biogeography of soil bacterial communities**

Proceedings of the National Academy of Sciences of the United States of America, 103 (2006), pp. 626-631

[CrossRefView Record in Scopus](#)

N. Fierer, M.A. Bradford, R.B. Jackson **Toward an ecological classification of soil bacteria**
Ecology, 88 (2007), pp. 1354-1364

[CrossRefView Record in Scopus](#)

J.A. Fuhrman **Microbial community structure and its functional implications**
Nature, 459 (2009), pp. 193-199

[CrossRefView Record in Scopus](#)

Y. Ge, J.Z. He, Y.G. Zhu, J.B. Zhang, Z.H. Xu, L.M. Zhang, Y.M. Zheng **Differences in soil bacterial diversity: driven by contemporary disturbances or historical contingencies?**
The ISME Journal, 2 (2008), pp. 254-264

[CrossRefView Record in Scopus](#)

E.S. Gómez-Acata, I. Valencia-Becerril, C. Valenzuela-Encinas, A.S. Velásquez-Rodríguez, Y.E. Navarro-Noya, N. Montoya-Ciriaco, M.C. Suárez-Arriaga, A. Rojas-Valdez, B.G. Reyes-Reyes, M. Luna-Guido, L. Dendooven **Deforestation and cultivation with maize (*Zea mays* L.) has a profound effect on the bacterial community structure in soil**
Land Degradation & Development, 27 (2016), pp. 1122-1130

[CrossRefView Record in Scopus](#)

S.C. Goslee, D.L. Urban **The ecodist package for dissimilarity-based analysis of ecological data**

Journal of Statistical Software, 22 (2007), pp. 1-19

[View Record in Scopus](#)

R.I. Griffiths, B.C. Thomson, P. James, T. Bell, M. Bailey, A.S. Whiteley **The bacterial biogeography of British soils**

Environmental Microbiology, 13 (2011), pp. 1642-1654

[CrossRefView Record in Scopus](#)

J.P. Grime **Plant Strategies and Vegetation Processes**
Wiley, Chichester (1979)

R. Guimera, L.A.N. Amaral **Functional cartography of complex metabolic networks**
Nature, 433 (2005), pp. 895-900

[CrossRefView Record in Scopus](#)

[Fierer et al., 2007](#)

[Fuhrman, 2009](#)

[Ge et al., 2008](#)

[Gómez-Acata et](#)

[Goslee and Urban](#)

[Griffiths et al., 2011](#)

[Grime, 1979](#)

[Guimera and Amaral](#)

[Hallam and McC](#)

S.J. Hallam, J.P. McCutcheon **Microbes don't play solitaire: how cooperation trumps isolation in the microbial world**

Environmental Microbiology Reports, 7 (2015), pp. 26-28

[CrossRefView Record in Scopus](#)

[Hanson et al., 20](#)

C.A. Hanson, J.A. Fuhrman, M.C. Horner-Devine, J.B.H. Martiny **Beyond biogeographic patterns: processes shaping the microbial landscape**

Nature Reviews Microbiology, 10 (2012), pp. 497-506

[View Record in Scopus](#)

[Hendershot et al](#)

J.N. Hendershot, Q.D. Read, J.A. Henning, N.J. Sanders, A.T. Classen **Consistently inconsistent drivers of patterns of microbial diversity and abundance at macroecological scales**

Ecology (2017), [10.1002/ecy.1829](#)

[He et al., 2017](#)

J.H. He, L.H. Tedersoo, A. Hu, C.H. Han, D. He, H. Wei, M. Jiao, S. Anslan, Y.X. Nie, Y.X. Jia, G. X. Zhang, G.R. Yu, S.R. Liu, W.J. Shen **Greater diversity of soil fungal communities and distinguishable seasonal variation in temperate deciduous forests compared with subtropical evergreen forests of eastern China**

FEMS Microbiology Ecology, 93 (2017)

[Horner-Devine e](#)

M.C. Horner-Devine, K.M. Carney, B.J.M. Bohannan **An ecological perspective on bacterial biodiversity**

Proceedings of the Royal Society B: Biological Sciences, 271 (2004), pp. 113-122

[CrossRefView Record in Scopus](#)

[Huston, 1979](#)

M. Huston **A general hypothesis of species diversity**

The American Naturalist, 113 (1979), pp. 81-101

[CrossRefView Record in Scopus](#)

[Jesus et al., 200](#)

E.D. Jesus, T.L. Marsh, J.M. Tiedje, F.M.D. Moreira **Changes in land use alter the structure of bacterial communities in Western Amazon soils**

The ISME Journal, 3 (2009), pp. 1004-1011

[View Record in Scopus](#)

[Jones and Willet](#)

D.L. Jones, V.B. Willett **Experimental evaluation of methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil**

Soil Biology and Biochemistry, 38 (2006), pp. 991-999

[ArticleDownload PDFView Record in Scopus](#)

[Kumada, 1988](#)

K. Kumada **Chemistry of Soil Organic Matter**
Elsevier (1988), p. 22

[Legendre et al., 1994](#)

P. Legendre, F.J. Lapointe, P. Casgrain **Modeling brain evolution from behavior - a permutational regression approach**
Evolution, 48 (1994), pp. 1487-1499

[CrossRef](#)

[Liang et al., 2015](#)

Y.T. Liang, L.Y. Wu, I.M. Clark, K. Xue, Y.F. Yang, J.D. Van Nostrand, Y. Deng, Z.L. He, S. McGrath, J. Storkey, P.R. Hirsch, B. Sun, J.Z. Zhou **Over 150 Years of long-term fertilization alters spatial scaling of microbial biodiversity**
mBio, 6 (2015)

[Liu et al., 2014](#)

J. Liu, Y. Sui, Z. Yu, Y. Shi, H. Chu, J. Jin, X. Liu, G. Wang **High throughput sequencing analysis of biogeographical distribution of bacterial communities in the black soils of northeast China**

Soil Biology and Biochemistry, 70 (2014), pp. 113-122

[ArticleDownload PDFView Record in Scopus](#)

[Ma et al., 2016](#)

B. Ma, H.Z. Wang, M. Dsouza, J. Lou, Y. He, Z.M. Dai, P.C. Brookes, J.M. Xu, J.A. Gilbert **Geographic patterns of co-occurrence network topological features for soil microbiota at continental scale in eastern China**

The ISME Journal, 10 (2016), pp. 1891-1901

[CrossRefView Record in Scopus](#)

[Martiny et al., 2006](#)

J.B.H. Martiny, B.J.M. Bohannan, J.H. Brown, R.K. Colwell, J.A. Fuhrman, J.L. Green, M.C. Horner-Devine, M. Kane, J.A. Krumins, C.R. Kuske, P.J. Morin, S. Naeem, L. Ovreas, A.L. Reysenbach, V.H. Smith, J.T. Staley **Microbial biogeography: putting microorganisms on the map**
Nature Reviews Microbiology, 4 (2006), pp. 102-112

[CrossRefView Record in Scopus](#)

[Martiny et al., 2011](#)

J.B.H. Martiny, J.A. Eisen, K. Penn, S.D. Allison, M.C. Horner-Devine **Drivers of bacterial beta-diversity depend on spatial scale**

Proceedings of the National Academy of Sciences of the United States of America, 108 (2011), pp. 7850-7854

[CrossRefView Record in Scopus](#)

[McKinney and Lomolino, 2001](#)

M.L. McKinney, J.L. Lockwood **Biotic homogenization: a few winners replacing many losers in the next mass extinction**

Trends in Ecology & Evolution, 14 (1999), pp. 450-453

[Article](#)[Download PDF](#)[View Record in Scopus](#)

[Morlon et al., 200](#)

H. Morlon, G. Chuyong, R. Condit, S. Hubbell, D. Kenfack, D. Thomas, R. Valencia, J.L. Green **A general framework for the distance-decay of similarity in ecological communities**

Ecology Letters, 11 (2008), pp. 904-917

[CrossRef](#)[View Record in Scopus](#)

[Morris et al., 201](#)

J.J. Morris, R.E. Lenski, E.R. Zinser **The black queen hypothesis: evolution of dependencies through adaptive gene loss**

mBio, 3 (2012)

[Mougi and Kond](#)

A. Mougi, M. Kondoh **Diversity of interaction types and ecological community stability**

Science, 337 (2012), pp. 349-351

[CrossRef](#)[View Record in Scopus](#)

[Naeem and Li, 1](#)

S. Naeem, S.B. Li **Biodiversity enhances ecosystem reliability**

Nature, 390 (1997), pp. 507-509

[CrossRef](#)[View Record in Scopus](#)

[Navarrete et al.,](#)

A.A. Navarrete, S.M. Tsai, L.W. Mendes, K. Faust, M. de

Hollander, N.A. Cassman, J. Raes, J.A. van Veen, E.E. Kuramae **Soil microbiome responses to the short-term effects of Amazonian deforestation**

Molecular Ecology, 24 (2015), pp. 2433-2448

[CrossRef](#)[View Record in Scopus](#)

[Nekola and Whit](#)

J.C. Nekola, P.S. White **The distance decay of similarity in biogeography and ecology**

Journal of Biogeography, 26 (1999), pp. 867-878

[CrossRef](#)[View Record in Scopus](#)

[Olden and Poff, 1](#)

J.D. Olden, N.L. Poff **Toward a mechanistic understanding and prediction of biotic homogenization**

The American Naturalist, 162 (2003), pp. 442-460

[CrossRef](#)[View Record in Scopus](#)

[Olesen et al., 2007](#)

J.M. Olesen, J. Bascompte, Y.L. Dupont, P. Jordano **The modularity of pollination networks**

Proceedings of the National Academy of Sciences of the United States of America, 104 (2007), pp. 19891-19896

[CrossRefView Record in Scopus](#)

[Peay et al., 2008](#)

K.G. Peay, P.G. Kennedy, T.D. Bruns **Fungal community ecology: a hybrid beast with a molecular master**

BioScience, 58 (2008), pp. 799-810

[CrossRefView Record in Scopus](#)

[Peay et al., 2013](#)

K.G. Peay, C. Baraloto, P.V.A. Fine **Strong coupling of plant and fungal community structure across western Amazonian rainforests**

The ISME Journal, 7 (2013), pp. 1852-1861

[CrossRefView Record in Scopus](#)

[Power et al., 1996](#)

M.E. Power, D. Tilman, J.A. Estes, B.A. Menge, W.J. Bond, L.S. Mills, G. Daily, J.C. Castilla, J. Lubchenco, R.T. Paine **Challenges in the quest for keystones**

BioScience, 46 (1996), pp. 609-620

[CrossRefView Record in Scopus](#)

[Ramette and Tiedje, 2007](#)

A. Ramette, J.M. Tiedje **Multiscale responses of microbial life to spatial distance and environmental heterogeneity in a patchy ecosystem**

Proceedings of the National Academy of Sciences of the United States of America, 104 (2007), pp. 2761-2766

[CrossRefView Record in Scopus](#)

[Ren et al., 2015](#)

D. Ren, J.S. Madsen, S.J. Sorensen, M. Burmolle **High prevalence of biofilm synergy among bacterial soil isolates in cocultures indicates bacterial interspecific cooperation**

The ISME Journal, 9 (2015), pp. 81-89

[CrossRefView Record in Scopus](#)

[Rodrigues et al., 2013](#)

J.L.M. Rodrigues, V.H. Pellizari, R. Mueller, K. Baek, E.D. Jesus, F.S. Paula, B. Mirza, G.S. Hamadou, S.M. Tsai, B. Feigl, J.M. Tiedje, B.J.M. Bohannan, K. Nusslein **Conversion of the Amazon rainforest to agriculture results in biotic homogenization of soil bacterial communities**

Proceedings of the National Academy of Sciences of the United States of America, 110 (2013), pp. 988-993

[CrossRefView Record in Scopus](#)

[Rousk et al., 2010](#)

J. Rousk, E. Baath, P.C. Brookes, C.L. Lauber, C. Lozupone, J.G. Caporaso, R. Knight, N. Fierer **Soil bacterial and fungal communities across a pH gradient in an arable soil**

The ISME Journal, 4 (2010), pp. 1340-1351

[CrossRefView Record in Scopus](#)

[Shi et al., 2016](#)

S.J. Shi, E.E. Nuccio, Z.J. Shi, Z.L. He, J.Z. Zhou, M.K. Firestone **The interconnected rhizosphere: high network complexity dominates rhizosphere assemblages**

Ecology Letters, 19 (2016), pp. 926-936

[CrossRefView Record in Scopus](#)

[Smart et al., 2006](#)

S.M. Smart, K. Thompson, R.H. Marrs, Duc M.G. Le, L.C. Maskell, L.G. Firbank **Biotic homogenization and changes in species diversity across human-modified ecosystems**

Proceedings of the Royal Society B: Biological Sciences, 273 (2006), pp. 2659-2665

[CrossRefView Record in Scopus](#)

[Soininen et al., 2007](#)

J. Soininen, R. McDonald, H. Hillebrand **The distance decay of similarity in ecological communities**

Ecography, 30 (2007), pp. 3-12

[CrossRefView Record in Scopus](#)

[Tian et al., 2018](#)

J. Tian, N. He, L. Hale, S. Niu, G. Yu, Y. Liu, E. Blagodatskava, Y. Kuzyakov, J.Z. Zhou **Soil organic matter availability and climate drive latitudinal patterns in bacterial diversity from tropical to cold-temperate forests**

Functional Ecology, 32 (2018), pp. 61-70

[CrossRefView Record in Scopus](#)

[Violle et al., 2010](#)

C. Violle, Z.C. Pu, L. Jiang **Experimental demonstration of the importance of competition under disturbance**

Proceedings of the National Academy of Sciences of the United States of America, 107 (2010), pp. 12925-12929

[CrossRefView Record in Scopus](#)

[Wang et al., 2017](#)

X.B. Wang, X.T. Lu, J. Yao, Z.W. Wang, Y. Deng, W.X. Cheng, J.Z. Zhou, X.G. Han **Habitat-specific patterns and drivers of bacterial beta-diversity in China's drylands**

The ISME Journal, 11 (2017), pp. 1345-1358

[CrossRefView Record in Scopus](#)

[Wood et al., 2017](#)

S.A. Wood, J.A. Gilbert, J.W. Leff, N. Fierer, H. D'Angelo, C. Bateman, S.M. Gedallovich, C.M. Gillikin, M.R. Gradoville, P. Mansor, A. Massmann, N. Yang, B.L. Turner, F.Q. Brearley, K.L. McGuire **Consequences of tropical forest conversion to oil palm on soil bacterial community and network structure**

Soil Biology and Biochemistry, 112 (2017), pp. 258-268

[ArticleDownload PDFView Record in Scopus](#)

[Xiong et al., 2012](#)

J.B. Xiong, Y.Q. Liu, X.G. Lin, H.Y. Zhang, J. Zeng, J.Z. Hou, Y.P. Yang, T.D. Yao, R. Knight, H.Y. Chu **Geographic distance and pH drive bacterial distribution in alkaline lake sediments across Tibetan Plateau**

Environmental Microbiology, 14 (2012), pp. 2457-2466

[CrossRefView Record in Scopus](#)

[Zhang et al., 2011](#)

J.J. Zhang, F. Hu, H.X. Li, Q. Gao, X.Y. Song, X.K. Ke, L.C. Wang **Effects of earthworm activity on humus composition and humic acid characteristics of soil in a maize residue amended rice-wheat rotation agroecosystem**

Applied Soil Ecology, 51 (2011), pp. 1-8

[ArticleDownload PDFCrossRefView Record in Scopus](#)

[Zhou et al., 2011](#)

J.Z. Zhou, Y. Deng, F. Luo, Z.L. He, Y.F. Yang **Phylogenetic molecular ecological network of soil microbial communities in response to elevated CO₂**

mBio, 2 (2011)

[Zhou et al., 2008](#)

J.Z. Zhou, S. Kang, C.W. Schadt, C.T. Garten **Spatial scaling of functional gene diversity across various microbial taxa**

Proceedings of the National Academy of Sciences of the United States of America, 105 (2008), pp. 7768-7773

[CrossRefView Record in Scopus](#)

[Zou et al., 1995](#)

X.M. Zou, C.P. Zucca, R.B. Waide, W.H. McDowell **Long-term influence of deforestation on tree species composition and litter dynamics of a tropical rain-forest in Puerto-Rico**

Forest Ecology and Management, 78 (1995), pp. 147-157

[ArticleDownload PDFView Record in Scopus](#)

[View Abstract](#)