

UC Irvine

UC Irvine Previously Published Works

Title

Comparison of Ca²⁺ puffs evoked by extracellular agonists and photoreleased IP₃

Permalink

<https://escholarship.org/uc/item/35s561vq>

Journal

Cell Calcium, 63

ISSN

0143-4160

Authors

Lock, Jeffrey T

Smith, Ian F

Parker, Ian

Publication Date

2017-05-01

DOI

10.1016/j.ceca.2016.11.006

Peer reviewed



Published in final edited form as:

Cell Calcium. 2017 May ; 63: 43–47. doi:10.1016/j.ceca.2016.11.006.

Comparison of Ca^{2+} puffs evoked by extracellular agonists and photoreleased IP_3

Jeffrey T. Lock^a, Ian F. Smith^a, and Ian Parker^{a,b}

^aDepartment of Neurobiology and Behavior, University of California, Irvine, CA92697

^bDepartment of Physiology and Biophysics, University of California, Irvine, CA92697

Abstract

The inositol trisphosphate (IP_3) signaling pathway evokes local Ca^{2+} signals (Ca^{2+} puffs) that arise from the concerted openings of clustered IP_3 receptor/channels in the ER membrane.

Physiological activation is triggered by binding of agonists to G-protein-coupled receptors (GPCRs) on the cell surface, leading to cleavage of phosphatidyl inositol bisphosphate and release of IP_3 into the cytosol. Photorelease of IP_3 from a caged precursor provides a convenient and widely employed means to study the final stage of IP_3 -mediated Ca^{2+} liberation, bypassing upstream signaling events to enable more precise control of the timing and relative concentration of cytosolic IP_3 . Here, we address whether Ca^{2+} puffs evoked by photoreleased IP_3 fully replicate those arising from physiological agonist stimulation. We imaged puffs in individual SH-SY5Y neuroblastoma cells that were sequentially stimulated by picospritzing extracellular agonist (carbachol, CCH or bradykinin, BK) followed by photorelease of a poorly-metabolized IP_3 analog, i- IP_3 . The centroid localizations of fluorescence signals during puffs evoked in the same cells by agonists and photorelease substantially overlapped (within $\sim 1 \mu\text{m}$), suggesting that IP_3 from both sources accesses the same, or closely co-localized clusters of IP_3Rs . Moreover, the time course and spatial spread of puffs evoked by agonists and photorelease matched closely. Because photolysis generates IP_3 uniformly throughout the cytoplasm, our results imply that IP_3 generated in SH-SY5Y cells by activation of receptors to CCH and BK also exerts broadly distributed actions, rather than specifically activating a subpopulation of IP_3Rs that are scaffolded in close proximity to cell surface receptors to form a signaling nanodomain.

Graphical abstract

Corresponding author: Ian Parker iparker@uci.edu.


Disclosures

All authors declare they have no competing financial interests.

Author contributions

Conception and design of the study by J.T.L., I.F.S., and I.P. Data were collected and analyzed by J.T.L. I.P. and I.F.S. directed the study. Manuscript was written by J.T.L. and I.P. All authors have read and approve the published manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Keywords

Ca²⁺ puffs; IP₃; GPCRs; calcium imaging; signaling nanodomains

1. Introduction

Numerous G-protein-coupled cell surface receptors (GPCRs) function by activating phospholipase C to cleave membrane phosphatidyl inositol bisphosphate, generating inositol trisphosphate (IP₃) that then diffuses in the cytosol to bind to IP₃ receptor/channels (IP₃R) in the membrane of the endoplasmic reticulum (ER) and causes them to open and release Ca²⁺ ions sequestered in the ER lumen [1]. Experimental studies of IP₃-mediated Ca²⁺ release in intact cells may be facilitated by circumventing the upstream components of this signaling pathway, utilizing photorelease of IP₃ from a caged precursor to enable better control of the timing and magnitude of [IP₃] transients than is possible by application of extracellular agonists[2–4]. Here, we address the question of whether GPCRs access the same population of IP₃Rs that are activated by photoreleased IP₃ by comparing the localizations and properties of subcellular Ca²⁺ puffs evoked by these different means. Our results further bear on the broader topic of whether particular GPCRs specifically interact with a subset of IP₃Rs by virtue of spatial proximity[5–9].

IP₃ is a small, soluble molecule that, based on the spread of radiolabeled IP₃ in extracts of oocyte cytoplasm, has long been thought to diffuse in the cytosol at a rate (diffusion coefficient ~280 μm² s⁻¹) nearly as fast as in free solution[10]. Consequently, it has been considered a ‘global’ messenger that would rapidly spread throughout a cell before being degraded[11]. However, we recently described that diffusion of IP₃ in neuroblastoma cells is much (30-fold) slower than this, so its range of action may be only a few μm[12]. At an even shorter scale, IP₃ has been reported to act as a highly localized signal in neurons and smooth muscle, where IP₃Rs may be tightly colocalized with cell surface GPCRs to generate specificity in receptor function[5–7].

To identify the sites where IP₃Rs are activated we imaged Ca²⁺ puffs in SH-SY5Y neuroblastoma cells. These transient, local events are generated by the concerted openings of small numbers of tightly clustered receptor/channels[13]. By determining the centroid locations of the fluorescence signals we could map Ca²⁺ puffs with sub-micron

precision[14], allowing us to compare the locations of IP₃R clusters activated in the same cell by sequential stimulation by GPCR agonists, followed by global photolysis of caged IP₃. SH-SY5Y cells endogenously express BK2 [15] and mACh3 [16] receptors, which we activated by extracellular application of bradykinin (BK) and carbachol (CCH).

2. Experimental methods

SH-SY5Y neuroblastoma cells were cultured and imaged as described previously[17]. In brief, cells were loaded by incubation with membrane-permeable esters of the Ca²⁺ indicator Cal-520; a caged precursor of a slowly metabolized IP₃ analog, i-IP₃ (SiChem; #cag-iso-2-145-10); and EGTA, a slow Ca²⁺ buffer introduced to inhibit propagation of cytosolic Ca²⁺ waves and sharpen the spatio-temporal profile of puffs[18,19]. Wide-field (WF) epi-fluorescence images of cells were acquired in a HEPES buffered salt solution containing 2 mM Ca²⁺ (HBSS) using a custom-built microscope equipped with a 60X objective (Olympus, NA 1.45) and electron-multiplied ccd camera (Photometrics Evolve; 220 frames s⁻¹, 2x2 binning for a final pixel size of 0.53 μm at the specimen). Ca²⁺ puffs were first evoked by picospritzing agonist (10 nM BK or 500 nM CCH) from a blunt (tip diameter ~1–2 μm) micropipette positioned above the cell, and subsequently by photoreleasing i-IP₃ by a flash of UV light (350–400 nm) from a Xenon arc lamp focused to evenly illuminate the same cell. Image stacks were analyzed using a custom algorithm[20] to locate the centroids of puffs with sub-pixel resolution and to derive traces of fluorescence over time from small regions centered on these sites. For a full description of the materials and procedures see Supplementary Methods online.

3. Results

Fig. 1 illustrates our experimental protocol. An SH-SY5Y cell was first stimulated by applying an agonist (500 nM CCH in this instance) from a micropipette positioned above the cell. The pressure, duration and timing of the pneumatic pulses were empirically adjusted to evoke local Ca²⁺ puffs without triggering global Ca²⁺ waves, and trials with fluorescein in the pipette solution showed that the plume of ejected solution engulfed the cell. Responses typically persisted for only a few seconds following a single pulse of agonist, so we delivered several pulses throughout a ~ 2 min recording period to generate repeated puffs that arose at several sites within the cell (Figs. 1A–C). After waiting ~ 5 min for agonist-evoked responses to fully cease, we then delivered a flash of UV light to uniformly photorelease i-IP₃ throughout the same cell, with the flash intensity and duration adjusted to again evoke puffs at multiple sites without triggering Ca²⁺ waves (Figs. 1E–G). Activity following a single flash persisted for several minutes (Fig. 1F) because the photoreleased i-IP₃ analog is poorly metabolized. For this reason switching the stimulant order, i.e. first stimulating with i-IP₃ followed by CCH, was impracticable. The same protocol was used to compare responses evoked by local application of bradykinin (10 nM) with those evoked by photoreleased i-IP₃.

3.1 Locations of puffs evoked by GPCR agonists and photoreleased IP₃

The stacks of raw fluorescence images were processed to form ratio ($\Delta F/F_0$) image sequences that were analyzed[20] to locate the centroid positions of all puffs evoked in a

given cell by agonist application and then by photoreleased i-IP₃, as illustrated in Figs. 1A,E. Fig 2A shows the same data, replotted to overlay the centroid locations of CCH and i-IP₃-evoked puffs, and Fig. 2B shows enlarged plots of puff localizations at sites where several repeated puffs were evoked by both stimuli. We had previously shown that localizations of IP₃-evoked puffs recorded over several minutes cluster tightly (within < 1 μm), indicating that the underlying cluster of IP₃Rs remains almost stationary[14]. Consistent with this, successive puffs evoked by CCH localized closely with one another (open circles, Fig. 2B), as did puffs evoked by photoreleased i-IP₃ (filled circles, Fig. 2B). Importantly, the locations of CCH-evoked puffs closely co-localized with those of i-IP₃-evoked puffs in the same cell (Fig. 2B).

To analyze these spatial distributions, we measured distances from the centroid localization of every puff to that of every other puff in that cell. Fig. 2C plots the pooled distribution of these all-neighbor distances for CCH-evoked puffs in 7 cells. The distribution showed a sharp peak at distances < 1 μm (shown on an expanded scale in the inset plot), which represents repeated puffs arising at the same, fixed sites, whereas the long 'tail' of the distribution represents distances between different puff sites (IP₃R clusters). A similar sharp peak was evident for i-IP₃-evoked puffs (Fig. 2D). Finally, Fig. 2E plots the distribution of distances between puffs evoked in the same cells by CCH and i-IP₃. Again, a sharp peak was evident at short distances, suggesting that both stimuli evoke Ca²⁺ release from the same clusters of IP₃Rs. Although the co-localization (width of the initial peak in the distribution) was slightly wider than for the self localizations (CCH to CCH and i-IP₃ to i-IP₃), that may have arisen from slight movement during the ~5 min recovery interval between trials with agonist application and photorelease.

Figs. 2F–J present a corresponding analysis of localizations of puffs evoked sequentially by BK and photorelease of i-IP₃. Similar to the results with CCH, puffs evoked by BK recurred at fixed, tightly localized sites (Fig. 2H), and co-localized with i-IP₃-evoked puffs at these sites (Fig. 2J); albeit with a wider spread than observed for co-localization between CCH- and i-IP₃-evoked puffs.

3.2 Spatiotemporal characteristics of puffs evoked by GPCR agonists and photoreleased IP₃

For each puff, the analysis software determined the peak amplitude of the fluorescence ratio change ($\Delta F/F_0$), the full duration at half maximum amplitude (FDHM), and the spatial spread of the fluorescence signal as full width at half maximal amplitude (FWHM), at the time of the peak signal[20]. Histograms showing the distributions of these parameters are plotted in Fig. 3 for puffs evoked by CCH and BK, in each case accompanied by the distributions of puffs evoked by photoreleased i-IP₃ in the same cells.

Puff amplitudes largely reflect the number of IP₃R channels that open during an event; in turn a function of the number of channels clustered at a puff site and the fraction of those that are activated by a given concentration of IP₃[21]. The amplitudes of puffs evoked by CCH (Fig. 3A) and BK (Fig. 3B) were, on average significantly smaller than those evoked by i-IP₃ (mean $\pm$ SD $\Delta F/F_0$ values, 0.60 ± 0.25 for CCH vs. 0.82 ± 0.33 for i-IP₃; 0.77 ± 0.55 BK vs. 1.05 ± 0.52 i-IP₃). However, these differences may largely be attributable to

differences in the resulting concentrations of IP₃, as the observed frequencies of puffs – a parameter that varies steeply with IP₃ concentration – were appreciably higher with photoreleased i-IP₃ (0.83 puffs per cell per s for CCH vs. 1.35 for i-IP₃; 0.54 for BK vs. 1.16 for i-IP₃).

The temporal durations (FDHM) and spatial widths (FWHM) of Ca²⁺ puffs evoked by stimulation with the GPCR agonists CCH or BK or by photorelease of i-IP₃ were closely similar.

4. Summary and Discussion

Our results indicate that local Ca²⁺ puffs evoked by photorelease of i-IP₃ from a caged precursor in SH-SY5Y neuroblastoma cells resemble closely those evoked in the same cells by activation of cell surface receptors by CCH and BK in terms of their locations, duration and spatial extent. Although GPCR-mediated cleavage of membrane phosphatidyl inositol bisphosphate by phospholipase C results in formation of diacylglycerol in addition to IP₃, we did not observe any differences in puff properties that might be attributable to that second messenger. We thus conclude that caged IP₃ provides an experimental system that mimics Ca²⁺ release evoked via GPCR-coupled formation of IP₃, while greatly facilitating the precise and reproducible control of the magnitude, timing and spatial localization of cytosolic IP₃.

The concentration of i-IP₃ generated by photorelease in our experiments would have been almost uniform throughout the cytosol. Therefore, the finding that puff sites (clusters of IP₃Rs) responding to CCH and BK map onto those seen with photoreleased i-IP₃ suggests that IP₃ formed by GPCR stimulation also distributes widely throughout the cytoplasm, and that the receptors to CCH and BK both access the same population of IP₃Rs. This conclusion differs from the notion of signaling nanodomains that have been proposed in neurons and other cell types, wherein highly localized gradients of IP₃ concentration activate IP₃Rs scaffolded in close proximity to GPCRs to confer selectivity and specificity in responses to different agonists [5,6,8,9]. Nevertheless, interactions between cell surface receptors and IP₃Rs may still be constrained even in cells lacking such nanodomains, albeit on a coarser scale. Although IP₃ has widely been considered to diffuse freely in the cytosol [10,11], we recently reported that its diffusion is strongly hindered in SH-SY5Y cells, reducing its range of action to a few μm [12]. Thus, different receptors might differentially access spatially separated populations of IP₃Rs if they were inhomogeneously distributed across the cell membrane on a scale of a few μm .

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by National Institutes of Health grants GM 100201 to I.F.S., and GM 048071 and GM 065830 to I.P.


References

1. Berridge MJ. Inositol trisphosphate and calcium signalling mechanisms. *Biochim Biophys Acta*. 2009; 1793:933–940. DOI: 10.1016/j.bbamcr.2008.10.005 [PubMed: 19010359]
2. Thomas D, Lipp P, Tovey SC, Berridge MJ, Li W, Tsien RY, Bootman MD. Microscopic properties of elementary Ca^{2+} release sites in non-excitable cells. *Curr Biol*. 2000; 10:8–15. [PubMed: 10660296]
3. Smith IF, Wiltgen SM, Parker I. Localization of puff sites adjacent to the plasma membrane: functional and spatial characterization of Ca^{2+} signaling in SH-SY5Y cells utilizing membrane-permeant caged IP_3 . *Cell Calcium*. 2009; 45:65–76. DOI: 10.1016/j.ceca.2008.06.001 [PubMed: 18639334]
4. Tovey SC, de Smet P, Lipp P, Thomas D, Young KW, Missiaen L, De Smedt H, Parys JB, Berridge MJ, Thuring J, Holmes A, Bootman MD. Calcium puffs are generic $\text{InsP}(3)$ -activated elementary calcium signals and are downregulated by prolonged hormonal stimulation to inhibit cellular calcium responses. *J Cell Sci*. 2001; 114:3979–3989. [PubMed: 11739630]
5. Delmas P, Wanaverbecq N, Abogadie FC, Mistry M, Brown DA, Sensoriellies I, Aiguiet CJ. Signaling Microdomains Define the Specificity of Receptor-Mediated InsP_3 Pathways in Neurons. 2002; 14:209–220.
6. Jacob SN, Choe C, Uhlen P, Degray B, Yeckel MF, Ehrlich BE. Trisphosphate-Mediated Intracellular Calcium Transients in Cultured Neurons. 2005; 25:2853–2864. DOI: 10.1523/JNEUROSCI.4313-04.2005
7. Olson ML, Sandison ME, Chalmers S, McCarron JG. Microdomains of muscarinic acetylcholine and $\text{Ins}(1,4,5)\text{P}(3)$ receptors create “ $\text{Ins}(1,4,5)\text{P}(3)$ junctions” and sites of $\text{Ca}(2)^+$ wave initiation in smooth muscle. *J Cell Sci*. 2012; 125:5315–5328. DOI: 10.1242/jcs.105163 [PubMed: 22946060]
8. Tovey SC, Dedos SG, Taylor EJA, Church JE, Taylor CW. Selective coupling of type 6 adenylyl cyclase with type 2 IP_3 receptors mediates direct sensitization of IP_3 receptors by cAMP. *J Cell Biol*. 2008; 183:297–311. DOI: 10.1083/jcb.200803172 [PubMed: 18936250]
9. Yuan Z, Cai T, Tian J, Ivanov AV, Giovannucci DR, Xie Z. Na/K-ATPase tethers phospholipase C and IP_3 receptor into a calcium-regulatory complex. *Mol Biol Cell*. 2005; 16:4034–4045. DOI: 10.1091/mbc.E05-04-0295 [PubMed: 15975899]
10. Allbritton NL, Meyer T, Stryer L. Range of messenger action of calcium ion and inositol 1,4,5-trisphosphate. *Science* (80–). 1992; 258:1812 LP–1815.
11. Sims CE, Allbritton NL. Metabolism of inositol 1,4,5-trisphosphate and inositol 1,3,4,5-tetrakisphosphate by the oocytes of *Xenopus laevis*. *J Biol Chem*. 1998; 273:4052–4058. [PubMed: 9461597]
12. Dickinson I, Ellefsen GD, Ponce-Dawson KL, Pearson S, Parker JE. Hindered cytoplasmic diffusion of inositol trisphosphate restricts its cellular range of action. *Sci Signal*. 2016; 9(453)
13. Smith IF, Parker I. Imaging the quantal substructure of single IP_3 R channel activity during Ca^{2+} puffs in intact mammalian cells. 2009; 106:3–8.
14. Wiltgen SM, Smith IF, Parker I. Superresolution localization of single functional IP_3 R channels utilizing Ca^{2+} flux as a readout. *Biophys J*. 2010; 99:437–446. DOI: 10.1016/j.bpj.2010.04.037 [PubMed: 20643061]
15. McDonald RL, Kaye DF, Reeve HL, Ball SG, Peers C, Vaughan PF. Bradykinin-evoked release of $[^3\text{H}]\text{noradrenaline}$ from the human neuroblastoma SH-SY5Y. *Biochem Pharmacol*. 1994; 48:23–30. [PubMed: 8043027]
16. Willets JM, Challiss RAJ, Nahorski SR. Endogenous G protein-coupled receptor kinase 6 Regulates M3 muscarinic acetylcholine receptor phosphorylation and desensitization in human SH-SY5Y neuroblastoma cells. *J Biol Chem*. 2002; 277:15523–15529. DOI: 10.1074/jbc.M111217200 [PubMed: 11856737]
17. Lock JT, Parker I, Smith IF. Cell Calcium A comparison of fluorescent Ca^{2+} indicators for imaging local Ca^{2+} signals in cultured cells. *Cell Calcium*. 2015; 58:638–648. DOI: 10.1016/j.ceca.2015.10.003 [PubMed: 26572560]

18. Callamaras N, Parker I. Phasic characteristic of elementary Ca^{2+} release sites underlies quantal responses to $\text{IP}(3)$. *EMBO J*. 2000; 19:3608–3617. DOI: 10.1093/emboj/19.14.3608 [PubMed: 10899115]
19. Dargan SL, Parker I. Buffer kinetics shape the spatiotemporal patterns of IP_3 -evoked Ca^{2+} signals. *J Physiol*. 2003; 553:775–788. DOI: 10.1113/jphysiol.2003.054247 [PubMed: 14555715]
20. Ellefsen KL, Settle B, Parker I, Smith IF. An algorithm for automated detection, localization and measurement of local calcium signals from camera-based imaging. *Cell Calcium*. 2014; 56:147–156. DOI: 10.1016/j.ceca.2014.06.003 [PubMed: 25047761]
21. Shuai J, Pearson JE, Foskett JK, Mak DOD, Parker I. A kinetic model of single and clustered IP_3 receptors in the absence of Ca^{2+} feedback. *Biophys J*. 2007; 93:1151–1162. DOI: 10.1529/biophysj.107.108795 [PubMed: 17526578]

Highlights

- GPCR agonists and photoreleased i-IP₃ evoke Ca²⁺ puffs at overlapping sites.
- Puffs evoked by GPCR agonists and photoreleased i-IP₃ have similar kinetics and spatial spread.
- Photoreleased i-IP₃ faithfully mimics puffs evoked by the physiological IP₃ pathway.
- Muscarinic and bradykinin receptors access the same IP₃ receptors.

**Fig. 1.**

Experimental procedure. **(A)** The background-subtracted resting Cal-520 fluorescence of a single SH-SY5Y cell is shown overlaid with dots indicating the centroid localizations of puffs evoked by application of CCH. **(B)** Traces showing fluorescence ratio measurements ($\Delta F/F_0$) from regions of interest ($\sim 1 \mu\text{m}$ square) centered on the puff sites numbered in A. Arrows indicate times at which repeated applications of CCH were delivered by pneumatic pressure pulses. **(C)** Superimposed examples of puffs evoked by CCH, shown on an expanded timescale and aligned by their rising phases. **(E-F)** Corresponding records of puffs evoked in the same cell about 5 min later by flash photorelease of $i\text{-IP}_3$ using UV light uniformly illuminating the entire cell. The lightning symbol marks the time of the photolysis flash.





Fig. 2. GPCR agonists and i-IP_3 evoke Ca^{2+} puffs at overlapping subcellular sites. **(A)** The dotted outline indicates the boundary of the same cell as in Fig. 1. Open circles mark the centroid localizations of puffs evoked by CCH, and filled circles mark localizations of puffs evoked by photoreleased i-IP_3 . **(B)** Examples of localizations of CCH-evoked puffs (open circles) and i-IP_3 -evoked puffs (filled circles) at four sites, shown on an expanded scale. **(C)** Distribution of all-neighbor distances between localizations of puffs evoked by CCH. The inset in this and other panels shows the same data re-plotted on an expanded distance scale. **(D)** Distribution of all-neighbor distances between localizations of puffs evoked by i-IP_3 . **(E)** Distribution of all-neighbor distances between localizations of puffs evoked by CCH and

puffs evoked in the same cell by $i\text{-IP}_3$. Data in C–E are from 7 cells; 581 puffs evoked by CCH and 946 evoked by $i\text{-IP}_3$. **(F,G)** Localizations of puffs evoked by BK and $i\text{-IP}_3$ in a different cell. **(H–J)** Corresponding distributions of all-neighbor distances for puffs evoked by BK **(H)**, $i\text{-IP}_3$ **(I)**, and between BK- and $i\text{-IP}_3$ -evoked puffs **(J)**. Data are from 7 cells; 383 puffs evoked by BK and 809 evoked by $i\text{-IP}_3$.

**Fig. 3.**

Comparison of Ca^{2+} puffs evoked by GPCR agonists with those evoked in the same cells by i-IP_3 . Histograms show the distributions of puff amplitudes (peak $\Delta F/F_0$: **A,B**), full-duration at half-maximal amplitude (**C,D**), and full spatial width at half-maximal amplitude (**E,F**). In each panel, the upper histogram shows the distribution of puffs evoked by CCH (open bars) or BK (grey bars) and the lower histogram shows the distribution of puffs evoked in the same cells by photoreleased i-IP_3 (black bars). The numbers of cells and puffs analyzed are as stated in the legend to Fig. 2.