UC San Diego

UC San Diego Previously Published Works

Title

Mental health impact of the COVID-19 pandemic in U.S. military veterans: a population-based, prospective cohort study.

Permalink

<https://escholarship.org/uc/item/3626s4rr>

Journal

Psychological medicine, 53(3)

ISSN

0033-2917

Authors

Hill, Melanie L
Nichter, Brandon
Na, Peter J
et al.

Publication Date

2021-06-01

DOI

10.1017/s0033291721002361

Peer reviewed

Original Article

Cite this article: Hill ML, Nichter B, Na PJ, Norman SB, Morland LA, Krystal JH, Pietrzak RH (2021). Mental health impact of the COVID-19 pandemic in U.S. military veterans: a population-based, prospective cohort study. *Psychological Medicine* 1–12. <https://doi.org/10.1017/S0033291721002361>

Received: 31 March 2021

Revised: 24 May 2021

Accepted: 26 May 2021

Key words:

Anxiety; COVID-19; depression; PTSD; psychiatric distress; veteran

Author for correspondence:

Melanie L. Hill, E-mail: mlhill@ucsd.edu

Mental health impact of the COVID-19 pandemic in U.S. military veterans: a population-based, prospective cohort study

Melanie L. Hill^{1,2} , Brandon Nichter¹ , Peter J. Na³ ,

Sonya B. Norman^{1,2,4,5} , Leslie A. Morland^{1,2,5,6} , John H. Krystal^{3,7}

and Robert H. Pietrzak^{3,7,8}

¹Department of Psychiatry, University of California, San Diego, CA, USA; ²VA San Diego Healthcare System, San Diego, CA, USA; ³Department of Psychiatry, Yale School of Medicine, New Haven, CT, USA; ⁴National Center for PTSD, White River Junction, VT, USA; ⁵VA Center of Excellence for Stress and Mental Health, San Diego, CA, USA; ⁶National Center for PTSD, Pacific Islands Division, Honolulu, HI, USA; ⁷National Center for PTSD, VA Connecticut Healthcare System, West Haven, CT, USA and ⁸Department of Social and Behavioral Sciences, Yale School of Public Health, New Haven, CT, USA

Abstract

Background. The coronavirus disease-2019 (COVID-19) pandemic has caused myriad health, social, and economic stressors. To date, however, no known study has examined changes in mental health during the pandemic in the U.S. military veteran population.

Methods. Data were analyzed from the 2019–2020 National Health and Resilience in Veterans Study, a nationally representative, prospective cohort survey of 3078 veterans. Pre-to-peri-pandemic changes in psychiatric symptoms were evaluated, as well as pre-pandemic risk and protective factors and pandemic-related correlates of increased psychiatric distress.

Results. The prevalence of generalized anxiety disorder (GAD) positive screens increased from pre- to peri-pandemic (7.1% to 9.4%; $p < 0.001$) and was driven by an increase among veterans aged 45–64 years (8.2% to 13.5%; $p < 0.001$), but the prevalence of major depressive disorder and posttraumatic stress disorder positive screens remained stable. Using a continuous measure of psychiatric distress, an estimated 13.2% of veterans reported a clinically meaningful pre-to-peri-pandemic increase in distress (mean = 1.1 standard deviation). Veterans with a larger pre-pandemic social network size and secure attachment style were less likely to experience increased distress, whereas veterans reporting more pre-pandemic loneliness were more likely to experience increased distress. Concerns about pandemic-related social losses, mental health COVID-19 effects, and housing stability during the pandemic were associated with increased distress, over-and-above pre-pandemic factors.

Conclusions. Although most U.S. veterans showed resilience to mental health problems nearly 1 year into the pandemic, the prevalence of GAD positive screens increased, particularly among middle-aged veterans, and one of seven veterans experienced increased distress. Clinical implications of these findings are discussed.

Introduction

The coronavirus disease-2019 (COVID-19) pandemic has caused widespread social, health, and economic challenges. Pandemic-related stressors including risk of virus exposure which could lead to severe illness or death, social distancing, and lockdown measures may negatively impact mental health, which has led to calls for research on the population-level mental health effects of COVID-19 (Holmes et al., 2020). These impacts might be especially pronounced in the United States, which has suffered the highest number of confirmed COVID-19 cases and deaths in the world (Dong, Du, & Gardner, 2020), as well as an economic recession that has persisted into 2021. Indeed, repeated cross-sectional surveys of nationally representative samples of American adults have found high prevalence of psychiatric distress during the pandemic – with roughly a quarter of respondents (*v.* 6.5–8.1% in 2019) reporting symptoms of anxiety and depression in one survey (Czeisler et al., 2020) and 13.6% (*v.* 3.9% in 2018) reporting serious psychological distress in another (McGinty, Presskreischer, Han, & Barry, 2020b) – raising concerns that the prevalence of psychiatric disorders such as major depressive disorder (MDD), generalized anxiety disorder (GAD), and posttraumatic stress disorder (PTSD) may have increased.

Despite these findings, only a few longitudinal studies to date have examined changes in psychiatric distress in the U.S. population during the pandemic (Breslau et al., 2021; Daly & Robinson, 2020; McGinty, Presskreischer, Anderson, Han, & Barry, 2020a), and several of these studies did not assess pre-pandemic distress (Daly & Robinson, 2020;

McGinty *et al.*, 2020a). This is an important gap, as repeated cross-sectional surveys rely on measurements from different samples, often using different methodologies. Longitudinal assessments of the same cohort over time are needed to understand whether mental health has worsened for individuals relative to their own pre-pandemic mental health. To our knowledge, only one previous longitudinal study has measured pre-to-peri-pandemic change in psychiatric distress in a nationally representative sample of U.S. adults (Breslau *et al.*, 2021). This study found that a similar proportion of the sample reported past-month serious psychological distress on the Kessler-6 (Kessler *et al.*, 2003) in May 2020 as reported past-year serious distress in February 2019 (10.9% *v.* 10.2%). Although past-month estimates were not directly compared, the authors interpreted this similarity as an increase in distress based on a previous finding from a nationally representative study that the past-year estimate of serious distress was approximately double the past-month estimate (Hedden *et al.*, 2012). Additional methodologically rigorous longitudinal studies of Americans, particularly in vulnerable subpopulations (Holmes *et al.*, 2020), are needed to better understand the mental health burden of the COVID-19 public health crisis. In addition, information about factors associated with pandemic-related psychiatric distress in these vulnerable groups is essential to identifying individuals most in need of support, and to inform targeted clinical and policy interventions.

One subpopulation of the United States that may be uniquely vulnerable to psychiatric distress during the pandemic is military veterans. Veterans have greater exposure to childhood adversity and combat trauma (Katon *et al.*, 2015), which are known to sensitize individuals to the negative effects of later stressors (Bandoli *et al.*, 2017; Nichter, Hill, Norman, Haller, & Pietrzak, 2020a; Smid *et al.*, 2015). In addition, veterans are more likely to have a history of psychiatric conditions, such as PTSD, suicidal behavior, and substance use disorders relative to the non-veteran population (Hoggatt, Lehavot, Krenek, Schweizer, & Simpson, 2017; Lehavot, Katon, Chen, Fortney, & Simpson, 2018; United States Department of Veterans Affairs Office of Mental Health & Suicide Prevention, 2019), which may be exacerbated by pandemic stressors (Murphy, Williamson, Baumann, Busuttill, & Fear, 2020). Indeed, one previous study of veterans with pre-existing psychiatric disorders found that pre-pandemic psychiatric symptom severity predicted suicide ideation during the pandemic (Na *et al.*, 2021). Finally, relative to civilians, veterans have higher rates of medical morbidities such as cardiovascular disease, obesity, and diabetes (Hoerster *et al.*, 2012; Lehavot, Hoerster, Nelson, Jakupcak, & Simpson, 2012), which may increase vulnerability to health-related stressors including risk of severe disease from COVID-19 infection and pandemic-related delays in obtaining routine health care (Földi *et al.*, 2020; Ssentongo, Ssentongo, Heilbrunn, Ba, & Chinchilli, 2020). Despite these characteristics no known study has examined pandemic-related changes in the mental health of the general U.S. veteran population.

To address this gap, we analyzed data from a nationally representative sample of U.S. military veterans who completed surveys before and nearly 1 year into the pandemic. To identify changes in specific disorders, we first measured the pre- and peri-pandemic prevalence of positive screens for common psychiatric conditions; then, following prior research (Breslau *et al.*, 2021), we used a continuous, transdiagnostic measure of psychiatric distress to assess variability in distress more sensitively and identify correlates of increased distress. We had three aims: (1) to examine changes in the prevalence of positive screens for MDD, GAD, and

PTSD from pre-to-peri-pandemic; (2) to identify pre-pandemic risk and protective factors that predicted a clinically meaningful increase [i.e. 0.5 standard deviation (S.D.) or higher; Norman, Sloan, and Wyrwich, 2003; Ready *et al.*, 2008; Schnurr *et al.*, 2003] in psychiatric distress (i.e. continuously assessed composite score of MDD, GAD, and PTSD symptoms); and (3) to evaluate associations between COVID-19-related infection stressors, general pandemic stressors, and pandemic-related changes in risk factors, and increased psychiatric distress.

With respect to aim 1, we hypothesized small but significant pre-to-peri-pandemic increases in positive screens for psychiatric disorders, based on general population findings (Breslau *et al.*, 2021) and greater pre-pandemic mental health problems among veterans (Hoglund & Schwartz, 2014). With respect to aim 2, we hypothesized that pre-pandemic factors that robustly predict psychiatric distress, such as prior trauma exposure, psychiatric and substance use problems, and loneliness (Bandoli *et al.*, 2017; Kuwert, Knaevelsrud, & Pietrzak, 2014; Seal *et al.*, 2011; Smid *et al.*, 2015), would positively predict increased distress during the pandemic, whereas psychological and social characteristics previously shown to protect mental health would negatively predict distress (Pietrzak & Cook, 2013). With respect to aim 3, we hypothesized that COVID-19 infection stressors, general pandemic stressors, and increases in pre-pandemic risk factors would be associated with increased psychiatric distress (Birditt, Turkelson, Fingerman, Polenick, & Oya, 2021; Hologue *et al.*, 2020; Holman, Thompson, Garfin, & Silver, 2020; McGinty *et al.*, 2020a; Shi *et al.*, 2020; Wang, Kala, & Jafar, 2020; Zheng *et al.*, 2021).

Methods

Participants

Data were analyzed from two waves (hereafter: pre-pandemic and peri-pandemic) of the 2019–2020 National Health and Resilience in Veterans Study (NHRVS), which surveyed a nationally representative sample of 4069 U.S. military veterans. Pre-pandemic data collection began before the first documented case of COVID-19 infection in China and ended before implementation of the first COVID-19-related lockdown in the USA (median completion date: 21 November 2019). Peri-pandemic data were collected from 3078 veterans (75.6% of the original sample) approximately 1 year later, during the 2020 fall and winter surge of COVID-19 cases (median completion date: 14 November 2020). Recruitment and sample details of the 2019–2020 NHRVS have been reported previously (Hill, Nichter, Loflin, Norman, & Pietrzak, 2021). Briefly, the sample was drawn from KnowledgePanel®, a survey research panel of more than 50 000 households maintained by Ipsos. Ipsos statisticians computed post-stratification weights to permit generalizability of results to the U.S. veteran population using benchmark distributions from the Veterans Supplement of the most contemporaneous U.S. Census Current Population Survey (Bureau of the Census, 2019). All participants provided informed consent. The study was approved by the Human Subjects Committee of the VA Connecticut Healthcare System and carried out in accordance with the Declaration of Helsinki.

The sample consisted of veterans aged 22–99 years (mean age = 63.2; S.D. = 14.7). The sample was mostly male (91.6%); 79.3% were non-Hispanic Caucasian, 10.3% non-Hispanic African American, 6.0% Hispanic, and 4.4% mixed/other race. The

sample included veterans who had served in all branches of the U.S. military [majority Army (47.3%), Navy (20.8%), or Air Force (18.9%)], 35.0% were combat veterans, 79.6% reported having enlisted in the military, and 20.1% reported utilizing VA as their primary source of healthcare.

Assessments

At the pre-pandemic assessment, sociodemographic and military characteristics, psychosocial risk and protective factors, and current psychiatric conditions were assessed. At the peri-pandemic assessment, psychosocial factors and psychiatric conditions were reassessed using the same measures, and COVID-19 infection stressors (related to actual infection of the self or close others) and pandemic stressors (related to general pandemic stressors such as risk of disease, social distancing, and lockdowns) were assessed. Table 1 presents detailed descriptions of all the measures used to assess mental health outcomes, as well as all predictors and correlates that were examined in relation to changes in psychiatric distress.

Data analysis

Statistical analyses were performed using SPSS version 25 and R statistical software. Raw unweighted frequencies are reported. Missing data (<3%) were multiply imputed using chained equations. When computing prevalence and inferential statistics, post-stratification sampling weights were applied to allow for generalizability to the entire population of U.S. military veterans (Ipsos, 2020).

Data analyses proceeded in six stages. First, descriptive statistics were computed to estimate the prevalence of positive screens for MDD, GAD, and PTSD at pre-pandemic and 1-year peri-pandemic, and McNemar's tests were computed to evaluate differences between prevalence estimates at each wave. Second, using standardized z scores of a composite measure of psychiatric distress (i.e. increased MDD, GAD, or PTSD symptoms), change in distress from the pre- to peri-pandemic assessment was computed ($z \text{ score}_{\text{peri-pandemic}} - z \text{ score}_{\text{pre-pandemic}}$), and then the sample was divided into veterans who did and did not experience a clinically meaningful increase in distress over this period using a minimum increase in distress of 0.5 $s.d.$. This threshold was selected based on evidence that a change of at least 0.5 $s.d.$ represents a minimally clinically important difference for a broad range of measures, including psychiatric symptoms (Norman et al., 2003; Ready et al., 2008; Schnurr et al., 2003). To further validate this threshold in the current sample, rates of mental health treatment initiation during the pandemic were compared between veterans with and without increased psychiatric distress. Third, independent samples' t tests for continuous variables and chi-squares for categorical variables were conducted to compare veterans with and without increased psychiatric distress with respect to sociodemographic and military characteristics, psychosocial risk and protective factors, pre-to-peri-pandemic changes in psychosocial risk factors, and COVID-19 infection and pandemic stressors. Fourth, a hierarchical logistic regression analysis was conducted to examine pre-pandemic predictors and peri-pandemic correlates of increased psychiatric distress. Variables that were statistically significantly associated with increased distress in bivariate analyses ($p < 0.05$) were entered as independent variables in this analysis. Step 1 included pre-pandemic sociodemographic, military, and psychosocial risk factors; step 2 included

pre-pandemic psychosocial protective factors; and step 3 included pre-to-peri-pandemic changes in risk factors and COVID-19 infection and pandemic stressors. Fifth, a series of planned post-hoc analyses were conducted to identify individual variables comprising pre- (e.g. social connectedness) and peri- (e.g. COVID-19-related disease worries) pandemic composite measures independently associated with increased psychiatric distress. Sixth, a relative importance analysis (Tonidandel & LeBreton, 2011) was conducted using the R statistical package *relaimpo* (R Core Team, 2019) to determine the relative contribution of each significant variable identified in the hierarchical regression model and planned post-hoc analyses. This analysis partitioned the explained variance in increased psychiatric distress that was attributable to each significant independent variable while accounting for intercorrelations among these variables.

Results

Table 2 presents the pre- and peri-pandemic prevalence estimates for MDD, GAD, and PTSD positive screens in the full sample and by age group. There was a statistically significant increase in the estimated prevalence of GAD from pre- to peri-pandemic in the full sample (7.1% to 9.4%), which was driven by an increase in GAD positive screens in veterans aged 45–64 years (8.2% to 13.5%).

A total of 389 (weighted 13.2%) veterans scored 0.5 $s.d.$ or higher on a composite measure of change in psychiatric distress at the 1-year peri-pandemic assessment (mean change in distress = 1.1 $s.d.$, $s.d.$ = 0.6). Supporting the clinical validity of the 0.5 $s.d.$ cut-off, veterans who met or exceeded this threshold were significantly more likely than those who did not to initiate mental health treatment during the pandemic (8.3% *v.* 3.4%, $\chi^2 = 17.29$, $p < 0.001$), and nearly 60% of these veterans screened positive for peri-pandemic MDD, GAD, and/or PTSD.

Table 3 presents bivariate comparisons of veterans with and without increased psychiatric distress on measures of sociodemographic and military characteristics, pre-pandemic psychosocial risk and protective factors, pre-to-peri-pandemic changes in risk factors, and COVID-19-related infection and pandemic stressors. Compared with veterans who did not experience an increase in distress (mean change in distress = -0.20 , $s.d.$ = 0.5), veterans who did were younger and more likely to have enlisted into the military. They were also more likely to report several pre-pandemic risk factors, including more adverse childhood experiences, lifetime diagnosis of MDD and/or PTSD, lifetime diagnosis of alcohol use disorder and/or drug use disorder, and greater severity of past-year alcohol problems, frequency of past-year non-prescription drug use, and loneliness. They also scored lower on pre-pandemic measures of protective psychosocial characteristics and social connectedness and were more likely to report increases in loneliness and alcohol-related problems during the pandemic. In addition, they were more likely to endorse exposure to a range of infection and pandemic stressors, including knowing someone infected with COVID-19 and experiencing COVID-19-related disease worries, social restriction stress, socioeconomic stress, and relationship difficulties.

Table 4 presents results of a hierarchical logistic regression analysis examining independent pre-pandemic predictors and COVID-19-related correlates of increased psychiatric distress. In step 1 of the model (pre-pandemic risk factors), fewer adverse childhood experiences and greater severity of past-year alcohol problems and loneliness predicted increased psychiatric distress.

Table 1. Measures of psychiatric, sociodemographic, military, and psychosocial variables and COVID-19 infection and pandemic stressors

Psychiatric conditions	
MDD	Score ≥ 3 on the two depression items of the Patient Health Questionnaire for Depression and Anxiety (PHQ-4; Kroenke, Spitzer, Williams, & Löwe, 2009) in the past 2 weeks
GAD	Score ≥ 3 on the two anxiety items of the PHQ-4 (Kroenke et al., 2009) in the past 2 weeks
PTSD	Past-month total score ≥ 33 on PTSD Checklist for DSM-5 (PCL-5; Weathers et al., 2013b); veterans reported on past-month PTSD symptoms in relation to their self-reported 'worst' Criterion A trauma on the Life Events Checklist for DSM-5 (Weathers et al., 2013a)
Sociodemographic characteristics	Age (continuous), sex (male, female), race/ethnicity (white, non-white), education (college graduate or higher, up to high school diploma), marital status (married/living with partner, not), household income (\$60 000 or more, less than \$60 000), retirement status (retired, not)
Military characteristics	Combat veteran status (combat exposure, not), primary source of healthcare (VA, other), enlistment status (enlisted into military, drafted into military, commissioned into military)
Pre-pandemic psychosocial risk factors	
Adverse childhood experiences	Adverse Childhood Experiences Questionnaire score (ACEQ; Finkelhor, Shattuck, Turner, & Hamby, 2015)
Total traumas	Items endorsed on Life Events Checklist for DSM-5 (LEC-5; Weathers et al., 2013a)
Lifetime MDD or PTSD	Lifetime MDD was assessed according to DSM-5 diagnostic criteria using the Mini International Neuropsychiatric Interview (MINI; Sheehan, 2016). Lifetime PTSD was defined as a score of 33+ (Bovin et al., 2016) on the PCL-5 (Weathers et al., 2013b), which was modified to include lifetime ratings of all PTSD symptoms in relation to veterans' self-reported 'worst' Criterion A trauma on the LEC-5 (Weathers et al., 2013a). Veterans who met criteria for either disorder were coded positive for lifetime MDD or PTSD
Lifetime AUD or DUD	Lifetime AUD and DUD were defined as meeting DSM-5 diagnostic criteria for AUD or DUD, respectively, as assessed using the MINI (Sheehan, 2016). Veterans who met criteria for either disorder were coded positive for lifetime AUD or DUD
Past-year alcohol use problem severity	Alcohol Use Disorders Identification Test (AUDIT) total score (Saunders, Aasland, Babor, De La Fuente, & Grant, 1993)
Past-year days of non-prescription drug use	Number of days reported in response to the following question from the Screen of Drug Use (Tiet et al., 2015): How many days in the past year have you used <i>non-prescription</i> drugs?
Loneliness	Score on 3-item measure adapted from the UCLA Loneliness Scale (Hughes et al., 2004)
Pre-pandemic psychosocial protective factors	
Protective psychosocial characteristics	Factor score using the following six indicators: score on Purpose in Life Test-Short Form (Schulenberg, Schnetzer, and Buchanan, 2011); score on Connor-Davidson Resilience Scale-10 (Campbell-Sills & Stein, 2007); rating (1 = strongly disagree to 7 = strongly agree) on single-item measure of optimism from Life Orientation Test-Revised ('In uncertain times, I usually expect the best') (Scheier, Carver, & Bridges, 1994); rating (1 = strongly disagree to 7 = strongly agree) on single-item measure of gratitude from Gratitude Questionnaire ('I have so much in life to be thankful for') (McCullough, Emmons, & Tsang, 2002); rating (1 = strongly disagree to 7 = strongly agree) on single-item measure of curiosity/exploration from Curiosity and Exploration Inventory-II ('I frequently find myself looking for new opportunities to grow as a person') (Kashdan et al., 2009); rating (1 = strongly disagree to 7 = strongly agree) on single item measuring perceived level of community integration ('I feel well integrated in my community')
Social connectedness	Factor score using the following three measures as indicators: score on 5-item Medical Outcomes Study Social Support Scale (Sherbourne & Stewart, 1991); size of social network as assessed by the question, 'About how many close friends and relatives do you have (people you feel at ease with and can talk to about what is on your mind)?'; secure attachment style as measured by the Attachment Style Questionnaire (ASQ; Hazan & Shaver, 1987)
Change in psychosocial risk factors from pre- to peri-pandemic	Increase in alcohol consumption (AUDIT consumption; Saunders et al., 1993), increase in alcohol use problem severity (AUDIT consequences; Saunders et al., 1993), increase in days of non-prescription drug use (Screen of Drug Use; Tiet et al., 2015), change in loneliness (UCLA Loneliness Scale; Hughes et al., 2004)
COVID-19 infection stressors	COVID-19 infection status (endorsement of: self-infected, know someone in household who was infected, know someone not in household who was infected, and know someone who died of COVID-19)
COVID-19 pandemic stressors	Questions from the Coronavirus Health Impact Survey (National Institute of Mental Health, 2020) were used to assess COVID-19-associated worries and concerns at the peri-pandemic assessment. Factor analysis revealed that these items loaded on five factors (eigen values = 1.01–4.94): COVID-19-related disease worries (e.g. 'In the past month, how worried have you been about being infected with coronavirus?'); COVID-19 social restriction stress (e.g. 'How stressful have these changes in social contacts been for you?'); COVID-19-associated socioeconomic stress (e.g. 'In the past month, to what degree have changes associated to the pandemic created financial problems for you or your family?'); COVID-19-associated relationship difficulties (e.g. 'Has the quality of the relationships between you and members of your family changed?'); and COVID-19-associated social engagement (e.g. 'In the past month, how many people, from outside of your household, have you had an in-person conversation with?')

COVID-19, coronavirus disease-2019; VA, veterans affairs; DSM-5, Diagnostic and Statistical Manual of Mental Disorders, 5th edition; MDD, major depressive disorder; GAD, generalized anxiety disorder; PTSD, posttraumatic stress disorder; AUD, alcohol use disorder; DUD, drug use disorder; UCLA, University of California, Los Angeles.

Table 2. Pre-pandemic to 1-year peri-pandemic prevalence of positive screens for MDD, GAD, and PTSD among U.S. military veterans

	Pre-pandemic N (weighted %)	1-Year peri-pandemic N (weighted %)	Test of difference	<i>p</i>
MDD	198 (7.6)	225 (8.6)	3.31	0.07
Age 18–44	31 (18.8)	33 (21.6)	1.41	0.24
Age 45–64	97 (9.7)	102 (10.5)	0.57	0.45
Age 65+	197 (3.4)	219 (4.0)	1.16	0.28
GAD	161 (7.1)	241 (9.4)	18.92	<0.001
Age 18–44	35 (21.7)	39 (23.0)	0.28	0.60
Age 45–64	76 (8.2)	123 (13.5)	22.26	<0.001
Age 65+	49 (2.5)	74 (3.3)	1.41	0.24
PTSD	140 (4.8)	113 (4.1)	2.16	0.14
Age 18–44	12 (16.6)	16 (13.3)	1.16	0.28
Age 45–64	59 (13.3)	64 (11.8)	0.65	0.42
Age 65+	31 (3.6)	30 (3.5)	0.00	1.00

MDD, major depressive disorder; GAD, generalized anxiety disorder; PTSD, posttraumatic stress disorder.
Statistically significant *p* values (*p* < 0.05) are bolded.

In step 2 of the model (pre-pandemic protective factors), veterans with higher levels of social connectedness were less likely to experience increased distress. In step 3 of the model (COVID-19-related changes and stressors), veterans with greater increases in loneliness and higher levels of COVID-19-related disease worries, social restriction and socioeconomic stress, and relationship difficulties were significantly more likely to experience increased psychiatric distress.

Results of planned post-hoc analyses revealed that veterans with a secure attachment style [Wald $\chi^2 = 8.02$, *p* = 0.005; odds ratio (OR) 0.65, 95% confidence interval (CI) 0.48–0.87] and larger social network (Wald $\chi^2 = 3.99$, *p* = 0.046; OR 0.98, 95% CI 0.96–0.99) pre-pandemic were significantly less likely to experience increased psychiatric distress 1 year into the pandemic. With regard to pandemic-related items, increased psychiatric distress was associated with greater concern about the stability of one's living situation (Wald $\chi^2 = 36.64$, *p* < 0.001; OR 1.67, 95% CI 1.42–1.98), worries about COVID-19 affecting one's mental health (Wald $\chi^2 = 11.90$, *p* = 0.001; OR 1.24, 95% CI 1.10–1.40), stress related to changes in social contacts (Wald $\chi^2 = 11.22$, *p* = 0.001; OR 1.31, 95% CI 1.12–1.53), worsening quality of one's relationships with friends (Wald $\chi^2 = 6.68$, *p* = 0.010; OR 1.30, 95% CI 1.06–1.58), and worries about COVID-19 infecting friends or family (Wald $\chi^2 = 6.49$, *p* = 0.011; OR 1.18, 95% CI 1.04–1.35).

As shown in Fig. 1, a relative importance analysis revealed that social network size [12.2% relative variance explained (RVE)] and loneliness (11.7% RVE) were the strongest pre-pandemic predictors of increased psychiatric distress. Stress related to changes in social contacts during the pandemic (15.8% RVE), worry about effects of the pandemic on one's mental health (15.3% RVE), and increased loneliness during the pandemic (13.6% RVE) were the strongest peri-pandemic correlates of this outcome.

Discussion

To our knowledge, this is the first study to examine pandemic-related mental health changes in a nationally

representative sample of U.S. military veterans. The current findings suggest that veterans evidenced considerable mental health resilience during the COVID-19 pandemic, but with substantial heterogeneity in individual vulnerability. Consistent with our first hypothesis, we found a pre-to-peri-pandemic increase in positive screens for GAD. This increase translates into substantial negative impact when extrapolated to the national population of nearly 18 million veterans (Vespa, 2020); using estimated population benchmarks (Bureau of the Census, 2019), more than 400 000 additional veterans may have screened positive for GAD 1-year into the pandemic. Elevated GAD was driven largely by veterans aged 45–64 years, among whom GAD positive screens increased by more than 65% from pre-to-peri pandemic (8.2% to 13.5%). This result was somewhat surprising given previous findings of greater pandemic-related distress among young adults (Breslau et al., 2021; Czeisler et al., 2020; Holman et al., 2020; McGinty et al., 2020b). However, data from previous studies came from early in the pandemic, and there is some evidence that age differences in psychiatric distress decreased over time (Banks, Fancourt, & Xu, 2021). Our results suggest that middle-aged veterans may have experienced unique challenges as the pandemic progressed, such as overlapping stressors related to the health and economic impacts of the pandemic. Indeed, post-hoc comparisons of the three age groups revealed that middle-aged veterans reported higher levels of COVID-19-related disease worries than younger veterans, and greater COVID-19-related socioeconomic stress than older veterans. As middle-aged individuals typically are not yet retired but are also of an age associated with more severe health impacts of COVID-19 (Fang et al., 2020), this group may have found it difficult to continuously self-isolate due to work-related responsibilities, and stressful to attend in-person work due to heightened disease risk. Moreover, this age cohort often balances work and care-taking responsibilities (Hopps, Iadeluca, McDonald, & Makinson, 2017), and they may have been disproportionately impacted by persistent school closures and remote learning (Garbe, Ogurlu, Logan, & Cook, 2020).

Although this increased anxiety is consistent with elevated distress observed in cross-sectional surveys (Czeisler et al., 2020;

Table 3. Characteristics of U.S. military veterans with and without increased psychological distress from pre- to peri-pandemic

	No increase in psychological distress composite score <i>N</i> = 2672 (86.8%)	Increase in psychological distress composite score <i>N</i> = 389 (13.2%)	Test of difference	<i>p</i>
	Mean (s.d.) or <i>N</i> (weighted %)	Mean (s.d.) or <i>N</i> (weighted %)		
Sociodemographic characteristics				
Age	63.6 (14.7)	61.5 (14.4)	2.59	0.010
Male sex	2391 (91.9)	327 (89.3)	3.01	0.08
White race/ethnicity	2210 (79.2)	317 (79.1)	0.00	0.96
College graduate or higher education	1225 (34.5)	176 (32.3)	0.71	0.40
Married/partnered	1931 (73.8)	280 (75.3)	0.41	0.52
Household income \$60k or higher	1620 (61.1)	224 (59.3)	0.48	0.49
Retired	1518 (47.2)	209 (44.5)	0.98	0.32
Military characteristics				
Combat veteran	916 (35.0)	130 (37.8)	1.17	0.28
VA primary source of healthcare	500 (20.3)	77 (19.8)	0.05	0.82
Enlisted into military	2004 (77.3)	316 (83.7)	8.26	0.016
Pre-pandemic psychosocial risk factors				
Baseline psychological distress	0.0 (0.6)	0.0 (0.9)	0.17	0.86
Adverse childhood experiences	1.4 (1.9)	1.7 (2.0)	−3.07	0.002
Total traumas	8.9 (8.3)	9.4 (8.2)	−1.11	0.27
Lifetime MDD and/or PTSD	466 (20.2)	123 (34.6)	38.89	<0.001
Lifetime AUD and/or DUD	1065 (41.8)	197 (50.5)	10.26	0.001
Past-year alcohol use problem severity	3.0 (4.4)	3.9 (4.7)	−3.48	0.001
Past-year days of non-prescription drug use	12.8 (60.4)	22.2 (79.5)	−2.18	0.030
Loneliness	4.4 (1.8)	5.3 (1.9)	−9.33	<0.001
Pre-pandemic psychosocial protective factors				
Protective psychosocial characteristics	0.1 (1.0)	−0.3 (0.9)	8.90	<0.001
Social connectedness	0.1 (1.0)	−0.4 (0.9)	8.34	<0.001
Change in risk factors from pre- to peri-pandemic				
Increase in alcohol consumption	451 (17.7)	72 (19.1)	0.45	0.50
Increase in alcohol-related problems	522 (7.7)	99 (14.9)	22.14	<0.001
Increase in non-prescription drug use	52 (2.2)	14 (3.5)	2.21	0.14
Change in loneliness	−0.1 (1.3)	0.2 (1.5)	−4.12	<0.001
COVID-19 infection stressors				
Infected with COVID-19	198 (8.0)	34 (9.8)	1.46	0.23
Someone in household infected with COVID-19	169 (7.4)	27 (8.3)	0.36	0.55
Know someone infected with COVID-19	198 (40.8)	34 (48.3)	7.82	0.005
Know someone who died of COVID-19	147 (5.5)	26 (5.6)	0.00	0.95

(Continued)

Table 3. (Continued.)

	No increase in psychological distress composite score $N = 2672$ (86.8%)	Increase in psychological distress composite score $N = 389$ (13.2%)		
	Mean (s.d.) or N (weighted %)	Mean (s.d.) or N (weighted %)	Test of difference	p
COVID-19 pandemic stressors				
COVID-19-related disease worries	-0.1 (1.0)	0.3 (1.0)	-6.94	<0.001
COVID-19-related social restriction stress	0.0 (1.0)	0.3 (1.0)	-5.64	<0.001
COVID-19-related socioeconomic stress	-0.1 (0.9)	0.3 (1.3)	-5.34	<0.001
COVID-19-related relationship difficulties	-0.1 (1.0)	0.3 (1.2)	-5.82	<0.001
Social engagement during pandemic	0.0 (1.0)	0.0 (1.1)	0.52	0.61
Hours of COVID-19-related media consumed per week	1.6 (2.1)	1.7 (2.1)	-1.11	0.27

s.d., standard deviation; VA, veterans affairs; MDD, major depressive disorder; PTSD, posttraumatic stress disorder; AUD, alcohol use disorder; DUD, drug use disorder; COVID-19, coronavirus disease 2019.

Increase in alcohol problems and increase in non-prescription drug use are binary variables; veterans with alcohol use problems or non-prescription drug use that increased by 0.5 s.d. or more from pre-pandemic to peri-pandemic were coded positive for respective increases in alcohol problems and drug use. Statistically significant p values ($p < 0.05$) are bolded.

Holman et al., 2020; McGinty et al., 2020b), the increase was smaller in the current study, and we observed notable stability in the prevalence of MDD and PTSD positive screens. This may be because the current study assessed the same cohort pre- and peri-pandemic, which is essential to accurately assessing pandemic-related changes over time. Indeed, our findings are consistent with a developing literature of longitudinal studies showing only small within-subject, population-level changes in mental health outcomes during the COVID-19 pandemic. For example, a recent rapid-review and meta-analysis of 25 longitudinal and natural experiment studies consisting of more than 70 000 participants from around the globe found small effects of COVID-19 lockdowns on anxiety ($g = 0.17$) and depression ($g = 0.15$), and null effects on general distress, positive functioning, and suicide risk (Prati & Mancini, 2021). Similarly, roughly 75% of respondents in the only previous U.S. population-based longitudinal study reported low psychiatric distress at both pre- and peri-pandemic assessments (Breslau et al., 2021). The current study extends this literature by demonstrating a similar pattern among U.S. military veterans, a population that has been characterized as high-risk for mental health impacts of the COVID-19 pandemic (Amsalem et al., 2021).

Despite considerable resilience, however, pandemic-related mental health changes among veterans were heterogeneous, mirroring research showing wide individual variability in responses to acute stress (Bonanno & Mancini, 2012). In fact, when we examined within-person changes in transdiagnostic symptoms of MDD, GAD, and PTSD, nearly 1 in 7 veterans experienced more than a 0.5 s.d. increase in distress relative to their pre-pandemic levels. This estimated 13.2% of veterans with increased distress is strikingly similar to the 12.8% of U.S. adults who reported increased distress from pre- to peri-pandemic in a previous longitudinal study (Breslau et al., 2021). As this increase was not fully accounted for by the increased prevalence of positive GAD screens, it appears that a significant proportion of the

distress that emerged during the pandemic was sub-diagnostic. However, the magnitude of this increase suggests a need for intervention to prevent increased distress from transitioning into more persistent psychiatric disorders. Thus, these findings highlight the importance of delineating risk factors for pandemic-related psychiatric distress to target interventions to vulnerable veterans.

Results of a relative importance analysis revealed that pre-pandemic social factors – specifically, social network size, loneliness, and attachment style – predicted increased psychiatric distress during the pandemic. Specifically, veterans with a larger pre-pandemic social network size and secure attachment style were less likely to experience an increase in distress, whereas veterans with greater pre-pandemic loneliness were more likely to experience such an increase. Collectively, pre-pandemic social factors accounted for roughly a third of the explained variance in increased distress. These findings, which are consistent with a literature showing that social support may help buffer the deleterious effects of stressors through psychobiological mechanisms (Ditzen & Heinrichs, 2014), align with calls for health care providers to regularly monitor social factors in populations at increased risk for pandemic-related distress (Killgore, Cloonan, Taylor, Lucas, & Dailey, 2020). Brief, validated measures such as the Three-Item Loneliness Scale (Hughes, Waite, Hawkey, & Cacioppo, 2004) could be used to screen veterans in primary and mental health care settings.

Results of a relative importance analysis also identified potentially modifiable pandemic-related stressors associated with increased psychiatric distress, including social losses – stress about changes in social contacts, increased loneliness, and worsening in the quality of friend relationships – worry about effects of the pandemic on mental health, and concern about housing stability during the pandemic. These findings support our third hypothesis and have several practical implications. First, associations of social loss with increased distress highlight the importance of prevention efforts such as outreach and messaging

Table 4. Results of hierarchical logistic regression analysis of predictors and correlates of increased psychological distress

	<i>B</i>	<i>S.E.</i>	Wald	<i>p</i>	AOR (95% CI)
Step 1: Pre-pandemic risk factors					
Age	0.00	0.00	0.10	0.76	1.00 (0.99–1.01)
Enlistment status					
Enlisted	–	–	–	–	1 (reference)
Drafted	–0.37	0.23	2.57	0.11	0.69 (0.44–1.08)
Commissioned	–0.16	0.22	0.50	0.48	0.86 (0.56–1.32)
Adverse childhood experiences	–0.09	0.03	6.96	0.008	0.91 (0.85–0.98)
Lifetime MDD and/or PTSD	–0.02	0.16	0.02	0.87	0.97 (0.71–1.34)
Lifetime AUD and/or DUD	–0.11	0.13	0.67	0.41	0.90 (0.69–1.17)
Alcohol use problem severity	0.03	0.01	6.03	0.014	1.03 (1.01–1.06)
Past-year non-prescription drug use	0.00	0.00	1.43	0.23	1.00 (1.00–1.00)
Loneliness	0.17	0.05	11.67	0.001	1.18 (1.07–1.30)
<i>R</i> ² : 0.06					
Step 2: Pre-pandemic protective factors					
Protective psychosocial characteristics	–0.11	0.08	2.25	0.13	0.89 (0.77–1.04)
Social connectedness	–0.21	0.09	5.65	0.017	0.81 (0.69–0.96)
<i>R</i> ² : 0.07					
<i>R</i> ² change: 0.01					
Step 3: COVID-19-related changes and stressors					
Increase in alcohol problems	0.24	0.14	2.76	0.10	1.27 (0.96–1.68)
Change in loneliness	0.25	0.05	26.37	<0.001	1.28 (1.16–1.40)
Know someone infected with COVID-19	0.03	0.13	0.07	0.80	1.03 (0.81–1.32)
COVID-19-related disease worries	0.37	0.06	35.66	<0.001	1.44 (1.28–1.62)
COVID-19-related social restriction stress	0.22	0.06	13.94	<0.001	1.25 (1.11–1.40)
COVID-19-related socioeconomic stress	0.19	0.05	13.00	<0.001	1.21 (1.09–1.34)
COVID-19-related relationship difficulties	0.22	0.06	14.97	<0.001	1.25 (1.12–1.40)
<i>R</i> ² : 0.16					
<i>R</i> ² change: 0.09					

AOR, adjusted odds ratio; CI, confidence interval; MDD, major depressive disorder; PTSD, posttraumatic stress disorder; AUD, alcohol use disorder; DUD, drug use disorder; COVID-19, coronavirus disease-2019.

Statistically significant *p* values (*p* < 0.05) are bolded.

campaigns to share strategies for virtual social connection and to destigmatize loneliness. Second, associations of distress with worry about pandemic effects on mental health underscore the need for public investment to reduce barriers to mental health care. As telehealth delivery is as effective as in-person delivery of evidence-based anxiety, depression, and PTSD treatments (Morland *et al.*, 2020; Tuerk, Keller, & Acerno, 2018), expansion of these services is warranted. Resources to provide smart tablets, as successfully piloted by the Department of Veterans Affairs (Zulman *et al.*, 2019), could expand access to low-income households where smart phone and laptop ownership are not universal (Nadkarni *et al.*, 2020). Brief video interventions that boost treatment-seeking intentions may also be helpful (Amsalem *et al.*, 2021), as many veterans even with severe psychological symptoms are not engaged in treatment (Nichter, Hill, Norman, Haller, & Pietrzak, 2020b).

Third, associations of distress with housing stability concerns highlight the need for continued government financial support for vulnerable individuals – perhaps particularly for middle-aged veterans. The high levels of financial stress and disease worries reported by this age cohort suggest greater fears of job instability and virus exposure, which may happen in work settings. This distress may signal a need for extension of unemployment insurance and direct payments introduced early in the pandemic to offset the impact of persistent job losses and concerns about the safety of returning to work. To abate housing concerns, expansion of housing-specific interventions such as rental and mortgage assistance and eviction diversion programs may be beneficial (Benfer *et al.*, 2021). For veterans specifically, this relief may come from the recent allocation of additional funding to the Housing and Urban Development-Veterans Affairs Supportive Housing program (HUD-VASH), which provides rental assistance and

Fig. 1. Results of relative importance analysis of significant correlates of increased psychological distress from pre- to peri-pandemic.

Note. COVID-19=coronavirus disease 2019; Error bars represent 95% confidence intervals.

Adverse childhood experiences (0.8% relative variance explained) and pre-pandemic alcohol use severity (2.8% relative variance explained) were not significant predictors of increased psychiatric distress in the relative importance analysis (both p 's > 0.05).

housing vouchers to veterans at risk of homelessness. Our findings support the expansion of this program's priorities during the pandemic to include a greater focus on homelessness prevention, which could include direct renter payments to veterans accumulating significant debt that may result in homelessness when eviction moratoria are lifted.

The current study has important strengths, including the use of a large, nationally representative sample of veterans, a pre-to-peri-pandemic longitudinal design, and assessment of the prevalence of psychiatric conditions in late 2020, after pandemic-related mental health effects had additional time to unfold. However, results must also be interpreted in the context of several limitations. First, the sample consisted of predominantly older, white, and male U.S. military veterans; distress might be higher among individuals underrepresented in the veteran population. As previous research has found greater pandemic-related distress among women (Breslau et al., 2021), ethnic/racial minorities (McGinty et al., 2020b), and young adults (Breslau et al., 2021; McGinty et al., 2020b), the current results may not generalize to civilians or veterans in these groups, or to veterans who are institutionalized or homeless. Second, although retention was relatively high (75.6%), veterans with a pre-pandemic diagnosis of MDD, GAD, and/or PTSD were less likely to complete the follow-up survey than veterans without these diagnoses (67.4% v. 76.7%). Therefore, it is possible that the prevalence of these disorders may have been underestimated due to retention or survivorship bias (Czeisler, Wiley, Czeisler, Rajaratnam, & Howard, 2021). Third, although we assessed peri-pandemic stress later in the pandemic than previous studies, mental health effects of the pandemic may be delayed and manifest over time, as economic, health, and social consequences of the pandemic compound. Fourth, screening instruments were used to assess psychiatric symptoms; studies using clinical interviews are needed to confirm the prevalence estimates reported in this study. In addition, we selected a 0.5 S.D. cut-off to indicate increased psychiatric distress, which, although identified in previous research as a valid indicator of clinical distress (e.g. Norman et al., 2003) and significantly associated with mental health

treatment-seeking in the current sample, may or may not indicate a transition to a clinically concerning level of symptoms; indeed, some studies have used higher thresholds to define adverse mental health outcomes (e.g. top 10% of the distribution of change scores; Chen and Hardy, 2009). However, we note that the group created using this cut-off experienced, on average, greater than a standard deviation increase in distress (mean change in distress = 1.1 S.D., S.D. = 0.6); in fact, the prevalence of positive screens for MDD, GAD, and/or PTSD in this group increased from 9.4% at pre-pandemic to 54.2% at peri-pandemic. Although it is possible that some of the increased distress we observed, particularly lower magnitude increases, reflected normal fluctuations in symptoms, we posit that even sub-diagnostic increases in distress are important to identify as they may portend more substantial problems. Fifth, COVID-19-related correlates of increased psychiatric distress were measured simultaneously with peri-pandemic distress, making the temporal direction of these associations unclear; it is possible that increased mental health symptoms caused veterans to report greater perceived stress related to the pandemic.

Conclusions

These limitations notwithstanding, the current study provides the first known nationally representative data on the mental health impact of the COVID-19 pandemic on U.S. military veterans. Our findings, which align with the broader literature showing high rates of adaptive functioning in the face of trauma, adversity, and disaster (Bonanno, 2005; Norris, Tracy, & Galea, 2009; North et al., 2002), reflect the remarkable human ability to adapt even to profound disruptions to daily life wrought by the pandemic. However, they also highlight disparities in the effects of the pandemic and the resultant need for targeted interventions to protect those at risk of deteriorating mental health, particularly middle-aged veterans and veterans with low social connectedness.

To better understand trajectories of risk and resilience, it will be important to continue tracking long-term changes in mental health, ideally using continuous measures of psychopathology,

which are more sensitive to detecting within-person changes in distress. Because of the ongoing nature of the pandemic and the continued presence of imminent stressors, it is unclear whether the elevated distress we observed will result in durable and functionally impairing psychiatric disorders. Prior research has found that population-level increases in distress following natural disasters often return to pre-disaster levels over time (Pietrzak et al., 2012). Although still early, emerging evidence about the persistence of distress during the pandemic in the USA is mixed; one study found an early spike in distress in spring of 2020 that returned to baseline levels by summer (Daly & Robinson, 2020), but another study found consistent high distress in April and July (McGinty et al., 2020a). Because data from the current study were collected in the fall of 2020, the elevated distress we observed may be an indicator of more persistent mental health problems. Post-pandemic follow-up will be essential to understanding the prevalence and determinants of acute *v.* long-term changes in mental health and functioning in veterans and the population at large.

Acknowledgements. The authors thank the veterans who participated in this study.

Financial support. The National Health and Resilience in Veterans Study is supported by the U.S. Department of Veterans Affairs National Center for Posttraumatic Stress Disorder.

Conflict of interest. The authors MH, BN, SN, LM, and RP have no conflicts of interest to declare. JK is a scientific advisor to Biohaven Pharmaceuticals, BioXcel Therapeutics, Inc., Cadent Therapeutics (Clinical Advisory Board), PsychoGenics, Inc., Stanley Center for Psychiatric research at the Broad Institute of MIT and Harvard, Lohocla Research Corporation. JK owns stock and/or stock options in Biohaven Pharmaceuticals, Sage Pharmaceuticals, Spring Care, Inc., BlackThorn Therapeutics, Inc., Terran Biosciences, Inc. JK reports income <\$10 000 per year from: AstraZeneca Pharmaceuticals, Biogen, Idex, MA, Biomedisyn Corporation, Bionomics, Limited (Australia), Boehringer Ingelheim International, Concert Pharmaceuticals, Inc., Epiodyne, Inc., Heptares Therapeutics, Limited (UK), Janssen Research & Development, L.E.K. Consulting, Otsuka America Pharmaceutical, Inc., Perception Neuroscience Holdings, Inc. Spring Care, Inc., Sunovion Pharmaceuticals, Inc., Takeda Industries, Taisho Pharmaceutical Co., Ltd. JK reports income >\$10 000 per year from Biological Psychiatry (Editor). JK received the drug, Saracatinib from AstraZeneca and Mavoglurant from Novartis for research related to NIAAA grant 'Center for Translational Neuroscience of Alcoholism' (CTNA-4) from AstraZeneca Pharmaceuticals. JK holds the following patents: (1) Seibyl JP, Krystal JH, Charney DS. Dopamine and noradrenergic reuptake inhibitors in treatment of schizophrenia. US Patent no. 5447948. 5 September 1995; (2) Vladimir, Coric, Krystal, John H, Sanacora, Gerard – Glutamate Modulating Agents in the Treatment of Mental Disorders US Patent No. 8778979 B2; Patent Issue Date: 15 July 2014. US Patent Application No. 15/695164; Filing Date: 5 September 2017; (3) Charney D, Krystal JH, Manji H, Matthew S, Zarate C, Intranasal Administration of Ketamine to Treat Depression United States Application No. 14/197767 filed on 5 March 2014; United States application or Patent Cooperation Treaty (PCT) International application No. 14/306382 filed on 17 June 2014; (4) Zarate, C, Charney, DS, Manji, HK, Mathew, Sanjay J, Krystal, JH, Department of Veterans Affairs 'Methods for Treating Suicidal Ideation', Patent Application No. 14/197767 filed on 5 March 2014 by Yale University Office of Cooperative Research; (5) Arias A, Petrakis I, Krystal JH. Composition and methods to treat addiction. Provisional Use Patent Application no. 61/973/961. 2 April 2014. Filed by Yale University Office of Cooperative Research; (6) Chekroud A, Gueorgieva R, Krystal JH. 'Treatment Selection for Major Depressive Disorder' (filing date 3 June 2016, USPTO docket number Y0087.70116US00). Provisional patent submission by Yale University; (7) Gihyun, Yoon, Petrakis I, Krystal JH Compounds, Compositions and

Methods for Treating or Preventing Depression and Other Diseases. U.S. Provisional Patent Application No. 62/444552, filed on 10 January 2017 by Yale University Office of Cooperative Research OCR 7088 US01; and (8) Abdallah C, Krystal JH, Duman R, Sanacora G. Combination Therapy for Treating or Preventing Depression or Other Mood Diseases. U.S. Provisional Patent Application No. 62/719935 filed on 20 August 2018 by Yale University Office of Cooperative Research OCR 7451 US01.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- Amsalem, D., Lazarov, A., Markowitz, J. C., Gorman, D., Dixon, L. B., & Neria, Y. (2021). Increasing treatment-seeking intentions of US veterans in the COVID-19 era: A randomized controlled trial. *Depression and Anxiety*, 38(6), 639–647. <https://doi.org/10.1002/da.23149>.
- Bandoli, G., Campbell-Sills, L., Kessler, R. C., Heeringa, S. G., Nock, M. K., Rosellini, A. J., ... Stein, M. B. (2017). Childhood adversity, adult stress, and the risk of major depression or generalized anxiety disorder in US soldiers: A test of the stress sensitization hypothesis. *Psychological Medicine*, 47(13), 2379–2392. <https://doi.org/10.1017/S0033291717001064>.
- Banks, J., Fancourt, D., & Xu, X. (2021). Mental health and the COVID-19 pandemic. In J. F. Helliwell, R. Layard, J. D. Sachs, J. E. De Neve, L. B. Aknin, & S. Wang (Eds.), pp. 107–130. New York, NY, USA: Sustainable Development Solutions Network. Retrieved from The World Happiness Report. <https://happiness-report.s3.amazonaws.com/2021/WHR+21.pdf>.
- Benfer, E. A., Vlahov, D., Long, M. Y., Walker-Wells, E., Pottenger, J. L., Gonsalves, G., ... Keene, D. E. (2021). Eviction, health inequity, and the spread of COVID-19: Housing policy as a primary pandemic mitigation strategy. *Journal of Urban Health*, 98(1), 1–12. <https://doi.org/10.1007/s11524-020-00502-1>.
- Birditt, K. S., Turkelson, A., Fingerman, K. L., Polenick, C. A., & Oya, A. (2021). Age differences in stress, life changes, and social ties during the COVID-19 pandemic: Implications for psychological well-being. *The Gerontologist*, 61(2), 205–216. <https://doi.org/10.1093/geront/gnaa204>.
- Bonanno, G. A. (2005). Resilience in the face of potential trauma. *Current Directions in Psychological Science*, 14(3), 135–138. <https://doi.org/10.1111/j.0963-7214.2005.00347.x>.
- Bonanno, G. A., & Mancini, A. D. (2012). Beyond resilience and PTSD: Mapping the heterogeneity of responses to potential trauma. *Psychological Trauma: Theory, Research, Practice, and Policy*, 4(1), 74–83. <https://doi.org/10.1037/a0017829>.
- Bovin, M. J., Marx, B. P., Weathers, F. W., Gallagher, M. W., Rodriguez, P., Schnurr, P. P., & Keane, T. M. (2016). Psychometric properties of the PTSD checklist for diagnostic and statistical manual of mental disorders-fifth edition (PCL-5) in veterans. *Psychological Assessment*, 28(11), 1379–1391. <https://doi.org/10.1037/pas0000254>.
- Breslau, J., Finucane, M. L., Locker, A. R., Baird, M. D., Roth, E. A., & Collins, R. L. (2021). A longitudinal study of psychological distress in the United States before and during the COVID-19 pandemic. *Preventive Medicine*, 143, 1–4. <https://doi.org/10.1016/j.ypmed.2020.106362>.
- Bureau of the Census. (2019). The Current Population Survey design and methodology (Technical Paper 66). Retrieved from <https://www2.census.gov/programs-surveys/cps/techdocs/cpsaug19.pdf>.
- Campbell-Sills, L., & Stein, M. B. (2007). Psychometric analysis and refinement of the Connor-Davidson Resilience Scale (CD-RISC): Validation of a 10-item measure of resilience. *Journal of Traumatic Stress*, 20(6), 1019–1028. <https://doi.org/10.1002/jts.20271>.
- Chen, L. Y., & Hardy, C. L. (2009). Alcohol consumption and health status in older adults: A longitudinal analysis. *Journal of Aging and Health*, 21(6), 824–847. <https://doi.org/10.1177/0898264309340688>.
- Czeisler, M., Lane, R., Petrosky, E., Wiley, J., Christensen, A., Njai, R., ... Rajaratnam, S. (2020). Mental health, substance use, and suicidal ideation during the COVID-19 pandemic – United States, June 24–30, 2020.

- Morbidity and Mortality Weekly Report, 69(32), 1049. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7440121/>.
- Czeisler, M. E., Wiley, J. F., Czeisler, C. A., Rajaratnam, S. M. W., & Howard, M. E. (2021). Uncovering survivorship bias in longitudinal mental health surveys during the COVID-19 pandemic. *Epidemiology and Psychiatric Sciences*, 30, 1–10. <https://doi.org/10.1017/S204579602100038X>.
- Daly, M., & Robinson, E. (2020). Psychological distress and adaptation to the COVID-19 crisis in the United States. *Journal of Psychiatric Research*, 136, 603–609. <https://doi.org/10.1016/j.jpsychires.2020.10.035>.
- Ditzen, B., & Heinrichs, M. (2014). Psychobiology of social support: The social dimension of stress buffering. *Restorative Neurology and Neuroscience*, 32(1), 149–162. <https://doi.org/10.3233/RNN-139008>.
- Dong, E., Du, H., & Gardner, L. (2020). An interactive web-based dashboard to track COVID-19 in real time. *The Lancet Infectious Diseases*, 20, 533–534. [https://doi.org/10.1016/S1473-3099\(20\)30120-1](https://doi.org/10.1016/S1473-3099(20)30120-1).
- Fang, X., Li, S., Yu, H., Wang, P., Zhang, Y., Chen, Z., ... Ma, X. (2020). Epidemiological, comorbidity factors with severity and prognosis of COVID-19: A systematic review and meta-analysis. *Aging*, 12(13), 12493–12503. <https://doi.org/10.18632/aging.103579>.
- Finkelhor, D., Shattuck, A., Turner, H., & Hamby, S. (2015). A revised inventory of adverse childhood experiences. *Child Abuse and Neglect*, 48, 13–21. <https://doi.org/10.1016/j.chiabu.2015.07.011>.
- Földi, M., Farkas, N., Kiss, S., Zádori, N., Váncsa, S., Szakó, L., ... Szentesi, A. (2020). Obesity is a risk factor for developing critical condition in COVID-19 patients: A systematic review and meta-analysis. *Obesity Reviews*, 21(10), e13095. <https://doi.org/10.1111/obr.13095>.
- Garbe, A., Ogurlu, U., Logan, N., & Cook, P. (2020). Parents' experiences with remote education during COVID-19 school closures. *American Journal of Qualitative Research*, 4(3), 45–65. <https://doi.org/10.29333/ajqr/8471>.
- Hazan, C., & Shaver, P. (1987). Romantic love conceptualized as an attachment process. *Journal of Personality and Social Psychology*, 52(3), 511–524. <https://doi.org/10.1037/0022-3514.52.3.511>.
- Hedden, S., Gfroerer, J., Barker, P., Smith, S., Pemberton, M. R., Saavedra, L. M., ... Novak, S. P. (2012). Comparison of NSDUH mental health data and methods with other data sources. In CBHSQ Data Review. Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK390286/>.
- Hill, M. L., Nichter, B., Loflin, M., Norman, S. B., & Pietrzak, R. H. (2021). Comparative associations of problematic alcohol and cannabis use with suicidal behavior in U.S. military veterans: A population-based study. *Journal of Psychiatric Research*, 135, 135–142. <https://doi.org/10.1016/j.jpsychires.2021.01.004>.
- Hoerster, K. D., Lehavot, K., Simpson, T., McFall, M., Reiber, G., & Nelson, K. M. (2012). Health and health behavior differences: U.S. military, veteran, and civilian men. *American Journal of Preventive Medicine*, 43(5), 483–489. <https://doi.org/10.1016/j.amepre.2012.07.029>.
- Hoggatt, K. J., Lehavot, K., Krennek, M., Schweizer, C. A., & Simpson, T. (2017). Prevalence of substance misuse among US veterans in the general population. *American Journal on Addictions*, 26(4), 357–365. <https://doi.org/10.1111/ajad.12534>.
- Hoglund, M. W., & Schwartz, R. M. (2014). Mental health in deployed and nondeployed veteran men and women in comparison with their civilian counterparts. *Military Medicine*, 179(1), 19–25. <https://doi.org/10.7205/MILMED-D-13-00235>.
- Holingue, C., Badillo-Goicoechea, E., Riehm, K. E., Veldhuis, C. B., Thrul, J., Johnson, R. M., ... Kalb, L. G. (2020). Mental distress during the COVID-19 pandemic among US adults without a pre-existing mental health condition: Findings from American trend panel survey. *Preventive Medicine*, 139, 1–8. <https://doi.org/10.1016/j.ypmed.2020.106231>.
- Holman, E., Thompson, R. R., Garfin, D. R., & Silver, R. C. (2020). The unfolding COVID-19 pandemic: A probability-based, nationally representative study of mental health in the United States. *Science Advances*, 6(42), 5390–5404. <https://doi.org/10.1126/sciadv.abd5390>.
- Holmes, E. A., O'Connor, R. C., Perry, V. H., Tracey, I., Wessely, S., Arseneault, L., ... Bullmore, E. (2020). Multidisciplinary research priorities for the COVID-19 pandemic: A call for action for mental health science. *The Lancet Psychiatry*, 7, 547–560. [https://doi.org/10.1016/S2215-0366\(20\)30168-1](https://doi.org/10.1016/S2215-0366(20)30168-1).
- Hopps, M., Iadeluca, L., McDonald, M., & Makinson, G. T. (2017). The burden of family caregiving in the United States: Work productivity, health care resource utilization, and mental health among employed adults. *Journal of Multidisciplinary Healthcare*, 10, 437–444. <https://doi.org/10.2147/JMDH.S135372>.
- Hughes, M. E., Waite, L. J., Hawkey, L. C., & Cacioppo, J. T. (2004). A short scale for measuring loneliness in large surveys: Results from two population-based studies. *Research on Aging*, 26(6), 655–672. <https://doi.org/10.1177/0164027504268574>.
- Ipsos. (2020). Ipsos public affairs project report for the 2019–2020 National Health and Resilience in Veterans Study. Ipsos Job Number: 19-073542-01.
- Kashdan, T. B., Gallagher, M. W., Silvia, P. J., Winterstein, B. P., Breen, W. E., Terhar, D., & Steger, M. F. (2009). The curiosity and exploration inventory-II: Development, factor structure, and psychometrics. *Journal of Research in Personality*, 43(6), 987–998. <https://doi.org/10.1016/j.jrp.2009.04.011>.
- Katon, J. G., Lehavot, K., Simpson, T. L., Williams, E. C., Barnett, S. B., Grossbard, J. R., ... Reiber, G. E. (2015). Adverse childhood experiences, military service, and adult health. *American Journal of Preventive Medicine*, 49(4), 573–582. <https://doi.org/10.1016/j.amepre.2015.03.020>.
- Kessler, R. C., Barker, P. R., Colpe, L. J., Epstein, J. F., Gfroerer, J. C., Hiripi, E., ... Zaslavsky, A. M. (2003). Screening for serious mental illness in the general population. *Archives of General Psychiatry*, 60(2), 184–189. <https://doi.org/10.1001/archpsyc.60.2.184>.
- Killgore, W. D. S., Cloonan, S. A., Taylor, E. C., Lucas, D. A., & Dailey, N. S. (2020). Loneliness during the first half-year of COVID-19 lockdowns. *Psychiatry Research*, 294, 1–2. <https://doi.org/10.1016/j.psychres.2020.113551>.
- Kroenke, K., Spitzer, R. L., Williams, J. B. W., & Löwe, B. (2009). An ultra-brief screening scale for anxiety and depression: The PHQ-4. *Psychosomatics*, 50(6), 613–621. [https://doi.org/10.1016/s0033-3182\(09\)70864-3](https://doi.org/10.1016/s0033-3182(09)70864-3).
- Kuwert, P., Knaevelsrud, C., & Pietrzak, R. H. (2014). Loneliness among older veterans in the United States: Results from the National Health and Resilience in Veterans Study. *American Journal of Geriatric Psychiatry*, 22(6), 564–569. <https://doi.org/10.1016/j.jagp.2013.02.013>.
- Lehavot, K., Hoerster, K. D., Nelson, K. M., Jakupcak, M., & Simpson, T. L. (2012). Health indicators for military, veteran, and civilian women. *American Journal of Preventive Medicine*, 42(5), 473–480. <https://doi.org/10.1016/j.amepre.2012.01.006>.
- Lehavot, K., Katon, J. G., Chen, J. A., Fortney, J. C., & Simpson, T. L. (2018). Post-traumatic stress disorder by gender and veteran status. *American Journal of Preventive Medicine*, 54(1), e1–e9. <https://doi.org/10.1016/j.amepre.2017.09.008>.
- McCullough, M. E., Emmons, R. A., & Tsang, J. A. (2002). The grateful disposition: A conceptual and empirical topography. *Journal of Personality and Social Psychology*, 82(1), 112–127. <https://doi.org/10.1037/0022-3514.82.1.112>.
- McGinty, E. E., Presskreischer, R., Anderson, K. E., Han, H., & Barry, C. L. (2020a). Psychological distress and COVID-19-related stressors reported in a longitudinal cohort of US adults in April and July 2020. *Journal of the American Medical Association*, 324(24), 2555–2557. <https://doi.org/10.1001/jama.2020.21231>.
- McGinty, E. E., Presskreischer, R., Han, H., & Barry, C. L. (2020b). Psychological distress and loneliness reported by US adults in 2018 and April 2020. *Journal of the American Medical Association*, 324(1), 93–94. <https://doi.org/10.1001/jama.2020.9740>.
- Morland, L. A., Wells, S. Y., Glassman, L. H., Greene, C. J., Hoffman, J. E., & Rosen, C. S. (2020). Advances in PTSD treatment delivery: Review of findings and clinical considerations for the use of telehealth interventions for PTSD. *Current Treatment Options in Psychiatry*, 7, 221–241. <https://doi.org/10.1007/s40501-020-00215-x>.
- Murphy, D., Williamson, C., Baumann, J., Busuttill, W., & Fear, N. T. (2020). Exploring the impact of COVID-19 and restrictions to daily living as a result of social distancing within veterans with pre-existing mental health difficulties. *BMJ Military Health*, 1–5. <https://doi.org/10.1136/bmjilitary-2020-001622>.
- Na, P. J., Tsai, J., Hill, M. L., Nichter, B., Norman, S. B., Southwick, S. M., & Pietrzak, R. H. (2021). Prevalence, risk and protective factors associated with suicidal ideation during the COVID-19 pandemic in U.S. military

- veterans with pre-existing psychiatric conditions. *Journal of Psychiatric Research*, 137, 351–359. <https://doi.org/10.1016/j.jpsychires.2021.03.021>.
- Nadkarni, A., Hasler, V., AhnAllen, C. G., Amonoo, H. L., Green, D. W., Levy-Carrick, N. C., & Mittal, L. (2020). Telehealth during COVID-19 – Does everyone have equal access? *American Journal of Psychiatry*, 177, 1093–1094. <https://doi.org/10.1176/appi.ajp.2020.20060867>.
- National Institute of Mental Health. (2020). The CoRonavIruS Health Impact Survey (CRISIS). Retrieved from National Institute of Mental Health Intramural Research Program Mood Spectrum Collaboration, Child Mind Institute of the NYS Nathan S. Kline Institute for Psychiatric Research. Retrieved from https://www.nlm.nih.gov/dr2/CRISIS_Adult_Self-Report_Baseline_Current_Form_V0.3.pdf.
- Nichter, B., Hill, M., Norman, S., Haller, M., & Pietrzak, R. H. (2020a). Associations of childhood abuse and combat exposure with suicidal ideation and suicide attempt in U.S. military veterans: A nationally representative study. *Journal of Affective Disorders*, 276, 1102–1108. <https://doi.org/10.1016/j.jad.2020.07.120>.
- Nichter, B., Hill, M., Norman, S., Haller, M., & Pietrzak, R. H. (2020b). Mental health treatment utilization among U.S. military veterans with suicidal ideation: Results from the National Health and Resilience in Veterans Study. *Journal of Psychiatric Research*, 130, 61–67. <https://doi.org/10.1016/j.jpsychires.2020.07.004>.
- Norman, G. R., Sloan, J. A., & Wyrwich, K. W. (2003). Interpretation of changes in health-related quality of life the remarkable universality of half a standard deviation. *Medical Care*, 41(5), 582–592. <https://doi.org/10.1097/00005650-200305000-00004>.
- Norris, F. H., Tracy, M., & Galea, S. (2009). Looking for resilience: Understanding the longitudinal trajectories of responses to stress. *Social Science and Medicine*, 68(12), 2190–2198. <https://doi.org/10.1016/j.socscimed.2009.03.043>.
- North, C. S., Tivis, L., McMillen, J. C., Pfefferbaum, B., Spitznagel, E. L., Cox, J., ... Smith, E. M. (2002). Psychiatric disorders in rescue workers after the Oklahoma City bombing. *American Journal of Psychiatry*, 159(5), 857–859. <https://doi.org/10.1176/appi.ajp.159.5.857>.
- Pietrzak, R. H., & Cook, J. M. (2013). Psychological resilience in older U. S. veterans: Results from the National Health and Resilience in Veterans Study. *Depression and Anxiety*, 30(5), 432–443. <https://doi.org/10.1002/da.22083>.
- Pietrzak, R. H., Tracy, M., Galea, S., Kilpatrick, D. G., Ruggiero, K. J., Hamblen, J. L., ... Norris, F. H. (2012). Resilience in the face of disaster: Prevalence and longitudinal course of mental disorders following Hurricane Ike. *PLoS ONE*, 7(6), 1–14. <https://doi.org/10.1371/journal.pone.0038964>.
- Prati, G., & Mancini, A. D. (2021). The psychological impact of COVID-19 pandemic lockdowns: A review and meta-analysis of longitudinal studies and natural experiments. *Psychological Medicine*, 51(2), 201–211. <https://doi.org/10.1017/S0033291721000015>.
- R Core Team. (2019). R Foundation for Statistical Computing. R: A language and environment for statistical computing.
- Ready, D. J., Thomas, K. R., Worley, V., Backscheider, A. G., Harvey, L. A. C., Baltzell, D., & Rothbaum, B. O. (2008). A field test of group based exposure therapy with 102 veterans with war-related posttraumatic stress disorder. *Journal of Traumatic Stress*, 21(2), 150–157. <https://doi.org/10.1002/jts.20326>.
- Saunders, J. B., Aasland, O. G., Babor, T. F., De La Fuente, J. R., & Grant, M. (1993). Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction*, 88(6), 791–804. <https://doi.org/10.1111/j.1360-0443.1993.tb02093.x>.
- Scheier, M. F., Carver, C. S., & Bridges, M. W. (1994). Distinguishing optimism from neuroticism (and trait anxiety, self-mastery, and self-esteem): A reevaluation of the Life Orientation Test. *Journal of Personality and Social Psychology*, 67(6), 1063–1078. <https://doi.org/10.1037/0022-3514.67.6.1063>.
- Schnurr, P. P., Friedman, M. J., Foy, D. W., Shea, M. T., Hsieh, F. Y., Lavori, P. W., ... Bernardy, N. C. (2003). Randomized trial of trauma-focused group therapy for posttraumatic stress disorder: Results from a Department of Veterans Affairs Cooperative Study. *Archives of General Psychiatry*, 60(5), 481–489. <https://doi.org/10.1001/archpsyc.60.5.481>.
- Schulenberg, S. E., Schnitzer, L. W., & Buchanan, E. M. (2011). The purpose in life test-short form: Development and psychometric support. *Journal of Happiness Studies*, 12(5), 861–876. <https://doi.org/10.1007/s10902-010-9231-9>.
- Seal, K. H., Cohen, G., Waldrop, A., Cohen, B. E., Maguen, S., & Ren, L. (2011). Substance use disorders in Iraq and Afghanistan veterans in VA healthcare, 2001–2010: Implications for screening, diagnosis and treatment. *Drug and Alcohol Dependence*, 116(1–3), 93–101. <https://doi.org/10.1016/j.drugalcdep.2010.11.027>.
- Sheehan, D. V. (2016). Mini international neuropsychiatric interview. English Version 7.0.2. For DSM-5. Copyright 1992–2016.
- Sherbourne, C. D., & Stewart, A. L. (1991). The MOS social support survey. *Social Science and Medicine*, 32(6), 705–714. [https://doi.org/10.1016/0277-9536\(91\)90150-B](https://doi.org/10.1016/0277-9536(91)90150-B).
- Shi, L., Lu, Z.-A., Que, J.-Y., Huang, X.-L., Liu, L., Ran, M.-S., ... Lu, L. (2020). Prevalence of and risk factors associated with mental health symptoms among the general population in China during the Coronavirus Disease 2019 Pandemic. *JAMA Network Open*, 3(7), e2014053–e2014053. <https://doi.org/10.1001/jamanetworkopen.2020.14053>.
- Smid, G. E., van Zuiden, M., Geuze, E., Kavelaars, A., Heijnen, C. J., & Vermetten, E. (2015). Cytokine production as a putative biological mechanism underlying stress sensitization in high combat exposed soldiers. *Psychoneuroendocrinology*, 51, 534–546. <https://doi.org/10.1016/j.psyneuen.2014.07.010>.
- Ssentongo, P., Ssentongo, A. E., Heilbrunn, E. S., Ba, D. M., & Chinchilli, V. M. (2020). Association of cardiovascular disease and 10 other pre-existing comorbidities with COVID-19 mortality: A systematic review and meta-analysis. *PLoS ONE*, 15(8), e0238215. <https://doi.org/10.1371/journal.pone.0238215>.
- Tiet, Q. Q., Leyva, Y. E., Moos, R. H., Frayne, S. M., Osterberg, L., & Smith, B. (2015). Screen of drug Use: Diagnostic accuracy of a new brief tool for primary care. *JAMA Internal Medicine*, 175(8), 1371–1377. <https://doi.org/10.1001/jamainternmed.2015.2438>.
- Tonidandel, S., & LeBreton, J. M. (2011). Relative importance analysis: A useful supplement to regression analysis. *Journal of Business and Psychology*, 26(1), 1–9. <https://doi.org/10.1007/s10869-010-9204-3>.
- Tuerk, P. W., Keller, S. M., & Acierno, R. (2018). Treatment for anxiety and depression via clinical videoconferencing: Evidence base and barriers to expanded access in practice. *FOCUS*, 16(4), 363–369. <https://doi.org/10.1176/appi.focus.20180027>.
- United States Department of Veterans Affairs Office of Mental Health and Suicide Prevention. (2019). 2019 National Veteran Suicide Prevention Annual Report. Retrieved from https://www.mentalhealth.va.gov/docs/data-sheets/2019/2019_National_Veteran_Suicide_Prevention_Annual_Report_508.pdf.
- Vespa, J. E. (2020). Those who served: America's veterans from World War II to the War on Terror. Womenveteransalliance.Org. Retrieved from www.census.gov/acs.
- Wang, Y., Kala, M. P., & Jafar, T. H. (2020). Factors associated with psychological distress during the coronavirus disease 2019 (COVID- 19) pandemic on the predominantly general population: A systematic review and meta-analysis. *PLoS ONE*, 15(12), e0244630. <https://doi.org/10.1371/journal.pone.0244630>.
- Weathers, F. J., Blake, D. D., Schnurr, P. P., Kaloupek, D. G., Marx, B. P., & Keane, T. M. (2013a). The life events checklist for DSM-5 (LEC-5). National Center for PTSD. Retrieved from <https://doi.org/10.1177/1073191104269954>.
- Weathers, F. W., Litz, B. T., Keane, T. M., Palmieri, P. A., Marx, B. P., & Schnurr, P. P. (2013b). The PTSD checklist for DSM-5 (PCL-5). National Center for PTSD. Retrieved from <https://doi.org/10.1037/t02622-000>.
- Zheng, J., Morstead, T., Sin, N., Klaiber, P., Umberson, D., Kamble, S., ... DeLongis, A. (2021). Psychological distress in North America during COVID-19: The role of pandemic-related stressors. *Social Science and Medicine*, 270, 1–8. <https://doi.org/10.1016/j.socscimed.2021.113687>.
- Zulman, D. M., Wong, E. P., Slightam, C., Gregory, A., Jacobs, J. C., Kimerling, R., ... Heyworth, L. (2019). Making connections: Nationwide implementation of video telehealth tablets to address access barriers in veterans. *JAMIA Open*, 2(3), 323–329. <https://doi.org/10.1093/jamiaopen/ooz024>.