

UC Berkeley

UC Berkeley Previously Published Works

Title

From Theory to Practice: Leveraging Chemical Principles To Improve the Performance of Peroxydisulfate-Based In Situ Chemical Oxidation of Organic Contaminants

Permalink

<https://escholarship.org/uc/item/3682d2b3>

Journal

Environmental Science and Technology, 58(1)

ISSN

0013-936X

Authors

McGachy, Lenka

Sedlak, David L

Publication Date

2024-01-09

DOI

10.1021/acs.est.3c07409

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

From Theory to Practice: Leveraging Chemical Principles To Improve the Performance of Peroxydisulfate-Based In Situ Chemical Oxidation of Organic Contaminants

Lenka McGachy* and David L. Sedlak*



Cite This: *Environ. Sci. Technol.* 2024, 58, 17–32



Read Online


ACCESS |

Metrics & More

Article Recommendations

ABSTRACT: In situ chemical oxidation (ISCO) using peroxydisulfate has become more popular in the remediation of soils and shallow groundwater contaminated with organic chemicals. Researchers have studied the chemistry of peroxydisulfate and the oxidative species produced upon its decomposition (i.e., sulfate radical and hydroxyl radical) for over five decades, describing reaction kinetics, mechanisms, and product formation in great detail. However, if this information is to be useful to practitioners seeking to optimize the use of peroxydisulfate in the remediation of hazardous waste sites, the relevant conditions of high oxidant concentrations and the presence of minerals and solutes that affect radical chain reactions must be considered. The objectives of this Review are to provide insights into the chemistry of peroxydisulfate-based ISCO that can enable more efficient operation of these systems and to identify research needed to improve understanding of system performance. By gaining a deeper understanding of the underlying chemistry of these complex systems, it may be possible to improve the design and operation of peroxydisulfate-based ISCO remediation systems.

KEYWORDS: *in situ chemical oxidation, peroxydisulfate, PDS, ISCO, organic contaminants*



1. INTRODUCTION

Starting in the mid-1970s, the discovery of widespread soil and groundwater contamination led to the investment of large sums of money in site remediation. By the early 1990s, this often resulted in the removal of concentrated contaminant sources using dig-and-haul and pump-and-treatment technologies.¹ The high cost of these remedial approaches and the desire to avoid long-term liability led to the development of active strategies, like biostimulation, permeable reactive barriers, and soil vapor extraction. Although these methods were effective, they often proved to be expensive or incapable of achieving the required cleanup goals. For this reason, more aggressive in situ remediation technologies, particularly those targeting source zones, started to be implemented in the early 2000s.²

Among the various new remediation approaches, in situ chemical oxidation (ISCO) became popular as a means of remediating soils and shallow aquifers contaminated with halogenated solvents,^{3,4} benzene, toluene, ethylbenzene, and xylenes (BTEX),^{5,6} petroleum hydrocarbons,⁷ polynuclear aromatic hydrocarbons (PAHs),^{8,9} pesticides,^{10–12} chlorobenzenes,^{13,14} and polychlorinated biphenyls (PCBs).¹⁵

ISCO typically involves the introduction of a solution containing a relatively high concentration (i.e., 0.05–1.25 M) of a strong oxidant, such as permanganate (MnO_4^-), hydrogen

peroxide (H_2O_2) or a peroxydisulfate salt, into an injection well. This is often accompanied by recovery of the injection fluid at one or more recovery wells.^{16–18} Fluids extracted from recovery wells are pumped into a holding tank and amended with a concentrated stock solution of the oxidant. Sufficient residence time is provided in the aquifer to ensure that only treated water is reinjected into the aquifer.¹⁹

The rate at which the oxidant decomposes in the subsurface is critical to the success of ISCO because it determines the delivery of reactive species to the target zone. For example, reactions of H_2O_2 with soil minerals and organic matter tend to take place over a period of minutes to hours in the subsurface,^{20,21} resulting in a limited transport distance for injected oxidants under many conditions. To slow these processes, stabilizers (e.g., KH_2PO_4 or sodium phytate) are used to keep H_2O_2 from decomposing too quickly.^{22,23} In addition, under certain circumstances (e.g., high

Received: September 8, 2023

Revised: November 28, 2023

Accepted: November 29, 2023

Published: December 18, 2023



concentrations of oxidant) ISCO can enhance the transfer of contaminants from soil to groundwater by oxidizing organic coatings on mineral surfaces and increasing groundwater temperature (H_2O_2 decomposition in soils is an exothermic reaction).^{24,25} Furthermore, the decomposition of H_2O_2 can produce O_2 bubbles ($2\text{H}_2\text{O}_2 \rightarrow 2\text{H}_2\text{O} + \text{O}_2$) that can increase the rate of transfer of volatile contaminants from the saturated zone to the vadose zone without oxidizing them.²⁶ In cases where this process releases high concentrations of contaminants, vapor recovery systems are used to minimize contaminant release.²⁶ Logistical consideration associated with well construction and the transfer of contaminants to the vadose zone often limits the use of ISCO to soils and shallow groundwater where injection and recovery is possible (typically depths less than 25 m²⁵).

Among the different oxidants employed for ISCO, solutions of peroxydisulfate salts (e.g., $\text{Na}_2\text{S}_2\text{O}_8$, $\text{K}_2\text{S}_2\text{O}_8$) became popular due to their relatively low cost, relative stability, and water solubility. After injection, peroxydisulfate anion ($\text{S}_2\text{O}_8^{2-}$) undergoes reactions that result in the formation of nonselective reactive oxidant species (ROS), primarily sulfate radicals ($\text{SO}_4^{\bullet-}$) that can be further converted into hydroxyl radicals ($\cdot\text{OH}$). Except for fully halogenated compounds (e.g., CCl_4 , hexachlorobenzene) and a few other contaminants, $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$ can oxidize most contaminants of concern.^{27,28}

$\text{S}_2\text{O}_8^{2-}$ -based ISCO has been a topic of considerable interest among researchers.²⁹ Numerous empirical studies have been conducted by remediation engineers (i.e., bench-scale treatability tests) as part of the process of determining appropriate $\text{S}_2\text{O}_8^{2-}$ doses and the need for additives (e.g., pH adjustment). Academic researchers also have studied the mechanisms of $\text{SO}_4^{\bullet-}$ -radical-based reactions to gain insight into kinetics and transformation products. However, academic research is rarely conducted under the conditions that are relevant to ISCO (i.e., high concentrations of oxidant, presence of high concentrations of mineral surfaces, and solutes that can affect radical chain reactions). The objective of this Review is to use the results of available studies and knowledge of the actual conditions employed in ISCO to gain insight into the chemistry of $\text{S}_2\text{O}_8^{2-}$ -based ISCO that can be used to operate these systems more efficiently.

2. PEROXYDISULFATE CHEMISTRY

Peroxydisulfate salts dissociate in water to form $\text{S}_2\text{O}_8^{2-}$, a species that is relatively stable in aqueous solutions. In the absence of other solutes, $\text{S}_2\text{O}_8^{2-}$ slowly oxidizes water through a series of radical reactions with the net stoichiometry shown in eq 1. In the absence of catalysts of other solutes, the half-life for $\text{S}_2\text{O}_8^{2-}$ at circumneutral pH values is around 14 months at 25 °C. $\text{S}_2\text{O}_8^{2-}$ -thermolysis (eq 2) exhibits a strong temperature dependence (i.e., the activation energy of the reaction is 128.9–140.2 kJ mol⁻¹³⁰). As a result, the rate of eq 2 can be accelerated by heating; at 50 °C the half-life of $\text{S}_2\text{O}_8^{2-}$ decreases to about 5 days. In weakly basic and neutral solutions, eq 2 is the main process contributing to the initiation of $\text{S}_2\text{O}_8^{2-}$ decomposition. Under acidic and strongly basic conditions, $\text{S}_2\text{O}_8^{2-}$ decomposes by two simultaneous reactions, one uncatalyzed (eq 2) and the other catalyzed by hydrogen (eq 3) and hydroxide (eq 8) ion, respectively.^{31,32} However, strong acidic (pH < 3) and basic (pH > 12) conditions are required to contribute significantly to $\text{S}_2\text{O}_8^{2-}$ decomposition (Figure 1).









Figure 1. Calculated kinetics of $\text{S}_2\text{O}_8^{2-}$ decomposition in aqueous solution as a function of the pH and temperature. The calculations are based on activation energies and pre-exponential factors of partial reactions (eqs 2, 3 and 8).³⁶

Under many conditions, the first step in the decomposition of $\text{S}_2\text{O}_8^{2-}$ is thermolysis to produce two $\text{SO}_4^{\bullet-}$ ions (eq 2). In the absence of high concentrations of contaminants, most of the $\text{SO}_4^{\bullet-}$ reacts with water (i.e., eq 4), hydroxide (eq 5) or $\text{S}_2\text{O}_8^{2-}$ (eq 6). The combined reactions of thermolysis (eq 2) and radical scavenging (eqs 4–6 and reactions with organic contaminants) result in low, steady-state $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$ concentrations during $\text{S}_2\text{O}_8^{2-}$ decomposition. The main factors responsible for the variations in $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$ concentrations are usually the concentration of $\text{S}_2\text{O}_8^{2-}$ and pH. Under conditions typical of ISCO, characterized by neutral or acidic pH levels, the majority of $\text{SO}_4^{\bullet-}$ does not undergo conversion into $\cdot\text{OH}$,³³ making $\text{SO}_4^{\bullet-}$ the predominant radical.



Organic contaminants typically do not become dominant sinks for the radicals until concentrations become high. The rate of $\text{S}_2\text{O}_8^{2-}$ decomposition may be altered by the presence of organic contaminants for several reasons. First, some compounds directly react with $\text{S}_2\text{O}_8^{2-}$, usually through 2-electron oxidation reactions (eq 7) to produce sulfate (SO_4^{2-}) and an oxidized organic compound. These reactions tend to be slow at temperatures typical of ISCO systems, but for electron-rich compounds, partial oxidation can occur through direct reactions. For example, the second-order rate constants for oxidation of phenols and anilines by $\text{S}_2\text{O}_8^{2-}$ at 30 °C typically range from 10^{-1} to 10^{-4} M⁻¹ s⁻¹.³⁴ In the presence of 0.1 M $\text{S}_2\text{O}_8^{2-}$ (i.e., a concentration typical of ISCO systems), the half-life of these compounds would range from approximately 1 min to 19 h. In addition, the strong temperature dependence of these reactions assures that this process is considerably faster at the higher temperatures used during heat activation of $\text{S}_2\text{O}_8^{2-}$.³⁵

The second reason that the rate of $\text{S}_2\text{O}_8^{2-}$ decomposition often changes in the presence of organic contaminants is related to reactions of organic compounds with $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$. Second-order rate constants of $\text{SO}_4^{\bullet-}$ reactions with organic contaminants range from 10^7 to 10^{10} M⁻¹ s⁻¹,³⁷ and $\cdot\text{OH}$ reacts with organic contaminants with second-order rate constants that range from 10^8 to 10^{11} M⁻¹ s⁻¹.³⁸ These reactions initiate a series of radical reactions that can be more important to

peroxydisulfate decomposition than the reaction pathways that dominate in the absence of solutes.



Although $\text{S}_2\text{O}_8^{2-}$ can be converted into $\text{SO}_4^{\cdot-}$ by thermolysis (eq 2), the rate of radical production is usually too slow to be useful for remediation under ambient conditions encountered in groundwater. As described in the following sections, technologies are available for heating the aquifer during ISCO. As an alternative, the rate at which $\text{S}_2\text{O}_8^{2-}$ is converted into $\text{SO}_4^{\cdot-}$ can be enhanced by raising the pH of the groundwater through the addition of a strong base (e.g., NaOH). The initial step of the alkaline activation process is the hydroxide-initiated fission of $\text{S}_2\text{O}_8^{2-}$ to form peroxymonosulfate (SO_5^{2-}) and SO_4^{2-} (eq 8). A similar fission of the S–O bond in SO_5^{2-} results in the formation of a hydroperoxyl anion (HO_2^-) (eq 8), followed by nucleophilic attack of HO_2^- on the peroxide (O–O) bond in $\text{S}_2\text{O}_8^{2-}$, resulting in the formation of $\text{SO}_4^{\cdot-}$ and superoxide ($\text{O}_2^{\cdot-}$) radical (eq 10).³⁹ Simultaneously, SO_5^{2-} decomposes to form SO_4^{2-} and O_2 (eq 11).⁴⁰



For the alkaline hydrolysis pathway to enhance the rate of conversion of $\text{S}_2\text{O}_8^{2-}$ to $\text{SO}_4^{\cdot-}$ and $\cdot\text{OH}$ (via eq 5) to rates that are useful for remediation, pH values greater than 12 must be achieved in the groundwater. It is challenging to maintain such high pH values in the subsurface due to the buffering by carbonates and other soil minerals, as well as the release of H^+ as $\text{S}_2\text{O}_8^{2-}$ is converted into SO_4^{2-} by various pathways (e.g., eq 1). As a result, alkaline activation typically requires the addition of large volumes of concentrated base into the subsurface, resulting in complicated logistics and possible health hazards to workers when the injection fluids are handled.^{41,42}

The addition of dissolved transition metals (e.g., Fe^{2+}) has been researched as a means of converting $\text{S}_2\text{O}_8^{2-}$ into $\text{SO}_4^{\cdot-}$. However, the use of Fe^{2+} for $\text{S}_2\text{O}_8^{2-}$ activation within ISCO appears to be impractical. Fe^{2+} reacts quickly with $\text{S}_2\text{O}_8^{2-}$ through a reaction analogous to the Fenton reaction, but unlike the Fenton-like processes that enable Fe to catalyze radical formation by cycling between the +II and +III oxidation states, $\text{S}_2\text{O}_8^{2-}$ and the radicals that it generates under neutral and low pH conditions do not result in a catalytic reaction. Thus, Fe^{2+} activation of $\text{S}_2\text{O}_8^{2-}$ would require the addition of relatively large quantities of iron. In addition, injected Fe^{2+} is converted into Fe^{3+} , which is insoluble at neutral pH.⁴³ It may be possible to avoid precipitation by adding a chelated form of Fe(II) into the aquifer.^{44,45} Nevertheless, chelating agents that are commonly used to prevent Fe(III) precipitation (e.g., citric acid, EDTA) are not Fe-specific and may complex with and mobilize other metals from the soil matrix.⁴⁶ Other methods for $\text{S}_2\text{O}_8^{2-}$ activation also appear to be impractical under conditions

relevant to ISCO. Carbon-based materials⁴⁷ and copper and cobalt oxides⁴⁸ can catalyze the conversion of $\text{S}_2\text{O}_8^{2-}$ into $\text{SO}_4^{\cdot-}$, but they are relatively expensive and cannot easily be adapted to deliver catalysts to porous media.^{49,50}

The homolytic cleavage of the O–O bond that produces two $\text{SO}_4^{\cdot-}$ (eq 2) can be accelerated by increasing the temperature in the subsurface.⁵¹ Although energy demanding, heat activation is considered one of the most promising methods for employing $\text{S}_2\text{O}_8^{2-}$ for ISCO.^{52–54} One benefit of thermal activation is that it allows engineers to determine the location in the aquifer where $\text{S}_2\text{O}_8^{2-}$ activation takes place.^{3,55} In addition, it avoids the need to handle additional chemicals (e.g., NaOH) and minimizes concerns about their impact on the subsurface environment.⁵⁵

To thermally activate $\text{S}_2\text{O}_8^{2-}$ in the field, various techniques have been employed, including in situ thermal treatment using steam, resistive heating and preceding $\text{S}_2\text{O}_8^{2-}$ addition by application of H_2O_2 , which generates heat as it is converted into O_2 and $\cdot\text{OH}$.^{18,56,57} Although these approaches have some advantages over base addition, they face their own set of challenges. For example, it is difficult to achieve uniform steam distribution in the subsurface, and steam generation can be complicated if the water used requires the addition of softeners to prevent the buildup of calcium, lime and rust in the steam lines.^{18,58} Electrical resistance heating provides better heat distribution in the subsurface, but it requires installation of electrodes, which can greatly increase project costs.^{59,60,53} Coupling $\text{S}_2\text{O}_8^{2-}$ application with H_2O_2 ⁵⁶ addition tends to be the least expensive and most practical approach for heating the subsurface because it uses much of the same infrastructure (i.e., injection well, piping) that is employed for $\text{S}_2\text{O}_8^{2-}$ -based ISCO. To thermally activate $\text{S}_2\text{O}_8^{2-}$, H_2O_2 is injected before $\text{S}_2\text{O}_8^{2-}$, and the recommended dosage is 1 mol of H_2O_2 per 1–2 mol of $\text{S}_2\text{O}_8^{2-}$ (the higher the H_2O_2 dosage, the shorter is the $\text{S}_2\text{O}_8^{2-}$ lifetime).⁶¹

The critical parameter of a thermally activated PDS-based ISCO is temperature. Most laboratory studies of $\text{S}_2\text{O}_8^{2-}$ use for oxidation of organic contaminants are conducted at temperatures of 30–90 °C.^{62–66} Higher temperatures result in rapid production of ROS as well as fast rates of contaminant oxidation. However, in the context of ISCO applications, lower temperatures (30–40 °C) may be more suitable because heating an aquifer to temperatures above 40 °C is expensive and rapid activation also limits the distance that $\text{S}_2\text{O}_8^{2-}$ can travel.^{52–54} As shown in Figure 1, the lifetime of peroxydisulfate is relatively short at elevated temperatures and can be even shorter in the presence of soil, which limits the persistence of PDS in the subsurface and thus the distance over which thermally activated PDS can be effective. Moreover, reducing the temperature of treatment decreases energy consumption.

3. DESIGN OF $\text{S}_2\text{O}_8^{2-}$ -BASED ISCO REMEDIATION SYSTEMS

Key factors determining the cost, complexity, and logistics of operating $\text{S}_2\text{O}_8^{2-}$ -based ISCO systems include oxidant volume and initial $\text{S}_2\text{O}_8^{2-}$ concentration, well spacing, depth and location, injection rates, and the duration, timing, and number of injection events. The design of ISCO systems is determined by the nature of the contamination, cleanup goals and uncertainties in subsurface conditions, including soil heterogeneity and properties affecting mass transfer rates.²⁵ The collection of extensive data on subsurface conditions can reduce some of these uncertainties. Before implementing $\text{S}_2\text{O}_8^{2-}$ -based ISCO, it is important to assess the potential for treatment to affect




Figure 2. Conceptual illustration of (A) direct injection process and (B) recirculation system.

downgradient water sources (e.g., by mobilizing toxic metals or other contaminants). This typically involves land assessment and bench treatability or pilot-scale treatability tests. These tests can offer insights into the risks of contaminant mobilization for the specific site, as well as design information regarding the required amount of $S_2O_8^{2-}$ and the rate at which the oxidant will be consumed in the subsurface.

The success of many ISCO projects depends on the ability of the injection system to distribute the oxidant into the subsurface in a uniform manner. Two approaches are typically used to deliver the $S_2O_8^{2-}$ solution into the subsurface: direct injection (Figure 2A) and recirculation (Figure 2B).⁵⁷ Direct injection involves mixing of solid $S_2O_8^{2-}$ with a specified volume of water from an external source above ground followed by injection into the aquifer without any effort to recover or recirculate the oxidant (Figure 2A). Because $S_2O_8^{2-}$ is injected at high concentrations, the possibility of acidity, metals mobility, and $S_2O_8^{2-}$ and/or SO_4^{2-} migration to nearby receptors should be assessed. Nonetheless, it is well documented that biotic and abiotic attenuation mechanisms reduce the impact of these processes to levels considered safe by regulatory agencies in a period of 1 to 4 months.^{18,26}

The direct injection process is mostly performed with temporary push points (Figure 2A); however, permanent injection wells can also be used. The typical direct push method involves using a machine with a hydraulic hammer that pushes small-diameter metal rods with an injection tool to the targeted treatment depth. This process can be performed using either a top-down and/or bottom-up injection sequence. As direct push technologies are mounted on a platform, the access of large or small vehicles to the injection area and the injection depth (typical maximum depth is 25 m) must be considered.

For recirculation systems, an $S_2O_8^{2-}$ -containing salt is dissolved in groundwater extracted from the contaminated zone. After mixing in a holding tank which also serves as an above-ground treatment unit, the treated groundwater is reinjected. The recirculation systems can be designed in several different configurations. Typically, as shown in Figure 2B, injection wells are installed in the upgradient portion of the

plume, and recovery wells are located downgradient of the plume. This strategy is particularly useful for treatment of contaminants that are not strongly sorbed to soil and aquifer materials. The mixing that occurs between $S_2O_8^{2-}$ and the groundwater in the above-ground treatment unit ensures that the contaminants are exposed to oxidants. In addition, recirculation enhances the distribution of an oxidant in the subsurface, provides hydraulic control in the treatment zone (e.g., develop a concentration gradient, containment), and ensures that any excess oxidants or products of oxidation reactions (e.g., acid, SO_4^{2-} , metals, and oxidation byproducts) are captured. Above-ground infrastructure consists mainly of metering pumps, prefiltration elements to remove sediments from extracted groundwater, a holding tank equipped with a mixer, a post-filtration element to remove solids from the injected fluid, system manifolds to distribute the flow and allow measurement and adjustment of injection/extraction pressures and flow rates, and hoses connecting the facility to the injection and recovery wells. When volatile contaminants (e.g., vinyl chloride) are treated, the tank is typically equipped with a blower to withdraw air from the headspace of the tank and discharge it through the activated carbon unit into the atmosphere. In general, water is reinjected without further additional treatment (e.g., acid neutralization), but in some situations (e.g., injection upgradient of a drinking water aquifer), additional treatment may be employed.

Although recirculation systems have numerous advantages over direct injection systems, the process often consumes more oxidant. It is also more complicated to operate and requires expensive above-ground infrastructure (e.g., hosing, tanks, electricity, and filtering of solids), making it less cost-effective just for small sites.¹⁶ Typically, higher volumes (e.g., 40–85% of a pore volume of the entire treatment area) and lower concentrations (0.05–0.2 M) of $S_2O_8^{2-}$ are used when injecting via recirculation, whereas lower volumes (e.g., ~1% of a pore volume of the entire treatment area) and higher $S_2O_8^{2-}$ concentrations (0.5–1.25 M) are applied when using direct injection.¹⁸ The number of pore volumes that will be injected or

recirculated depends on site-specific factors such as oxidant dispersion/diffusion.⁵⁷

Another factor affecting the cost and design of $S_2O_8^{2-}$ -based ISCO systems is the duration of deployment. The direct injection typically takes place over a period of 4 h to 2 days with $S_2O_8^{2-}$ lasting in the ground for between 30 and 90 days, depending upon the volume and injection rates, while recirculation is typically completed over a period of 4–8 weeks.^{18,67} Due to the difficulty associated with achieving contact between $S_2O_8^{2-}$ and contaminants adsorbed onto clay or organic-rich particles, multiple injection events (redosing) are often necessary to achieve cleanup goals. The number of redosing events is typically determined by data collected after the injection event.²⁵ ISCO is considered one of the most rapid in situ remediation technologies, with times required to remediate a site on average 1–3 years.⁶⁸ In general, remediation of simple sites (e.g., sandy aquifers, small sites without strongly adsorbed contaminants) is easier compared to sites with low permeability formations or sites with extensive contamination.⁶⁷

Although solid $S_2O_8^{2-}$ is a very stable material, excessive heat or moisture can result in decomposition of the stored material (e.g., material at the bottom of a pallet or in the center of a drum). As heat generated may stimulate self-accelerating decomposition, $S_2O_8^{2-}$ should be stored in a cool, clean, dry place away from point sources of heat (e.g., radiant heaters) and moisture (e.g., rain). When working with caustic chemicals (e.g., NaOH) as an activator, it is imperative to apply increased precautionary measures. Additionally, materials such as soft metals (e.g., copper) can be degraded by concentrated $S_2O_8^{2-}$ solutions.⁵⁷ Therefore, the materials used for the delivery system should be compatible with long-term PDS exposure (e.g., stainless steel, high-density polyethylene, and polyvinyl chloride). Similarly, at sites where large volumes of $S_2O_8^{2-}$ are injected, daylighting (surfacing of groundwater) can occur. Thus, a spill containment plan also should be developed as part of the remedial action work plan.⁶⁹

The cost of $S_2O_8^{2-}$ -based ISCO is highly site-specific, as it is influenced by the various parameters mentioned above. The total cost of remediation can be divided into two components. The first component is the capital cost, which is a fixed project expense and includes costs associated with site preparation (e.g., utility lines and concrete pad installation), mobilization/demobilization (including transportation to the site, storage, fabrication, assembly, setup, and dismantling), equipment acquisition, and labor (for transportation and setup on site if not already included in mobilization). The second component is operating cost, which is variable and includes mainly expenditure on equipment purchase/rental (e.g., storage tanks and pumps), chemicals/reagents, energy, waste disposal and labor required to operate the system. In general, capital costs increase with site size and range from approximately \$170,000 for smaller sites (with a target aquifer volume of approximately 900 m³) to approximately \$1,300,000 for larger sites (with a target aquifer volume of approximately 34,700 m³).¹⁸ Overall, these costs are competitive with alternative remediation techniques and offer the benefit of being completed more quickly (i.e., successful ISCO projects can be completed in several months, whereas pump-and-treat and bioremediation often require several years to decades to complete).

4. CHALLENGES AFFECTING THE PERFORMANCE OF $S_2O_8^{2-}$ -BASED ISCO SYSTEMS

Although laboratory studies show that $S_2O_8^{2-}$ can be used to oxidize contaminants under well-controlled conditions, the effectiveness of this technology under field conditions is often limited by several practical issues. By understanding the practical limitations of using $S_2O_8^{2-}$ -based ISCO, it may be possible for researchers to develop more effective ways to predict when this approach will work, increase its effectiveness, and develop approaches to minimize the unintended consequences of its use.

4.1. Delivering $S_2O_8^{2-}$ to Low Permeability Zones.

Irrespective of the oxidant employed, the success of all ISCOs often depends upon subsurface geology. Due to the heterogeneity in the subsurface and the presence of layers or lenses of low-permeable minerals (e.g., clay), dead-end pores or stagnant zones,⁷⁰ oxidants may not reach contaminated zones that require treatment. Under these conditions, oxidants will preferentially flow through areas of least resistance (e.g., highly permeable sand layers), resulting in incomplete oxidation of contaminants.⁷¹ This is especially problematic for hydrophobic contaminants that tend to associate with particulate organic matter and clays,⁷² especially under conditions in which contaminants have been present for several decades.⁷³ To ensure that oxidants have the opportunity to come into contact with contaminants in these difficult-to-reach zones, efforts are needed to maximize the longevity of the oxidant in the subsurface.^{27,74}

Failure of ISCO systems to deliver oxidants to contaminated zones often results in the rebound effect, as contaminants gradually diffuse out of clay- and organic-matter-rich zones after oxidant application ends. In response, researchers have developed techniques to deliver $S_2O_8^{2-}$ -containing fluids to low-permeability zones using electrokinetic techniques.^{55,54,75,76} Although these techniques have the potential to help with the delivery of $S_2O_8^{2-}$ to low-permeability zones, their widespread application in these approaches has not made substantial progress.

4.2. Loss of $S_2O_8^{2-}$ through Reactions with Aquifer Solids.

Researchers who have studied the fate of H_2O_2 in ISCO systems have found that transition metal oxides and soil organic matter can cause rapid loss of oxidant.^{77,78} Consequently, they hypothesized that the same phenomenon could lead to the loss of $S_2O_8^{2-}$ in ISCO systems. However, laboratory studies and field observations indicate that $S_2O_8^{2-}$ is much less susceptible to loss by reactions with aquifer solids. Nevertheless, there are situations in which oxidant loss occurs through reactions with geologic materials.

$S_2O_8^{2-}$ is very stable in pure sand, having a half-life of more than 1 year.^{27,81} However, as shown in Figure 3, the rate of $S_2O_8^{2-}$ decomposition accelerates in the presence of more reactive minerals and surfaces. Half-lives of $S_2O_8^{2-}$ decrease to less than a year when Fe(III) and Mn(IV) oxides comprise more than about 1% of the aquifer solids.^{27,80–82} In addition, the loss of $S_2O_8^{2-}$ increases significantly when the oxidant comes into contact with pure minerals (e.g., ferrihydrite, pyrolusite).²⁷ Metal oxides rarely account for more than 10% mass of aquifer solids on a mass basis. Therefore, the rates of reactivity of $S_2O_8^{2-}$ with pure minerals exaggerate the importance of this loss mechanism.

Similarly, if the organic matter in the soil is more than about 1% of the mass of aquifer solids, the decomposition rate of $S_2O_8^{2-}$ also increases (Figure 3), with half-lives dropping to less




Figure 3. Observed first-order kinetic rate constants of $S_2O_8^{2-}$ decomposition in different soils and minerals²⁷ under different thermal conditions and experimental designs (batch, column, push–pull) compared with calculated kinetics of $S_2O_8^{2-}$ decay in aqueous solution at pH 5.^{36,27,36,79–81} OSR is oxidant-to-soil mass ratio.

than ~30 days.⁸¹ However, the fraction of soil organic matter in aquifer solids tends to be much less than 1%, typically <0.01%.^{83–85} Under these conditions, the loss of $S_2O_8^{2-}$ by direct reactions with organic matter is likely negligible.

In addition, increasing the $S_2O_8^{2-}$ concentration from low concentrations typically used in laboratory experiments (e.g., 0.05 M⁸¹) to the high concentrations employed in the field (e.g., 0.2 M) results in an increase of half-life by more than an order of magnitude⁷⁹ (Figure 3, push–pull).

4.3. Biogeochemical Impacts and Mobilization of Metals of $S_2O_8^{2-}$ Injection. Application of $S_2O_8^{2-}$ -based ISCO alters the geochemistry of the subsurface, mainly due to the production of acid and the release of sulfate.³² Although large quantities of acid are generated as the injection fluid decomposes, the pH of groundwater after treatment typically ranges from 2 to 5 due to the neutralization of protons by minerals (e.g., $CaCO_{3(s)}$) associated with the aquifer material.^{18,26} Furthermore, the neutralization of acidity is not instantaneous; pH values typically increase over several days after treatment ends. In well-buffered systems, pH values eventually return to preinjection pH values. However, at sites with a paucity of minerals capable of neutralizing persistent acidity, pH values below 2 have been observed.¹⁸

The drop in pH often results in the release of toxic trace elements and/or the formation of new mineral species. Elevated concentrations of trace elements such as arsenic, chromium and vanadium have been reported in treatment zone wells where $S_2O_8^{2-}$ -ISCO was employed even after pH values returned to preinjection levels.¹⁸ This has been a concern with respect to chromium because the relatively nontoxic Cr(III) can be converted into its carcinogenic form, Cr(VI), during the oxidation process.⁸⁶


Acid produced during the decomposition of $S_2O_8^{2-}$ can also lead to the formation of minerals that alter the permeability of the subsurface. For example, Ca(II) released when $CaCO_{3(s)}$ dissolves under acidic conditions can react with SO_4^{2-} to produce gypsum ($CaSO_4 \cdot 2H_2O$), which is relatively insoluble.⁸⁷ Under acidic conditions, Fe(III) and Mn(IV) oxides also will dissolve. Because these minerals often play a role in maintaining soil structure and can precipitate when the acidity is neutralized, the permeability of the subsurface can change after $S_2O_8^{2-}$ -ISCO.

In addition to producing acidic conditions, the high concentrations of SO_4^{2-} and other dissolved ions (e.g., Na^+ , Ca^{2+}) produced during the decomposition of $S_2O_8^{2-}$ and subsequent acid-catalyzed dissolution processes can degrade water quality. $S_2O_8^{2-}$ -based injection fluids typically contain over 1000 mg/L of SO_4^{2-} , which is considerably higher than the secondary drinking water standard for SO_4^{2-} of 250 mg/L. In addition, under anaerobic conditions SO_4^{2-} can be reduced to sulfide (HS^-), compromise water quality and result in the precipitation of sulfide minerals.

Changes in pH and redox potential induced by the injection of $S_2O_8^{2-}$ -based fluids can also affect microbial communities. Microorganisms that are sensitive to oxidative treatments, including anaerobic organisms such as organohalide-respiring bacteria (e.g., *Dehalococcoides mccartyi*, *Dehalobacter*, *Geobacter*, and *Desulfitobacterium*), will experience a decrease in abundance and activity.⁸⁸ In contrast, the population and activity of microbes that are less sensitive to the alteration in redox potential (e.g., facultative anaerobes) might remain unchanged or even exhibit positive responses.^{89–91} For example, it has been demonstrated that acidification (pH < 3) and an increase in redox potential (>500 mV) associated with $S_2O_8^{2-}$ treatment disrupt the microbial community, with full recovery of anaerobic bacteria such as *Dehalococcoides mccartyi* taking longer than half a year.⁹⁰ The recovery is likely due to the recolonization of the previously oxidized area by organisms moving in groundwater from the upgradient direction or the reintroduction of microorganisms into a treated area, potentially originating from blind pockets or low-permeability zones.^{92,93} Additionally, the observed increase in biomass following the application of $S_2O_8^{2-}$ oxidant could be linked to the increased presence of bioavailable oxidation byproducts.⁹⁴

4.4. Effect of Sorption on Contaminant Transformation Rates during $S_2O_8^{2-}$ -Based ISCO. Observations based upon laboratory studies and field applications indicate that $S_2O_8^{2-}$ -based ISCO is less effective in the treatment of organic contaminants that are strongly sorbed to solids.^{95,96} The inefficiency of the oxidation processes is related to the fact that most $SO_4^{\bullet-}$ or OH^{\bullet} are produced in solution (e.g., when thermal activation is employed) or at the mineral–water interfaces (e.g., when $S_2O_8^{2-}$ is activated by processes taking place on mineral surfaces). Particle-associated organic contaminants tend to exhibit low reactivity with aqueous $SO_4^{\bullet-}$ or OH^{\bullet} radicals because the oxidants are consumed by other

Table 1. Important Reactions Involving $\text{SO}_4^{\bullet-}$ and Dissolved Species Typically Encountered during ISCO

Reaction	Second order reaction rate constant [$\text{M}^{-1}\text{s}^{-1}$]	No.	Ref.
$\text{SO}_4^- + \text{Cl}^- \leftrightarrow \text{SO}_4^{2-} + \text{Cl}^\cdot$	2.7×10^{8a} / 2.5×10^{8b}	Eq. 12	110, 119
$\text{Cl}^\cdot + \text{Cl}^- \leftrightarrow \text{Cl}_2^{\cdot-}$	8.5×10^{9a} / 6.0×10^{4b}	Eq. 13	120
$\text{Cl}^\cdot + \text{H}_2\text{O} \rightarrow \text{HOCl}^\cdot$	2.5×10^5	Eq. 14	121
$\text{SO}_4^- + \text{HCO}_3^- \rightarrow \text{SO}_4^{2-} + \text{CO}_3^{\cdot-} + \text{H}^+$	9.1×10^6	Eq. 15	122
	$6.38 \pm 0.53 \times 10^{6c}$ - $3.68 \pm 0.34 \times 10^{7c}$	Eq. 16	117

^aForward. ^bBackward. ^c $\text{M}_C^{-1} \text{s}^{-1}$.

dissolved species (i.e., natural organic matter, HCO_3^- , and $\text{S}_2\text{O}_8^{2-}$) before they can diffuse to the interfaces where particle-associated organic contaminants are located.⁹⁷ In addition, uncharged organic contaminants often partition into particulate organic matter and organic coatings on minerals; this polymeric organic material would compete with contaminants for reactive oxidant species like $\text{SO}_4^{\bullet-}$. Thus, rates of transformation of absorbed contaminants will often be limited by the rates at which they are released to solution to reestablish equilibrium partitioning after dissolved forms of the compounds are depleted.

The rate of desorption of neutral organic contaminants from the solid phase depends upon the hydrophobicity of the compound as well as the amount of organic carbon and surface area of the material from which the solute is associated.^{98,99} Although elevated temperatures encountered during heat-activated $\text{SO}_4^{\bullet-}$ treatment result in increased concentrations of hydrophobic compounds in the aqueous phase and faster desorption kinetics, this effect tends to be relatively unimportant. For example, an increase of temperature from 25 to 50 °C will only increase the equilibrium concentration of dissolved pyrene by about 45% assuming an enthalpy of sorption (ΔH_{sorp}) of -11.8 kJ/mol .¹⁰⁰

To enhance the efficacy of ISCO in the treatment of hydrophobic organic compounds, researchers have employed surfactants.^{101–103} However, this approach has not been widely adopted in practice due to a variety of factors. First, most surfactants react with $\text{SO}_4^{\bullet-}$ or OH^\cdot ; competition for radicals can slow the rate of contaminants oxidation. In addition, there is evidence that some surfactants (e.g., Brij 35, Triton X-100, and Tween 80) slow the rate of $\text{S}_2\text{O}_8^{2-}$ activation by interrupting radical chain reactions that are important to the oxidation process.^{104–106} Finally, residual surfactants remaining after treatment can compromise the water quality.

4.5. Effect of Dissolved Solutes on the Kinetics of $\text{S}_2\text{O}_8^{2-}$ -Based ISCO. Much of the power of ISCO treatment with oxidants can be traced to the occurrence of radical chain reactions, which can be divided into three phases. After the initial production of primary radicals (e.g., $\text{SO}_4^{\bullet-}$), a radical propagation phase is initiated as stable molecules (e.g., organic contaminants and inorganic solutes) are converted into radicals. These secondary radicals (e.g., OH^\cdot , carbon-centered radicals)

eventually react with $\text{S}_2\text{O}_8^{2-}$ to propagate the radical chain or react with other radicals to form a stable nonradical adduct and terminate the chain reactions.

The kinetics of $\text{S}_2\text{O}_8^{2-}$ activation as well as the efficiency through which contaminants are transformed are affected by dissolved solutes, such as chloride (Cl^-), bicarbonate (HCO_3^-), and natural organic matter. The way in which these species affect the efficacy of the treatment processes depends upon their concentrations, which, in turn, is often determined by local conditions (e.g., pH, mineral types, and presence of cocontaminants).

Cl^- reacts with $\text{SO}_4^{\bullet-}$ quickly (Table 1, eq 12) to form chlorine radical (Cl^\cdot) as the primary product, which in turn initiates a chain of propagation reactions (e.g., eqs 13 and eq 14 in Table 1).^{107,108} However, the rate constant for the reverse of eq 13 is also fast, which can push the reaction backward. Therefore, there is usually little loss of $\text{SO}_4^{\bullet-}$ through this process unless chloride concentrations are very high.^{109,110} If Cl^- are present at an increased concentration compared to the initial $\text{S}_2\text{O}_8^{2-}$ (e.g., $[\text{Cl}^-] > 0.3 \text{ M}$, $[\text{S}_2\text{O}_8^{2-}] > 0.023 \text{ M}$), the forward reaction (eq 13) would be significant and would lead to a greater level of sulfate radical scavenging.¹¹¹ Generally, Cl^- concentration in groundwater is less than 0.015 M ,¹¹² and $\text{S}_2\text{O}_8^{2-}$ is used at substantially higher concentrations during ISCO. Therefore, Cl^- usually will not be a significant sink for $\text{SO}_4^{\bullet-}$ during $\text{S}_2\text{O}_8^{2-}$ -based ISCO treatment.

Similarly, HCO_3^- also can scavenge $\text{SO}_4^{\bullet-}$ (e.g., eq 15 in Table 1) and impact reaction pathways, kinetics, and the overall efficiency of oxidative treatment.^{113,114} The concentrations of HCO_3^- in groundwater depend on the nature of minerals through which the groundwater flows (e.g., typically around 1 mM in granitic rock or sandy aquifers and $\sim 10 \text{ mM}$ in limestone formations).¹¹⁵ Under the low pH conditions typical of $\text{S}_2\text{O}_8^{2-}$ -based ISCO (i.e., < 5), H_2CO_3^* will be the predominant form of dissolved inorganic carbon.¹¹⁶ Thus, the scavenging of $\text{SO}_4^{\bullet-}$ by carbonate species will be negligible. If base activation is used for $\text{S}_2\text{O}_8^{2-}$ activation, scavenging could be more significant.

In addition, dissolved organic matter presented in groundwater can react with $\text{SO}_4^{\bullet-}$ (eq 16 in Table 1). If dissolved organic matter is presented at high concentrations compared to the initial molarity of $\text{S}_2\text{O}_8^{2-}$ (e.g., $25 \text{ mg}_C/\text{L}$, $0.005 \text{ M S}_2\text{O}_8^{2-}$), its scavenging of $\text{SO}_4^{\bullet-}$ can become important.¹¹⁷ However,




Figure 4. Oxidation of organic contaminants by $\text{SO}_4^{\bullet-}$ and product radicals. The process is initiated by homolytic (Δ) or heterolytic (e^-) cleavage of the O–O bond in $\text{S}_2\text{O}_8^{2-}$ and generation of $\text{SO}_4^{\bullet-}$. Subsequently, $\text{SO}_4^{\bullet-}$ reacts with organic compounds to produce secondary organic radicals. $\text{SO}_4^{\bullet-}$ can react with water or hydroxide to generate $\cdot\text{OH}$, which can oxidize many contaminants of concern. Due to eq 5, the conversion of $\text{SO}_4^{\bullet-}$ into $\cdot\text{OH}$ is especially significant at high pH values (e.g., during base activation). However, under neutral or acidic pH, which are typical of ISCO, most of $\text{SO}_4^{\bullet-}$ reacts with other solutes before it can be converted to $\cdot\text{OH}$. Additionally, the reaction of $\text{SO}_4^{\bullet-}$ with $\text{S}_2\text{O}_8^{2-}$ produces $\text{S}_2\text{O}_8^{\bullet-}$ that together with other radicals such as α -hydroxyalkyl radicals can in the absence of O_2 reduce highly chlorinated compounds. Created with BioRender.com.

typical dissolved organic matter concentrations in groundwater (i.e., 1–2 mg_C/L ¹¹⁸) are considerably lower than the amount of $\text{S}_2\text{O}_8^{2-}$ employed in ISCO.

5. TRANSFORMATION OF ORGANIC CONTAMINANTS



There are three main mechanisms through which organic contaminants are transformed during $\text{S}_2\text{O}_8^{2-}$ -based ISCO: (1) direct reaction with $\text{S}_2\text{O}_8^{2-}$ (i.e., two-electron transfer), (2) reactions with $\text{SO}_4^{\bullet-}$, and (3) reactions with secondary radicals.

5.1. Two-Electron Transfer Mechanism. Due to its strong electrophilic character, $\text{S}_2\text{O}_8^{2-}$ is capable of oxidizing electron-rich moiety organic contaminants (e.g., substituted phenols, arylamines) without the occurrence of radical chain reactions (Elbs and Boyland–Sims oxidation).^{123,124} The two-electron transport reaction usually occurs via two steps: (i) attack of $\text{S}_2\text{O}_8^{2-}$ at the carbon atom located ortho/para to the electron-donating moiety, accompanied by deprotonation, resulting in formation of phenolate/arylamine sulfate ions, followed by (ii) hydrolysis to form hydroxylated aromatics and sulfate.³⁴ Based on the Behrman's mechanism, $\text{S}_2\text{O}_8^{2-}$ reacts with phenols and arylamines in two ways depending on whether or not the compound is protonated.¹²⁵ Under alkaline conditions (e.g., pH \approx 13), *o*-phenolsulfates/*o*-aminoarylsulfates are formed, while under acidic conditions phenols and amines polymerize to form polyphenols/polyanilines. Recent studies, however, suggest the formation of polyarylamine precipitates as well as soluble *o*-aminoarylsulfates over a broader pH range (i.e., >5).^{126,127}

5.2. Transformation of Organic Contaminants by $\text{SO}_4^{\bullet-}$ and Secondary Radicals. $\text{SO}_4^{\bullet-}$ radicals react with organic molecules and solutes by three mechanisms:¹²⁸ (i) hydrogen atom abstraction, (ii) single-electron transfer, and (iii) addition to double or triple bonds, leading to the formation of secondary radicals that undergo a suite of reactions leading to a variety of different products (Figure 4). Furthermore, these reactions may occur simultaneously.

For example, $\text{SO}_4^{\bullet-}$ reacts with alkylated benzenes (i.e., BTEX) by single-electron transfer from the attacked species to $\text{SO}_4^{\bullet-}$ radical, resulting in the formation of the benzene radical cations and SO_4^{2-} .^{129–131} The next step involves the addition of OH^- to the benzene-derivative radical cation that then forms a hydroxycyclohexadienyl radical.¹³² This radical then can react with O_2 to produce hydroxycyclohexadienyl peroxy radicals, which then can eliminate hydroperoxyl radical (HO_2^{\bullet}) to produce hydroxylated products (eq 17).¹³³ Alternatively, hydrogen abstraction can produce hydroquinone radical and then hydroquinone (eq 18).^{134,135} In an alternative pathway, the hydroxycyclohexadienyl peroxy radical can also undergo oxygen addition to a double bond to form a bicyclic peroxy radical, which in the absence of any oxidant may be disproportionated to produce two endoperoxides. Both products are unstable and can undergo ring opening and production of 5- and 6-membered unsaturated aldehydes and ketones (eq 17),^{136,137} which can be further oxidized to carboxylic acid and finally CO_2 . Similar reactions can also be triggered by the reaction of aromatic compounds with $\cdot\text{OH}$.^{138–140} In addition, $\text{SO}_4^{\bullet-}$ also reacts with

aromatic compounds through radical addition (Figure 4). Although the addition process is usually considered a secondary pathway,^{141,142} it leads to the formation of organosulfates, which may be of environmental concern due to their higher mobility in the subsurface compared to the hydrophobic parent compounds as well as their elevated toxicity.^{143,144} However, organosulfates may undergo further transformation, such as further reaction with $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$ or biotransformation, resulting in complete mineralization (i.e., conversion to CO_2).^{137,145}



Similarly, polyaromatic hydrocarbons react with $\text{SO}_4^{\bullet-}$ predominantly by single-electron transfer, leading to the formation of oxygen transformation products (i.e., aromatic ketones, aromatic quinones, and aliphatic or aromatic acids), which can further react with $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$ ⁸ or undergo biotransformation.¹⁴⁶

Alternatively, hydrogen abstraction of a carbon-bond hydrogen atom is believed to be the main mechanism by which $\text{SO}_4^{\bullet-}$ reacts with tertiary dialkyl ethers (methyl and ethyl *tert*-butyl ether, and methyl *tert*-amyl ether) to generate carbon centered radicals, which then react with O_2 , forming peroxy radicals that undergo an acid-catalyzed hydrolysis reaction to yield tertiary alcohols and carboxylic acids.⁶⁵




Similarly, haloalkanes (e.g., $\text{CH}_2\text{Cl}-\text{CHCl}_2$) react with $\text{SO}_4^{\bullet-}$ via abstraction of a carbon-bond hydrogen atom, resulting in the formation of alkyl-like radicals (e.g., $\text{CH}_3-\text{C}^{\bullet}\text{Cl}_2$).¹¹⁴ These radicals react with O_2 to produce peroxy radicals (e.g., $\text{CH}_3-\text{C}(\text{Cl}_2)\text{OO}^{\bullet}$) that may hydrolyze to corresponding carboxylic acid, HCl, and superoxide radical ($\text{O}_2^{\bullet-}$).¹⁴⁷

$\text{S}_2\text{O}_8^{2-}$ -based ISCO is not a very effective means of treating highly halogenated organic contaminants (e.g., polychlorinated benzenes, polyhalogenated C_1 and C_2 compounds) under oxic condition.^{10,64,148} This is related to the low reactivity of carbon-halogen bonds with electrophilic radicals (e.g., $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$).^{149,150} However, highly halogenated contaminants more readily undergo reductive dehalogenation reactions, creating the possibility that reducing radicals produced during the chain decomposition of $\text{S}_2\text{O}_8^{2-}$ could produce transformation products that react with $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$.

$\text{SO}_4^{\bullet-}$ reacts with $\text{S}_2\text{O}_8^{2-}$ to produce a persulfate radical ($\text{S}_2\text{O}_8^{\bullet-}$) (eq 6). In the absence of O_2 , $\text{S}_2\text{O}_8^{\bullet-}$ reacts with $\text{S}_2\text{O}_8^{2-}$, initiates radical chain reactions (eq 19) and reduces highly chlorinated contaminants like hexachloroethane¹⁵¹ and DDT.¹² In addition, the consumption of one molecule of $\text{S}_2\text{O}_8^{2-}$ by $\text{S}_2\text{O}_8^{\bullet-}$ produces one molecule of $\text{SO}_4^{\bullet-}$, which again can react with $\text{S}_2\text{O}_8^{2-}$ to produce $\text{S}_2\text{O}_8^{\bullet-}$. However, in the presence of O_2 , the reaction between $\text{S}_2\text{O}_8^{\bullet-}$ and O_2 terminates the reaction ($4\text{S}_2\text{O}_8^{\bullet-} + \text{O}_2 + 2\text{H}_2\text{O} \rightarrow 8\text{HSO}_5^- + 4\text{H}^+$). Another approach involves the use of carbon-centered radicals that are produced when $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$ radicals react with short chain length aliphatic compounds (e.g., methanol, ethanol). It has been reported that these radicals can reduce highly halogenated

compounds such as carbon tetrachloride or hexachloroethane.^{151,152} However, O_2 competes with contaminant carbon-centered radicals (e.g., $\cdot\text{CH}(\text{CH}_3)\text{OH} + \text{O}_2 \rightarrow \text{CH}_3\text{CHO} + \text{HO}_2^{\bullet}$).¹⁵³ HO_2^{\bullet} and its conjugate base, $\text{O}_2^{\bullet-}$, then quickly undergo bimolecular dismutation to produce H_2O_2 under acidic conditions.^{154–156} Since $\text{S}_2\text{O}_8^{2-}$ decomposition produces O_2 (eq 1), it is necessary to use sufficient quantities of alcohol to ensure simultaneous reduction of halogenated contaminants and initiate chain decomposition of $\text{S}_2\text{O}_8^{2-}$ that produces additional carbon-centered radicals (eq 20) and consumes O_2 . In this first phase the reactions of carbon-centered radicals result in the formation of partially dehalogenated products that contain electron-rich double bonds or carbon-hydrogen bonds (Figure 4) and are less susceptible to further reduction but are prone to subsequent oxidation by $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$.^{151,157} The following oxidation reactions require sufficient concentrations of $\text{S}_2\text{O}_8^{2-}$ as well as a means of accelerating the rate of $\text{S}_2\text{O}_8^{2-}$ activation.¹⁵²

Additionally, it has been reported that perfluoroalkyl acids which are resistant to oxidation by $\cdot\text{OH}$ treatment react with $\text{SO}_4^{\bullet-}$ to form shorter-chain perfluoroalkyl carboxylic acids, which may be further degraded and eventually mineralized.^{66,158} The multistep mechanism responsible for the transformation of perfluorooctanoic acid to perfluoroheptanoic acid can be described by eq 21. The subsequent oxidation of perfluoroheptanoic acid to produce perfluorohexanoic acid follows an analogous mechanism. The transformation of perfluorooctanoic acid to shorter-chain perfluorocarboxylic acids by $\text{SO}_4^{\bullet-}$ can only be carried out under strongly acidic conditions ($\text{pH} \leq 3$).¹⁵⁹



5.3. Transformation Products. The oxidation of organic contaminants by a high concentration of $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$ often results in mineralization (i.e., contaminants are converted into water, carbon dioxide, Cl^- and other relatively innocuous products). However, the incomplete oxidation of contaminants could lead to the formation of transformation products that are more toxic and/or mobile than the compounds targeted for remediation, especially if those compounds are less reactive with $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$ than the parent compounds.¹⁶⁰ For example, reactions of $\text{SO}_4^{\bullet-}$ and $\cdot\text{OH}$ with benzene and alkylbenzenes produce toxic ring-cleavage products such as α,β -unsaturated aldehydes or ketones (e.g., acrolein).^{161,137} Additionally, incomplete transformation of haloalkanes can produce harmful haloalkenes, such as dichloromethane,¹⁶² which is a possible human carcinogen.¹⁶³ Similarly, if $\text{S}_2\text{O}_8^{2-}$ treatment of perfluoroalkylcarboxylic acids does not lead to complete mineralization, the resulting shorter-chain perfluoroalkylcarboxylic acids may be more mobile than the parent PFAS compounds¹⁶⁴ and more difficult to remove by sorption technologies.¹⁶⁵

Additionally, the oxidation of halides (X^-) in the soil or groundwater by $SO_4^{\bullet-}$ can initiate a cascade of radical chain reactions (e.g., eqs 12–14), leading to the production of secondary radicals (X^\bullet , $X_2^{\bullet-}$, and XHO^\bullet) and free halogens (X_2/HXO).¹⁶⁶ These species are extremely reactive electrophiles that can halogenate electron-rich substances, such as NOM¹⁶⁷ and benzene derivatives.¹⁶⁸ This process may lead to the production of halogenated byproducts, such as trihalomethanes and haloacetic acids,^{169–171} which raises concerns due to their well documented human health effects.^{172–175} Field measurements made during ISCO treatment have revealed elevated concentrations of chloromethane and methylene chloride, which are precursors to trihalomethanes; however, chloroform or other trihalomethanes have not been detected at elevated concentrations after the application of $S_2O_8^{2-}$.¹⁸ Therefore, more research is necessary to assess the formation of halogenated byproducts under the conditions typical for $S_2O_8^{2-}$ -based ISCO. Another concern is the potential production of chlorate (ClO_3^-); due to sequential oxidation of Cl^- , concentrations that could pose a health hazard have been reported under conditions similar to those employed for ISCO.^{108,176}

$SO_4^{\bullet-}$ also reacts with ammonia (NH_4^+) and nitrate (NO_2^-) to produce nitrogen dioxide (NO_2^\bullet) and monoxide (NO^\bullet) radicals, which can react with electron-rich compounds such as PAHs or NOM to produce toxic nitro-byproducts.^{177–181} For example, nitro-PAHs are known to possess higher mutagenicity and carcinogenicity compared to their parent PAHs.^{182,183} While PAHs need initial enzymatic activation before they can act as mutagens or carcinogens,¹⁸⁴ their nitro-byproducts are direct acting mutagens.¹⁸⁵ Similarly, nitro-aromatics pose potential hazards to ecosystems and human health, including the risk of cancer.¹⁸⁶ In addition, the introduction of a nitro group into the molecule could enhance the mobility of the compound by reducing its tendency to partition onto surfaces (i.e., aquifer solids).

$SO_4^{\bullet-}$ also has the potential to form organosulfonates through addition reactions. These compounds as well exhibit greater mobility in the subsurface and higher toxicity than their parent compounds.¹⁴⁴ For example, *p*-cresol sulfate, a relatively polar transformation product of benzene treatment by $SO_4^{\bullet-}$, is associated with renal toxicity and kidney damage.¹⁸⁷

Most of the transformation products underwent further oxidative treatment with $SO_4^{\bullet-}$. However, they can also undergo biodegradation. For example, nitro-aromatics can be completely mineralized via catabolic processes without the need for additional external substrates in an oxic environment.¹⁸⁸ The primary mechanism of aerobic degradation of nitroaromatics is initiated by the addition of a molecular oxygen atom to the benzene ring catalyzed by oxygenase, with subsequent release of a nitro group and ring cleavage, as in the biodegradation of 4-nitrophenol.¹⁸⁹ Desulfurization of aromatic sulfonates proceeds by a similar pathway.¹⁹⁰ However, evidence for the biodegradation of nitro-PAHs is lacking. So far, only a few microorganisms have been identified that are able to degrade nitro-PAHs, such as nitroanthracene.¹⁹¹ Nitro-PAHs are also more resistant to further oxidation by $SO_4^{\bullet-}$ than their parent compounds.¹⁸⁰ Therefore, additional research is needed to better identify the formation of toxic transformation products and develop appropriate treatment methods.

6. IMPLICATIONS AND PROSPECTS

As the use of $S_2O_8^{2-}$ -based ISCO continues to advance, it is becoming a well-established and extensively studied field of research. This progress is evident with the increasing number of academic studies and the implementation of full-scale applications in soil and groundwater remediation. However, there are still challenges that limit the performance of this technology in field applications.

$S_2O_8^{2-}$ is a relatively stable oxidant; however, there is a trade-off associated with this intrinsic stability due to its slow rate of reaction with most contaminants. Therefore, $S_2O_8^{2-}$ needs to be activated to generate $SO_4^{\bullet-}$ in order to oxidize most recalcitrant contaminants. Among various activation methods, thermal activation appears to be the most viable option, but the optimal temperature range (30–40 °C) must balance energy consumption and reaction efficiency. H_2O_2 represents a suitable method to thermally activate $S_2O_8^{2-}$. However, because of its very high soil oxidant demand, using H_2O_2 may not be feasible for sites with a high clay content and low permeability. Heat can be introduced into the subsurface also by steam injection or with electrodes; however, these approaches are suitable mainly for smaller sites (as the size increases, so does the cost). Nevertheless, challenges still exist with heat activation, such as achieving uniform heat distribution. Hence, further research and engineering improvements are crucial to maximize the effectiveness of the heat-activated $S_2O_8^{2-}$ -based ISCO.

In cases where thermal activation is not feasible, alkaline activation can serve as an alternative approach. This method involves the injection of a strong base (e.g., NaOH) that raises the pH to above 12 and activates $S_2O_8^{2-}$. However, there are several important considerations associated with this approach. First, careful attention must be paid to the use and safe handling of strong bases throughout the design and implementation of the project. Another critical consideration relates to pH adjustment, which consists of two key components. The first component pertains to the amount needed to adjust the pH of the soil and groundwater to the desired level. The second component involves the quantity required to counteract the impact of the acid formed during $S_2O_8^{2-}$ decomposition. Therefore, both of these components must be carefully considered when determining the amount of pH modifier.

Although other methods, like employing heterogeneous catalyst or dissolved transition metals, are actively being researched, their applicability to ISCO remains limited. Solid phase catalysts can activate $S_2O_8^{2-}$,^{192,193} but their use during in situ treatment is restricted by the challenges associated with delivering them to the source zone and concerns about their toxicity. Research in this area is limited, with most studies being conducted as batch, column, or box experiments.^{194–196} There is also a lack of information regarding pilot- or full-scale implementation. Concerning dissolved transition metals, only Fe(II) is a viable option. However, research²⁷ suggests that this process is essentially one-way, meaning that Fe(III) does not get reduced back to Fe(II) to create a catalytic loop. Therefore, the introduction of Fe(II) into the aquifer, followed by the addition of $S_2O_8^{2-}$, results in a burst of $SO_4^{\bullet-}$ production, but a certain portion is consumed in oxidizing Fe(II), making it an impractical approach. It might be possible to prevent Fe(III) precipitation by introducing a chelated form of Fe(II) into the aquifer. However, common chelating agents used to prevent Fe(III) precipitation, such as citric acid and EDTA, are not specific to

iron and may form complexes with other metals and mobilize them from the soil matrix.

In practice, the $S_2O_8^{2-}$ -based ISCO for contaminant remediation faces several limitations under field conditions. The heterogeneity of subsurface geology and the presence of low-permeability zones can hinder the delivery of $S_2O_8^{2-}$ to contaminated areas requiring treatment. To increase the potential for treating difficult-to-reach low-permeability zones, methods may be needed to enhance the lifetime of the oxidant as it diffuses into these regions. Additionally, the sorption of contaminants to organic matter can limit the transformation rates during $S_2O_8^{2-}$ -based ISCO, especially for strongly sorbed organic contaminants.

Furthermore, potential loss of $S_2O_8^{2-}$ through reactions with aquifer solids, such as Fe(III)- and Mn(IV) oxides, as well as the presence of high concentrations of organic matter and dissolved solutes, such as Cl^- , O_2 or organic cocontaminants, can impact the kinetics and efficiency of $S_2O_8^{2-}$ -based oxidation reactions. However, the use of a high concentration of $S_2O_8^{2-}$ during the field application can effectively counteract these inhibitory effects.

Acidic conditions and exposure of minerals and solutes to large amounts of oxidants can alter the geochemistry of the subsurface, leading to changes in the solubility of certain inorganic species and the mobilization of metals. Furthermore, the $S_2O_8^{2-}$ -based ISCO can partly or fully inhibit certain microbial species, especially those that are sensitive to oxygen and acidity. Since natural attenuation often is expected to play an important role in the breakdown of residual contaminants at sites where $S_2O_8^{2-}$ -based ISCO is planned, permanent inhibition in microbial activity is undesirable. Therefore, treatability tests should be conducted on bench- and pilot-scale to assess the potential impact of $S_2O_8^{2-}$ on geochemistry and microbial activity.

$SO_4^{\bullet-}$ reacts with organic molecules through various mechanisms, leading to the formation of secondary radicals and a wide range of transformation products. Despite the success of $S_2O_8^{2-}$ -based ISCO in the treatment of various contaminants, the technology is not very effective when highly halogenated contaminants are present. One promising approach is to employ reductive radicals, such as carbon-centered radicals produced during chain decomposition of $S_2O_8^{2-}$. However, further research is necessary to assess the impact of field application conditions (e.g., presence of aquifer solids).

Incomplete oxidation of contaminants can result in the formation of potentially more toxic or mobile transformation byproducts. Similarly, reactions with solutes may lead to the formation of products such as ClO_3^- or chlorinated and nitro-containing byproducts that may pose health risks to consumers of groundwater. Furthermore, $SO_4^{\bullet-}$ has the potential to form organosulfates through radical addition that may also be more toxic and mobile than the parent compounds. These byproducts may undergo substantial oxidation or be transformed by microorganisms. However, certain byproducts, such as nitro-PAHS, are resistant also to biodegradation. The formation of these difficult to treat byproducts should be carefully assessed prior to $S_2O_8^{2-}$ -based ISCO application.

Despite all these challenges, $S_2O_8^{2-}$ -based ISCO is a promising technology that can be very useful in the treatment of contaminated subsurface. We believe that understanding the practical limitations of $S_2O_8^{2-}$ -based ISCO reviewed in this paper can guide researchers and remediation engineers in developing more effective approaches, predicting its applic-

ability, enhancing its effectiveness, and minimizing unintended consequences when it is applied in the field.

AUTHOR INFORMATION

Corresponding Authors

Lenka McGachy – Department of Environmental Chemistry, University of Chemistry and Technology Prague, 16628 Prague, Czech Republic; orcid.org/0000-0002-2251-1798; Email: lenka.mc.gachy@vscht.cz

David L. Sedlak – Department of Civil and Environmental Engineering, University of California, Berkeley, California 94720, United States; orcid.org/0000-0003-1686-8464; Email: sedlak@berkeley.edu

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.est.3c07409>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The financial support of the J. William Fulbright Commission in the Czech Republic and Czech Science Foundation (Grant 23-05901S) is gratefully acknowledged. Additional support was provided by the U.S. National Institute for Environmental Health Sciences Superfund Research Program (Grant P42 ES004705) at the University of California, Berkeley.

REFERENCES

- (1) Chen, Q.; Fan, G.; Na, W.; Liu, J.; Cui, J.; Li, H. Past, Present, and Future of Groundwater Remediation Research: A Scientometric Analysis. *International Journal of Environmental Research and Public Health* **2019**, *16* (20), 3975.
- (2) Siegrist, R. L.; Crimi, M.; Brown, R. A. In Situ Chemical Oxidation: Technology Description and Status. In *In Situ Chemical Oxidation for Groundwater Remediation*, Siegrist, R. L., Crimi, M., Simpkin, T. J., Eds.; Springer: New York, 2011; pp 1–32.
- (3) Waldemer, R. H.; Tratnyek, P. G.; Johnson, R. L.; Nurmi, J. T. Oxidation of Chlorinated Ethenes by Heat-Activated Persulfate: Kinetics and Products. *Environ. Sci. Technol.* **2007**, *41* (3), 1010–1015.
- (4) Costanza, J.; Otaño, G.; Callaghan, J.; Pennell, K. D. PCE Oxidation by Sodium Persulfate in the Presence of Solids. *Environ. Sci. Technol.* **2010**, *44* (24), 9445–9450.
- (5) Sra, K. S.; Thomson, N. R.; Barker, J. F. Persulfate injection into a gasoline source zone. *Journal of Contaminant Hydrology* **2013**, *150*, 35–44.
- (6) Liang, C.; Huang, C.-F.; Chen, Y.-J. Potential for activated persulfate degradation of BTEX contamination. *Water Res.* **2008**, *42* (15), 4091–4100.
- (7) Lominchar, M. A.; Santos, A.; de Miguel, E.; Romero, A. Remediation of aged diesel contaminated soil by alkaline activated persulfate. *Science of The Total Environment* **2018**, *622–623*, 41–48.
- (8) Liao, X.; Zhao, D.; Yan, X.; Huling, S. G. Identification of persulfate oxidation products of polycyclic aromatic hydrocarbon during remediation of contaminated soil. *Journal of Hazardous Materials* **2014**, *276*, 26–34.
- (9) Zhao, D.; Liao, X.; Yan, X.; Huling, S. G.; Chai, T.; Tao, H. Effect and mechanism of persulfate activated by different methods for PAHS removal in soil. *Journal of Hazardous Materials* **2013**, *254–255*, 228–235.
- (10) Usman, M.; Tascone, O.; Faure, P.; Hanna, K. Chemical oxidation of hexachlorocyclohexanes (HCHs) in contaminated soils. *Science of The Total Environment* **2014**, *476–477*, 434–439.
- (11) Zhang, X. H.; Liu, X. T.; Zhao, J. G.; Sun, W. J.; Li, X.; Li, J. W.; Zhang, J.; Jing, Z. B. NaOH-activated persulfate-assisted mechanochemical mechanism and removal of lindane from contaminated soil. *Journal of Environmental Chemical Engineering* **2021**, *9* (4), 105391.

- (12) Zhu, C.; Wang, D.; Zhu, F.; Liu, S.; Fang, G.; Gao, J.; Zhou, D. Rapid DDTs degradation by thermally activated persulfate in soil under aerobic and anaerobic conditions: Reductive radicals vs. oxidative radicals. *Journal of Hazardous Materials* **2021**, *402*, No. 123557.
- (13) Sedlak, D. L.; Andren, A. W. Oxidation of chlorobenzene with Fenton's reagent. *Environ. Sci. Technol.* **1991**, *25* (4), 777–782.
- (14) Barbash, A. M.; Hoag, G. E.; Nadim, F. Oxidation and removal of 1,2,4-trichlorobenzene using sodium persulfate in a sorption-desorption experiment. *Water Air Soil Pollut.* **2006**, *172* (1–4), 67–80.
- (15) Fang, G.-D.; Dionysiou, D. D.; Zhou, D.-M.; Wang, Y.; Zhu, X.-D.; Fan, J.-X.; Cang, L.; Wang, Y.-J. Transformation of polychlorinated biphenyls by persulfate at ambient temperature. *Chemosphere* **2013**, *90* (5), 1573–1580.
- (16) Krembs, F. J.; Siegrist, R. L.; Crimi, M. L.; Furrer, R. F.; Petri, B. G. ISCO for Groundwater Remediation: Analysis of Field Applications and Performance. *Groundwater Monitoring & Remediation* **2010**, *30* (4), 42–53.
- (17) Tsitonaki, A.; Petri, B.; Crimi, M.; Mosbaek, H.; Siegrist, R. L.; Bjerg, P. L. In Situ Chemical Oxidation of Contaminated Soil and Groundwater Using Persulfate: A Review. *Critical Reviews in Environmental Science and Technology* **2010**, *40* (1), 55–91.
- (18) Rosansky, S.; Dindal, A. *Cost and Performance Report for Persulfate Treatability Studies*; TR-2333-ENV; NAVFAC Engineering Service Center: 2010.
- (19) Lowe, K. S.; Gardner, F. G.; Siegrist, R. L.; Houck, T. C. *Field Pilot Test of In Situ Chemical Oxidation through Recirculation Using Vertical Wells at the Portsmouth Gaseous Diffusion Plant*. EPA Office of Research and Development: Washington, DC, 1999.
- (20) Cooper, W. J.; Zepp, R. G. Hydrogen Peroxide Decay in Waters with Suspended Soils: Evidence for Biologically Mediated Processes. *Canadian Journal of Fisheries and Aquatic Sciences* **1990**, *47* (5), 888–893.
- (21) Petigara, B. R.; Blough, N. V.; Mignerey, A. C. Mechanisms of Hydrogen Peroxide Decomposition in Soils. *Environ. Sci. Technol.* **2002**, *36* (4), 639–645.
- (22) Kakarla, P. K. C.; Watts, R. J. Depth of Fenton-Like Oxidation in Remediation of Surface Soil. *J. Environ. Eng.* **1997**, *123* (1), 11–17.
- (23) Watts, R. J.; Finn, D. D.; Cutler, L. M.; Schmidt, J. T.; Teel, A. L. Enhanced stability of hydrogen peroxide in the presence of subsurface solids. *Journal of Contaminant Hydrology* **2007**, *91* (3), 312–326.
- (24) Nyer, E. K.; Vance, D. Hydrogen Peroxide Treatment: The Good, the Bad, the Ugly. *Groundwater Monitoring Remediation* **1999**, *19* (3), 54–57.
- (25) Siegrist, R.; Crimi, M.; Petri, B.; Simpkin, T.; Palaia, T.; Krembs, F.; Munakata-Marr, J.; Illangasekare, T.; Ng, G.; Singletary, M. *In Situ Chemical Oxidation for Groundwater Remediation: Site Specific Engineering and Technology Application. ER-0623 Final Report, CD-ROM, Version PRv1. 01, October 29, 2010; ESTCP: Arlington, VA, 2010.*
- (26) Huling, S. G.; Pivetz, B. E. *In-situ chemical oxidation*; U.S. Environmental Protection Agency, Office Of Water: Washington, DC, 2006.
- (27) Liu, H.; Bruton, T. A.; Doyle, F. M.; Sedlak, D. L. In Situ Chemical Oxidation of Contaminated Groundwater by Persulfate: Decomposition by Fe(III)- and Mn(IV)-Containing Oxides and Aquifer Materials. *Environ. Sci. Technol.* **2014**, *48* (17), 10330–10336.
- (28) Liang, C. J.; Chen, C. Y. Characterization of a Sodium Persulfate Sustained Release Rod for in Situ Chemical Oxidation Groundwater Remediation. *Ind. Eng. Chem. Res.* **2017**, *56* (18), 5271–5276.
- (29) Zhang, B.-T.; Zhang, Y.; Teng, Y.; Fan, M. Sulfate Radical and Its Application in Decontamination Technologies. *Critical Reviews in Environmental Science and Technology* **2015**, *45* (16), 1756–1800.
- (30) House, D. A. Kinetics and Mechanism of Oxidations by Peroxydisulfate. *Chem. Rev.* **1962**, *62* (3), 185–203.
- (31) Behrman, E. J.; Edwards, J. O. The thermal decomposition of peroxydisulfate ions. *Reviews in Inorganic Chemistry* **1980**, *2*, 179–206.
- (32) Kolthoff, I. M.; Miller, I. K. The Chemistry of Persulfate. I. The Kinetics and Mechanism of the Decomposition of the Persulfate Ion in Aqueous Medium. *J. Am. Chem. Soc.* **1951**, *73* (7), 3055–3059.
- (33) Liang, C.; Su, H.-W. Identification of Sulfate and Hydroxyl Radicals in Thermally Activated Persulfate. *Ind. Eng. Chem. Res.* **2009**, *48* (11), 5558–5562.
- (34) Behrman, E. J. The Persulfate Oxidation of Phenols and Arylamines (The Elbs and the Boyland–Sims Oxidations). *Organic Reactions* **1988**, 421–511.
- (35) Behrman, E. J. Studies on the reaction between peroxydisulfate ions and aromatic amines. Boyland-Sims oxidation. *J. Am. Chem. Soc.* **1967**, *89* (10), 2424–2428.
- (36) Johnson, R. L.; Tratnyek, P. G.; Johnson, R. O. B. Persulfate Persistence under Thermal Activation Conditions. *Environ. Sci. Technol.* **2008**, *42* (24), 9350–9356.
- (37) Wojnárovits, L.; Takács, E. Rate constants of sulfate radical anion reactions with organic molecules: A review. *Chemosphere* **2019**, *220*, 1014–1032.
- (38) Parsons, S. *Advanced oxidation processes for water and wastewater treatment*; IWA Publishing: 2004.
- (39) Furman, O. S.; Teel, A. L.; Watts, R. J. Mechanism of Base Activation of Persulfate. *Environ. Sci. Technol.* **2010**, *44* (16), 6423–6428.
- (40) Chandra Singh, U.; Venkatarao, K. Decomposition of peroxodisulphate in aqueous alkaline solution. *Journal of Inorganic and Nuclear Chemistry* **1976**, *38* (3), 541–543.
- (41) Santos, A.; Fernandez, J.; Rodriguez, S.; Dominguez, C. M.; Lominchar, M. A.; Lorenzo, D.; Romero, A. Abatement of chlorinated compounds in groundwater contaminated by HCH wastes using ISCO with alkali activated persulfate. *Science of The Total Environment* **2018**, *615*, 1070–1077.
- (42) Huang, M. Q.; Wang, X. L.; Zhu, C. Y.; Zhu, F. X.; Liu, P.; Wang, D. X.; Fang, G. D.; Chen, N.; Gao, S. X.; Zhou, D. M. Efficient chlorinated alkanes degradation in soil by combining alkali hydrolysis with thermally activated persulfate. *Journal of Hazardous Materials* **2022**, *438*, 129571.
- (43) Stumm, W.; Morgan, J. J. *Aquatic chemistry: chemical equilibria and rates in natural waters*; John Wiley & Sons: 2012.
- (44) Hou, K. J.; Pi, Z. J.; Yao, F. B.; Wu, B.; He, L.; Li, X. M.; Wang, D. B.; Dong, H. R.; Yang, Q. A critical review on the mechanisms of persulfate activation by iron-based materials: Clarifying some ambiguity and controversies. *Chemical Engineering Journal* **2021**, *407*, 127078.
- (45) Škarohlíd, R.; McGachy, L.; Martinec, M.; Rošková, Z. Removal of PCE/TCE from groundwater by peroxydisulfate activated with citric acid chelated ferrous iron at 13 °C. *Environmental Technology & Innovation* **2020**, *19*, No. 101004.
- (46) Bennedsen, L. R.; Krischker, A.; Jørgensen, T. H.; Søgaard, E. G. Mobilization of metals during treatment of contaminated soils by modified Fenton's reagent using different chelating agents. *Journal of Hazardous Materials* **2012**, *199–200*, 128–134.
- (47) Kumar, M.; Xiong, X. N.; Sun, Y. Q.; Yu, I. K. M.; Tsang, D. C. W.; Hou, D. Y.; Gupta, J.; Bhaskar, T.; Pandey, A. Critical Review on Biochar-Supported Catalysts for Pollutant Degradation and Sustainable Biorefinery. *Advanced Sustainable Systems* **2020**, *4* (10), 2070020.
- (48) Zawadzki, P. Persulfate activation by organic compounds: advancements and challenges. *Current Opinion in Chemical Engineering* **2022**, *37*, No. 100837.
- (49) Pearce, D. A.; Sherman, F. Toxicity of copper, cobalt, and nickel salts is dependent on histidine metabolism in the yeast *Saccharomyces cerevisiae*. *J. Bacteriol.* **1999**, *181* (16), 4774–9.
- (50) Guo, M.; Song, W.; Tian, J. Biochar-Facilitated Soil Remediation: Mechanisms and Efficacy Variations. *Frontiers in Environmental Science* **2020**, *8*, 521512.
- (51) Arvaniti, O. S.; Ioannidi, A. A.; Mantzavinos, D.; Frontistis, Z. Heat-activated persulfate for the degradation of micropollutants in water: A comprehensive review and future perspectives. *Journal of Environmental Management* **2022**, *318*, No. 115568.
- (52) Wen, D.; Liu, H.; Zhang, Y.; Fu, R. Electrokinetically-delivered persulfate and pulsed direct current electric field induced transport, mixing and thermally activated in situ for remediation of PAHs contaminated soil. *Journal of Hazardous Materials* **2023**, *444*, No. 130414.

- (53) Li, J.; Wang, L.; Peng, L.; Deng, Y.; Deng, D. A combo system consisting of simultaneous persulfate recirculation and alternating current electrical resistance heating for the implementation of heat activated persulfate ISCO. *Chemical Engineering Journal* **2020**, *385*, No. 123803.
- (54) Head, N. A.; Gerhard, J. I.; Inglis, A. M.; Nunez Garcia, A.; Chowdhury, A. I. A.; Reynolds, D. A.; de Boer, C. V.; Sidebottom, A.; Austrins, L. M.; Eimers, J.; O'Carroll, D. M. Field test of electrokinetically-delivered thermally activated persulfate for remediation of chlorinated solvents in clay. *Water Res.* **2020**, *183*, No. 116061.
- (55) Chowdhury, A. I. A.; Gerhard, J. I.; Reynolds, D.; O'Carroll, D. M. Low Permeability Zone Remediation via Oxidant Delivered by Electrokinetics and Activated by Electrical Resistance Heating: Proof of Concept. *Environ. Sci. Technol.* **2017**, *51* (22), 13295–13303.
- (56) Thompson, S.; Riggenbach, J.; Brown, R.; Hines, J.; Haselow, J. In *Catalyzed persulfate remediation of chlorinated and recalcitrant compounds in soil*, Proceedings of the 5th International Conference on Remediation of Chlorinated and Recalcitrant Compounds, Monterey, CA; Battelle Press: Columbus, OH, 2006.
- (57) ITRC. *Technical and regulatory guidance for in situ chemical oxidation of contaminated soil and groundwater*; Interstate Technology & Regulatory Council: Washington, DC, 2005.
- (58) Davis, E. L. *How heat can enhance in-situ soil and aquifer remediation*; United States Environmental Protection Agency, Office of Research and Development, Office of Solid Waste and Emergency Response: Washington, DC, 1997.
- (59) Beyke, G.; Fleming, D. In situ thermal remediation of DNAPL and LNAPL using electrical resistance heating. *Remediation Journal* **2005**, *15* (3), 5–22.
- (60) Triplett Kingston, J. L.; Dahlen, P. R.; Johnson, P. C. State-of-the-Practice Review of In Situ Thermal Technologies. *Groundwater Monitoring & Remediation* **2010**, *30* (4), 64–72.
- (61) Block, P. A.; Brown, R. A.; Robinson, D. In *Novel activation technologies for sodium persulfate in situ chemical oxidation*, Proceedings of the Fourth International Conference on the remediation of chlorinated and recalcitrant compounds; Battelle Press: Columbus, OH, 2004; pp 24–27.
- (62) Lee, Y.; Lo, S.; Kuo, J.; Hsieh, C. Decomposition of perfluorooctanoic acid by microwave-activated persulfate: Effects of temperature, pH, and chloride ions. *Frontiers of Environmental Science & Engineering* **2012**, *6* (1), 17–25.
- (63) Qi, C.; Liu, X.; Zhao, W.; Lin, C.; Ma, J.; Shi, W.; Sun, Q.; Xiao, H. Degradation and dechlorination of pentachlorophenol by microwave-activated persulfate. *Environmental Science and Pollution Research* **2015**, *22* (6), 4670–4679.
- (64) Xu, M.; Gu, X.; Lu, S.; Qiu, Z.; Sui, Q.; Miao, Z.; Zang, X.; Wu, X. Degradation of carbon tetrachloride in aqueous solution in the thermally activated persulfate system. *Journal of Hazardous Materials* **2015**, *286*, 7–14.
- (65) Huang, K.-C.; Couttenye, R. A.; Hoag, G. E. Kinetics of heat-assisted persulfate oxidation of methyl tert-butyl ether (MTBE). *Chemosphere* **2002**, *49* (4), 413–420.
- (66) Bruton, T. A.; Sedlak, D. L. Treatment of Aqueous Film-Forming Foam by Heat-Activated Persulfate under Conditions Representative of in Situ Chemical Oxidation. *Environ. Sci. Technol.* **2017**, *51* (23), 13878–13885.
- (67) Desrosiers, R. J.; Rach, B. ISCO Optimization in a Low Permeability Formation Using Groundwater Recirculation. In *Battelle Conference on the Remediation of Chlorinated and Recalcitrant Compounds 2022*; Battelle: Palm Springs, CA, 2022.
- (68) Baciocchi, R.; D'Aprile, L.; Innocenti, I.; Massetti, F.; Verginelli, I. Development of technical guidelines for the application of in-situ chemical oxidation to groundwater remediation. *Journal of Cleaner Production* **2014**, *77*, 47–55.
- (69) Rosansky, S.; Condit, W.; Sirabian, R. *Best Practices for Injection and Distribution of Amendments*; Battelle: Port Hueneme, CA, 2013.
- (70) Muller, K. A.; Johnson, C. D.; Bagwell, C. E.; Truex, M. J. Methods for Delivery and Distribution of Amendments for Subsurface Remediation: A Critical Review. *Groundwater Monitoring & Remediation* **2021**, *41* (1), 46–75.
- (71) Pac, T. J.; Baldock, J.; Brodie, B.; Byrd, J.; Gil, B.; Morris, K. A.; Nelson, D.; Parikh, J.; Santos, P.; Singer, M.; Thomas, A. In situ chemical oxidation: Lessons learned at multiple sites. *Remediation Journal* **2019**, *29* (2), 75–91.
- (72) Calvet, R. Adsorption of organic chemicals in soils. *Environ. Health Perspect* **1989**, *83*, 145–77.
- (73) Essaid, H. I.; Bekins, B. A.; Cozzarelli, I. M. Organic contaminant transport and fate in the subsurface: Evolution of knowledge and understanding. *Water Resour. Res.* **2015**, *51* (7), 4861–4902.
- (74) Zahasky, C.; Benson, S. M. Preferential Solute Transport in Low Permeability Zones During Spontaneous Imbibition in Heterogeneous Porous Media. *Water Resour. Res.* **2022**, *58* (1), No. e2020WR029460.
- (75) Wei, K.-H.; Ma, J.; Xi, B.-D.; Yu, M.-D.; Cui, J.; Chen, B.-L.; Li, Y.; Gu, Q.-B.; He, X.-S. Recent progress on in-situ chemical oxidation for the remediation of petroleum contaminated soil and groundwater. *Journal of Hazardous Materials* **2022**, *432*, No. 128738.
- (76) Yukselen-Aksoy, Y.; Reddy, K. R. Effect of soil composition on electrokinetically enhanced persulfate oxidation of polychlorobiphenyls. *Electrochim. Acta* **2012**, *86*, 164–169.
- (77) Pham, A. L.-T.; Doyle, F. M.; Sedlak, D. L. Inhibitory Effect of Dissolved Silica on H₂O₂ Decomposition by Iron(III) and Manganese(IV) Oxides: Implications for H₂O₂-Based In Situ Chemical Oxidation. *Environ. Sci. Technol.* **2012**, *46* (2), 1055–1062.
- (78) Baciocchi, R.; Boni, M. R.; D'Aprile, L. Application of H₂O₂ lifetime as an indicator of TCE Fenton-like oxidation in soils. *Journal of Hazardous Materials* **2004**, *107* (3), 97–102.
- (79) Sra, K. S.; Thomson, N. R.; Barker, J. F. Persistence of Persulfate in Uncontaminated Aquifer Materials. *Environ. Sci. Technol.* **2010**, *44* (8), 3098–3104.
- (80) Oliveira, F. C.; Freitas, J. G.; Furquim, S. A. C.; Rollo, R. M.; Thomson, N. R.; Alleoni, L. R. F.; Nascimento, C. A. O. Persulfate Interaction with Tropical Soils. *Water, Air, & Soil Pollution* **2016**, *227* (9), 343.
- (81) Fang, G.; Chen, X.; Wu, W.; Liu, C.; Dionysiou, D. D.; Fan, T.; Wang, Y.; Zhu, C.; Zhou, D. Mechanisms of Interaction between Persulfate and Soil Constituents: Activation, Free Radical Formation, Conversion, and Identification. *Environ. Sci. Technol.* **2018**, *52* (24), 14352–14361.
- (82) Sra, K. S.; Thomson, N. R.; Barker, J. F. Stability of Activated Persulfate in the Presence of Aquifer Solids. *Soil Sediment Contam* **2014**, *23* (8), 820–837.
- (83) Lebron, C. A.; Heron, G.; Kueper, B.; LaChance, J.; Leeson, A.; Brogan, D.; Nielsen, S. G.; Phelan, D.; Rodriguez, D. *Dense non aqueous phase liquid (DNAPL) removal from fractured rock using thermal conductive heating (TCH)*; Environmental Security Technology Certification Program: Alexandria, VA, 2013.
- (84) Barber, L. B. Sorption of Chlorobenzenes to Cape Cod Aquifer Sediments. *Environ. Sci. Technol.* **1994**, *28* (5), 890–897.
- (85) Steventon-Barnes, H. *Solid organic matter in UK aquifers: its role in sorption of organic contaminants*; University of London, University College London: United Kingdom, 2000.
- (86) Kaur, K.; Crimi, M. Release of Chromium from Soils with Persulfate Chemical Oxidation. *Groundwater* **2014**, *52* (5), 748–755.
- (87) Goetschl, K. E.; Purgstaller, B.; Dietzel, M.; Mavromatis, V. Effect of sulfate on magnesium incorporation in low-magnesium calcite. *Geochim. Cosmochim. Acta* **2019**, *265*, 505–519.
- (88) Honetschlägerová, L.; Martinec, M.; Škarohlid, R. Coupling in situ chemical oxidation with bioremediation of chloroethenes: a review. *Reviews in Environmental Science and Bio/Technology* **2019**, *18* (4), 699–714.
- (89) McGachy, L.; Škarohlid, R.; Martinec, M.; Roskova, Z.; Smrhova, T.; Strejcek, M.; Uhlík, O.; Marek, J. Effect of chelated iron activated peroxydisulfate oxidation on perchloroethene-degrading microbial consortium. *Chemosphere* **2021**, *266*, No. 128928.
- (90) Sutton, N. B.; Atashgahi, S.; van der Wal, J.; Wijn, G.; Grotenhuis, T.; Smidt, H.; Rijnaarts, H. H. M. Microbial Dynamics During and After

In Situ Chemical Oxidation of Chlorinated Solvents. *Groundwater* **2015**, *53* (2), 261–270.

(91) Chang, Y.-C.; Chen, T.-Y.; Tsai, Y.-P.; Chen, K.-F. Remediation of trichloroethene (TCE)-contaminated groundwater by persulfate oxidation: a field-scale study. *RSC Adv.* **2018**, *8* (5), 2433–2440.

(92) Kastner, J. R.; Domingo, J. S.; Denham, M.; Molina, M.; Brigmon, R. Effect of Chemical Oxidation on Subsurface Microbiology and Trichloroethene (TCE) Biodegradation. *Bioremediation Journal* **2000**, *4* (3), 219–236.

(93) Sercu, B.; Jones, A. D. G.; Wu, C. H.; Escobar, M. H.; Serlin, C. L.; Knapp, T. A.; Andersen, G. L.; Holden, P. A. The Influence of In Situ Chemical Oxidation on Microbial Community Composition in Groundwater Contaminated with Chlorinated Solvents. *Microbial Ecology* **2013**, *65* (1), 39–49.

(94) Azadpour-Keeley, A.; Wood, L. A.; Lee, T. R.; Mravik, S. C. Microbial responses to in situ chemical oxidation, six-phase heating, and steam injection remediation technologies in groundwater. *Remediation Journal: The Journal of Environmental Cleanup Costs, Technologies & Techniques* **2004**, *14* (4), 5–17.

(95) Dugan, P. J.; Siegrist, R. L.; Crimi, M. L. Coupling surfactants/cosolvents with oxidants for enhanced DNAPL removal: A review. *Remediation Journal* **2010**, *20* (3), 27–49.

(96) Teel, A. L.; Cutler, L. M.; Watts, R. J. Effect of sorption on contaminant oxidation in activated persulfate systems. *Journal of Environmental Science and Health, Part A* **2009**, *44* (11), 1098–1103.

(97) Sedlak, D. L.; Andren, A. W. The effect of sorption on the oxidation of polychlorinated biphenyls (PCBs) by hydroxyl radical. *Water Res.* **1994**, *28* (5), 1207–1215.

(98) Karickhoff, S. W.; Brown, D. S.; Scott, T. A. Sorption of hydrophobic pollutants on natural sediments. *Water Res.* **1979**, *13* (3), 241–248.

(99) Chiou, C. T.; Peters, L. J.; Freed, V. H. A Physical Concept of Soil-Water Equilibria for Nonionic Organic Compounds. *Science* **1979**, *206* (4420), 831–832.

(100) Hiller, E.; Jurkovič, L.; Bartal, M. Effect of temperature on the distribution of polycyclic aromatic hydrocarbons in soil and sediment. *Soil and Water Research* **2008**, *3* (4), 231–240.

(101) Tsai, T. T.; Kao, C. M.; Hong, A. Treatment of tetrachloroethylene-contaminated groundwater by surfactant-enhanced persulfate/BOF slag oxidation—A laboratory feasibility study. *Journal of Hazardous Materials* **2009**, *171* (1), 571–576.

(102) Pan, L. W.; Siegrist, R. L.; Crimi, M. Effects of In Situ Remediation Using Oxidants or Surfactants on Subsurface Organic Matter and Sorption of Trichloroethene. *Groundwater Monitoring & Remediation* **2012**, *32* (2), 96–105.

(103) Zheng, F.; Gao, B.; Sun, Y.; Shi, X.; Xu, H.; Wu, J.; Gao, Y. Removal of tetrachloroethylene from homogeneous and heterogeneous porous media: Combined effects of surfactant solubilization and oxidant degradation. *Chemical Engineering Journal* **2016**, *283*, 595–603.

(104) Peluffo, M.; Pardo, F.; Santos, A.; Romero, A. Use of different kinds of persulfate activation with iron for the remediation of a PAH-contaminated soil. *Science of The Total Environment* **2016**, *563*–564, 649–656.

(105) Wang, L.; Wu, H.; Deng, D. Role of surfactants in accelerating or retarding persulfate decomposition. *Chemical Engineering Journal* **2020**, *384*, No. 123303.

(106) Wang, L.; Peng, L.; Xie, L.; Deng, P.; Deng, D. Compatibility of Surfactants and Thermally Activated Persulfate for Enhanced Subsurface Remediation. *Environ. Sci. Technol.* **2017**, *51* (12), 7055–7064.

(107) Liou, S.-Y.; Dodd, M. C. Evaluation of hydroxyl radical and reactive chlorine species generation from the superoxide/hypochlorous acid reaction as the basis for a novel advanced oxidation process. *Water Res.* **2021**, *200*, No. 117142.

(108) Lutze, H. V.; Kerlin, N.; Schmidt, T. C. Sulfate radical-based water treatment in presence of chloride: Formation of chlorate, interconversion of sulfate radicals into hydroxyl radicals and influence of bicarbonate. *Water Res.* **2015**, *72*, 349–360.

(109) Huie, R. E.; Clifton, C. L. Rate constants for hydrogen abstraction reactions of the sulfate radical, SO₄^{•−}. Alkanes and ethers. *International Journal of Chemical Kinetics* **1989**, *21* (8), 611–619.

(110) McElroy, W. J. A laser photolysis study of the reaction of sulfate(1[−]) with chloride and the subsequent decay of chlorine(1[−]) in aqueous solution. *J. Phys. Chem.* **1990**, *94* (6), 2435–2441.

(111) Liang, C.; Wang, Z.-S.; Mohanty, N. Influences of carbonate and chloride ions on persulfate oxidation of trichloroethylene at 20 °C. *Science of The Total Environment* **2006**, *370* (2), 271–277.

(112) Mullaney, J. R.; Lorenz, D. L.; Arntson, A. D. *Chloride in groundwater and surface water in areas underlain by the glacial aquifer system, northern United States*; U.S. Geological Survey: Reston, VA, 2009; Vol. 2009.

(113) Saha, S.; Reza, A. H. M. S.; Roy, M. K. Hydrochemical evaluation of groundwater quality of the Tista floodplain, Rangpur, Bangladesh. *Applied Water Science* **2019**, *9* (8), 198.

(114) Peyton, G. R. The free-radical chemistry of persulfate-based total organic carbon analyzers. *Marine Chemistry* **1993**, *41* (1), 91–103.

(115) Schwarzenbach, R. P.; Gschwend, P. M.; Imboden, D. M. *Environmental organic chemistry*; John Wiley & Sons: 2016.

(116) Al-Rawajfeh, A. E.; Al-Amairah, M. N. The influence of CO₂ injection on the carbonate chemistry and scaling in multiple-effect distillers. *Desalin Water Treat* **2009**, *7* (1–3), 191–197.

(117) Lei, X.; Lei, Y.; Guan, J.; Westerhoff, P.; Yang, X. Kinetics and Transformations of Diverse Dissolved Organic Matter Fractions with Sulfate Radicals. *Environ. Sci. Technol.* **2022**, *56* (7), 4457–4466.

(118) Goody, D.; Hinsby, K. Organic quality of groundwaters. *Natural groundwater quality* **2008**, 59–70.

(119) Huie, R. E.; Clifton, C. L.; Neta, P. Electron transfer reaction rates and equilibria of the carbonate and sulfate radical anions. *International Journal of Radiation Applications and Instrumentation. Part C. Radiation Physics and Chemistry* **1991**, *38* (5), 477–481.

(120) Buxton, G. V.; Bydder, M.; Salmon, G. A.; Williams, J. E. The reactivity of chlorine atoms in aqueous solution. Part III. The reactions of Cl• with solutes. *Phys. Chem. Chem. Phys.* **2000**, *2* (2), 237–245.

(121) Buxton, G. V.; Bydder, M.; Salmon, G. A. Reactivity of chlorine atoms in aqueous solution Part I The equilibrium ClMNSbd+Cl-Cl2. *Journal of the Chemical Society, Faraday Transactions* **1998**, *94* (5), 653–657.

(122) Ross, A. B.; Neta, P. *Rate Constants for Reactions of Inorganic Radicals in Aqueous Solution*; American Institute of Physics: 1979.

(123) Elbs, K. Ueber Nitrohydrochinon. *Journal für Praktische Chemie* **1893**, *48* (1), 179–185.

(124) Boyland, E.; Manson, D.; Sims, P. 729. The preparation of o-aminophenyl sulphates. *Journal of the Chemical Society (Resumed)* **1953**, *0*, 3623–3628.

(125) Behrman, E. J. Comment on “Revised Mechanism of Boyland–Sims Oxidation. *J. Phys. Chem. A* **2011**, *115* (26), 7863–7864.

(126) Marjanović, B.; Juranić, I.; Ćirić-Marjanović, G. Revised Mechanism of Boyland–Sims Oxidation. *J. Phys. Chem. A* **2011**, *115* (15), 3536–3550.

(127) Surwade, S. P.; Dua, V.; Manohar, N.; Manohar, S. K.; Beck, E.; Ferraris, J. P. Oligoaniline intermediates in the aniline-peroxydisulfate system. *Synth. Met.* **2009**, *159* (5), 445–455.

(128) Wang, J.; Wang, S. Reactive species in advanced oxidation processes: Formation, identification and reaction mechanism. *Chemical Engineering Journal* **2020**, *401*, No. 126158.

(129) Luo, S.; Wei, Z.; Dionysiou, D. D.; Spinney, R.; Hu, W.-P.; Chai, L.; Yang, Z.; Ye, T.; Xiao, R. Mechanistic insight into reactivity of sulfate radical with aromatic contaminants through single-electron transfer pathway. *Chemical Engineering Journal* **2017**, *327*, 1056–1065.

(130) Olmez-Hanci, T.; Arslan-Alaton, I. Comparison of sulfate and hydroxyl radical based advanced oxidation of phenol. *Chemical Engineering Journal* **2013**, *224*, 10–16.

(131) Mora, V. C.; Rosso, J. A.; Mártire, D. O.; Gonzalez, M. C. Phenol depletion by thermally activated peroxydisulfate at 70 °C. *Chemosphere* **2011**, *84* (9), 1270–1275.

(132) Pan, X.-M.; Schuchmann, M. N.; von Sonntag, C. Oxidation of benzene by the OH radical. A product and pulse radiolysis study in

oxygenated aqueous solution. *Journal of the Chemical Society, Perkin Transactions 2* **1993**, No. 3, 289–297.

(133) Pari, S.; Wang, I. A.; Liu, H.; Wong, B. M. Sulfate radical oxidation of aromatic contaminants: a detailed assessment of density functional theory and high-level quantum chemical methods. *Environmental Science: Processes & Impacts* **2017**, *19* (3), 395–404.

(134) Liu, H.; Bruton, T. A.; Li, W.; Buren, J. V.; Prasse, C.; Doyle, F. M.; Sedlak, D. L. Oxidation of Benzene by Persulfate in the Presence of Fe(III)- and Mn(IV)-Containing Oxides: Stoichiometric Efficiency and Transformation Products. *Environ. Sci. Technol.* **2016**, *50* (2), 890–898.

(135) Arslan-Alaton, I.; Aytac, E.; Kusk, K. O. Effect of Fenton treatment on the aquatic toxicity of bisphenol A in different water matrices. *Environmental Science and Pollution Research* **2014**, *21* (21), 12122–12128.

(136) Li, Y.-T.; Zhang, J.-J.; Li, Y.-H.; Chen, J.-L.; Du, W.-Y. Treatment of soil contaminated with petroleum hydrocarbons using activated persulfate oxidation, ultrasound, and heat: A kinetic and thermodynamic study. *Chemical Engineering Journal* **2022**, 428, No. 131336.

(137) Van Buren, J.; Prasse, C.; Marron, E. L.; Skeel, B.; Sedlak, D. L. Ring-Cleavage Products Produced during the Initial Phase of Oxidative Treatment of Alkyl-Substituted Aromatic Compounds. *Environ. Sci. Technol.* **2020**, *54* (13), 8352–8361.

(138) Pan, X.-M.; Sonntag, C. v. Notizen: OH-Radical-Induced Oxidation of Benzene in the Presence of Oxygen: $R+O_2 \rightleftharpoons RO_2$. Equilibria in Aqueous Solution. A Pulse Radiolysis Study. *Zeitschrift für Naturforschung B* **1990**, *45* (9), 1337–1340.

(139) Fang, X.; Pan, X.; Rahmann, A.; Schuchmann, H.-P.; von Sonntag, C. Reversibility in the Reaction of Cyclohexadienyl Radicals with Oxygen in Aqueous Solution. *Chemistry A European Journal* **1995**, *1* (7), 423–429.

(140) Dorfman, L. M.; Taub, I. A.; Bühler, R. E. Pulse Radiolysis Studies. I. Transient Spectra and Reaction-Rate Constants in Irradiated Aqueous Solutions of Benzene. *J. Chem. Phys.* **1962**, *36* (11), 3051–3061.

(141) Caregnato, P.; David Gara, P. M.; Bosio, G. N.; Gonzalez, M. C.; Russo, N.; Michelini, M. d. C.; Mártire, D. O. Theoretical and Experimental Investigation on the Oxidation of Gallic Acid by Sulfate Radical Anions. *J. Phys. Chem. A* **2008**, *112* (6), 1188–1194.

(142) Luo, S.; Wei, Z.; Spinney, R.; Villamena, F. A.; Dionysiou, D. D.; Chen, D.; Tang, C.-J.; Chai, L.; Xiao, R. Quantitative structure–activity relationships for reactivities of sulfate and hydroxyl radicals with aromatic contaminants through single–electron transfer pathway. *Journal of Hazardous Materials* **2018**, *344*, 1165–1173.

(143) Swartz, C. H.; Reddy, S.; Benotti, M. J.; Yin, H.; Barber, L. B.; Brownawell, B. J.; Rudel, R. A. Steroid Estrogens, Nonylphenol Ethoxylate Metabolites, and Other Wastewater Contaminants in Groundwater Affected by a Residential Septic System on Cape Cod, MA. *Environ. Sci. Technol.* **2006**, *40* (16), 4894–4902.

(144) Van Buren, J.; Cuthbertson, A. A.; Ocasio, D.; Sedlak, D. L. Ubiquitous Production of Organosulfates during Treatment of Organic Contaminants with Sulfate Radicals. *Environmental Science & Technology Letters* **2021**, *8* (7), 574–580.

(145) Cunningham, J. A.; Rahme, H.; Hopkins, G. D.; Lebron, C.; Reinhard, M. Enhanced In Situ Bioremediation of BTEX-Contaminated Groundwater by Combined Injection of Nitrate and Sulfate. *Environ. Sci. Technol.* **2001**, *35* (8), 1663–1670.

(146) Chen, K. F.; Chang, Y. C.; Chiou, W. T. Remediation of diesel-contaminated soil using in situ chemical oxidation (ISCO) and the effects of common oxidants on the indigenous microbial community: a comparison study. *J. Chem. Technol. Biotechnol.* **2016**, *91* (6), 1877–1888.

(147) Lal, M.; Mönig, J.; Asmus, K.-D. Acid formation in the radical-mediated degradation of chlorinated ethanes in aqueous environment. A radiation chemical study. *Journal of the Chemical Society, Perkin Transactions 2* **1987**, No. 11, 1639–1644.

(148) Cao, J.; Zhang, W.-X.; Brown, D. G.; Sethi, D. Oxidation of Lindane with Fe(II)-Activated Sodium Persulfate. *Environmental Engineering Science* **2008**, *25* (2), 221–228.

(149) Minisci, F.; Citterio, A.; Giordano, C. Electron-transfer processes: peroxydisulfate, a useful and versatile reagent in organic chemistry. *Acc. Chem. Res.* **1983**, *16* (1), 27–32.

(150) Li, S.; Elliott, D. W.; Spear, S. T.; Ma, L.; Zhang, W.-X. Hexachlorocyclohexanes in the Environment: Mechanisms of Dechlorination. *Critical Reviews in Environmental Science and Technology* **2011**, *41* (19), 1747–1792.

(151) Zhu, C.; Zhu, F.; Liu, C.; Chen, N.; Zhou, D.; Fang, G.; Gao, J. Reductive Hexachloroethane Degradation by $S_2O_8^{\bullet-}$ with Thermal Activation of Persulfate under Anaerobic Conditions. *Environ. Sci. Technol.* **2018**, *52* (15), 8548–8557.

(152) Kim, T.-K.; Sedlak, D. L. Mineralization of a Fully Halogenated Organic Compound by Persulfate under Conditions Relevant to in Situ Reduction and Oxidation: Reduction of Hexachloroethane by Ethanol Addition Followed by Oxidation. *Environ. Sci. Technol.* **2023**, *57* (36), 13691–13698.

(153) Popov, E.; Mametkulyev, M.; Santoro, D.; Liberti, L.; Eloranta, J. Kinetics of UV–H₂O₂ Advanced Oxidation in the Presence of Alcohols: The Role of Carbon Centered Radicals. *Environ. Sci. Technol.* **2010**, *44* (20), 7827–7832.

(154) Bielski, B. H. J. Reevaluation of the Spectral and Kinetic Properties of HO₂ and O₂⁻ Free Radicals. *Photochem. Photobiol.* **1978**, *28* (4-5), 645–649.

(155) Bielski, B. H. J.; Allen, A. O. Mechanism of the disproportionation of superoxide radicals. *J. Phys. Chem.* **1977**, *81* (11), 1048–1050.

(156) Bielski, B. H. J.; Cabelli, D. E.; Arudi, R. L.; Ross, A. Reactivity of HO₂/O₂⁻ Radicals in Aqueous Solution. *J. Phys. Chem. Ref. Data* **1985**, *14*, 1041–1100.

(157) Butler, E. C.; Hayes, K. F. Effects of Solution Composition and pH on the Reductive Dechlorination of Hexachloroethane by Iron Sulfide. *Environ. Sci. Technol.* **1998**, *32* (9), 1276–1284.

(158) Vecitis, C. D.; Park, H.; Cheng, J.; Mader, B. T.; Hoffmann, M. R. Treatment technologies for aqueous perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA). *Frontiers of Environmental Science & Engineering in China* **2009**, *3* (2), 129–151.

(159) Bruton, T. A.; Sedlak, D. L. Treatment of perfluoroalkyl acids by heat-activated persulfate under conditions representative of in situ chemical oxidation. *Chemosphere* **2018**, *206*, 457–464.

(160) Wang, W.; Chen, M.; Wang, D.; Yan, M.; Liu, Z. Different activation methods in sulfate radical-based oxidation for organic pollutants degradation: Catalytic mechanism and toxicity assessment of degradation intermediates. *Science of The Total Environment* **2021**, 772, No. 145522.

(161) LoPachin, R. M.; Gavin, T. Molecular Mechanisms of Aldehyde Toxicity: A Chemical Perspective. *Chem. Res. Toxicol.* **2014**, *27* (7), 1081–1091.

(162) Gu, X.; Lu, S.; Li, L.; Qiu, Z.; Sui, Q.; Lin, K.; Luo, Q. Oxidation of 1,1,1-Trichloroethane Stimulated by Thermally Activated Persulfate. *Ind. Eng. Chem. Res.* **2011**, *50* (19), 11029–11036.

(163) Dekant, W.; Jean, P.; Arts, J. Evaluation of the carcinogenicity of dichloromethane in rats, mice, hamsters and humans. *Regul. Toxicol. Pharmacol.* **2021**, *120*, No. 104858.

(164) Guelfo, J. L.; Higgins, C. P. Subsurface Transport Potential of Perfluoroalkyl Acids at Aqueous Film-Forming Foam (AFFF)-Impacted Sites. *Environ. Sci. Technol.* **2013**, *47* (9), 4164–4171.

(165) Appleman, T. D.; Higgins, C. P.; Quiñones, O.; Vanderford, B. J.; Kolstad, C.; Zeigler-Holady, J. C.; Dickenson, E. R. V. Treatment of poly- and perfluoroalkyl substances in U.S. full-scale water treatment systems. *Water Res.* **2014**, *51*, 246–255.

(166) Zhang, K.; Parker, K. M. Halogen Radical Oxidants in Natural and Engineered Aquatic Systems. *Environ. Sci. Technol.* **2018**, *52* (17), 9579–9594.

(167) Korshin, G. V.; Benjamin, M. M.; Xiao, H.-B. Interactions of Chlorine with Natural Organic Matter and Formation of Intermediates:

- Evidence by Differential Spectroscopy. *Acta hydrochimica et hydrobiologica* **2001**, *28* (7), 378–384.
- (168) Alegre, M. L.; Geronés, M.; Rosso, J. A.; Bertolotti, S. G.; Braun, A. M.; Mártire, D. O.; Gonzalez, M. C. Kinetic Study of the Reactions of Chlorine Atoms and Cl₂•⁻ Radical Anions in Aqueous Solutions. 1. Reaction with Benzene. *J. Phys. Chem. A* **2000**, *104* (14), 3117–3125.
- (169) Lu, J.; Benjamin, M. M.; Korshin, G. V.; Gallard, H. Reactions of the Flavonoid Hesperetin with Chlorine: A Spectroscopic Study of the Reaction Pathways. *Environ. Sci. Technol.* **2004**, *38* (17), 4603–4611.
- (170) Lei, Y.-J.; Tian, Y.; Fang, C.; Zhan, W.; Duan, L.-C.; Zhang, J.; Zuo, W.; Kong, X.-W. Insights into the oxidation kinetics and mechanism of diesel hydrocarbons by ultrasound activated persulfate in a soil system. *Chemical Engineering Journal* **2019**, *378*, No. 122253.
- (171) Lei, Y.; Lei, X.; Westerhoff, P.; Zhang, X.; Yang, X. Reactivity of Chlorine Radicals (Cl• and Cl₂•⁻) with Dissolved Organic Matter and the Formation of Chlorinated Byproducts. *Environ. Sci. Technol.* **2021**, *55* (1), 689–699.
- (172) Kumari, M.; Gupta, S. K. Occurrence and Exposure to Trihalomethanes in Drinking Water: A Systematic Review and Meta-analysis. *Exposure and Health* **2022**, *14* (4), 915–939.
- (173) Pals, J. A.; Ang, J. K.; Wagner, E. D.; Plewa, M. J. Biological Mechanism for the Toxicity of Haloacetic Acid Drinking Water Disinfection Byproducts. *Environ. Sci. Technol.* **2011**, *45* (13), 5791–5797.
- (174) Bove, G. E.; Rogerson, P. A.; Vena, J. E. Case-Control Study of the Effects of Trihalomethanes on Urinary Bladder Cancer Risk. *Archives of Environmental & Occupational Health* **2007**, *62* (1), 39–47.
- (175) Rahman, M. B.; Driscoll, T.; Cowie, C.; Armstrong, B. K. Disinfection by-products in drinking water and colorectal cancer: a meta-analysis. *International Journal of Epidemiology* **2010**, *39* (3), 733–745.
- (176) Hou, S.; Ling, L.; Dionysiou, D. D.; Wang, Y.; Huang, J.; Guo, K.; Li, X.; Fang, J. Chlorate Formation Mechanism in the Presence of Sulfate Radical, Chloride, Bromide and Natural Organic Matter. *Environ. Sci. Technol.* **2018**, *52* (11), 6317–6325.
- (177) Ji, Y.; Wang, L.; Jiang, M.; Lu, J.; Ferronato, C.; Chovelon, J.-M. The role of nitrite in sulfate radical-based degradation of phenolic compounds: An unexpected nitration process relevant to groundwater remediation by in-situ chemical oxidation (ISCO). *Water Res.* **2017**, *123*, 249–257.
- (178) Yang, P.; Ji, Y.; Lu, J.; Huang, Q. Formation of Nitrophenolic Byproducts during Heat-Activated Peroxydisulfate Oxidation in the Presence of Natural Organic Matter and Nitrite. *Environ. Sci. Technol.* **2019**, *53* (8), 4255–4264.
- (179) Yang, P.; Qian, L.; Cheng, Y.; Ji, Y.; Lu, J.; Kong, D. Formation of nitrophenolic byproducts in soils subjected to sulfate radical oxidation. *Chemical Engineering Journal* **2021**, *403*, No. 126316.
- (180) Chu, L.; Cang, L.; Sun, Z.; Wang, X.; Chen, H.; Fang, G.; Gao, J. Mechanism of nitro-byproducts formation during persulfate-based electrokinetic in situ oxidation for remediation of anthracene contaminated soil. *Journal of Hazardous Materials* **2023**, *453*, No. 131396.
- (181) Yang, P.; Ji, Y.; Lu, J. Transformation of ammonium to nitrophenolic byproducts by sulfate radical oxidation. *Water Res.* **2021**, *202*, No. 117432.
- (182) Perrini, G.; Tomasello, M.; Librando, V.; Minniti, Z. Nitrated Polycyclic Aromatic Hydrocarbons in the Environment: Formation, Occurrences and Analysis. *Annali di Chimica* **2005**, *95* (7–8), 567–577.
- (183) Lewtas, J.; Nishioka, M. G. Nitroarenes: Their Detection, Mutagenicity and Occurrence in the Environment. In *Nitroarenes: Occurrence, Metabolism, and Biological Impact*; Howard, P. C., Hecht, S. S., Beland, F. A., Eds.; Springer US: Boston, MA, 1990; pp 61–72.
- (184) Pashin, Y. V.; Bakhitova, L. M. Mutagenic and carcinogenic properties of polycyclic aromatic hydrocarbons. *Environ. Health Persp* **1979**, *30*, 185–189.
- (185) Lee, Y.-Y.; Hsieh, Y.-K.; Huang, B.-W.; Mutuku, J. K.; Chang-Chien, G.-P.; Huang, S. An Overview: PAH and Nitro-PAH Emission from the Stationary Sources and their Transformations in the Atmosphere. *Aerosol and Air Quality Research* **2022**, *22* (7), No. 220164.
- (186) Kovacic, P.; Somanathan, R. Nitroaromatic compounds: Environmental toxicity, carcinogenicity, mutagenicity, therapy and mechanism. *Journal of Applied Toxicology* **2014**, *34* (8), 810–824.
- (187) Vanholder, R.; Schepers, E.; Pletinck, A.; Nagler, E. V.; Glorieux, G. The Uremic Toxicity of Indoxyl Sulfate and p-Cresyl Sulfate: A Systematic Review. *Journal of the American Society of Nephrology* **2014**, *25* (9), 1897–1907.
- (188) Zhu, C.; Huang, H.; Chen, Y. Recent advances in biological removal of nitroaromatics from wastewater. *Environ. Pollut.* **2022**, *307*, No. 119570.
- (189) Pimviriyakul, P.; Pholert, P.; Somjitt, S.; Choowongkamon, K. Role of conserved arginine in HadA monooxygenase for 4-nitrophenol and 4-chlorophenol detoxification. *Proteins: Struct., Funct., Bioinf.* **2022**, *90* (6), 1291–1302.
- (190) Kertesz, M. A. Riding the sulfur cycle – metabolism of sulfonates and sulfate esters in Gram-negative bacteria. *FEMS Microbiol Rev.* **2000**, *24* (2), 135–175.
- (191) Cambria, M. T.; Minniti, Z.; Librando, V.; Cambria, A. Degradation of Polycyclic Aromatic Hydrocarbons by *Rigidoporus lignosus* and its Laccase in the Presence of Redox Mediators. *Appl. Biochem. Biotechnol.* **2008**, *149* (1), 1–8.
- (192) Huang, W. Q.; Xiao, S.; Zhong, H.; Yan, M.; Yang, X. Activation of persulfates by carbonaceous materials: A review. *Chemical Engineering Journal* **2021**, *418*, 129297.
- (193) Wang, J.; Wang, S. Activation of persulfate (PS) and peroxymonosulfate (PMS) and application for the degradation of emerging contaminants. *Chemical Engineering Journal* **2018**, *334*, 1502–1517.
- (194) Guo, R.; Xi, B.; Hu, Y.; Yuan, Y.; Chen, H.; Gong, Y.; Ge, X.; Lv, N. Enhancing persulfate activation by carbon-based Mn-Fe nanoparticles for groundwater ISCO treatment: Efficiency, stability and mechanism. *Chemical Engineering Journal* **2023**, *476*, No. 146827.
- (195) Rayaroth, M. P.; Oh, D.; Lee, C.-S.; Kang, Y.-G.; Chang, Y.-S. In situ chemical oxidation of contaminated groundwater using a sulfidized nanoscale zerovalent iron-persulfate system: Insights from a box-type study. *Chemosphere* **2020**, *257*, No. 127117.
- (196) Karim, A. V.; Jiao, Y. L.; Zhou, M. H.; Nidheesh, P. V. Iron-based persulfate activation process for environmental decontamination in water and soil. *Chemosphere* **2021**, *265*, 129057.