

UC Irvine

UC Irvine Previously Published Works

Title

Towards an integrated view of HCN channel role in epilepsy

Permalink

<https://escholarship.org/uc/item/36b7s1p8>

Journal

Current Opinion in Neurobiology, 21(6)

ISSN

0959-4388

Authors

Noam, Yoav
Bernard, Christophe
Baram, Tallie Z

Publication Date

2011-12-01

DOI

10.1016/j.conb.2011.06.013

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Published in final edited form as:

Curr Opin Neurobiol. 2011 December ; 21(6): 873–879. doi:10.1016/j.conb.2011.06.013.

Towards an integrated view of HCN channel role in epilepsy

Yoav Noam^{1,2}, Christophe Bernard^{3,*}, and Tallie Z. Baram^{1,4,*}

¹Department of Pediatrics, University of California-Irvine, Irvine, CA 92697-4475, USA ²SILS-Center for Neuroscience, University of Amsterdam, Amsterdam, The Netherlands ³INSERM-U751, Université de la Méditerranée, 27, Bd Jean Moulin, 13005 Marseille, France ⁴Departments of Anatomy/Neurobiology and Neurology, University of California-Irvine, Irvine, CA 92697-4475, USA

Abstract

Epilepsy is the third most common brain disorder and affects millions of people. Epilepsy is characterized by the occurrence of spontaneous seizures, i.e., bursts of synchronous firing of large populations of neurons. These are believed to result from abnormal regulation of neuronal excitability that favors hypersynchrony. Among the intrinsic conductances that govern neuronal excitability, the hyperpolarization-activated current (I_h) plays complex and important roles in the fine-tuning of both cellular and network activity. Not surprisingly, *dys*regulation of I_h and/or of its conducting ion-channels (HCN) has been strongly implicated in various experimental models of epilepsy, as well as in human epilepsy. Here we provide an overview of recent findings on the distinct physiological roles played by I_h in specific contexts, and the cellular mechanisms that underlie these functions, including the subunit make-up of the channels. We further discuss current knowledge of dysregulation of I_h and HCN channels in epilepsy in light of the multifaceted functions of I_h in the brain.

The h-current (I_h): a versatile regulator of cellular and network excitability

Among the many ionic conductances that shape neuronal excitability, the hyperpolarization-activated current I_h has long been a subject of interest and debate among researchers. Unlike other voltage-gated currents, I_h is activated upon relative *hyper*-polarization of the cell membrane. Conducted by a mixed cationic current with reversal potential values around -30 mV, I_h activation counteracts the hyperpolarization that triggered it [1,2]. Because I_h is partially activated at physiological, “resting” conditions, it provides constant depolarization of the membrane potential [3-5]. Thus, I_h is thought to function as a stabilizing negative-feedback loop that responds to alterations in membrane potential [1,2]. Alongside its depolarizing role, I_h exerts a shunting effect on excitable cells: being open at subthreshold potentials, I_h reduces the input resistance of the membrane (R_{in}), thus dampening the ability of incoming inputs to alter membrane voltage.

The net effect of the depolarizing and shunting properties of I_h on excitability is combinatorial and depends on many factors (Figure 1), as illustrated by the roles of I_h in

© 2011 Elsevier Ltd. All rights reserved.

Corresponding authors: Baram, Tallie Z. (tallie@uci.edu, Tel:+1 949 824 1131, Fax: +1 949 824 1106) Bernard, Christophe (christophe.bernard@univmed.fr).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

CA1 hippocampal pyramidal neurons. In these neurons, the density of I_h along apical dendrites increases with the distance from the soma [6]. A physiological consequence of this heterogeneous distribution (mediated by the effect of I_h on the input resistance and membrane time constant) is that EPSP time course is increasingly shortened with the distance from the soma [6]. As a result, temporal summation of synaptic inputs at the soma is similar regardless of whether these inputs occur at proximal or distal sites [6] (but see [7]). Another physiological action of I_h , mediated by its tonic depolarizing effect, is to increase steady-state inactivation of low-threshold, voltage-gated calcium channels [8], and thus restrict the genesis of dendritic calcium spikes. The effects of I_h on excitability involve interactions with other intrinsic conductances: whereas I_h increases the peak voltage amplitude of *weak* EPSPs, it inhibits the peak amplitude of responses to *strong* stimuli, and the net effect on EPSPs depends upon an intricate interaction between the I_h -mediated tonic depolarization and the K^+ -mediated conductance, I_M [9]. Regulation of synaptic signaling by I_h is not limited to excitatory inputs. I_h suppresses inhibitory (GABA_A-mediated) post-synaptic potentials (IPSPs) through interactions with other active conductances and cellular passive electric properties [10,11]. The relative depolarization generated by I_h in distal dendrites can alter GABA_A signaling from shunting to hyperpolarizing, and thus shortens the time-window for coincidence-detection [12].

From the above, it is clear that I_h has multiple effects in regulating dendritic excitability, including through interactions with other intrinsic conductances. Yet how do these different effects of I_h on inhibitory, excitatory and intrinsic signals integrate in a physiological context? This question was recently addressed in subthalamic nucleus (STN) neurons [13]. In these neurons (which express different HCN channel subunits compared with CA1 pyramidal neurons), dendritic I_h was activated only upon strong hyperpolarizing input onto the dendrite, and thus served a homeostatic role in counteracting GABA_A-mediated signals. In an analogous manner to that described in CA1 pyramidal neurons [8], activation of I_h by inhibitory input facilitated steady-state inactivation of low-threshold voltage-gated calcium channels, leading to inhibition of dendritic calcium spikes. Whereas theoretically I_h could suppress also the temporal summation of *excitatory* inputs, the authors demonstrated that the same inhibitory input that activated I_h shunted the effects of I_h on EPSP summation, and concluded that, in STN neuronal dendrites, I_h selectively regulated inhibitory signaling [13].

I_h is not limited to somato-dendritic subcellular distribution, and presynaptic I_h has been reported in various classes of neurons [14-17]. The presynaptic functions of I_h in mammalian brain have remained elusive [1, 16], but, in axon terminals of layer 3 entorhinal cortex neurons, I_h -mediated depolarization was recently found to restrict the activity of T-type Ca^{2+} channels, leading to reduced calcium influx and inhibition of synaptic release [17]. It remains to be seen whether similar mechanisms exist in presynaptic terminals of *interneurons* [14,15] or whether I_h serves an opposite, facilitatory role in these neuronal populations [14].

In addition to its roles in dendritic integration, membrane potential stabilization and regulation of synaptic transmission, I_h is an important modulator of oscillatory activity at both cellular and network levels. At cellular levels, I_h is critical for theta resonance (the preferential response of a neuron to oscillating inputs at specific frequencies). The slow kinetics and sub-threshold activation of I_h enable this current to filter out inputs at low frequencies (<3 Hz). This high-pass filtering property, in combination with low-pass filtering properties provided by the membrane capacitance, render cells with a strong I_h particularly responsive to inputs in the theta range (3-12 Hz) [18,19], a property also found in specific types of interneurons [20]. In cell-types with high I_h density at distal dendrites, dendritic I_h -dependent resonance is more pronounced, and preferentially filters signals that propagate from the soma to dendrites [19]. The filtering properties of I_h may contribute to

regulating *in vivo* theta rhythms [4], and other network oscillations (e.g., delta and gamma) and rhythmic firing (reviewed by [1]).

I_h diversity originates from exquisite control of HCN channels at multiple levels and enables physiological plasticity

The functional diversity and multiple roles of I_h in different physiological settings (discussed above) require tight regulation of the pore-forming proteins that conduct I_h, namely, HCN channels. HCN channel regulation involves many processes that take place at three levels: (1) regulation of the biophysical properties of the channels (such as voltage-activation profile and kinetics); (2) regulation of the number of channels expressed on the plasma membrane; and (3) localization of the channels to distinct subcellular compartments.

A critical determinant of the biophysical properties of HCN channels is their molecular make-up. Four different channel isoforms exist (HCN1-4) that can assemble in different combinations to yield homo- or hetero-tetrameric complexes with different properties [1,2,21]. Indeed, variability in I_h properties among cell populations and during different developmental stages is often associated with distinct expression profiles of specific HCN channel isoforms [22-25]. The conductive properties of HCN channels are further diversified by discrete interactions with other molecules, notably cyclic AMP (cAMP): cAMP binds a sequence on the C terminus of the channel, and influences HCN channel function by accelerating its kinetics and shifting its voltage-activation curve to more depolarized values [1,2]. The sensitivity of HCN channels to cAMP is isoform-specific: HCN4 > HCN2 >> HCN1 [1,2]. Thus, isoform-specific interaction of HCN channels with regulating molecules provides another dimension to the diversity of I_h. HCN channel function is modified by other 'small' molecules such as PIP(2) [26,27], as well as by interacting/auxiliary proteins (reviewed in [28]). Finally, phosphorylation of different HCN channel isoforms at multiple sites influences their biophysical properties (e.g. [29-31]).

The *number* of surface-expressed HCN channels influences the magnitude and properties of I_h (via changes in relative abundance of the constituents of the channel). Mechanisms controlling the number of surface HCN channels act at different time-scales. At a relatively long time scale (hours to days), transcriptional regulation determines the total amount of HCN channel protein in the cell [1]. This regulation is isoform-specific, acts via specific transcription factors [32], and depends on network activity [33]. Post-translational mechanisms also modulate HCN channels. For example, these channels are heavily glycosylated in the mammalian brain [34,35] and glycosylation influences both the total number of HCN channels in the membrane and their heteromerization (and hence properties) in a subunit-specific manner [35,36]. Modulation of HCN channel surface expression at shorter time-scales can occur via local regulation of channel membrane insertion, internalization and recycling [37,38]. Importantly, the dynamics of HCN channel trafficking and surface expression are activity-dependent [38], with implications to neuroplasticity and disease. Auxiliary proteins interact with HCN channels to regulate their surface expression: a family of splice variants of TRIP8b (an auxiliary protein that interacts directly with HCN channels) can either up- or down-regulate HCN channel surface expression [39,40]. Other candidate auxiliary proteins have been implicated in surface regulation of HCN channels (reviewed in [28]).

The targeting of HCN channels to distinct sub-cellular domains influences their location-dependent roles in regulation of excitability. Distinct distribution patterns of HCN channels exist in somata, dendrites and axons of specific cell types and brain regions, as well as during development [16,17,23,41,42]. Whereas ample information exists on the distribution of HCN channels, little is known of the molecular mechanisms that underlie their targeting

to sub-cellular domains. Time-lapse live imaging in hippocampal neurons indicates that vesicular trafficking dynamics of HCN channels are isoform-specific [38], and neuronal activity influences both trafficking and long-term subcellular distribution [38,43]. Because the HCN-interacting protein TRIP8b colocalizes with HCN1 channels and the distribution of the protein is disrupted in HCN1 knockout mice, TRIP8b is a candidate for dendritic targeting of HCN channels [44,45]. In addition to a heterogeneous distribution in the different cell compartments, HCN channels appear to be differently distributed within the same brain region. Thus, I_h decreases along the dorsal-ventral axis in entorhinal cortex stellate cells, providing these neurons with different integrative properties [46,47].

Together, the diverse regulatory mechanisms described above enable exquisite functional plasticity of I_h , such as seen during development [16, 22] or when synaptic plasticity is triggered [18,38]. When the mechanisms of HCN channel plasticity are disrupted, the resulting dysregulation of I_h might contribute to neurological disorders including epilepsy.

(3) HCN channel dysregulation in epilepsy: the wrong amount in the wrong place, at the wrong time

Since the original implication of HCN channels in epilepsy in 2001 [48], many studies have linked these channels to the epileptogenic process. In resected hippocampi from patients with temporal-lobe (limbic) epilepsy, enhanced levels of HCN1 channel expression and dendritic localization were found in granule cells of the dentate gyrus [49], and recent work has identified a mutation in the *HCN2* gene and augmentation of I_h in patients with genetic epilepsy with febrile seizures plus (GEFS+) [50]. Deletion of the *HCN1* gene in mice results in increased excitability and seizure susceptibility [51,52], and reduction or deletion of the *HCN2* isoform leads to spontaneous ‘absence’ seizures [53,54].

In accord with the diverse regulatory mechanisms and versatile functions of I_h in the normal brain, the *dysregulation* of I_h and HCN channels in epilepsy is dynamic and intricate (Figure 2). HCN channel abnormalities in the epileptic brain can manifest as altered mRNA and protein expression [55-60], sub-cellular distribution [44,61] or biophysical properties [48,58,62]. The causal relationship between aberrant HCN channel regulation and the epileptic process is further complicated by the fact that alterations in HCN channel expression, localization and function vary across animal models of epilepsy. Both early and late changes affecting diverse isoforms in distinct spatial patterns have been reported. For example, in the pilocarpine model, progressive reduction in HCN1 and HCN2 protein levels results in diminished I_h amplitude in dendrites of CA1 pyramidal neurons, leading to the disruption of theta resonance [58,63]. In contrast, following hyperthermia-induced seizures, the same population of cells exhibit *enhanced* dendritic I_h accompanied by altered gating properties [48,62], likely mediated by isoform-specific transcriptional regulation and increased HCN1/HCN2 heteromerization [56]. These seemingly conflicting results from different animal models demonstrate that augmented I_h can be associated with both increased and decreased excitability, depending on the physiological context [64], and potential interaction with other conductances.

The *temporal* patterns of HCN dysregulation in the epileptic brain are complex, and both transient and long-lasting alterations in hippocampal HCN/ I_h have been reported in pilocarpine, KA and febrile seizures models [55,56,58,60,63]. Interestingly, alterations in I_h gating properties in CA1 pyramidal neurons were reported within hours following induction of seizure-like-activity *in vitro*, attributable to activation of the phosphatase calcineurin and inhibition of p38 MAPK [65].

Spatial selectivity of HCN-channel pathology further contributes to the complex involvement of I_h in epilepsy. In acquired hippocampal epilepsy, HCN channel expression levels vary across different hippocampal regions in isoform-specific and region-specific fashions [55,57]. Region-specific alterations have been found in the WAG/Rij rat model of genetic absence epilepsy, where *reduced* HCN1 channel expression in layer 5 cortical neurons was reported to increase excitability via enhanced calcium electrogenesis [66], while *increased* HCN1 levels in thalamocortical neurons impaired their firing pattern via reducing the responsiveness of I_h to cAMP [59]. Reduced response of I_h to cAMP (triggered by imbalance of HCN subunit expression) was found also in the GAERS rat model of absence epilepsy [25].

Epilepsy may also involve abnormal subcellular distribution of HCN channels. Dendritic HCN1 localization was augmented in hippocampi resected from patients with epilepsy [49]. In animal models, transient upregulation of HCN1 surface expression occurred in CA1 pyramidal neurons 1-2 days after an epilepsy-provoking insult, followed by a *downregulation* 4 weeks later with mislocalization of the channels from distal dendritic domains to somata [44]. Recent work in freeze-lesion models of cortical dysplasia-provoked epilepsy found reduced HCN1 channel presence in distal dendrites of layer 5 cortical neurons [61].

In summary, multiple mechanisms, including transcriptional control, trafficking and channel modification act at different temporal and spatial scales to modulate I_h in the epileptic brain. In general, such changes might contribute to epileptogenesis, i.e., ‘cause’ epilepsy, or be a result of the epilepsy. The occurrence of several of these HCN changes early after the insult that triggers epilepsy, and/ or prior to the onset of spontaneous seizures [32,55,66] suggests a causal role. To fully grasp the contribution of these alterations to the epileptogenic process, physiological, molecular and cellular approaches should be integrated (Box 1). Recent discoveries have greatly advanced our understanding of the protean roles that I_h plays in the normal brain in different contexts, and their underlying cellular mechanisms. These, and future studies, will ultimately lead to an integrated view on the role of HCN channel pathology in epilepsy, at levels of analysis spanning single molecules to cellular and network domains. This understanding, in turn, will promote the design of new strategies for better treatment of the epilepsies.

Acknowledgments

We thank Barbara Cartwright for editorial assistance. The authors are supported by NIH grants R37 NS35439 (TZB); T32 NS04550 (YN); INSERM, ANR ANTARES and MINOS grants (CB), and a grant from the Dutch Epilepsy Fund (NEF 08.01, YN).

References

- *1. Biel M, Wahl-Schott C, Michalakis S, Zong X. Hyperpolarization-activated cation channels: from genes to function. *Physiol Rev.* 2009; 89:847–885. A comprehensive review of HCN channel structure, function and regulation. [PubMed: 19584315]
2. Robinson RB, Siegelbaum SA. Hyperpolarization-activated cation currents: from molecules to physiological function. *Annual Review of Physiology.* 2003; 65:453–480.
3. Lupica CR, Bell JA, Hoffman AF, Watson PL. Contribution of the hyperpolarization-activated current (I_h) to membrane potential and GABA release in hippocampal interneurons. *J Neurophysiol.* 2001; 86:261–268. [PubMed: 11431507]
4. Nolan MF, Malleret G, Dudman JT, Buhl DL, Santoro B, Gibbs E, Vronskaya S, Buzsaki G, Siegelbaum SA, Kandel ER, Morozov A. A behavioral role for dendritic integration: HCN1 channels constrain spatial memory and plasticity at inputs to distal dendrites of CA1 pyramidal neurons. *Cell.* 2004; 119:719–732. [PubMed: 15550252]

5. Meuth SG, Kanyshkova T, Meuth P, Landgraf P, Munsch T, Ludwig A, Hofmann F, Pape HC, Budde T. Membrane resting potential of thalamocortical relay neurons is shaped by the interaction among TASK3 and HCN2 channels. *J of Neurophysiol.* 2006; 96:1517–1529. [PubMed: 16760342]
6. Magee JC. Dendritic Ih normalizes temporal summation in hippocampal CA1 neurons. *Nature Neuroscience.* 1999; 2:508–514.
7. Angelo K, London M, Christensen SR, Hausser M. Local and global effects of I(h) distribution in dendrites of mammalian neurons. *J Neurosci.* 2007; 27:8643–8653. [PubMed: 17687042]
8. Tsay D, Dudman JT, Siegelbaum SA. HCN1 channels constrain synaptically evoked Ca²⁺ spikes in distal dendrites of CA1 pyramidal neurons. *Neuron.* 2007; 56:1076–1089. [PubMed: 18093528]
- **9. George MS, Abbott LF, Siegelbaum SA. HCN hyperpolarization-activated cation channels inhibit EPSPs by interactions with M-type K(+) channels. *Nat Neurosci.* 2009; 12:577–584. Using experimental and modeling approaches, the authors assess the relative contribution of I_h-mediated depolarization and shunting to dendritic excitability, and the important interplay between HCN channels and other dendritic channels. [PubMed: 19363490]
10. Hardie JB, Pearce RA. Active and passive membrane properties and intrinsic kinetics shape synaptic inhibition in hippocampal CA1 pyramidal neurons. *J Neurosci.* 2006; 26:8559–8569. [PubMed: 16914682]
11. Williams SR, Stuart GJ. Voltage- and site-dependent control of the somatic impact of dendritic IPSPs. *J Neurosci.* 2003; 23:7358–7367. [PubMed: 12917370]
12. Pavlov I, Scimemi A, Savtchenko L, Kullmann DM, Walker MC. I(h)-mediated depolarization enhances the temporal precision of neuronal integration. *Nat Commun.* 2011; 2:199. [PubMed: 21326231]
- *13. Atherton JF, Kitano K, Baufreton J, Fan K, Wokosin D, Tkatch T, Shigemoto R, Surmeier DJ, Bevan MD. Selective participation of somatodendritic HCN channels in inhibitory but not excitatory synaptic integration in neurons of the subthalamic nucleus. *J Neurosci.* 2010; 30:16025–16040. A comprehensive analysis of the role of HCN channel in regulating dendritic excitability in subthalamic nucleus neurons, showing complex, context-dependent roles mediated by interaction of I_h with GABA(A) receptors and low-threshold calcium channels. [PubMed: 21106841]
14. Aponte Y, Lien CC, Reisinger E, Jonas P. Hyperpolarization-activated cation channels in fast-spiking interneurons of rat hippocampus. *J Physiol.* 2006; 574:229–243. [PubMed: 16690716]
15. Lujan R, Albasanz JL, Shigemoto R, Juiz JM. Preferential localization of the hyperpolarization-activated cyclic nucleotide-gated cation channel subunit HCN1 in basket cell terminals of the rat cerebellum. *Eur J Neurosci.* 2005; 21:2073–2082. [PubMed: 15869503]
16. Bender RA, Kirschstein T, Kretz O, Brewster AL, Richichi C, Ruschenschmidt C, Shigemoto R, Beck H, Frotscher M, Baram TZ. Localization of HCN1 channels to presynaptic compartments: novel plasticity that may contribute to hippocampal maturation. *J Neurosci.* 2007; 27:4697–4706. [PubMed: 17460082]
- *17. Huang Z, Lujan R, Kadurin I, Uebele VN, Renger JJ, Dolphin AC, Shah MM. Presynaptic HCN1 channels regulate Ca(V)_{3.2} activity and neurotransmission at select cortical synapses. *Nat Neurosci.* 2011; 14:478–486. A pioneering study of the role of axonal HCN channels in synaptic transmission. [PubMed: 21358644]
18. Narayanan R, Johnston D. Long-term potentiation in rat hippocampal neurons is accompanied by spatially widespread changes in intrinsic oscillatory dynamics and excitability. *Neuron.* 2007; 56:1061–1075. [PubMed: 18093527]
19. Hu H, Vervaeke K, Graham LJ, Storm JF. Complementary theta resonance filtering by two spatially segregated mechanisms in CA1 hippocampal pyramidal neurons. *J Neurosci.* 2009; 29:14472–14483. [PubMed: 19923281]
20. Zemankovics R, Kali S, Paulsen O, Freund TF, Hajos N. Differences in subthreshold resonance of hippocampal pyramidal cells and interneurons: the role of h-current and passive membrane characteristics. *J Physiol.* 2010; 588:2109–2132. [PubMed: 20421280]
21. Santoro B, Baram TZ. The multiple personalities of h-channels. *Trends Neurosci.* 2003; 26:550–554. [PubMed: 14522148]

22. Bender RA, Baram TZ. Hyperpolarization activated cyclic-nucleotide gated (HCN) channels in developing neuronal networks. *Prog Neurobiol.* 2008; 86:129–140. [PubMed: 18834920]
23. Brewster AL, Chen Y, Bender RA, Yeh A, Shigemoto R, Baram TZ. Quantitative analysis and subcellular distribution of mRNA and protein expression of the hyperpolarization-activated cyclic nucleotide-gated channels throughout development in rat hippocampus. *Cereb Cortex.* 2007; 17:702–712. [PubMed: 16648453]
24. Kanyshkova T, Pawlowski M, Meuth P, Dube C, Bender RA, Brewster AL, Baumann A, Baram TZ, Pape HC, Budde T. Postnatal expression pattern of HCN channel isoforms in thalamic neurons: relationship to maturation of thalamocortical oscillations. *J Neurosci.* 2009; 29:8847–8857. [PubMed: 19587292]
25. Kuisle M, Wanaverbecq N, Brewster AL, Frere SG, Pinault D, Baram TZ, Luthi A. Functional stabilization of weakened thalamic pacemaker channel regulation in rat absence epilepsy. *J Physiol.* 2006; 575:83–100. [PubMed: 16728450]
26. Zolles G, Klocker N, Wenzel D, Weisser-Thomas J, Fleischmann BK, Roeper J, Fakler B. Pacemaking by HCN channels requires interaction with phosphoinositides. *Neuron.* 2006; 52:1027–1036. [PubMed: 17178405]
27. Pian P, Bucci A, Decostanzo A, Robinson RB, Siegelbaum SA. Modulation of cyclic nucleotide-regulated HCN channels by PIP(2) and receptors coupled to phospholipase C. *Pflugers Arch.* 2007; 455:125–45. [PubMed: 17605039]
28. Lewis AS, Estep CM, Chetkovich DM. The fast and slow ups and downs of HCN channel regulation. *Channels (Austin).* 2010; 4
29. Li CH, Zhang Q, Teng B, Mustafa SJ, Huang JY, Yu HG. Src tyrosine kinase alters gating of hyperpolarization-activated HCN4 pacemaker channel through Tyr531. *Am J Physiol Cell Physiol.* 2008; 294:C355–362. [PubMed: 17977941]
30. Liao Z, Lockhead D, Larson ED, Proenza C. Phosphorylation and modulation of hyperpolarization-activated HCN4 channels by protein kinase A in the mouse sinoatrial node. *J Gen Physiol.* 2010; 136:247–258. [PubMed: 20713547]
31. Hammelmann V, Zong X, Hofmann F, Michalakis S, Biel M. The cGMP-dependent protein kinase II is an inhibitory modulator of the hyperpolarization-activated HCN2 channel. *PLoS One.* 2011; 6:e17078. [PubMed: 21347269]
32. McClelland S, Flynn C, Dubé C, Richichi C, Zha QQ, Ghestem A, Esclapez M, Bernard C. NRSF-mediated HCN channelopathy in experimental temporal lobe epilepsy. *Ann Neurol.* 2011 in print.
33. Richichi C, Brewster AL, Bender RA, Simeone TA, Zha Q, Yin HZ, Weiss JH, Baram TZ. Mechanisms of seizure-induced ‘transcriptional channelopathy’ of hyperpolarization-activated cyclic nucleotide gated (HCN) channels. *Neurobiol Dis.* 2008; 29:297–305. [PubMed: 17964174]
34. Much B, Wahl-Schott C, Zong X, Schneider A, Baumann L, Moosmang S, Ludwig A, Biel M. Role of subunit heteromerization and N-linked glycosylation in the formation of functional hyperpolarization-activated cyclic nucleotide-gated channels. *J Biol Chem.* 2003; 278:43781–43786. [PubMed: 12928435]
35. Zha Q, Brewster AL, Richichi C, Bender RA, Baram TZ. Activity-dependent heteromerization of the hyperpolarization-activated, cyclic-nucleotide gated (HCN) channels: role of N-linked glycosylation. *J Neurochem.* 2008; 105:68–77. [PubMed: 17988239]
36. Hogle AP, Nazzari H, Roth A, Angoli D, Accili EA. Evolutionary emergence of N-glycosylation as a variable promoter of HCN channel surface expression. *Am J Physiol Cell Physiol.* 2010; 298:C1066–1076. [PubMed: 20130205]
37. Hardel N, Harmel N, Zolles G, Fakler B, Klocker N. Recycling endosomes supply cardiac pacemaker channels for regulated surface expression. *Cardiovasc Res.* 2008; 79:52–60. [PubMed: 18326556]
- *38. Noam Y, Zha Q, Phan L, Wu RL, Chetkovich DM, Wadman WJ, Baram TZ. Trafficking and surface expression of hyperpolarization-activated cyclic nucleotide-gated channels in hippocampal neurons. *J Biol Chem.* 2010; 285:14724–14736. The original characterization of HCN channel trafficking dynamics in live neurons, using time-lapse imaging techniques. Activity-dependent channel trafficking and membrane expression point out roles in plasticity and pathology. [PubMed: 20215108]

39. Lewis AS, Schwartz E, Chan CS, Noam Y, Shin M, Wadman WJ, Surmeier DJ, Baram TZ, Macdonald RL, Chetkovich DM. Alternatively spliced isoforms of TRIP8b differentially control h channel trafficking and function. *J Neurosci*. 2009; 29:6250–6265. [PubMed: 19439603]
40. Santoro B, Piskorowski RA, Pian P, Hu L, Liu H, Siegelbaum SA. TRIP8b splice variants form a family of auxiliary subunits that regulate gating and trafficking of HCN channels in the brain. *Neuron*. 2009; 62:802–813. [PubMed: 19555649]
41. Lorincz A, Notomi T, Tamas G, Shigemoto R, Nusser Z. Polarized and compartment-dependent distribution of HCN1 in pyramidal cell dendrites. *Nat Neurosci*. 2002; 5:1185–1193. [PubMed: 12389030]
42. Notomi T, Shigemoto R. Immunohistochemical localization of Ih channel subunits, HCN1–4, in the rat brain. *J Comp Neurol*. 2004; 471:241–276. [PubMed: 14991560]
43. Shin M, Chetkovich DM. Activity-dependent regulation of h channel distribution in hippocampal CA1 pyramidal neurons. *J Biol Chem*. 2007; 282:33168–33180. [PubMed: 17848552]
44. Shin M, Brager D, Jaramillo TC, Johnston D, Chetkovich DM. Mislocalization of h channel subunits underlies h channelopathy in temporal lobe epilepsy. *Neurobiol Dis*. 2008; 32:26–36. [PubMed: 18657617]
45. Santoro B, Wainger BJ, Siegelbaum SA. Regulation of HCN channel surface expression by a novel C-terminal protein-protein interaction. *J Neurosci*. 2004; 24:10750–10762. [PubMed: 15564593]
46. Giocomo LM, Zilli EA, Fransen E, Hasselmo ME. Temporal frequency of subthreshold oscillations scales with entorhinal grid cell field spacing. *Science*. 2007; 315:1719–1722. [PubMed: 17379810]
47. Nolan MF, Dudman JT, Dodson PD, Santoro B. HCN1 channels control resting and active integrative properties of stellate cells from layer II of the entorhinal cortex. *J Neurosci*. 2007; 27:12440–12451. [PubMed: 18003822]
48. Chen K, Aradi I, Thon N, Eghbal-Ahmadi M, Baram TZ, Soltesz I. Persistently modified h-channels alter complex febrile seizures convert the seizure-induced enhancement of inhibition to hyperexcitability. *Nat Med*. 2001; 7:331–337. [PubMed: 11231632]
49. Bender RA, Soleymani SV, Brewster AL, Nguyen ST, Beck H, Mathern GW, Baram TZ. Enhanced expression of a specific hyperpolarization-activated cyclic nucleotide-gated cation channel (HCN) in surviving dentate gyrus granule cells of human and experimental epileptic hippocampus. *J Neurosci*. 2003; 23:6826–6836. [PubMed: 12890777]
- *50. Dibbens LM, Reid CA, Hodgson B, Thomas EA, Phillips AM, Gazina E, Cromer BA, Clarke AL, Baram TZ, Scheffer IE, et al. Augmented currents of an HCN2 variant in patients with febrile seizure syndromes. *Ann Neurol*. 2010; 67:542–546. This study provides evidence for the possible involvement of HCN channel mutations in familial forms of human epilepsy. A triple amino acid deletion HCN2 gene was found in a subset of human subjects with febrile seizures-related syndromes and was associated with augmented I_h . [PubMed: 20437590]
51. Huang Z, Walker MC, Shah MM. Loss of dendritic HCN1 subunits enhances cortical excitability and epileptogenesis. *J Neurosci*. 2009; 29:10979–10988. [PubMed: 19726656]
52. Santoro B, Lee JY, Englot DJ, Gildersleeve S, Piskorowski RA, Siegelbaum SA, Winawer MR, Blumenfeld H. Increased seizure severity and seizure-related death in mice lacking HCN1 channels. *Epilepsia*. 2010; 51:1624–1627. [PubMed: 20384728]
53. Ludwig A, Budde T, Stieber J, Moosmang S, Wahl C, Holthoff K, Langebartels A, Wotjak C, Munsch T, Zong X, et al. Absence epilepsy and sinus dysrhythmia in mice lacking the pacemaker channel HCN2. *EMBO J*. 2003; 22:216–224. [PubMed: 12514127]
54. Chung WK, Shin M, Jaramillo TC, Leibel RL, LeDuc CA, Fischer SG, Tzilianos E, Gheith AA, Lewis AS, Chetkovich DM. Absence epilepsy in apathetic, a spontaneous mutant mouse lacking the h channel subunit, HCN2. *Neurobiol Dis*. 2009; 33:499–508. [PubMed: 19150498]
55. Brewster A, Bender RA, Chen Y, Dube C, Eghbal-Ahmadi M, Baram TZ. Developmental febrile seizures modulate hippocampal gene expression of hyperpolarization-activated channels in an isoform- and cell-specific manner. *J Neurosci*. 2002; 22:4591–4599. [PubMed: 12040066]
56. Brewster AL, Bernard JA, Gall CM, Baram TZ. Formation of heteromeric hyperpolarization-activated cyclic nucleotide-gated (HCN) channels in the hippocampus is regulated by developmental seizures. *Neurobiol Dis*. 2005; 19:200–207. [PubMed: 15837575]

57. Powell KL, Ng C, O'Brien TJ, Xu SH, Williams DA, Foote SJ, Reid CA. Decreases in HCN mRNA expression in the hippocampus after kindling and status epilepticus in adult rats. *Epilepsia*. 2008; 49:1686–1695. [PubMed: 18397293]
58. Jung S, Jones TD, Lugo JN Jr, Sheerin AH, Miller JW, D'Ambrosio R, Anderson AE, Poolos NP. Progressive dendritic HCN channelopathy during epileptogenesis in the rat pilocarpine model of epilepsy. *J Neurosci*. 2007; 27:13012–13021. [PubMed: 18032674]
59. Budde T, Caputi L, Kanyshkova T, Staak R, Abrahamczik C, Munsch T, Pape HC. Impaired regulation of thalamic pacemaker channels through an imbalance of subunit expression in absence epilepsy. *J Neurosci*. 2005; 25:9871–9882. [PubMed: 16251434]
60. Shah MM, Anderson AE, Leung V, Lin X, Johnston D. Seizure-induced plasticity of h channels in entorhinal cortical layer III pyramidal neurons. *Neuron*. 2004; 44:495–508. [PubMed: 15504329]
61. Hablitz JJ, Yang J. Abnormal pyramidal cell morphology and HCN channel expression in cortical dysplasia. *Epilepsia*. 2010; 51(Suppl 3):52–55. [PubMed: 20618401]
62. Dyhrfjeld-Johnsen J, Morgan RJ, Foldy C, Soltesz I. Upregulated H-current in hyperexcitable CA1 dendrites after febrile seizures. *Front Cell Neurosci*. 2008; 2:2. [PubMed: 18946517]
- *63. Marcelin B, Chauviere L, Becker A, Migliore M, Esclapez M, Bernard C. H Channel-Dependent Deficit of Theta Oscillation Resonance and Phase Shift in Temporal Lobe Epilepsy. *Neurobiol Dis*. 2009; 33:436–447. This study links I_h dysregulation and network excitability in epilepsy by demonstrating reduced hippocampal theta resonance due to down-regulation of HCN channel expression and I_h . [PubMed: 19135151]
- **64. Dyhrfjeld-Johnsen J, Morgan RJ, Soltesz I. Double Trouble? Potential for Hyperexcitability Following Both Channelopathic up- and Downregulation of $I(h)$ in Epilepsy. *Front Neurosci*. 2009; 3:25–33. A treatise on the possible implications of up- and down-regulation of I_h in epilepsy. The authors challenge unilateral views of I_h roles in epilepsy, and suggest a complex paradigm whereby either of these changes can result in hyperexcitability. [PubMed: 19753094]
65. Jung S, Bullis JB, Lau IH, Jones TD, Warner LN, Poolos NP. Downregulation of dendritic HCN channel gating in epilepsy is mediated by altered phosphorylation signaling. *J Neurosci*. 2010; 30:6678–6688. [PubMed: 20463230]
66. Kole MH, Brauer AU, Stuart GJ. Inherited cortical HCN1 channel loss amplifies dendritic calcium electrogenesis and burst firing in a rat absence epilepsy model. *J Physiol*. 2007; 578:507–525. [PubMed: 17095562]

BOX 1. Factors that influence the potential contribution of HCN channel dysregulation to hyperexcitability and epilepsy

- **Brain region and neuronal population:** In what cell populations are I_h /HCN channels altered? (Principal cells? Interneurons? Which brain region?); What is the specific role of I_h in this population and how does it influence network excitability?
- **Sub-cellular distribution:** Do changes in I_h /HCN channels take place in dendrites, soma, axon terminals?
- **Quantitative properties:** Is I_h upregulated or reduced?
- **Qualitative properties:** How is the molecular makeup of the channels altered? How are biophysical properties changed (voltage-activation, kinetics, cAMP gating)?
- **Temporal pattern:** Are changes in I_h dynamic or static? Activity dependent? Are they a result of other cellular and network changes?
- **Interplay with active and passive properties:** Are I_h /HCN changes accompanied by changes in other intrinsic conductances, synaptic inputs or the passive properties of the cell? Do changes in I_h lead to alterations of other channels?

Highlights

- The hyperpolarization-activated current I_h plays complex and important roles in the fine-tuning of both cellular and network activity
- The distinct physiological roles of I_h depend on factors including HCN subunit composition, interacting proteins and molecules, subcellular localization and the context of synaptic activity
- I_h dysregulation in epilepsy occurs at multiple regulatory levels and time frames, with complex effects on neuronal excitability
- A full understanding of the physiological and pathological roles of I_h will clarify the role of HCN channels as molecular targets in epilepsy therapy




Figure 1. The multiple functions of I_h on neuronal activity are context-and location driven

A schematized neuron depicting principal functions of I_h in distinct subcellular domains. In dendrites, I_h reduces dendritic summation (inset from [1], with permission); the current also represses dendritic calcium spikes (inset from [8], with permission), and converts the effect of IPSPs on the membrane potential from shunting to hyperpolarizing (inset modified from [12], with permission). In axon terminals, I_h reduces synaptic release through interaction with calcium channels (inset from [17], with permission). In both soma and dendrites, I_h augments theta-frequency resonance (inset modified from [19], with permission). In neurons that possess increased dendritic I_h density, the effects on theta-frequency resonance are more pronounced in dendrites compared to the soma.




Figure 2. Abnormal HCN channel regulation and function in epilepsy

Dysregulation of HCN channels/ I_h in epilepsy occurs at multiple levels. Seizure-induced alterations of the biophysical properties of I_h include either up- or down- regulation of current amplitude and modification of gating. The sub-cellular distribution of HCN channels along the somato-dendritic axis is altered both short-term, e.g., in hippocampal CA1 neurons in the pilocarpine model (insert modified, with permission, from [44]), and in dentate gyrus granule cells of human epileptic patients (insert from [49], with permission). Seizure-induced alterations in HCN channel mRNA and protein levels occur in many epilepsy models, and are both region- and channel isoform- specific (insert from [55], with permission). Altered synthesis of specific HCN channel isoforms may drive, at least in part, the increased, seizure-induced HCN1/2 heteromerization in hippocampus [55,56].