

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

THE EFFECT OF THE PHENYL SUBSTITUENT ON THE NITROGEN INVERSION BARRIER IN A 2-PHENYLOXAZIRIDINE

### Permalink

<https://escholarship.org/uc/item/36k8p1x8>

### Authors

Ono, Howard  
Splitter, Janet S.  
Calvin, Melvin.

### Publication Date

1973-08-01

THE EFFECT OF THE PHENYL SUBSTITUENT ON THE  
NITROGEN INVERSION BARRIER IN A  
2-PHENYLOXAZIRIDINE

Howard Ono, Janet S. Splitter, and Melvin Calvin


August 14, 1973

Prepared for the U. S. Atomic Energy Commission  
under Contract W-7405-ENG-48

RECEIVED  
LAWRENCE  
RADIATION LABORATORY

JAN 25 1974

LIBRARY AND  
DOCUMENTS SECTION



## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THE EFFECT OF THE PHENYL SUBSTITUENT ON THE NITROGEN INVERSION BARRIER  
IN A 2-PHENYLOXAZIRIDINE

Howard Ono<sup>1</sup>, Janet S. Splitter and Melvin Calvin

Department of Chemistry and the Laboratory of Chemical Biodynamics  
University of California, Berkeley, California, 94720

Comparisons of the pyramidal inversion barrier<sup>2</sup> with changes in substituent have recently been made for amines and arsines relative to phosphines<sup>3</sup> as shown in Figure 1. The slope of the correlation line was used as a measure of the sensitivity of the inverting center to substituent effects.<sup>3</sup> From data in the literature for aziridines,<sup>4</sup> in which the nitrogen atom is subjected to the barrier raising constraints of the three membered ring, a similar correlation line shows a much greater sensitivity to substituent than for the amines (Figure 1) and approximately the same sensitivity to substituent as for the phosphines. However, the aziridine inversion barriers are  $\sim 17$  kcal/mol less than for the phosphines.

The nitrogen inversion barriers of 2-methyl-3,3-disubstituted oxaziridines<sup>5</sup> are





Fig. 1. Correlation lines: -----, as determined in Ref. 3; ———, determined from data in Ref. 4 and this work (see text).

$\sim 14$  kcal/mol greater than those of 1-methyl-2,2-dimethylaziridines<sup>4d</sup> due to the barrier raising effect of the electronegative oxygen atom in the ring. No inversion data have been available for other kinds of substituents on the nitrogen atom in oxaziridines primarily because many of these oxaziridines undergo other thermal reactions. Until this time the known 2-aryloxaziridines were thermally too unstable to observe for nitrogen inversion.<sup>6</sup> We have now prepared 2,3-diphenyl-3-methyloxaziridine in predominantly the *cis* form and have observed its equilibration to 70% *trans*:30% *cis*-oxaziridine before rearrangement reactions occurred.<sup>7</sup>



The nmr spectrum of nitrene I<sup>8</sup> in carbon tetrachloride showed a single methyl resonance at  $\tau$  7.47 (3H,s) and aromatic proton resonances at  $\tau$  2.95 (10H,m), indicative of a single isomer, the cis-nitrene I.<sup>10</sup> A solution of 50 mg of I in 100 ml methylene chloride was irradiated for 4 minutes in a Pyrex immersion apparatus equipped with a 450 watt Hanovia medium pressure mercury lamp. Removal of the solvent in vacuo afforded an oil, II:  $\text{uv}_{\text{max}}^{\text{CH}_3\text{CN}}$ , end absorption at 230 nm ( $\epsilon$  9400). Its ir showed no carbonyl or imine bands and iodometric titration gave a 90% active oxygen content. Reflux of II in acetonitrile for 24 hours resulted in a 2:1 mixture of N-methylbenzanilide and acetyldiphenylamine. These properties were consistent with the structure of 2,3-diphenyl-3-methyloxaziridine for II.

An nmr spectrum of II in carbon tetrachloride taken as soon as possible after preparation showed two methyl resonances at  $\tau$  8.08 and  $\tau$  8.46 in a ratio of about 80%:20%.<sup>13</sup> Peak integration also showed the expected 10:3 ratio between aromatic and methyl hydrogens.

On standing at room temperature, the  $\tau$  8.08 resonance slowly decreased while the  $\tau$  8.46 resonance increased until an equilibrium ratio of 30%:70% was obtained. The ir, uv, and




Fig. 2. Arrhenius plot for the inversion of cis-II to trans-II in  $\text{CCl}_4$ .

active oxygen measurements remained unchanged during the isomerization. Oxaziridines prepared by the irradiation of the corresponding nitrones have been shown to have the same stereochemistry as the nitrene.<sup>11</sup> Therefore, the  $\tau$  8.08 resonance in the oxaziridine was assigned to cis-oxaziridine II and the  $\tau$  8.46 resonance to trans-oxaziridine II. This assignment is consistent with the more stable oxaziridine being the trans-oxaziridine.<sup>14</sup>

The first order rate constants for the isomerization (inversion) of cis-II to trans-II were determined at several temperatures.<sup>15</sup> In Figure 2 is shown an Arrhenius plot from which

$E_a = 23.2 \pm 1$  kcal/mol and  $\Delta S^\ddagger = -3 \pm 3$  eu were calculated. The  $\Delta S^\ddagger$  values for other pyramidal inversions have been found to be relatively small<sup>2,4</sup> and valid comparisons have been shown with  $\Delta H^\ddagger$  values when the same value of  $\Delta S^\ddagger$  was used for a series of the same type of compounds.<sup>16</sup> When  $\Delta S^\ddagger = 0$ , which is a reasonable assumption for pyramidal inversions,  $\Delta H^\ddagger = \Delta G^\ddagger$ .

The  $\Delta G^\ddagger$  of 23.4 kcal/mol for the inversion of oxaziridine II was calculated from the rate constants using the Eyring equation. From this value of  $\Delta G^\ddagger$  and the  $\Delta G^\ddagger$  values for 2-methyl-3,3-disubstituted-oxaziridines reported previously,<sup>5</sup> an approximate correlation line was drawn for the oxaziridines as shown in Figure 1. A comparison of the slopes shows that the oxaziridines have a greater sensitivity to substituent than the aziridines, which in turn have a greater sensitivity than the amines.

#### REFERENCES AND FOOTNOTES

1. Present address: Dept. of Chemistry, California State University, Fresno, Calif. 93710.
2. For reviews see: (a) A. Rauk, L.C. Allen and K. Mislow, *Angew.Chem., Int.Ed.Engl.*, 9, 400 (1970); (b) J.M. Lehn, *Fortschr.Chem.Forsch.*, 15, 311 (1970); (c) S.J. Brois, *Trans.N.Y.Acad. Sci.*, 31, 931 (1969); (d) J.B. Lambert, *Topics in Stereochem.* 6, 19 (1971).
3. R.D. Baechler, J.D. Andose, J. Stackhouse and K. Mislow, *J. Amer. Chem. Soc.*, 94, 8060 (1972).
4. Representative substituents were placed on the abscissa according to the magnitude of their calculated  $(CH_3)_2P$ -inversion barriers (slope = 1)<sup>3</sup>. The correlation line for the corresponding 1-substituted aziridines was drawn utilizing the following values for  $\Delta G^\ddagger$  (kcal/mol) adjusted to 25°C assuming  $\Delta S^\ddagger = 0$ . The aziridines were chosen, where possible, for structural similarities to 3,3-disubstituted oxaziridines. The solvents are given in parentheses. (a) 1-acetyl-, <5.6 (vinyl chloride); F.A. Anet and J.M. Osyany, *J. Amer. Chem. Soc.*, 89, 352 (1967). (b) 1-phenyl-2,2-dimethyl-, 11.1 ( $CF_2Cl_2$ ); J.D. Andose, J.M. Lehn, K. Mislow and J. Wagner, *J. Amer. Chem. Soc.*, 92, 4050 (1970). (c) 1-methylsulfonyl-2,2-dimethyl-, 13.2 ( $CDCl_3$ ); J.M. Lehn and J. Wagner, *Chem. Commun.*, 1298 (1968). (d) 1-methyl-2,2-dimethyl-, 17.6 (toluene); T. Drakenberg and J.M. Lehn, *J. Chem. Soc., Perkin II*, 532 (1972). (e) 7-chloro-7-azabicyclo[4.1.0]heptane, 23.6 ( $C_6H_6$ ); D. Felix and A. Eschenmoser, *Angew. Chem., Int. Ed. Engl.*, 7, 224 (1968).
5. (a)  $\Delta G^\ddagger_{25^\circ} = 32.2$  kcal/mol, calculated from rates given in F. Montanari, I. Moretti and G. Torre, *Chem. Commun.* 1086 (1969). (b)  $\Delta G^\ddagger_{25^\circ} = 32.2$  kcal/mol; A. Mammshreck, J. Linss and W. Seitz, *Ann.*, 727, 224 (1969).
6. J.S. Splitter and M. Calvin, *J. Org. Chem.*, 30, 3427 (1965).
7. Similar results were found with 3-(4-methoxyphenyl)-2,3-diphenyloxaziridine.
8. I was prepared by reacting acetophenone hydrazone (9a) first with silver oxide and then with nitrosobenzene in petroleum ether (9b) and collecting the resulting precipitate; mp 93-96°;

uv  $\text{Cl}_3\text{CN}$   
max 291 nm ( $\epsilon$  8,800); ms (70 ev) m/e (rel. int.) 211 ( $\text{M}^+$ , 27), 210(8) 195 (10), 180 (18), 91 (100), 77 (40). It was thermally unstable, hydrolyzing to acetophenone after several days at room temperature.

9. (a) G. Lock and K. Stach, Chem. Ber., 77B, 293 (1944); (b) H.G. Aurich, *ibid.*, 98, 3917 (1965).
10. The nmr spectra of trans- $\alpha$ -arylnitrones exhibit downfield shifts of the ortho protons. This characteristic is absent in the cis isomers<sup>11</sup> or when the ortho protons are substituted with methyl groups.<sup>12</sup>
11. J.S. Splitter, T. Su, H. Ono and M. Calvin, J. Amer. Chem. Soc., 93, 4075 (1971).
12. K. Koyano and H. Suzuki, Bull. Chem. Soc. Japan, 42, 3306 (1969).
13. Since considerable time was involved in the preparation of the nmr sample, the original ratio of the two isomers after irradiation is probably even greater than 4:1. In a series of isolable 2,3-diaryloxaziridines, nmr measurements in  $\text{CCl}_4$  at room temperature showed only a single  $\alpha$ -hydrogen resonance which remained unchanged with time: 2-(4-cyanophenyl)-3-phenyl oxaziridine,  $\tau$  5.55(s); 2-(4-nitrophenyl)-3-phenyloxaziridine,  $\tau$  5.37(s); 3-styryl-2-phenyloxaziridine,  $\tau$  5.70(d). This indicated either rapid nitrogen inversion or, more probably, the existence of a single isomeric species.
14. The downfield shift of the cis-3-methyl group is analogous to the downfield shifts of the cis-3-H relative to the trans-3-H-oxaziridines; D.M. Jerina, D.R. Boyd, L. Paolillo, and E.D. Becker, Tetrahedron Lett., 1483 (1970).
15. Kinetics were measured in a Varian A-60 nmr spectrometer equipped with a constant temperature probe. The 1° kinetics were done in an ice bath, the sample being transferred to the nmr spectrometer at various intervals. The rate constants ( $\text{sec}^{-1}$ ) found were: at 35°,  $13.6 \times 10^{-5}$ ; at 27°,  $4.37 \times 10^{-5}$ ; at 24°,  $3.80 \times 10^{-5}$ ; at 1°,  $0.15 \times 10^{-5}$ .
16. J. Lehn and J. Wagner, Tetrahedron, 26, 4227 (1970).

LEGAL NOTICE

*This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.*

TECHNICAL INFORMATION DIVISION  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720