

UCLA

UCLA Previously Published Works

Title

Optimal anesthetic conduct regarding immediate and short-term outcomes after liver transplantation - Systematic review of the literature and expert panel recommendations

Permalink

<https://escholarship.org/uc/item/3701w0kf>

Journal

Clinical Transplantation, 36(10)

ISSN

0902-0063

Authors

Chadha, Ryan
Patel, Dhupal
Bhangui, Pooja
et al.

Publication Date



2022-10-01

DOI

10.1111/ctr.14613

Peer reviewed

Optimal anesthetic conduct regarding immediate and short-term outcomes after liver transplantation – Systematic review of the literature and expert panel recommendations

Ryan Chadha¹  | Dhupal Patel² | Pooja Bhangui³ | Annabel Blasi⁴ | Victor Xia⁵  | Matteo Parotto⁶ | Christopher Wray⁵ | James Findlay⁷ | Michael Spiro^{8,10} | Dimitri Aristotle Raptis^{9,10} | the ERAS4OLT.org Working Group: Claus Niemann, San Francisco, CA, USA, Joerg-Matthias Pollok, London, UK, Marina Berenguer, Valencia, Spain, Pascale Tinguely, London, UK, Ka Hay Fan, London, UK, Giselle Farrer, London, UK.

¹Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Jacksonville, USA

²Department of Anesthesia and Intensive Care Medicine, Addenbrooke's Hospital, Cambridge, UK

³Department of Anesthesiology, Medanta Liver Institute, Gurgaon, India

⁴Department of Anesthesiology, Hospital Clinic BarcelonaInstitut d'Investigacio Biomèdica Pi I Suner (IDIBAPS), Spain

⁵Department of Anesthesiology, University of California, Los Angeles, USA

⁶Department of Anesthesiology and Interdepartmental Division of Critical Care Medicine, Toronto General Hospital, University of Toronto, Canada

⁷Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, USA

⁸Department of Anesthesia and Intensive Care Medicine, Royal Free Hospital, London, UK

⁹Clinical Service of HPB Surgery and Liver Transplantation, Royal Free Hospital, London, UK

¹⁰Division of Surgery & Interventional Science, University College London, UK

Correspondence

Ryan Chadha, Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, 4500 San Pablo Road, Jacksonville, FL 32224, USA.
 Email: chadha.ryan@mayo.edu

Funding information

International Liver Transplant Society (ILTS)

Abstract

Background: In the era of enhanced recovery after surgery, there is significant discussion regarding the impact of intraoperative anesthetic management on short-term outcomes following liver transplantation (LT), with no clear consensus in the literature.

Objectives: To identify whether or not intraoperative anesthetic management affects short-term outcomes after liver transplantation.

Data Sources: Ovid MEDLINE, Embase, Scopus, Google Scholar, and Cochrane Central.

Methods: A systematic review following PRISMA guidelines was undertaken. The systematic review was registered on PROSPERO (CRD42021239758). An international expert panel made recommendations for clinical practice using the GRADE approach.

Results: After screening, 14 studies were eligible for inclusion in this systematic review. Six were prospective randomized clinical trials, three were prospective nonrandomized clinical trials, and five were retrospective studies. These manuscripts were reviewed to look at five questions regarding anesthetic care and its impact on short term outcomes following liver transplant. After review of the literature, the quality of evidence according to the following outcomes was as follows: intraoperative and postoperative

morbidity and mortality (low), early allograft dysfunction (low), and hospital and ICU length of stay (moderate).

Conclusions: For optimal short term outcomes after liver transplantation, the panel recommends the use of volatile anesthetics in preference to total intravenous anesthesia (TIVA) (Level of Evidence: Very low; Strength of Recommendation: Weak) and minimum alveolar concentration (MAC) versus bispectral index (BIS) for depth of anesthesia monitoring (Level of Evidence: Very low; Strength of Recommendation: Weak). Regarding ventilation and oxygenation, the panel recommends a restrictive oxygenation strategy targeting a PaO₂ of 70–120 mmHg (10–14 kPa), a tidal volume of 6–8 ml/kg ideal body weight (IBW), administration of positive end expiratory pressure (PEEP) tailored to patient intraoperative physiology, and recruitment maneuvers. (Level of evidence: Very low; Strength of Recommendation: Strong). Finally, the panel recommends the routine use of antiemetic prophylaxis. (Level of evidence: low; Strength of Recommendation: Strong).

KEYWORDS

bispectral index (BIS), depth of anesthesia, liver transplantation, minimum alveolar concentration (MAC), oxygen management, positive end-expiratory pressure (PEEP), total intravenous anesthesia, ventilation, volatile or inhalational anesthesia

1 | INTRODUCTION

Enhanced recovery after surgery (ERAS) programs have been developed to improve patient outcomes and recovery following surgery through improvements in the quality of peri-operative care. Anesthetic conduct during the intraoperative period is a well-recognized component of ERAS pathways.¹ Several studies have focused on understanding the importance of different components of routine anesthetic care including type, depth and monitoring of anesthesia,^{2–4} and intraoperative mechanical ventilation settings.⁵ Reviews as well as specific recommendations have been published for certain types of abdominal surgery.^{6,7}

In this study, we provide an updated review of the literature focused on understanding the optimal anesthetic conduct regarding immediate and short-term outcomes specifically for patients undergoing liver transplantation (LT). In particular, we aim to answer the following questions:

1. Does the type of anesthesia (Total Intravenous Anesthetic (TIVA) vs. Volatile) affect immediate and short-term outcomes after liver transplantation?
2. Does the type of depth of anesthesia assessment (EEG based awareness monitoring such as Bispectral Index (BIS) vs. Minimum Alveolar Concentration (MAC)) affect immediate and short term outcomes after liver transplantation?
3. Does a liberal versus restrictive O₂ management affect immediate and short-term outcomes after liver transplantation?
4. Do intraoperative ventilation strategies (positive end-expiratory pressure (PEEP), tidal volumes, recruitment manoeuvres) affect immediate and short-term outcomes after liver transplantation?
5. Does antiemetic prophylaxis affect immediate and short-term outcomes after liver transplantation?

Based on the findings of this literature review, we provide expert panel recommendations for the optimal anesthetic conduct in patients undergoing liver transplantation.

This work was conducted in preparation for the ILTS – ERAS4OLT.org Consensus Conference on Enhanced Recovery for Liver Transplantation, January 2022, Valencia, Spain.

2 | METHODS

2.1 | Protocol and registration

The PROSPERO protocol ID is CRD42021239758.

2.2 | Eligibility criteria

The population being studied in this review was adults with end-stage liver disease who were listed for and underwent deceased donor orthotopic liver transplantation. Split-livers and patients undergoing living donor liver transplantation were excluded. The interventions and controls being studied include: TIVA versus volatile anesthesia, depth of anesthesia assessment using BIS versus MAC, liberal versus restrictive oxygen administration, use of intraoperative ventilatory strategies such as low tidal volume ventilation, PEEP, and recruitment manoeuvres, and use or avoidance of antiemetic prophylaxis.

2.3 | Information sources and search

The following bibliographic databases were searched: Ovid MEDLINE, Embase, Scopus, Google Scholar, Clinical.Trials.gov and the Cochrane

Central Register of Controlled Trials. Any study published prior to March 7, 2021 was eligible for inclusion. There were no language limitations. Studies reporting on pediatric populations as well as case reports or conference abstracts were excluded. The search strategy involved searching the terms liver or hepatic and transplantation with the following keywords: (1) total intravenous and anesthesia or anesthesia, (2) TIVA, propofol, isoflurane, sevoflurane, desflurane, fentanyl, remifentanyl, sufentanyl, volatile, inhalational (3) depth of anesthesia, bispectral index, BIS, minimum alveolar concentration, minimal alveolar concentration, and (4) oxygen management, ventilation, end-tidal concentration, positive end-expiratory pressure, PEEP. The search was performed by expert University of Zurich librarians and Junior Research Committee members.

2.4 | Study selection

Types of studies that were reviewed include both comparative and single cohort studies, retrospective or prospective, describing outcomes in patients that received different anesthetics conducts during liver transplantation.

2.5 | Quality of studies and recommendations grading

The “Grading of Recommendations Assessment, Development and Evaluation” (GRADE) approach was used for grading quality of evidence and strength of recommendations.⁸ The GRADE system was designed to provide a comprehensive and structured approach to rating the quality of evidence (QOE) for systematic reviews, and to grade the strength of recommendations for development of guidelines in health care. We applied the modified GRADE approach for QOE assessment derived from systematic reviews using estimates summarized narratively.⁹ The QOE was rated separately for each outcome. The direction and strength of recommendation was assessed individually by all authors and disagreements resolved by consensus.^{10,11}

3 | RESULTS

3.1 | Study selection

Initial search of the databases revealed 3374 records, which was reduced to 1670 after duplicates were removed. After screening for eligibility, 23 records remained, with nine excluded due to lack of outcome data, lack of data regarding operative data, studies related to living donors rather than recipients, or being a case presentation or abstract. This resulted in 14 records which were taken forward for analysis (Figure 1).

3.2 | Study characteristics

The characteristics of the included studies are available in Table 1.

3.3 | Results of individual studies

The outcomes of the included studies are listed in Table 2.

3.4 | Quality of evidence

After review of the literature, we assessed its quality based on the questions presented in the introductions with specific focus at the following short-term outcomes: intraoperative morbidity and mortality, postoperative morbidity and mortality, early allograft dysfunction, length of intensive care unit (ICU) and hospital stay.

The summary of findings for the main outcomes, including the quality of evidence (QOE) assessment according to the GRADE approach are summarized in Table 3.

3.5 | Recommendations

Despite moderate to low QOE across the main outcomes, the expert working group issued the following recommendations for optimal short-term outcomes following LT (Table 4 for evidence framework):

1. The use of volatile anesthesia with consideration of TIVA in patients at risk of, or with raised intracranial pressure.
2. The use of MAC rather than EEG based awareness monitoring such as BIS to measure depth of anesthesia during LT when using volatile agents.
3. A restrictive oxygen strategy titrated to PaO₂ 70–110 mmHg (10–14 kPa).
4. Ventilatory strategies: tidal volumes 6–8 ml/kg IBW, PEEP titrated to patient physiology, recruitment maneuvers.
5. The routine use of antiemetic prophylaxis.

4 | DISCUSSION

This systematic review focuses on establishing the literature and expert panel opinion supporting intraoperative anesthesia technique and its association with intermediate and short term outcomes after LT.

When comparing the use of TIVA versus volatile in LT, one study showed a faster wake-up time with Propofol versus Desflurane, but no other outcome impact.¹⁸ An additional study showed a lower




FIGURE 1 Study flow diagram

postreperfusion MDA (marker of ischemic reperfusion injury) levels in the Propofol versus the non-Propofol groups, but no other effects on hemodynamics.²³ Propofol may also be beneficial in the acute liver failure, acute on chronic liver failure, or other transplant populations at risk for increased ICP.³⁰ Conversely, the panel noted the impact of hepatic and renal function on Propofol metabolism. Although there are no studies specific to LT, the increased cost of certain volatile agents coupled with their negative environmental impact due to increased greenhouse gas emissions is well known.³¹

Based on this information reviewed, the panel can make a weak recommendation supporting the use of volatile anesthesia, with a secondary recommendation to consider total intravenous anesthetics in populations at risk for increased ICP. There is no recommendation regarding the use of a particular volatile anesthetic as the panel feels appropriate intraoperative utilization can be left to the provider.

In the comparison of EEG based monitors versus MAC as a marker of anesthetic depth, one paper showed no difference in incidence of post-

operative cognitive dysfunction when utilizing BIS versus control¹³; one reference assessed BIS versus MAC in patients with hepatic encephalopathy, however no difference in time to return of consciousness was found.²² While there are some referenced potential benefits to EEG based anesthetic depth monitoring in the surgical population²⁶ its practice may be limited by availability and cost internationally. It should be noted as well that EEG awareness monitoring is not a standard monitor as recommended by the American Society of Anesthesiologists.³² It is the opinion of this panel for a weak recommendation supporting MAC utilization over EEG based anesthetic depth monitoring.

There is also no evidence in the liver transplant literature reviewed regarding liberal versus restrictive O₂ use, specific ventilatory strategies, and antiemetic prophylaxis. However, the panel felt despite the lack of evidence, strong recommendations could be made in these specific domains based on established literature from other surgical arenas. Regarding the use of liberal versus restrictive O₂ use in

TABLE 1 Study characteristics

Author, year (Reference)	Study type	Number of participants inclusion and exclusion criteria	Main outcomes assessed
Najafi et al. (2019) ¹²	Single-center, randomized	78 Propofol: 39 Sevoflurane: 39	<ul style="list-style-type: none"> Hemodynamic parameters, coagulation, hepatic and renal function
Cao et al. (2017) ¹³	Combined prospective and retrospective case control	60 EEG based awareness monitoring (BIS): 33 Control: 27	<ul style="list-style-type: none"> incidence of POCD Day 7 post LT
Beck-Schimmer et al. (2015) ¹⁴	Multi-center, double-blinded, randomized	98 Propofol: 48 Sevoflurane: 50	<ul style="list-style-type: none"> Peak postop AST EAD In-hospital complications Hospital and ICU LOS Secondary outcomes: postop ALT, early allograft dysfunction (INR > / = 1.6, ALT > 2000, bilirubin > / = 10 mg/dl at day 7 post-LT), in-hospital complications (Clavien-Dindo complication score), ICU length of stay, hospital length of stay
Gucyetmez (2016) ¹⁵	Single center, randomized, blinded	70 Control: 35 Treatment group: 35	<ul style="list-style-type: none"> Postoperative tramadol requirements ICU and hospital LOS
Gajate Martin (2016) ¹⁶	Single center, retrospective	201 Propofol: 143 Sevoflurane: 58	<ul style="list-style-type: none"> Primary graft dysfunction
Ali Sahmeddini (2012) ¹⁷	Single center, randomized, blinded	40 Norepinephrine: 20 Acupuncture: 20	<ul style="list-style-type: none"> Haemodynamic parameters
Ali Sahmeddini, 2012	Randomized blinded unicentric 20 acupuncture 20 NE	40	MAP in hepatectomy-anhepatic-neohepatic
Lu (2016) ¹⁸	Single center, retrospective	111 total patients TIVA: 66 Desflurane: 45	<ul style="list-style-type: none"> Haemodynamic parameters Extubation time INR, ALT 24 h Hospital and ICU LOS.
Mangus (2018) ¹⁹	Single-center retrospective study	1291 total cases Isoflurane: 797 Desflurane: 102 Sevoflurane: 392	<ul style="list-style-type: none"> IRI ALT and bilirubin up to Day7 Peri-operative mortality and graft loss 1-year mortality ad graft survival EAD Hospital LOS
Toprak (2011) ²⁰	Single center, prospective, non-randomized,	50 patients	Primary outcome: End-tidal Isoflurane concentration at the 3 phases of LT <ul style="list-style-type: none"> Haemodynamic parameters
Pan (2016) ²¹	Single center, prospective, non-randomized, un-blinded	53 patients MELD < 9: 32 MELD > 10: 21	<ul style="list-style-type: none"> Haemodynamic parameters Haemodynamic instability

(Continues)

TABLE 1 (Continued)

Author, year (Reference)	Study type	Number of participants inclusion and exclusion criteria	Main outcomes assessed
Kim (2019) ²²	Single center, retrospective, observational	64 Non-severe HE, Grade 1-2): 26 Severe HE, Grade 3-4: 38	<ul style="list-style-type: none"> • Postop recovery of consciousness
Tsai (2012) ²³	Single center, prospective, randomized, unblinded	20 Control: 10 Propofol group: 10	<ul style="list-style-type: none"> • IRI
Sorensen (2014) ²⁴	Single center, retrospective	49	<ul style="list-style-type: none"> • Cerebral oxygenation and ET_{CO}₂ during LT
Jowkar (2020) ²⁵	Single center, randomized	100	<ul style="list-style-type: none"> • Changes in liver and renal biochemistry and coagulation from day 1–7 post LT • Extubation time • Hospital and ICU LOS

Abbreviations: ALP, alkaline phosphatase; ALT, alanine transaminase; AST, aspartate transaminase; BIS, bispectral index; BUN, blood urea nitrogen; CO, cardiac output; CVP, central venous pressure; EAD, early allograft dysfunction; ET_{CO}₂, end tidal carbon dioxide; HE: hepatic encephalopathy, GA: general anesthesia, HR: heart rate, INR: international normalized ratio, IRI: ischemic reperfusion injury; MAP, mean arterial pressure; MDA, malondialdehyde; MPAP, mean pulmonary artery pressure; NSE, neuron specific enolase; LOC, loss of consciousness; LOS, length of stay; LT, liver transplant; MELD, model for end stage liver disease; POCD, postoperative cognitive dysfunction; PT, prothrombin time; PTT partial thrombin time; S100B, calcium-binding protein B; TCI, target controlled infusion; TIVA, total intravenous anesthesia.

abdominal surgery, there are studies showing increased risk of mortality, increased cancer incidence, increased incidence of atelectasis, and increased cardiac injury markers with utilization of higher O₂ concentrations perioperatively.^{33–36} There appears to be minimal evidence for higher O₂ concentrations contributing to an increased risk of surgical site infections.^{37–39} From a resources standpoint, reducing the amount of oxygen needed per patient could lead to improved costs with reduced ventilator and ICU time in liver transplantation patients, with the caveat that data has been extrapolated. Looking at the harms of liberal O₂ and potential benefits of lower oxygen concentrations, the panel feels it can propose a strong recommendation for restrictive oxygen administration with a FiO₂ to maintain a PaO₂ of 70–110 mmHg (10–14 kPa).

Considering specific ventilatory strategies, there is significant literature supporting low-tidal volume ventilation in abdominal surgery in overall improved mortality and reduction in pulmonary complications.^{40–42} Similarly, there are multiple studies showing improved oxygenation and reduction in postoperative hypoxemic events when PEEP is utilized.^{43,44} While PEEP itself is recommended, the panel cannot recommend a specific PEEP value or range of values. Instead, intraoperative PEEP should be tailored and optimized to individual patient requirements dependent on their body habitus, oxygen requirements, and where possible flow volume loops. Along similar lines, the use of recruitment maneuvers has very little evidence in abdominal surgery and liver transplantation,⁴⁵ but their benefit is well proven to improve oxygenation and lung function,

particularly in the ICU population. As a result, the panel feels a strong recommendation can be proposed for the use of low tidal volume ventilation of 6–8 ml/kg ideal body weight (IBW), use of PEEP tailored to patient physiology, and recruitment maneuvers in liver transplantation.

Regarding antiemetic prophylaxis, there is well supported consensus opinion regarding the management of postoperative nausea and vomiting.⁴⁶ The utility of routine of antiemetic prophylaxis may be low given that a large number of centers do not routinely extubate in the operating room, but within a specified number of hours postoperatively. However, with increasing implementation of fast-track liver transplantation pathways and the favorable benefit to risk ratio of antiemetic prophylaxis (an important risk being the potential QT prolongation associated with most antiemetics, which may provide an issue in the setting of cardiomyopathy), the panel feels that a strong recommendation can be proposed for the use of antiemetic prophylaxis during liver transplantation.

5 | LIMITATIONS

This review has several limitations. Firstly, there is a distinct lack of evidence for intraoperative anesthetic care in liver transplantation and short term outcomes, therefore extrapolation from other high-quality studies not specific to liver transplantation as well as a heavy reliance

TABLE 2 Study outcomes

Author, year (Reference)	Intra-operative morbidity and mortality	Post-operative morbidity and mortality	Early allograft dysfunction	ICU and hospital length of stay
Najafi et al. (2019)	-	No significant difference between 2 groups in hemodynamic parameters, coagulation, hepatic and renal function	-	-
Cao et al. (2017)	EEG based awareness monitoring (BIS) guided anesthesia could avoid hypotension during surgery through reduced consumption of Propofol	No significant difference in incidence of POCD between the 2 groups (p-value = 0.089)	-	-
Beck-Schimmer et al. (2015)	-	No significant difference in liver biochemistry (ALT, AST, bilirubin) between both groups No significant difference in major in-hospital complications between both groups (p-value = 0.23) No significant difference in overall morbidity rate between both groups (p-value = 0.42)	No significant difference in incidence of EAD in both groups (p-value = 0.45)	No significant difference in ICU LOS (p-value = 0.64) or hospital LOS (p-value = 0.77) between both groups
Gucyetmez (2016)	-	24-h total tramadol requirement lower in the study group 4.1 ± 0.07 versus 3.7 ± 0.05 mg/kg/h Time to first tramadol longer in the study group 3.2 ± 0.7 versus 17.5 ± 2.5 h Reduced duration of mechanical ventilation in the study group in multivariate analysis (p-value < 0.001)	-	Similar ICU and hospital LOS between both groups
Gajate Martin (2016)	-	-	No significant difference in incidence of primary graft dysfunction between both groups. (OR 0.76, CI 0.42-1.47; p-value = 0.45).	-
Ali Sahmeddini (2012)	No differences in hemodynamic variables (MAP, HR, CVP) during LT between both groups	-	-	-
Lu (2016)	Statistically significantly lower dose of norepinephrine during reperfusion in TIVA versus Desflurane group (p-value = 0.012) No significant difference in haematological (INR) and biochemical (ALT) markers 24 h post LT	Awake time was significantly faster in the TIVA group versus Desflurane (p-value = 0.034) No significant differences in extubation time	-	No significant difference in ICU or total hospital LOS
Mangus (2018)	No significant differences in perioperative death	No significant differences in death at 1 year post LT Greatest perioperative IRI occurred in Isoflurane group	No significant difference in EAD	No significant difference in hospital LOS between all 3 groups

(Continues)

TABLE 2 (Continued)

Author, year (Reference)	Intra-operative morbidity and mortality	Post-operative morbidity and mortality	Early allograft dysfunction	ICU and hospital length of stay
Toprak (2011)	Significant changes in MAP, MPAP, ETCO ₂ , EEG based awareness monitoring (BIS), and body temperature across all 3 phases of LT (all values remained clinically acceptable) No difference in HR across the 3 phases of LT.	-	-	-
Pan (2016)	Significant difference in hypotension (MAP) seen at 30 min (<i>p</i> -value = 0.027) No significant difference in bradycardia No significant difference in quantity of vasoactive agents (atropine and phenylephrine) used	-	-	-
Kim (2019)	-	Time to recovery of consciousness significantly longer in severe HE group vs. non-severe group (2.8 days vs. 1.2 days, <i>p</i> -value = 0.002) Multivariate analysis: MELD score and GCS at 24 h were associated with increased consciousness recovery time: (MELD: hazard ratio 1.09, <i>p</i> -value = 0.02) (GCS: hazard ratio 0.692, <i>p</i> -value < 0.001) -	-	-
Tsai (2012)	No significant difference in perioperative hemodynamic data between groups No significant difference in mean MDA levels between both groups before reperfusion Mean MDA significantly lower in Propofol group versus control group at 3, 5, 0, and 60 min after reperfusion (all time points: <i>p</i> -value < 0.05)	-	-	-
Sorensen (2014)	Cerebral oxygenation and ETCO ₂ reduced by 4.3% and 0.3 kPa respectively in the anhepatic phase (<i>p</i> -value < 0.0001) Cerebral oxygenation and ETCO ₂ increased by 5.5% and 0.7 kPa respectively during reperfusion of the liver	-	-	-

(Continues)

TABLE 2 (Continued)

Author, year (Reference)	Intra-operative morbidity and mortality	Post-operative morbidity and mortality	Early allograft dysfunction	ICU and hospital length of stay
Jowkar (2020)	-	No significant difference in postoperative liver biochemistry (AST, INR, bilirubin) Patients who received remifentanyl had statistically significant, but clinically insignificant, changes in ALT No change in renal function over 7 days post LT No significant difference in extubation time	-	No significant difference in ICU LOS (p-value = 0.75) and hospital LOS (p-value = 0.23)

Outcomes are presented in the same order as in Table 1.

Abbreviations: ALP, alkaline phosphatase; ALT, alanine transaminase; AST, aspartate transaminase; BIS, bispectral index; BUN, blood urea nitrogen; CI, confidence interval; CO, cardiac output; CVP, central venous pressure; EAD, early allograft dysfunction; ETCO₂, end tidal carbon dioxide; FFP, fresh frozen plasma; HE, hepatic encephalopathy; GA, general anesthesia; GCS, Glasgow coma scale; HR, heart rate; INR, international normalized ratio; IRI, ischemic reperfusion injury; LT, liver transplant; MAC, minimum alveolar concentration; MAP, mean arterial pressure; MDA, malondialdehyde; MPAP, mean pulmonary artery pressure; NSE, neuron specific enolase; LOC, loss of consciousness; LOS, length of stay; LT, liver transplant; MELD, model for end stage liver disease; OR, odds ratio; PCC, prothrombin complex concentrate; POCD, postoperative cognitive dysfunction; PT, prothrombin time; PTT, partial thrombin time; S100B, calcium-binding protein B; TCI, target controlled infusion; TIVA, total intravenous.

TABLE 3 Summary of findings leading to the quality of evidence assessment according to the GRADE approach

Summary of findings								
Number of studies								
RCT	Observational comparative	Observational non-comparative	Limitations	Inconsistency	Indirectness	Imprecision	Publication bias	Quality of evidence (GRADE)
Outcome 1: Intraoperative morbidity and mortality								
2	5	1	Serious	Not serious	Serious	Serious	Not likely	Low ●●○○
Outcome 2: Postoperative morbidity and mortality								
4	3	0	Serious	Not serious	Serious	Serious	Not likely	Low ●●○○
Outcome 3: Early allograft dysfunction								
1	2	0	Not serious	Not serious	Not serious	Serious	Not likely	Low ●●○○
Outcome 4: ICU length of stay								
3	1	0	Not serious	Not serious	Not serious	Serious	Not likely	Moderate ●●●○
Outcome 5: Hospital length of stay								
3	2	0	Not serious	Not serious	Not serious	Serious	Not likely	Moderate ●●●○

on expert opinion has been utilized to make recommendations. Secondly, the questions have several financial, resource, and environmental implications which pose a challenge to making strong recommendations that are widely applicable, given that LT is a worldwide procedure occurring in environments with different resource constraints. For example, certain anesthetic resources like EEG based awareness monitoring such as BIS, or target controlled propofol infusions are asso-

ciated with increased costs and subsequently variable access; volatile agents have been shown to have a detrimental effect on the environment and are expensive, but remain widely used. Thirdly, an element of conflict exists within subsequent questions, as demonstrated by the first question which compares type of anesthesia (TIVA vs. volatile) and the second question which examines MAC, implying the use of volatile anesthesia, versus EEG based awareness monitoring.

TABLE 4 Evidence to recommendation framework according to the GRADE approach

Question 1: Does the type of anaesthesia (TIVA vs. Volatile) affect immediate and short-term outcomes after liver transplantation?			
Decision domain	Judgement		Reason for Judgement
	Yes	No	
Balance between desirable and undesirable outcomes (estimated effects), with consideration of values and preferences (estimated typical)	✓		<p>The literature reviewed showed a small amount of evidence supporting faster wakeup and markers of decreased ischemic reperfusion injury in patients receiving TIVA when compared to volatile anaesthesia</p> <p>In general, when assessing volatile agents as an anesthesia type, their desirability lies in their ability to provide amnesia at a measurable level and lack of dependence on hepatic or renal metabolism, which outweighs their undesirable effects such as postoperative nausea and vomiting, and environmental impact particularly with desflurane</p> <p>When generally assessing TIVA as an anesthetic type, its desirability as an agent is due to more rapid emergence, improved profile in patients with increased intracranial pressure (ICP), and lower incidence of PONV are outweighed by the need for metabolism by the liver and non-validated means of confirming amnesia intraprocedurally, especially in the setting of rapid blood loss.</p>
Confidence in the magnitude of estimates of effect of the interventions on important outcomes (overall quality of evidence for outcomes)		✓	The evidence is low when looking at TIVA versus volatile anesthesia in terms of their contribution to important outcomes post-liver transplantation
Confidence in Values and Preference, and their Variability		✓	There is inconsistency between the studies analyzed in the manner in which TIVA versus volatile was administered. In certain studies, target controlled infusions for TIVA were utilized versus anesthesiologist titration to BIS or discretion, which was also similar to the volatile administration parameters
Resource implications <i>Are the resources worth the expected net benefit from following the recommendation?</i>	✓		There are increased costs and potential environmental effects depending on the agent used. For TIVA, resource implications include the cost of propofol vials, as well as the resources needed for target controlled infusions and non-validated depth of anaesthesia monitoring techniques technologies
Overall Quality of Evidence: Very Low			
Recommendation of the Panel: Weak for the use of volatile anesthesia for use in liver transplantation for optimal short term outcomes, with consideration of TIVA in patients at risk for or with increased intracranial pressure.			
Question 2 : Does the type of depth of anaesthesia assessment (EEG based awareness monitoring vs. MAC) affect immediate and short term outcomes after liver transplantation?			
Decision domain	Judgement		Reason for Judgement
	Yes	No	
Balance between desirable and undesirable outcomes (estimated effects), with consideration of values and preferences (estimated typical)	✓		In general, there is some data supporting the use of EEG based awareness monitoring for reduction of anesthetic delivered, improved hemodynamics, and improved survival, and has been shown to be of similar reference value in cirrhotic patients ²⁶
Confidence in the magnitude of estimates of effect of the interventions on important outcomes (overall quality of evidence for outcomes)			The evidence is low regarding the use of EEG based awareness monitoring versus MAC contributing to important outcomes post-liver transplantation

(Continues)

TABLE 4 (Continued)

Question 2 : Does the type of depth of anaesthesia assessment (EEG based awareness monitoring vs. MAC) affect immediate and short term outcomes after liver transplantation?			
Decision domain	Judgement		Reason for Judgement
	Yes	No	
Confidence in Values and Preference, and their Variability	✓		BIS values have been shown to be consistent in the cirrhotic population, with MAC requirements shown to be reduced in cirrhotics undergoing liver transplantation ^{27,28}
Resource implications	✓		There is an increased cost associated with the utilization of EEG based awareness monitoring monitoring relative to the use of MAC during the intraoperative phase of case
Overall Quality of Evidence: Very Low			
Recommendation: Weak for the use of MAC as a measure of anesthetic depth during liver transplantation for optimal short term outcomes			
Question 3: Does a liberal versus restrictive O₂ management affect immediate and short-term outcomes after liver transplantation?			
Decision domain	Judgement		Reason for judgement
	Yes	No	
Balance between desirable and undesirable outcomes (estimated effects), with consideration of values and preferences (estimated typical)	✓		In the literature reviewed, there is no data supporting varying oxygen concentrations. However, there is significant data supporting the use of lower oxygen concentrations as well as data showing the harms of liberal oxygen use in other surgical populations
Confidence in the magnitude of estimates of effect of the interventions on important outcomes (overall quality of evidence for outcomes)	✓		The evidence is low to support different levels of oxygen administration for important outcomes after liver transplantation
Confidence in Values and Preference, and their Variability		✓	The concentration of O ₂ for restrictive O ₂ concentrations is not well defined. Liberal O ₂ concentration is defined as a FiO ₂ of 80–100%
Resource implications	✓		There is a cost associated with oxygen to centers, so there is a benefit with restrictive O ₂ management techniques
Overall Quality of Evidence: Very Low			
Recommendation: Strong for restrictive O₂ administration titrated to a PaO₂ of 70–110 mmHg (10–14 kPa) should be utilized during liver transplantation for optimal short term outcomes.			
Question 4: Do intraoperative ventilation strategies (PEEP, tidal volumes, recruitment manoeuvres) affect immediate and short-term outcomes after liver transplantation?			
Decision domain	Judgement		Reason for Judgement
	Yes	No	
Balance between desirable and undesirable outcomes (estimated effects), with consideration of values and preferences (estimated typical)	✓		Low tidal volumes (TV) as defined 6–8 ml/kg IBW, use of PEEP, and recruitment maneuvers all are beneficial to patients with minimal undesirable outcomes associated with their utilization
Confidence in the magnitude of estimates of effect of the interventions on important outcomes (overall quality of evidence for outcomes)	✓		There is low-quality evidence supporting the use of low tidal volume ventilation, PEEP, and recruitment maneuvers for improving outcomes postliver transplantation

(Continues)

TABLE 4 (Continued)

Question 4: Do intraoperative ventilation strategies (PEEP, tidal volumes, recruitment manoeuvres) affect immediate and short-term outcomes after liver transplantation?			
Decision domain	Judgement		Reason for Judgement
	Yes	No	
Confidence in Values and Preference, and their Variability	✓		The definition of low TV is well defined as 6–8 ml/kg. Optimal PEEP has not been defined and varies by patient factors. The definition of recruitment maneuver is also varied depending on literature ²⁹
Resource implications	✓		Although not defined in the evidence, there could be a potential resource saving in reduced need for mechanical ventilation or ICU days
Overall Quality of Evidence: Very Low			
Recommendation: Strong for the use of tidal volumes of 6–8 ml/kg IBW, PEEP titrated to patient intraoperative physiology, and recruitment maneuvers in liver transplantation for optimal short-term outcomes.			
Question 5: Does antiemetic prophylaxis affect immediate and short-term outcomes after liver transplantation?			
Decision domain	Judgement		Reason for Judgement
	Yes	No	
Balance between desirable and undesirable outcomes (estimated effects), with consideration of values and preferences (estimated typical)	✓		The benefits of antiemesis outweigh the risk of QT prolongation associated with most common antiemetic agents.
Confidence in the magnitude of estimates of effect of the interventions on important outcomes (overall quality of evidence for outcomes)	✓		The evidence is low for the support of antiemetic prophylaxis contributing to important outcomes in liver transplantation
Confidence in Values and Preference, and their Variability		✓	n/a
Resource implications?	✓		No resource difference in patients receiving PONV prophylaxis
Overall Quality of Evidence: Very low			
Recommendation: Strong for the use of antiemetic prophylaxis in liver transplantation for optimal short term outcomes			

6 | CONCLUSION

When considering optimal anesthetic management and its contribution to short-term outcomes after liver transplantation, the panel was able to come to the above conclusions about the five proposed questions. However, more importantly, this review demonstrates significant deficiencies in the anesthesia literature. More outcome focused research is needed to identify the most beneficial anesthetic techniques for liver transplant patients in the perioperative period, such research should also consider the resource implications of specific techniques.

Panel suggestions for key areas of future research:

1. A core outcome set (<https://www.comet-initiative.org>) to define key outcomes that should be reported in any study focusing on intraoperative anesthetic management in liver transplantation.
2. Volatile Anesthetics: studies comparing the use of different volatiles on short term outcomes after LT including: time to extubation, EAD, comparison of cost relative to hospital and ICU length of stay.
3. International multicenter RCTs comparing volatile agents versus TIVA focusing on:
 - a. Cardiovascular stability and intraoperative vasopressor, inotrope, and fluid requirements
 - b. Ability to extubate early (on-table and within a specified number of hours postoperatively on ICU)
 - c. Postoperative cognitive dysfunction
 - d. Early Allograft Dysfunction
4. Multi-center RCTs comparing liberal versus restrictive oxygenation strategies focusing on:
 - a. Ischaemia-reperfusion injury (given the theory that IRI is partly driven by an increase in oxygen free radicals)

- b. Postoperative pulmonary complications such as atelectasis, pneumonia, and pulmonary edema.
- c. Early allograft dysfunction
5. Explanatory studies to assess: Depth of anesthesia monitoring: correlation of MAC with EEG based awareness monitoring in LT recipients

ACKNOWLEDGEMENTS

Claus Niemann, San Francisco, CA, USA, Joerg-Matthias Pollok, London, UK, Marina Berenguer, Valencia, Spain, Pascale Tinguely, London, UK, Ka Hay Fan, London, UK, Giselle Farrer, London, UK. This manuscript was prepared for the ERAS4OLT.org Consensus Conference 2022, which is partially funded by the International Liver Transplant Society (ILTS). This work was conducted in preparation for the ILTS – ERAS4OLT.org Consensus Conference on Enhanced Recovery for Liver Transplantation, January 2022, Valencia, Spain.

AUTHORSHIP

All authors qualify for authorship as per the International Committee of Medical Journal Editors (ICMJE) guidelines.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Ryan Chadha  <https://orcid.org/0000-0002-7463-6912>

Victor Xia  <https://orcid.org/0000-0003-1182-3179>

REFERENCES

1. Feldheiser A, Aziz O, Baldini G, et al. Enhanced recovery after surgery (ERAS) for gastrointestinal surgery, part 2: consensus statement for anaesthesia practice. *Acta Anaesthesiol Scand*. 2015;60(3):289-334. <http://doi.org/10.1111/aas.12651>.
2. Myles P, Leslie K, Chan M, et al. The safety of addition of nitrous oxide to general anaesthesia in at-risk patients having major non-cardiac surgery (ENIGMA-II): a randomised, single-blind trial. *Lancet North Am Ed*. 2014;384(9952):1446-1454. [http://doi.org/10.1016/s0140-6736\(14\)60893](http://doi.org/10.1016/s0140-6736(14)60893).
3. Punjasawadwong Y, Phongchiewboon A, Bunchungmongkol N. Bispectral index for improving anaesthetic delivery and postoperative recovery. *Cochrane Database Syst Rev*. 2014;2014(6):CD003843. <http://doi.org/10.1002/14651858.CD003843.pub3>.
4. Chan M, Cheng B, Lee T, Gin T. BIS-guided anesthesia decreases postoperative delirium and cognitive decline. *J Neurosurg Anesthesiol*. 2013;25(1):33-42. <http://doi.org/10.1097/ana.0b013e3182712fba>.
5. Futier E, Constantin JM, Paugam-Burtz C, et al, IMPROVE Study Group. A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *N Engl J Med*. 2013;369(5):428-437. <http://doi.org/10.1056/NEJMoa1301082>.
6. Gustafsson UO, Scott MJ, Hubner M, et al. Guidelines for perioperative care in elective colorectal surgery: enhanced recovery after surgery (ERAS®) society recommendations: 2018. *World J Surg*. 2019;43(3):659-695. <http://doi.org/10.1007/s00268-018-4844-y>.
7. Farges O, Kokudo N, Vauthey JN, Clavien PA, Demartines N. Guidelines for perioperative care for liver surgery: enhanced recovery after surgery (ERAS) society recommendations. *World J Surg*. 2016;40(10):2425-2440. <http://doi.org/10.1007/s00268-016-3700-1>.
8. Guyatt G, Oxman A, Akl E, et al. GRADE guidelines: 1. Introduction—GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. 2011;64(4):383-394. <http://doi.org/10.1016/j.jclinepi.2010.04.026>.
9. Murad M, Mustafa R, Schünemann H, Sultan S, Santesso N. Rating the certainty in evidence in the absence of a single estimate of effect. *Evid Based Med*. 2017;22(3):85-87. <http://doi.org/10.1136/ebmed-2017-110668>.
10. Andrews J, Guyatt G, Oxman A, et al. GRADE guidelines: 14. Going from evidence to recommendations: the significance and presentation of recommendations. *J Clin Epidemiol*. 2013;66(7):719-725. <http://doi.org/10.1016/j.jclinepi.2012.03.013>.
11. Andrews J, Schünemann H, Oxman A, et al. GRADE guidelines: 15. Going from evidence to recommendation—determinants of a recommendation's direction and strength. *J Clin Epidemiol*. 2013;66(7):726-735. <http://doi.org/10.1016/j.jclinepi.2013.02.003>.
12. Najafi A, Hussain Khan Z, Faez Abdunabi S, Mohammadpour Z, Barzin G, Zebardast J. Comparing the effect of propofol and sevoflurane on hemodynamics and coagulation status during liver transplant anesthesia and hepatic and renal function of the patients after liver transplant. *Cochrane Database Syst Rev*. 2019;9:CD003843. PMID: 24937564; PMCID: PMC6483694. <http://doi.org/10.18502/aacc.v5i4.1453>.
13. Cao Y, Chi P, Zhao Y, Dong X. Effect of bispectral index-guided anesthesia on consumption of anesthetics and early postoperative cognitive dysfunction after liver transplantation. *Medicine (Baltimore)*. 2017;96(35):e7966. <http://doi.org/10.1097/md.0000000000007966>.
14. Beck-Schimmer B, Bonvini J, Schadde E, et al. Conditioning with sevoflurane in liver transplantation. *Transplantation*. 2015;99(8):1606-1612. <http://doi.org/10.1097/tp.0000000000000644>.
15. Gucyetmez B, Atalan H, Aslan S, Yazar S, Polat K. Effects of intraoperative magnesium sulfate administration on postoperative tramadol requirement in liver transplantation: a prospective, double-blind study. *Transplant Proc*. 2016;48(8):2742-2746. <http://doi.org/10.1016/j.transproceed.2016.08.033>.
16. Gajate Martín L, González C, Ruiz Torres I, et al. Effects of the hypnotic agent on primary graft dysfunction after liver transplantation. *Transplant Proc*. 2016;48(10):3307-3311. <http://doi.org/10.1016/j.transproceed.2016.08.048>.
17. Sahmeddini M, Eghbal M, Khosravi M, Ghaffaripour S, Janatmakan F, Shokrizade S. Electro-acupuncture stimulation at acupoints reduced the severity of hypotension during anesthesia in patients undergoing liver transplantation. *J Acupunct Meridian Stud*. 2012;5(1):11-14. <http://doi.org/10.1016/j.jams.2011.11.001>.
18. Lu C, Yeh C, Huang Y, et al. Hemodynamic and biochemical changes in liver transplantation: a retrospective comparison of desflurane and total intravenous anesthesia by target-controlled infusion under auditory evoked potential guide. *Acta Anaesthesiol Taiwan*. 2014;52(1):6-12. <http://doi.org/10.1016/j.aat.2014.05.004>.
19. Mangus R, Kinsella S, Farar D, Fridell J, Woolf L, Kubal C. Impact of volatile anesthetic agents on early clinical outcomes in liver transplantation. *Transplant Proc*. 2018;50(5):1372-1377. <http://doi.org/10.1016/j.transproceed.2018.03.001>.
20. Toprak H, Sener A, Gedik E, et al. bispectral index monitoring to guide end-tidal isoflurane concentration at three phases of operation in patients with end-stage liver disease undergoing orthotopic liver transplantation. *Transplant Proc*. 2011;43(3):892-895. <http://doi.org/10.1016/j.transproceed.2010.11.023>.

21. Pan J, Cai J, Zhou S, et al. Pharmacodynamic analysis of target-controlled infusion of propofol in patients with hepatic insufficiency. *Biomed Rep.* 2016;5(6):693-698. [10.3892/br.2016.786](https://doi.org/10.3892/br.2016.786).
22. Kim D, Shin B, Song I, et al. Relationship between intraoperative bispectral index and consciousness recovery in patients with hepatic encephalopathy undergoing liver transplant: a retrospective analysis. *Transplant Proc.* 2019;51(3):798-804. <http://doi.org/10.1016/j.transproceed.2018.10.031>.
23. Tsai Y, Lin C, Lee W, Yu H. Propofol attenuates ischemic reperfusion-induced formation of lipid peroxides in liver transplant recipients. *Transplant Proc.* 2012;44(2):376-379. <http://doi.org/10.1016/j.transproceed.2012.01.013>.
24. SÅyrensen H, Grocott H, Niemann M, et al. Ventilatory strategy during liver transplantation: implications for near-infrared spectroscopy-determined frontal lobe oxygenation. *Front Physiol.* 2014;5. <http://doi.org/10.3389/fphys.2014.00321>.
25. Jowkar S, Khosravi M, Sahmeddini M, Eghbal M, Samadi K. Preconditioning effect of remifentanyl versus fentanyl in prevalence of early graft dysfunction in patients after liver transplant: a randomized clinical trial. *Exp Clin Transplant.* 2020;18(5):598-604. <http://doi.org/10.6002/ect.2019.0014>.
26. Myles PS, Leslie K, McNeil J, Forbes A, Chan MT, B-Aware Trial Group. Bispectral index monitoring to prevent awareness during anaesthesia: the B-Aware randomised controlled trial. *Lancet North Am Ed.* 2004;363(9423):1757-1763.
27. Wang CH, Chen CL, Cheng KW, et al. Bispectral index monitoring in healthy, cirrhotic, and end-stage liver disease patients undergoing hepatic operation. *Transplantation Proceedings*, 1 October 2008 (Vol. 40, No. 8, pp. 2489-2491). Elsevier.
28. Kang JG, Ko JS, Kim GS, Gwak MS, Kim YR, Lee SK, The relationship between inhalational anesthetic requirements and the severity of liver disease in liver transplant recipients according to three phases of liver transplantation. *Transplantation Proceedings*, 1 April 2010 (Vol. 42, No. 3, pp. 854-857). Elsevier.
29. Santos RS, Silva PL, Pelosi P, Rocco PR. Recruitment maneuvers in acute respiratory distress syndrome: the safe way is the best way. *World J Crit Care Med.* 2015;4(4):278.
30. Stravitz RT, Kramer AH, Davern T, et al. Intensive care of patients with acute liver failure: recommendations of the US acute liver failure study group. *Crit Care Med Baltimore.* 2007;35(11):2498. Nov 1.
31. Meyer MJ. MD desflurane should des-appear: global and financial rationale. *Anesth Anal.* 2020;131(4):1317-1322. <http://doi.org/10.1213/ANE.0000000000005102>.
32. <https://www.asahq.org/standards-and-guidelines/standards-for-basic-anesthetic-monitoring>
33. Meyhoff C, Jorgensen L, Wetterslev J, Christensen K, Rasmussen L. Increased long-term mortality after a high perioperative inspiratory oxygen fraction during abdominal surgery. *Anesth Anal.* 2012;115(4):849-854. <http://doi.org/10.1213/ane.0b013e3182652a51>.
34. Meyhoff C, Jorgensen L, Wetterslev J, Siersma V, Rasmussen L. Risk of new or recurrent cancer after a high perioperative inspiratory oxygen fraction during abdominal surgery. *Br J Anaesth.* 2014;113:i74-i81. <http://doi.org/10.1093/bja/aeu110>.
35. Park M, Jung K, Sim W, et al. Perioperative high inspired oxygen fraction induces atelectasis in patients undergoing abdominal surgery: a randomized controlled trial. *J Clin Anesth.* 2021;72:110285. <http://doi.org/10.1016/j.jclinane.2021.110285>.
36. Reiterer C, Kabon B, Taschner A, et al. Perioperative supplemental oxygen and NT-proBNP concentrations after major abdominal surgery – a prospective randomized clinical trial. *J Clin Anesth.* 2021;73:110379. <http://doi.org/10.1016/j.jclinane.2021.110379>.
37. Meyhoff C, Wetterslev J, Jorgensen L, et al. Effect of high perioperative oxygen fraction on surgical site infection and pulmonary complications after abdominal surgery. *JAMA.* 2009;302(14):1543. <http://doi.org/10.1001/jama.2009.1452>.
38. Thibon P, Borgey F, Boutreux S, Hanouz J, Le Coutour X, Parienti J. Effect of perioperative oxygen supplementation on 30-day surgical site infection rate in abdominal, gynecologic, and breast surgery. *Anesthesiology.* 2012;117(3):504-511. <http://doi.org/10.1097/aln.0b013e3182632341>.
39. Patel S, Coughlin S, Malthaner R. High-concentration oxygen and surgical site infections in abdominal surgery: a meta-analysis. *Can J Surg.* 2013;56(4):E82-E90. <http://doi.org/10.1503/cjs.001012>.
40. Futier E, Constantin J, Paugam-Burtz C, et al. A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *N Engl J Med.* 2013;369(5):428-437. <http://doi.org/10.1056/nejmoa1301082>.
41. Severgnini P, Selmo G, Lanza C, et al. Protective mechanical ventilation during general anesthesia for open abdominal surgery improves postoperative pulmonary function. *Anesthesiology.* 2013;118(6):1307-1321. <http://doi.org/10.1097/aln.0b013e31829102de>.
42. Park S. Perioperative lung-protective ventilation strategy reduces postoperative pulmonary complications in patients undergoing thoracic and major abdominal surgery. *Kor J Anesthesiol.* 2016;69(1):3. <http://doi.org/10.4097/kjae.2016.69.1.3>.
43. Wetterslev J, Hansen E, Roikjaer O, Kanstrup I, Heslet L. Optimizing perioperative compliance with PEEP during upper abdominal surgery: effects on perioperative oxygenation and complications in patients without preoperative cardiopulmonary dysfunction. *Eur J Anaesthesiol.* 2001;18(6):358-365. <http://doi.org/10.1097/00003643-200106000-00003>.
44. Mini G, Ray B, Anand R, et al. Effect of driving pressure-guided positive end-expiratory pressure (PEEP) titration on postoperative lung atelectasis in adult patients undergoing elective major abdominal surgery: a randomized controlled trial. *Surgery.* 2021;170(1):277-283. <http://doi.org/10.1016/j.surg.2021.01.047>.
45. Cui Y, Cao R, Li G, Gong T, Ou Y, Huang J. The effect of lung recruitment maneuvers on post-operative pulmonary complications for patients undergoing general anesthesia: a meta-analysis. *PLoS One.* 2019;14(5):e0217405. <http://doi.org/10.1371/journal.pone.0217405>.
46. Gan T, Diemunsch P, Habib A, et al. Consensus guidelines for the management of postoperative nausea and vomiting. *Anesth Anal.* 2014;118(1):85-113. <http://doi.org/10.1213/ane.0000000000000002>.

How to cite this article: Chadha R, Patel D, Bhangui P, et al. Optimal anesthetic conduct regarding immediate and short-term outcomes after liver transplantation – Systematic review of the literature and expert panel recommendations. *Clin Transplant.* 2022;e14613. <https://doi.org/10.1111/ctr.14613>