

# UC Davis

## UC Davis Previously Published Works

### Title

Draft Genome Sequence of the Multidrug-Resistant *Citrobacter freundii* 132-2 Strain Isolated from a Domestic Duck in Bangladesh.

### Permalink

<https://escholarship.org/uc/item/37n5752m>

### Journal

Microbiology Resource Announcements, 12(7)

### Authors

Ahmed, Tarana

Islam, Md Saiful

Nuruzzaman, Mohammad

et al.

### Publication Date

2023-07-18




### DOI

10.1128/mra.00378-23

Peer reviewed



# Draft Genome Sequence of the Multidrug-Resistant *Citrobacter freundii* 132-2 Strain Isolated from a Domestic Duck in Bangladesh

 Tarana Ahmed,<sup>a,b</sup>  M. Saiful Islam,<sup>a</sup> Mohammad Nuruzzaman,<sup>c</sup> Mohammad Sadekuzzaman,<sup>d</sup> S. M. Lutful Kabir,<sup>a</sup>  
 M. Tanvir Rahman,<sup>a</sup> M. Shahidur Rahman Khan<sup>a</sup>

<sup>a</sup>Department of Microbiology and Hygiene, Bangladesh Agricultural University, Mymensingh, Bangladesh

<sup>b</sup>Department of Livestock Services, Ministry of Fisheries & Livestock, Government of the Peoples Republic of Bangladesh, Dhaka, Bangladesh

<sup>c</sup>Ministry of Public Administration, Dhaka, Bangladesh

<sup>d</sup>Central Disease Investigation Laboratory (CDIL), Department of Livestock Services, Ministry of Fisheries & Livestock, Government of the Peoples Republic of Bangladesh, Dhaka, Bangladesh

**ABSTRACT** We sequenced a multidrug-resistant strain of *Citrobacter freundii*, 132-2, isolated from a cloacal swab sample of a domestic duck. The whole genome of the *C. freundii* 132-2 strain had a length of 5,097,592 bp, 62 contigs, two plasmids, and an average G+C content of 51.85%, with a 105.0× genome coverage.

The extensive and inappropriate use of antimicrobial agents has resulted in the emergence of antimicrobial resistance in bacteria, leading to various multidrug-resistant (MDR) clones, which have become a major global public health concern (1, 2). Ducks have the potential to harbor antimicrobial-resistant and MDR pathogens that could transfer to humans due to their interactions with them (3). *Citrobacter freundii* is commonly found in the environment, soil, water, human clinical samples, and intestinal tracts of animals (4). *C. freundii* can be transmitted from ducks to humans through various means, such as contact with infected eggs, raw or undercooked meat, and the handling of duck carcasses at the slaughterhouse (3, 5). In humans, *C. freundii* isolates can cause bacteremia and severe neurological symptoms such as extreme cognitive impairment, seizures, and hemiparesis, and, in some cases, even death in children, so that it represents a considerable threat to human health (6, 7).

From January 2020 to January 2022, cloacal swab samples of domestic ducks (*Anas platyrhynchos domesticus*) were collected from the Kishoreganj district of Bangladesh and transported to our laboratory (24.7196°N, 90.4267°E). The specimens were then cultivated in nutrient broth (HiMedia, India), streaked on a xylose-lysine deoxycholate agar (HiMedia, India) plate, and incubated at 37°C for one night. Afterward, pure colonies were selected and subjected to staining and biochemical tests to isolate *Citrobacter* spp. (8). Finally, *C. freundii* was identified using a matrix-assisted laser desorption ionization–time of flight mass spectrometry assay (9). The MDR *C. freundii* isolate (resistant to >7 antimicrobial classes by disk diffusion method [10]) was incubated overnight in nutrient broth (HiMedia, India) at 37°C. The DNA was extracted from the collected culture using the Qiagen DNA minikit (Qiagen, Hilden, Germany). We used a NanoDrop 2000 UV-visible (UV-Vis) spectrophotometer (Thermo Fisher, Waltham, MA, USA) to determine the concentration of DNA present and its level of purity. To create the DNA libraries, the Nextera DNA flex library prep kit (Illumina, San Diego, CA, USA) was employed. The Illumina NextSeq2000 platform was used for whole-genome sequencing (WGS) (read length, 2 by 150 bp). The genome was assembled using Unicycler v0.4.9 (11). Before that, the raw paired-end reads ( $n = 7,855,310$ ) were subjected to trimming using Trimmomatic v0.39 (12) and underwent quality assessment with FastQC v0.11.7 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). The annotation of the genome was performed with Prokka v1.14.6 (13) and PATRIC (14). The PathogenFinder

**Editor** Simon Roux, DOE Joint Genome Institute

**Copyright** © 2023 Ahmed et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Tarana Ahmed, tarana\_ahmed07@yahoo.com, or M. Shahidur Rahman Khan, msrkhan001@yahoo.com.

The authors declare no conflict of interest.

**Received** 5 May 2023

**Accepted** 15 June 2023

**Published** 26 June 2023

1.1 (15), MLST 2.0 (16), PlasmidFinder-2.0 (17), CARD (18), VFDB (19), and RAST.v.2 (20) databases were utilized to identify the pathogenicity index, sequence type, plasmids, antimicrobial resistance genes (ARGs), virulence factor genes (VFGs), and metabolic functional features, respectively, in our assembled genome. All tools were run with default parameters.

The genome coverage of *Citrobacter freundii* 132-2 was 105.0 $\times$ , and a total of 62 contigs were obtained. Our assembled genome had a total length of 5,097,592 bp, a GC content of 51.85%, and an  $N_{50}$  value of 536,796 bp. This assembled genome identified two important plasmid replicons, i.e., IncFIB(pHCM2) (110,089 bp with a 98.51% identity to [AL513384](#)) and IncR (16,190 bp with a 100% identity to [DQ449578](#)). Moreover, this genome consists of a sequence type of ST18 (using MLST 2.0) and a probability of being a pathogen for humans of 89.6% (using PathogenFinder 1.1). The strain carried 45 predicted ARGs under 19 antimicrobial categories and 83 predicted virulence genes under more than 12 predicted virulence factors. Moreover, *Citrobacter freundii* 132-2 contained 395 subsystems (having 2,298 genes) with 33% coverage, 5,024 protein-coding sequences, and 81 RNA genes.

The Animal Welfare and Experimentation Ethical Committee, which is the institutional ethics committee of Bangladesh Agricultural University in Mymensingh, granted approval for the procedures and protocols associated with this research under the reference number of AWEEC/BAU/2020(10).

**Data availability.** The whole-genome sequencing (WGS) shotgun study for *Citrobacter freundii* 132-2 was submitted to NCBI/GenBank with accession number [JAPQWA000000000](#), and the raw reads were deposited under SRA accession number [SRR24848516](#) (BioProject accession number [PRJNA907483](#)). The current version referred to this paper is identified as [JAPQWA010000000](#).

## ACKNOWLEDGMENTS

We acknowledge the Bangladesh Agricultural Research Council for providing funds through the National Agricultural Technology Program Phase II project for the in-country Ph.D. scholarship (P149553) to facilitate this study.

We extend our appreciation to the Department of Livestock Services, the Ministry of Fisheries and Livestock of Bangladesh, for their assistance in conducting this research.

## REFERENCES

- Islam MS, Nayeem MMH, Sobur MA, Levy S, Islam MA, Rahman S, Kafi MA, Ashour HM, Rahman MT. 2021. Virulence determinants and multidrug resistance of *Escherichia coli* isolated from migratory birds. *Antibiotics* 10:190. <https://doi.org/10.3390/antibiotics10020190>.
- Islam MS, Paul A, Talukder M, Roy K, Sobur MA, Levy S, Nayeem MMH, Rahman S, Nazir KNH, Hossain MT, Rahman MT. 2021. Migratory birds travelling to Bangladesh are potential carriers of multi-drug resistant *Enterococcus* spp., *Salmonella* spp., and *Vibrio* spp. *Saudi J Biol Sci* 28: 5963–5970. <https://doi.org/10.1016/j.sjbs.2021.06.053>.
- Ahmed T, Islam MS, Haider N, Elton L, Hasan B, Nuruzzaman M, Rahman MT, Kabir SL, Khan MSR. 2023. Phenotypic and genotypic characteristics of antimicrobial resistance in *Citrobacter freundii* isolated from domestic ducks (*Anas platyrhynchos domesticus*) in Bangladesh. *Antibiotics* 12: 769. <https://doi.org/10.3390/antibiotics12040769>.
- Liu L, Lan R, Liu L, Wang Y, Zhang Y, Wang Y, Xu J. 2017. Antimicrobial resistance and cytotoxicity of *Citrobacter* spp. in Maanshan Anhui Province, China. *Front Microbiol* 8:1357. <https://doi.org/10.3389/fmicb.2017.01357>.
- Lessenger JE. 2006. Diseases from animals, poultry, and fish, p 367–382. In Lessenger JE (ed), *Agricultural medicine: a practical guide*. Springer, New York, NY.
- Doran TI. 1999. The role of *Citrobacter* in clinical disease of children. *Clin Infect Dis* 28:384–394. <https://doi.org/10.1086/515106>.
- Liu L, Chen D, Liu L, Lan R, Hao S, Jin W, Sun H, Wang Y, Liang Y, Xu J. 2018. Genetic diversity, multidrug resistance, and virulence of *Citrobacter freundii* from diarrheal patients and healthy individuals. *Front Cell Infect Microbiol* 8:233. <https://doi.org/10.3389/fcimb.2018.00233>.
- Brenner DJ, O'Hara CM, Grimont PA, Janda JM, Falsen E, Aldova E, Ageron E, Schindler J, Abbott SL, Steigerwalt AG. 1999. Biochemical identification of *Citrobacter* species defined by DNA hybridization and description of *Citrobacter gillenii* sp. nov. (formerly *Citrobacter* genomospecies 10) and *Citrobacter murlinae* sp. nov. (formerly *Citrobacter* genomospecies 11). *J Clin Microbiol* 37:2619–2624. <https://doi.org/10.1128/JCM.37.8.2619-2624.1999>.
- Kolínská R, Španělová P, Dřevínek M, Hrabák J, Žemličková H. 2015. Species identification of strains belonging to genus *Citrobacter* using the biochemical method and MALDI-TOF mass spectrometry. *Folia Microbiol (Praha)* 60:53–59. <https://doi.org/10.1007/s12223-014-0340-4>.
- Bauer AW, Kirby WMM, Sherris JC, Turck M. 1966. Antibiotic susceptibility testing by a standardized single diffusion method. *Am J Clin Pathol* 45:493–496. [https://doi.org/10.1093/ajcp/45.4\\_ts.493](https://doi.org/10.1093/ajcp/45.4_ts.493).
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Wattam AR, Abraham D, Dalay O, Disz TL, Driscoll T, Gabbard JL, Gillespie JJ, Gough R, Hix D, Kenyon R, Machi D, Mao C, Nordberg EK, Olson R, Overbeek R, Pusch GD, Shukla M, Schulman J, Stevens RL, Sullivan DE, Vonstein V, Warren A, Will R, Wilson MJC, Yoo HS, Zhang C, Zhang Y, Sobral BW. 2014. PATRIC, the bacterial bioinformatics database and analysis resource. *Nucleic Acids Res* 42: D581–D591. <https://doi.org/10.1093/nar/gkt1099>.
- Cosentino S, Voldby Larsen M, Møller Aarestrup F, Lund O. 2013. PathogenFinder—distinguishing friend from foe using bacterial whole genome sequence data. *PLoS One* 8:e77302. <https://doi.org/10.1371/journal.pone.0077302>.
- Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Pontén T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol* 50:1355–1361. <https://doi.org/10.1128/JCM.06094-11>.

17. Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller Aarestrup F, Hasman H. 2014. *In silico* detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <https://doi.org/10.1128/AAC.02412-14>.
18. Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Boucharde M, Edalatmand A, Huynh W, Nguyen A-LV, Cheng AA, Liu S, Min SY, Miroshnichenko A, Tran H-K, Werfalli RE, Nasir JA, Oloni M, Speicher DJ, Florescu A, Singh B, Faltyn M, Hernandez-Koutoucheva A, Sharma AN, Bordeleau E, Pawlowski AC, Zubyk HL, Dooley D, Griffiths E, Maguire F, Winsor GL, Beiko RG, Brinkman FSL, Hsiao WWL, Domselaar GV, McArthur AG. 2020. CARD 2020: antibiotic resistance surveillance with the comprehensive antibiotic resistance database. *Nucleic Acids Res* 48:D517–D525. <https://doi.org/10.1093/nar/gkz935>.
19. Liu B, Zheng D, Zhou S, Chen L, Yang J. 2022. VFDB 2022: a general classification scheme for bacterial virulence factors. *Nucleic Acids Res* 50:D912–D917. <https://doi.org/10.1093/nar/gkab1107>.
20. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.