

UC Davis

UC Davis Previously Published Works

Title

Glycoprotein In Vitro N-Glycan Processing Using Enzymes Expressed in E. coli

Permalink

<https://escholarship.org/uc/item/38d4h19q>

Journal

Molecules, 28(6)

ISSN

1431-5157

Authors

Zhang, Libo

Li, Yanhong

Li, Riyao

et al.

Publication Date

2023

DOI

10.3390/molecules28062753

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Article

Glycoprotein In Vitro N-Glycan Processing Using Enzymes Expressed in *E. coli*

Libo Zhang [†], Yanhong Li [†], Riyao Li [†], Xiaohong Yang, Zimin Zheng, Jingxin Fu, Hai Yu and Xi Chen ^{*†} 

Department of Chemistry, University of California, Davis, CA 95616, USA

^{*} Correspondence: xiichen@ucdavis.edu; Tel.: +1-530-754-6037[†] These authors contributed equally to this work.

Abstract: Protein N-glycosylation is a common post-translational modification that plays significant roles on the structure, property, and function of glycoproteins. Due to N-glycan heterogeneity of naturally occurring glycoproteins, the functions of specific N-glycans on a particular glycoprotein are not always clear. Glycoprotein in vitro N-glycan engineering using purified recombinant enzymes is an attractive strategy to produce glycoproteins with homogeneous N-glycoforms to elucidate the specific functions of N-glycans and develop better glycoprotein therapeutics. Toward this goal, we have successfully expressed in *E. coli* glycoside hydrolases and glycosyltransferases from bacterial and human origins and developed a robust enzymatic platform for in vitro processing glycoprotein N-glycans from high-mannose-type to α 2-6- or α 2-3-disialylated biantennary complex type. The recombinant enzymes are highly efficient in step-wise or one-pot reactions. The platform can find broad applications in N-glycan engineering of therapeutic glycoproteins.

Keywords: N-glycan; glycoprotein; glycan engineering; glycosyltransferase; mannosidase; *E. coli* expression



Citation: Zhang, L.; Li, Y.; Li, R.; Yang, X.; Zheng, Z.; Fu, J.; Yu, H.; Chen, X. Glycoprotein In Vitro N-Glycan Processing Using Enzymes Expressed in *E. coli*. *Molecules* **2023**, *28*, 2753. <https://doi.org/10.3390/molecules28062753>

Academic Editor: Kiichiro Totani

Received: 2 February 2023

Revised: 5 March 2023

Accepted: 16 March 2023

Published: 18 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Protein N-glycosylation is an important post-translational modification that affects the structure, property, and function of glycoproteins including folding, solubility, stability, localization, trafficking, molecular recognition, and interactions, etc. Glycoprotein N-glycans are attached via the innermost N-acetylglucosamine residue to the L-asparagine residue (GlcNAc β 1-Asn) in the Asn-X-Ser/Thr sequon (where X is an amino acid that is not an L-proline) of the protein with a β -linked N-glycosidic bond. All eukaryotic glycoprotein N-glycans share a trimannosyl chitobiose (Man₃GlcNAc₂) core and can be classified as high-mannose, complex, and hybrid types based on the glycan structures extended from the terminal mannose residues on the core [1,2].

Many therapeutic proteins and enzymes are N-glycosylated. The level of N-glycosylation and the structure of their N-glycans can directly affect their solubility, stability, safety, function, efficacy, delivery, pharmacokinetics, immunogenicity, and dose frequency [1,3–9]. Therefore, N-glycosylation is a critical quality attribute (CQA) of glycoprotein therapeutics considered by regulatory authorities [3,5,10]. Homogeneous glycoproteins with preferred N-glycoforms are highly desirable for their pharmaceutical applications and for exploring the fundamental understanding of their functions at the molecular level [7].

Nevertheless, N-glycosylated glycoproteins are intrinsically heterogeneous with variations on the N-glycan location, site occupancy of N-glycosylation, and N-glycan structures at individual N-glycosylation sites [11–13], and the N-glycosylation process is influenced by many factors [3,14]. Cells of different origins have been used to produce N-glycosylated glycoproteins [4]. Numerous strategies including cell line engineering, as well as the addition of inhibitors for metabolic glycosylation and glycosidase have been developed to reduce glycoprotein N-glycan heterogeneity [15,16]. Synthetic methods including total

synthesis, semi-synthesis, and chemoenzymatic synthesis have been developed to obtain homogeneous glycoproteins [17–20]. It is, however, still challenging to achieve glycoprotein N-glycan homogeneity [15], especially in large-scale productions. For example, monoclonal IgG1-type antibodies are an important type of therapeutic glycoproteins that have seen significant increased clinical applications. The desired N-glycans at the conserved N-glycosylation site of the Fc domain of these antibodies are biantennary complex-type fully sialylated with α 2–6-linked N-acetylneuraminic acid (Neu5Ac) without a core fucose [21]. Nevertheless, commercial therapeutic monoclonal antibodies have undesired high levels of high mannose-type N-glycans that are challenging to control during manufacturing processes [22]. In vitro enzymatic conversion of the high-mannose-type N-glycans on the glycoproteins to the preferred complex biantennary-type is a potential solution to overcome the challenge.

Among the methods developed for producing glycoproteins with structurally defined homogeneous N-glycans [4,19], the strategy of using recombinant glycoside hydrolases and glycosyltransferases for in vitro modifying N-glycan of glycoproteins is attractive [16,23]. Success in kilogram-scale N-glycan engineering of a recombinant human immunoglobulin G (hIgG) antibody using a purified recombinant β 1–4-galactosyltransferase 1 (B β 4GalT1) [24] highlights the feasibility of applying enzymatic glycoprotein in vitro N-glycan processing for large-scale production of therapeutic glycoproteins with structurally defined N-glycans. Nevertheless, the scope of several reported examples [25–28] is limited to the involvement of one or two glycosyltransferases with or without glycoside hydrolases. The utilization of multiple carbohydrate-active enzymes including glycoside hydrolases and glycosyltransferases for glycoprotein in vitro N-glycan processing has been traditionally underexplored, but has shown a high potential in several recent examples [23,29]. In general, glycosyltransferases that are expressed in mammalian [24,27] or insect cells [25] have been commonly used for glycoprotein N-glycan engineering due to the challenges in obtaining their active forms from common *E. coli* expression systems [29,30]. As many of these enzymes expressed in eukaryotic cells are N-glycosylated glycoproteins themselves, their N-glycans may complicate N-glycan analysis of target glycoproteins, a potential problem that deserves consideration and investigation, but has not been paid sufficient attention.

We believe that glycoprotein in vitro N-glycan processing and engineering using purified recombinant enzymes expressed in *E. coli* is an efficient platform that can be combined conveniently with different glycoprotein production systems to produce glycoproteins with homogeneous N-glycoforms. Using enzymes expressed in *E. coli* has the advantage of not introducing external N-glycans that may interfere with N-glycan analysis of target glycoproteins. *E. coli* expression system is also considered as the most convenient and economical system for protein production [29,31].

Using ribonuclease B (RNase B), a glycoprotein with high-mannose-type N-glycans at a single N-glycosylation site [32,33], as a model glycoprotein, we demonstrate here the success of processing glycoprotein N-glycans from high-mannose type to α 2–3 or α 2–6-linked disialylated complex biantennary N-glycans, using recombinant enzymes expressed in *E. coli* in both step-wise and one-pot reactions.

2. Results and Discussion

2.1. Glycoprotein In Vitro N-Glycan Processing Route Design and Enzyme Selection

To establish an efficient platform for processing glycoprotein N-glycans from high-mannose-type to disialylated complex biantennary N-glycans, commercially available bovine pancreatic RNase B was chosen as a model. It is a relatively small glycoprotein of 124 amino acids with a single N-glycosylation site at Asn34, which is attached with high mannose-type N-glycans (Man_{5,9}GlcNAc₂) containing five to nine mannose residues [32–34]. As eukaryotic glycosyltransferases involved in N-glycan processing are well known and the corresponding bacterial alternatives have not been fully identified, attempts to express the former in *E. coli* are one of the focuses of our enzymatic glycoprotein

in vitro N-glycan processing strategy. Meanwhile, bacterial alternatives of eukaryotic N-glycan processing mannosidases and glycosyltransferases that have already been identified with desired functions are great choices for developing the efficient glycoprotein in vitro N-glycan processing platform. As shown in Figure 1, the heterogeneous $\text{Man}_{5-9}\text{GlcNAc}_2$ N-glycans on RNase B can be processed to a homogeneous $\text{Man}_5\text{GlcNAc}_2$ N-glycan by removing all $\alpha 1-2$ -linked mannose residues, using a recombinant *Enterococcus faecalis* $\alpha 1-2$ -mannosidase (EfMan-I) expressed in *E. coli* that we reported previously [34]. EfMan-I belongs to the carbohydrate active enzyme (CAZY) glycoside hydrolase [35,36] family 92 (GH92) and requires a divalent metal cation, such as Ca^{2+} or Mg^{2+} for activity [34]. After EfMan-I treatment, the natural N-glycan processing steps [2] can be followed in vitro using recombinant glycosyltransferases and glycosidases from different origins to form target homogeneous glycoproteins with the desired disialylated biantennary complex-type N-glycans.




Figure 1. Glycoprotein in vitro N-glycan processing from high-mannose to disialylated biantennary complex-type N-glycans. The conserved trimannosyl chitobiose core ($\text{Man}_3\text{GlcNAc}_2$) of eukaryotic glycoprotein N-glycans is highlighted in a red rectangle.

Human N-acetylglucosaminyltransferase I (hGnT-I or hMGAT1) in the CAZY glycosyltransferase [37,38] family 13 (GT13) was chosen to add an N-acetylglucosamine (GlcNAc) residue $\beta 1-2$ -linked to the $\alpha 1-3$ -linked mannose residue on the trimannosyl chitobiose core of the $\text{Man}_5\text{GlcNAc}_2$ N-glycan on the EfMan-I-treated RNase B to form a hybrid-type $\text{GlcNAcMan}_5\text{GlcNAc}_2$ N-glycan. The hGnT-I uses uridine-5'-diphosphate GlcNAc (UDP-GlcNAc) as the donor substrate and requires a divalent metal cation, such as Mn^{2+} as a cofactor. It is highly selective toward $\text{Man}_5\text{GlcNAc}_2$ N-glycan with dramatically decreased activity for $\text{Man}_3\text{GlcNAc}_2$ and other high-mannose-type N-glycans [39]. This acceptor substrate preference is beneficial for our one-pot multienzyme (OPME) N-glycan processing approach described below.

To process the $\text{GlcNAcMan}_5\text{GlcNAc}_2$ N-glycan on glycoproteins further to form $\text{GlcNAcMan}_3\text{GlcNAc}_2$, the reported *Bacteroides thetaiotaomicron* $\alpha 1-6$ -mannosidase Bt3994 and $\alpha 1-3$ -mannosidase Bt1769 [40] were chosen. Similar to EfMan-I, they are Ca^{2+} -dependent CAZY GH92 bacterial mannosidases [40]. In contrast to EfMan-I, which is an $\alpha 1-2$ -mannosidase, Bt3994 was reported as an $\alpha 1-6$ -mannosidase to catalyze the cleavage of the terminal $\alpha 1-6$ -mannosidic linkage in $\text{Man}_5\text{GlcNAc}_2$ and $\text{Man}_3\text{GlcNAc}_2$ N-glycans

to form $\text{Man}_4\text{GlcNAc}_2$ and $\text{Man}_2\text{GlcNAc}_2$, respectively, while Bt1769 was reported as an $\alpha 1$ -3-mannosidase to catalyze the cleavage of the terminal unbranched $\alpha 1$ -3-linked mannose of the $\text{Man}_4\text{GlcNAc}_2$ N-glycan generated by Bt3994 to form $\text{Man}_3\text{GlcNAc}_2$ [40]. Based on the reported activities of Bt3994 and Bt1769, we hypothesized that they can coordinate with each other to convert $\text{GlcNAcMan}_5\text{GlcNAc}_2$ N-glycan on glycoproteins to $\text{GlcNAcMan}_3\text{GlcNAc}_2$.

To generate the second antenna in the glycoprotein biantennary complex N-glycans, human N-acetylglucosaminyltransferase II (hGnT-II or hMGAT2, CAZy family GT16) was chosen. It requires a divalent metal cation for catalyzing the transfer of GlcNAc from UDP-GlcNAc to form a $\beta 1$ -2-linkage to the $\alpha 1$ -6-linked terminal mannose on $\text{GlcNAcMan}_3\text{GlcNAc}_2$ to produce $\text{GlcNAc}_2\text{Man}_3\text{GlcNAc}_2$. hGnT-II has been shown to have a substrate binding pocket that interacts with both the $\alpha 1$ -6-linked terminal mannose as the glycosylation site and the other $\text{GlcNAc}\beta 1$ -2 $\text{Man}\alpha 1$ -3 $\text{Man}\beta$ branch as the additional “recognition arm” [41]. This high acceptor substrate selectivity, again, is advantageous for the OPME N-glycan engineering approach presented below.

The biantennary $\text{GlcNAc}_2\text{Man}_3\text{GlcNAc}_2$ complex-type N-glycan on RNase B generated from the hGnT-II reaction can be $\beta 1$ -4-galactosylated using a well-known bovine $\beta 1$ -4-galactosyltransferase 1 (B β 4GalT1) [42–45], a CAZy GT7 family enzyme, to form RNase B containing $\text{Gal}_2\text{GlcNAc}_2\text{Man}_3\text{GlcNAc}_2$ N-glycan using uridine 5'-diphosphate galactose (UDP-Gal) as the donor substrate.

The in vitro N-glycan processing can be completed by a final sialylation step using a suitable sialyltransferase in the presence of CMP-sialic acid to form RNase B containing homogeneous $\alpha 2$ -3- or $\alpha 2$ -6-sialylated biantennary complex-type N-glycans ($\text{Sia}_2\text{Gal}_2\text{GlcNAc}_2\text{Man}_3\text{GlcNAc}_2$). Different sialic acid forms can be introduced in the enzymatic sialylation step [46] and the most common sialic acid form, N-acetylneuraminic acid (Neu5Ac), is introduced as an example in our study presented here.

2.2. Enzyme Cloning and Expression

To facilitate the purification of enzymes that will be used for glycoprotein in vitro N-glycan processing, an His₆-tag was introduced at the C-terminus of each recombinant enzyme to allow its easy purification by Ni²⁺-affinity columns. Furthermore, we found that fusing a maltose binding protein (MBP) at the N-terminus of the target recombinant protein and removing the N-terminal transmembrane domain of mammalian glycosyltransferases by truncation worked well to improve their soluble expression in *E. coli* [47]. These were the strategies that guided our general design to construct the plasmids for expressing target recombinant enzymes. In addition, *E. coli* Origami B (DE3) strain harboring pGro7 for chaperon expression was found to be a better choice for expressing mammalian enzymes than *E. coli* BL21 (DE3) strain, which was used to express recombinant enzymes from bacterial origins.

As we reported previously [34], EfMan-I was expressed as a C-terminal His₆-tagged soluble and active enzyme in *E. coli* BL21 (DE3) cells with an expression level of 85 mg/L LB culture (Table 1).

Although *Bacteroides thetaiotaomicron* mannosidases Bt3994 and Bt1769 were cloned previously in pET21a vector and expressed in *E. coli* Turner or B834 cells as C-terminal His₆-tagged recombinant proteins [40], their expression levels were not reported. We found that removing Bt3994 N-terminal 24 amino acid residues and Bt1769 N-terminal 18 amino acid residues significantly improved their soluble expression levels in *E. coli* BL21 (DE3) cells. Both $\Delta 24\text{Bt3994}$ -His₆ and $\Delta 18\text{Bt1769}$ -His₆ were expressed at a level of around 55 mg/L LB media (Table 1) as soluble and active proteins with the expected molecular weights of 82 and 83 kDa, respectively (Figure 2A,B).

Table 1. Enzymes obtained for glycoprotein in vitro N-glycan processing and their expression levels in *E. coli* BL21 (DE3) or Origami B (DE3) cells.

Enzyme	CAZY Family	Source	Expression Host	Expression Level (mg/L LB)
EfMan-I-His ₆	GH92	<i>Enterococcus faecalis</i> V583	BL21 (DE3)	85 [a]
Δ24Bt3994-His ₆	GH92	<i>Bacteroides thetaiotaomicron</i>	BL21 (DE3)	55
Δ18Bt1769-His ₆	GH92	<i>Bacteroides thetaiotaomicron</i>	BL21 (DE3)	55
MBP-Δ28hGnT-I-His ₆	GT13	Human	Origami B (DE3)/pGro7	5
MBP-Δ27hGnT-II-His ₆	GT16	Human	Origami B (DE3)/pGro7	~1
MBP-Δ128Bβ4GalT1-His ₆	GT7	Bovine	Origami B (DE3)/pGro7	60
MBP-Δ89ST6GAL-I-His ₆	GT29	Human	Origami B (DE3)/pGro7	30
MBP-CjCst-IΔ145-His ₆	GT42	<i>Campylobacter jejuni</i>	BL21 (DE3)	60

[a] As reported in Ref. [34].




Figure 2. SDS-PAGE analysis results for enzyme expression and purification. (A) Δ24Bt3994-His₆, (B) Δ18Bt1769-His₆, (C) MBP-Δ28hGnT-I-His₆, (D) MBP-Δ27hGnT-II-His₆, (E) MBP-Δ128Bβ4GalT1-His₆, (F) MBP-Δ89ST6GAL-I-His₆, (G) MBP-CjCst-IΔ145-His₆. Lanes: BI, before induction; AI, after induction; Lys, lysate; PP, purified protein; M, PageRuler™ Plus Prestained Protein Ladder, 10 to 250 kDa. Target proteins are marked with an arrow on the right.

To obtain active and soluble recombinant hGnT-I and hGnT-II from *E. coli*, their N-terminal sequences containing the putative transmembrane domains were removed, and the truncated sequences were expressed as fusion proteins with an N-terminal maltose binding protein (MBP) and a C-terminal His₆-tag. The resulting MBP-Δ28hGnT-I-His₆ and MBP-Δ27hGnT-II-His₆ were expressed at a level of 5 and 1 mg/L, respectively (Table 1), with the expected molecular weights of 91 and 92 kDa (Figure 2C,D).

Bβ4GalT1 expressed in *E. coli* has been purified from inclusion bodies [42,44,48]. We previously cloned and expressed an N-terminal 128 amino acid-truncated Bβ4GalT1 in pET15b as an N-terminal His₆-tagged fusion protein (His₆-Δ128Bβ4GalT1) in *E. coli* BL21 (DE3) cells, which had a relatively low soluble expression (<1 mg/L culture) [45]. We redesigned the construct to express a protein with both an N-terminal MBP fusion and C-terminal His₆-tag. The resulting MBP-Δ128Bβ4GalT1-His₆ was expressed in a dramatically improved 60 mg/L yield (Table 1) with an expected molecular weight at 75 kDa (Figure 2E).

For sialyltransferases, in addition to the recombinant sialyltransferases that we previously expressed in *E. coli* including bacterial α 2–3-sialyltransferases, such as *Pasteurella multocida* α 2–3-sialyltransferase 1 (PmST1) [49] and its mutant PmST1_M144D [50], *Pasteurella multocida* α 2–3-sialyltransferase 3 (PmST3) [51], as well as α 2–6-sialyltransferases, such as *Photobacterium damsela* α 2–6-sialyltransferase (Pd2,6ST) [52] and its mutant Pd2,6ST_A200Y/S232Y [53], *Photobacterium* species α 2–6-sialyltransferase (Psp2,6ST) [54] and its mutant Psp2,6ST_A366G [55], we cloned and expressed a recombinant human α 2–6-sialyltransferase hST6GAL-I and a recombinant *Campylobacter jejuni* α 2–3-sialyltransferase CjCst-I.

Similar to other eukaryotic sialyltransferases, hST6GAL-I is a CAZy GT29 enzyme [56]. It has been shown to selectively α 2–6-sialylate glycoprotein N-glycans [57] and has been successfully expressed in *E. coli* as a soluble and active fusion protein with MBP at its N-terminus [31]. The soluble expression of the N-terminal MBP-fused and C-terminal His₆-tagged N-terminal truncated hST6GAL-I (MBP- Δ 89hST6GAL-I-His₆) that we constructed reached 30 mg/L LB culture (Table 1) with an expected molecular weight at 80 kDa (Figure 2F).

Campylobacter jejuni CjCst-I is an α 2–3-sialyltransferase belonging to CAZy GT42 family [58,59]. It has been shown to utilize Gal β 1–3/4OR as acceptors. The 145 residues at the C terminus of CjCst-I were removed and MBP was fused at the N terminus to form MBP-CjCst-I Δ 145-His₆. It was successfully expressed in *E. coli* as a soluble and active fusion protein at a level of 60 mg/L LB culture with an expected molecular weight at 77 kDa (Figure 2G).

2.3. Step-Wise Reactions and Enzyme Activity Determination Using Glycoprotein Substrates

With all enzymes in hand, their activities and applications for glycoprotein in vitro N-glycan processing were tested using step-wise enzymatic reactions and the product formation was monitored by matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry (MS) analysis. Reaction conditions were optimized by varying the types of the buffers used, pH, temperature, ion strength, incubation time, etc.

As shown in Figure 3, treating RNase B (5 mg/mL) (Figure 3A) with 3% (*w/w*) EfMan-I-His₆ at 30 °C for 2 h, the high-mannose-type N-glycans Man_{5–9}GlcNAc₂ on RNase B were completely trimmed down to Man₅GlcNAc₂ (Figure 3B). Treating the resulting RNase B sample with 3% (*w/w*) MBP- Δ 28hGnT-I in the presence of MnCl₂ (2 mM) and UDP-GlcNAc (1 mM) at 30 °C for 2 h completed the reaction for the formation of RNase B with a homogeneous GlcNAc₁Man₅GlcNAc₂ N-glycan (Figure 3C). Incubation of the resulting RNase B with 4% (*w/w*) Δ 24Bt3994-His₆ and 3% (*w/w*) Δ 18Bt1769-His₆ in the presence of 2 mM CaCl₂ at 30 °C for 2 h completed the cleavage of the terminal α 1–6- and α 1–3-linked mannose residues to form RNase B with a homogeneous GlcNAc₁Man₃GlcNAc₂ N-glycan, which was not cleaved further in the presence of both Δ 24Bt3994-His₆ and Δ 18Bt1769-His₆ (Figure 3D). The resulting reaction mixture was incubated with MBP- Δ 27hGnT-II-His₆ (10% *w/w*) in the presence of 2 mM MnCl₂ and 1 mM UDP-GlcNAc at 30 °C overnight to form RNase B containing homogeneous GlcNAc₂Man₃GlcNAc₂ N-glycan (Figure 3E). MBP- Δ 128B β 4GalT1-His₆ (3% *w/w*) was then used to process the N-glycan in the resulting RNase B to form RNase B containing homogeneous Gal₂GlcNAc₂Man₃GlcNAc₂ N-glycan by incubating at 30 °C in 2 h in the presence of 5 mM MnCl₂ and 2 mM UDP-Gal (Figure 3F). Notably, all enzymes used for RNase B N-glycan processing were active in the presence of Tris-HCl (100 mM, pH 7.5).




Figure 3. MALDI-TOF MS analysis results for RNase B with heterogeneous high-mannose N-glycans (A) and RNase B sequentially treated with *EfMan-I-His₆* (B), *MBP-Δ28hGnT-I-His₆* (C), *Δ24Bt3994-His₆* and *Δ18Bt1769-His₆* (D), *MBP-Δ27hGnT-II-His₆* (E), and *MBP-Δ128Bp4GalT1-His₆* (F) in step-wise reactions. The peak marked with an asterisk (*) in each figure is for RNase A, the non-glycosylated ribonuclease that is present in the commercially obtained RNase B sample, which was used as the internal standard. The schematic illustrations of the corresponding reactions are shown. The number above each peak represents the m/z value of the expected product + 2Na species.

2.4. Multi-Step OPME N-Glycan Processing

Due to the high specificity of the acceptor substrate preference of mammalian glycosyltransferases and the high efficiency of the bacterial mannosidases used in the glycoprotein *in vitro* N-glycan processing described above, we hypothesized that the step-by-step process was not necessary, and one-pot approaches were possible and could be more efficient. To test this hypothesis, a series of one-pot multienzyme (OPME) reactions were carried out and the resulting RNase B samples were analyzed directly by MALDI-TOF MS assays after dialysis.

Indeed, reactions with *EfMan-I-His₆* (3% *w/w*) alone, one-pot two-enzyme (OP2E) reactions containing *EfMan-I-His₆* (3% *w/w*) and *MBP-Δ28hGnT-I-His₆* (3% *w/w*), as well

as one-pot four-enzyme (OP4E) containing EfMan-I-His₆ (3% *w/w*), MBP- Δ 28hGnT-I-His₆ (3% *w/w*), Δ 18Bt1769-His₆ (3% *w/w*), and Δ 24Bt3994-His₆ (4% *w/w*) were very efficient. The RNase B products containing the expected N-glycan structures were all obtained in 2 h at 30 °C (Figure 4A–C). In comparison, one-pot five-enzyme (OP5E) reactions containing EfMan-I-His₆ (3% *w/w*), MBP- Δ 28hGnT-I-His₆ (3% *w/w*), Δ 18Bt1769-His₆ (3% *w/w*), Δ 24Bt3994-His₆ (4% *w/w*), and MBP- Δ 27hGnT-II-His₆ (10% *w/w*) were slower, and the RNase B product containing the target N-glycan was obtained after incubation at 30 °C overnight (~18 h) (Figure 4D). This indicated that the addition of the second GlcNAc catalyzed by MBP- Δ 27hGnT-II-His₆ was the rate limiting step of the process in this system. It is worth mentioning that further treatment of the OP4E product with Δ 24Bt3994-His₆ (5% *w/w*) or Δ 18Bt1769-His₆ (3% *w/w*) at 30 °C overnight did not lead to further cleavage of the N-glycan (data not shown), highlighting the applicable combination of Δ 18Bt1769-His₆ and Δ 24Bt3994-His₆ in glycoprotein N-glycan processing from high-mannose N-glycans without the concern of removing extra mannose residues.




Figure 4. MALDI-TOF MS analysis results for RNase B treated with EfMan-I-His₆ for 2 h (A), OP2E containing EfMan-I-His₆, MBP- Δ 28hGnT-I-His₆ for 2 h (B), OP4E containing EfMan-I-His₆, MBP- Δ 28hGnT-I-His₆, Δ 24Bt3994-His₆, and Δ 18Bt1769-His₆ for 2 h (C), OP5E containing EfMan-I-His₆, MBP- Δ 28hGnT-I-His₆, Δ 24Bt3994-His₆, Δ 18Bt1769-His₆, and MBP- Δ 27hGnT-II-His₆ for 18 h (D), and OP5E for 18 h followed by incubation with MBP- Δ 128B β 4GalT1-His₆ for additional 2 h (E). All reactions were carried out at 30 °C in Tris-HCl (100 mM, pH 7.5) containing CaCl₂ (2 mM), MgCl₂ (2 mM), and MnCl₂ (2 mM). The peak marked with an asterisk (*) in each figure is for RNase A, the non-glycosylated ribonuclease that is present in the commercially obtained RNase B sample, which was used as the internal standard. The symbol representations of the expected products are shown.

The N-glycan of the RNase B product of the OP5E reaction was released by peptide:N-glycosidase F (PNGase F) and analyzed by MALDI-TOF MS assays. The results (Figure 5) were consistent with those obtained with the intact RNase B samples. Only the target N-glycan GlcNAc₂Man₃GlcNAc₂ (+Na, $m/z = 1339.737$) was observed without the presence of other N-glycans, confirming the efficiency of the OP5E reactions and the N-glycan homogeneity of the RNase B glycoprotein obtained.




Figure 5. MALDI-TOF MS analysis result of the N-glycan released from the RNase B product of OP5E reactions; m/z calculated for the sodium adduct of the product GlcNAc₂Man₃GlcNAc₂ [M + Na]⁺ was 1339.476, found 1339.737.

Incubating the reaction mixture of the OP5E reaction with UDP-Gal (2 mM) and MBP- $\Delta 128\beta 4\text{GalT1-His}_6$ (3% w/w) at 30 °C for 2 h completed the formation of RNase B with Gal₂GlcNAc₂Man₃GlcNAc₂ N-glycan (Figure 4E).

The formation of RNase B with disialylated biantennary complex N-glycan was accomplished by incubating with MBP- $\Delta 89\text{hST6GAL-I-His}_6$ (3% w/w) or MBP-CjCst- $\Delta 145\text{-His}_6$ (3% w/w) in the presence of cytidine 5'-monophosphate-Neu5Ac (CMP-Neu5Ac, 5 mM) for the formation of $\alpha 2-6$ or $\alpha 2-3$ -sialyl linkage, respectively. It was observed that sialic acids were cleaved from sialosides during MALDI-TOF analysis. Therefore, high-resolution mass spectrometry (HRMS) analysis was used to analyze the N-glycans released from RNase B sialylation products by PNGase F digestion and purified by Cotton HILIC SPE microtips [60]. We found that in situ generation of CMP-Neu5Ac from Neu5Ac and CTP by *Neisseria meningitidis* CMP-sialic acid synthetase (NmCSS) [61] during enzymatic sialylation process in a one-pot two-enzyme system [62] further improved the efficiency of sialylation. As shown in Figure 6, the ionized species of disialyl biantennary N-glycans released from RNase B sialylation products (Neu5Ac₂Gal₂GlcNAc₂Man₃GlcNAc₂, m/z value expected 1110.3842; m/z values observed were 1110.3849 and 1110.3848 for the double-charged $\alpha 2-6$ and $\alpha 2-3$ -linked species, respectively) were clearly observed. Ionized species for monosialylated N-glycan (Neu5Ac₁Gal₂GlcNAc₂Man₃GlcNAc₂, m/z value expected: 1930.6803) and N-glycan released from the RNase B substrate for sialylation reactions (Gal₂GlcNAc₂Man₃GlcNAc₂, m/z value expected 1663.5819, or 1675.5610 for its Cl adduct) were not observed. This indicated that the di-sialylation reactions went to completion.




Figure 6. HRMS (negative mode) assay results for the N-glycans released from the RNase B products produced by a one-pot two-enzyme (OP2E) sialylation reaction containing *Neisseria meningitidis* CMP-sialic acid synthetase (NmCSS) and MBP- Δ 89ST6GAL-I-His₆ (**A**) or MBP-CjCst-I Δ 145-His₆ (**B**). The m/z value expected for the ionized disialyl N-glycan (Neu5Ac₂Gal₂GlcNAc₂Man₃GlcNAc₂) was 1110.3842. The ionized substrate (m/z value expected 1663.5819), substrate + Cl (m/z value expected 1675.5610), or monosialylated N-glycan (m/z value expected 1930.6803) was not observed.

3. Conclusions

Using RNase B as a model glycoprotein substrate, we have successfully established a glycoprotein in vitro N-glycan processing platform for the production of glycoproteins containing homogeneous α 2-6- or α 2-3-linked disialylated biantennary complex N-glycans using enzymes expressed in *E. coli*. Several mammalian glycoprotein N-glycan processing glycosyltransferases including hGnT-I, hGnT-II, B β 4GalT1, and hST6GAL-I have been successfully expressed in *E. coli* Origami B (DE3) cells as N-terminal MBP-fused and C-terminal His₆-tagged fusion proteins in a soluble and active form. In addition, bacterial mannosidases including EfManI, Bt3994, and Bt1769, as well as a bacterial sialyltransferase CjCst-I have been successfully expressed as His₆-tagged proteins. These enzymes can be easily purified using a single Ni²⁺-column. The in vitro processing of the high-mannose-type N-glycans in glycoprotein RNase B has been successfully achieved with the combination of these enzymes used in sequential step-wise reactions or in one-pot reactions. The platform developed can find broad applications for producing glycoproteins with homogeneous N-glycoforms.

4. Materials and Methods

4.1. Materials and Instruments

Chemicals were purchased and used as received without further purification. Matrix-assisted laser desorption/ionization mass spectra were obtained using Bruker UltraFlex-treme MALDI-TOF/TOF (Bruker, Billerica, MA, USA) and high-resolution mass spectrometry results were obtained using Thermo Q-Exactive HF (Thermo Fisher Scientific, Asheville, NC, USA) at the University of California-Davis Campus Mass Spectrometry Facilities. Proteins were purified using NGC medium-pressure liquid chromatography systems (Bio-Rad, Hercules, CA, USA). Peptide N-glycosidase F (PNGase F) was obtained from R&D Systems, Minneapolis, Minnesota, USA. Vector plasmids pMAL-c4X and pMAL-c2X were from New England Biolabs, Inc., Ipswich, MA, USA. Vector plasmids pET15b and pET22b(+) were from Novagen, Madison, WI, USA. Profinity and Nuvia IMAC Ni-charged resins, and mini Nuvia IMAC cartridges (5 mL) were from Bio-Rad, Hercules, California, USA. Galactose, *N*-acetylglucosamine (GlcNAc), LB media, isopropyl-1-thio-D-galactopyranoside (IPTG), and ampicillin were from Fisher Scientific, Inc., Hampton, NH, USA. Kanamycin sulfate was from AMRESCO, Fairlawn, NJ, USA. Chloramphenicol was from ALDRICH, St. Louis, MO, USA. Tetracycline hydrochloride was from Calbiochem, La Jolla, CA, USA. *N*-Acetylneuraminic acid (Neu5Ac) was from Inalco (Milano, Italy). Adenosine 5'-triphosphate (ATP), cytosine 5'-triphosphate (CTP), and uridine 5'-triphosphate (UTP) were purchased from Hangzhou Meiya Pharmaceutical Co. Ltd. (Hangzhou, China). DH5 α competent cells, BL21 (DE3) competent cells, restriction enzymes, GeneJET Gel Extraction Kit, GeneJET Plasmid Miniprep Kit, PhusionTM Plus DNA Polymerase, T4 ligase, and 2,5-dihydroxybenzoic acid (2,5-DHB) were purchased from Thermo Fisher Scientific, Asheville, NC, USA. Sinapinic (or sinapic) acid was from TCI America, Portland, OR, USA. Origami B (DE3) competent cells were from Novagen and pGro7 was from Takara Bio USA Inc, Mountain View, CA, USA. Herculase-enhanced DNA polymerase was from Agilent Technologies, Santa Clara, California, USA. UDP-GlcNAc [63], UDP-Gal [64], and CMP-Neu5Ac [61] were prepared as described previously.

4.2. Cloning

All polymerase chain reactions (PCRs) were carried out with Phusion[®] HF DNA polymerase by following the standard protocol provided by the manufactory unless noted. Briefly, the reaction was performed in a reaction mixture (50 μ L) containing a template (10 ng plasmid or synthetic DNA, or 1 μ g of genomic DNA), 10 \times Phusion[®] HF buffer (5 μ L), dNTP mixture (1 mM each), and 5 U (1 μ L) of Phusion[®] HF DNA polymerase, forward and reverse primers (1 μ M each). The reaction mixture was subjected to 30 cycles of amplification. The primers and the annealing temperature (T_a) used for each PCR reaction are listed in Table 2. The PCR products were purified by GeneJET Gel Extraction Kit and digested with two restriction enzymes at 37 $^{\circ}$ C for 2 h. The digested products were purified by GeneJET Gel Extraction Kit and ligated with vector plasmid pre-digested with the same restriction enzymes and similarly gel extraction purified. The ligation was carried out at 16 $^{\circ}$ C overnight using T4 DNA ligase. The ligated product was transformed into the chemical competent *E. coli* DH5 α cells. Plasmids were purified using GeneJET Plasmid Miniprep Kit and sequences were confirmed by DNA sequencing (See Supplementary Materials). Positive plasmids were selected and transferred to chemically competent *E. coli* BL21 (DE3) or Origami B (DE3) cells for expression.

number: XP_019821962.1) as an N-terminal MBP-fused and C-terminal His₆-tagged fusion protein MBP- Δ 128B β 4GalT1-His₆.

MBP- Δ 89hST6GAL-I-His₆: A synthetic gene of an N-terminal 189 amino acid truncated human ST6GAL-I (hST6GAL-I, GenBank accession number: NP_001340845.1) with codon optimized for *E. coli* expression was cloned in pMAL-c2X vector to construct plasmid pMAL-c2X- Δ 89hST6GAL-I for expressing MBP- Δ 89hST6GAL-I-His₆.

MBP-CjCst-I Δ 145-His₆: A synthetic gene of C-terminal 145 amino acids truncated *Campylobacter jejuni* Cst-I (CjCst-I, GenBank accession number: AAF13495.1) with codon optimized for *E. coli* expression was cloned into pMAL-c4X to construct plasmid pMAL-c4X-CjCst-I Δ 145 for expressing MBP-CjCst-I Δ 145-His₆.

4.3. Enzyme Expression and Purification

To express enzymes in *E. coli* BL21 (DE3) expression system, cells harboring the plasmid of interest were cultured in LB media (10 g/L tryptone, 5 g/L yeast extract, and 10 g/L NaCl) supplemented with ampicillin (100 μ g/mL). When the OD_{600 nm} of the culture reached 0.6–0.8, isopropyl-1-thio-D-galactopyranoside (IPTG, 0.1 mM) was added and the culture was incubated at 20 °C for 20 h. To express enzymes in *E. coli* Origami B (DE3) harboring pGro7 and the plasmid of interest, cells were cultured in LB media supplemented with ampicillin (50 mg/mL), tetracycline (5 mg/L), chloramphenicol (17 mg/L), kanamycin (25 mg/L), and L-arabinose (1 g/L, for chaperon expression). When the OD_{600 nm} of the culture reached 0.6–0.8, IPTG (0.1 mM) was added and the culture was incubated at 16 °C for 48 h.

After the expression was completed, cells were harvested by centrifugation (6000 \times g) at 4 °C for 30 min and re-suspended in lysis buffer (50 mM Tris-HCl, pH 8.0, 300 mM NaCl, 0.1% Triton X-100). The cell resuspension was subjected to the homogenizer (EmulsiFlex-C3) to break the cells. The cell lysate was obtained as the supernatant after centrifugation (9016 \times g) at 4 °C for 60 min and purified by a Ni²⁺-NTA affinity column, such as a mini Nuvia IMAC cartridge (5 mL), on a Bio-Rad NGC system. The column was pre-equilibrated with 6 column volumes of binding buffer containing Tris-HCl buffer (50 mM, pH 8.0), NaCl (300 mM). It was washed with 10 column volumes of binding buffer, followed by washing with 10 column volumes of 10% elute buffer, and 10 column volumes of 20% elute buffer, and then eluted with elute buffer containing Tris-HCl (50 mM, pH 8.0), NaCl (300 mM), imidazole (250 mM). The fractions containing the purified protein were combined for dialysis against a dialysis buffer (Tris-HCl, 50 mM, pH 7.5, 250 mM NaCl) or for concentration using a protein concentrator (10 kDa cut-off). Finally, 20% glycerol (for MBP- Δ 128B β 4GalT1-His₆) or 10% glycerol (for other enzymes) was added before storing the samples at –20 °C.

4.4. Stepwise Enzymatic Reaction Using RNase B as The Substrate

RNase B (5 mg/mL, 330 μ M) was incubated with EfMan-I-His₆ (150 μ g/mL, 1.8 μ M) in Tris-HCl (100 mM, pH 7.5) at 30 °C for 2 h. The resulting mixture was then dialyzed against ddH₂O and subjected to MALDI-TOF MS analysis. It was also used for the assays described below.

The dialyzed EfMal-I-His₆-treated RNase B (~330 μ M) was used as the acceptor substrate. It was incubated with UDP-GlcNAc (1 mM) and MBP- Δ 27hGnT-I-His₆ (1.6 μ M) in Tris-HCl (100 mM, pH 7.5) containing MnCl₂ (2 mM) at 30 °C for 2 h. The weight ratio of MBP- Δ 27hGnT-I-His₆ to RNase B was 3% (*w/w*). The resulting mixture was then dialyzed against ddH₂O and subjected to MALDI-TOF MS analysis. It was also used for the assays described below.

The RNase B obtained above was incubated with Δ 24Bt3994-His₆ (1.9 μ M) and Δ 18Bt1769-His₆ (1.8 μ M) in Tris-HCl (100 mM, pH 7.5) containing CaCl₂ (2 mM) at 30 °C for 2 h. The weight ratios of Δ 24Bt3994-His₆ and Δ 18Bt1769-His₆ to RNase B were 4% (*w/w*) and 3% (*w/w*), respectively. The resulting mixture was then dialyzed against ddH₂O and subjected to MALDI-TOF MS analysis. It was also used for the assays described below.

The RNase B obtained above was incubated with UDP-GlcNAc (2 mM) and MBP- Δ 27hGnT-II-His₆ (3.2 μ M) in Tris-HCl (100 mM, pH 7.5) containing MnCl₂ (2 mM) at 30 °C for 2 h. The weight ratio of Δ 27hGnT-II to RNase B was 10% (*w/w*). The resulting mixture was then dialyzed against ddH₂O and subjected to MALDI-TOF MS analysis. It was also used for the assays described below.

The RNase B obtained above was incubated with UDP-Gal (2 mM) and MBP- Δ 128B β 4GalT1-His₆ (3.9 μ M) in Tris-HCl (100 mM, pH 7.5) containing MnCl₂ (2 mM) at 30 °C for 2 h. The weight ratio of MBP- Δ 128B β 4GalT1-His₆ to RNase B was 10% *w/w*. The resulting mixture was then dialyzed against ddH₂O and subjected to MALDI-TOF MS analysis.

Dialysis was not necessary for step-wise reactions, but was preferred before MALDI-TOF MS analysis of the samples.

4.5. OPME Reactions for N-Glycan Processing of RNase B

One-pot two-enzyme (OP2E) reactions were carried out in a 0.5 mL microcentrifuge tube by incubating RNase B (330 μ M), UDP-GlcNAc (2 mM), EfMan-I-His₆ (1.8 μ M, 3% *w/w*), and MBP- Δ 28hGnT-I-His₆ (1.6 μ M, 3% *w/w*), in Tris-HCl (100 mM, pH 7.5) containing MnCl₂ (2 mM), CaCl₂ (2 mM), MnCl₂ (2 mM) at 30 °C for 2 h.

One-pot two-enzyme (OP2E) reactions were carried out similarly as described above for OP2E reactions, except for the fact that two more enzymes including Δ 24Bt3994-His₆ (1.9 μ M, 4% *w/w*) and Δ 18Bt1769-His₆ (1.8 μ M, 3% *w/w*) were added.

One-pot two-enzyme (OP5E) reactions were carried out similarly to OP4E reactions except for the fact that one more enzyme MBP- Δ 27hGnT-II-His₆ (3.2 μ M, 10% *w/w*) was added and the reaction was continued overnight.

Galactosylation was accomplished by adding UDP-Gal (2 mM) and MBP- Δ 128B β 4GalT1-His₆ (3.9 μ M, 10% *w/w*) to the reaction mixture after OP5E reaction followed by incubation at 30 °C for 2 h.

The final step α 2–6-sialylation was completed by adding CMP-Neu5Ac (5 mM) and MBP- Δ 89hST6GAL-I-His₆ (1.9 μ M, 3% *w/w*) to the reaction mixture after galactosylation followed by incubation at 30 °C overnight.

Similarly, α 2–3-sialylation was completed by adding CMP-Neu5Ac (5 mM) and MBP-CjCst-I Δ 145-His₆ (1.9 μ M, 3% *w/w*) to the reaction mixture after galactosylation followed by incubation at 30 °C for a shorter time of 2 h.

For sialylation reactions with in situ generation of the sialyltransferase donor, CMP-Neu5Ac (5 mM) was replaced by Neu5Ac (5 mM), CTP (7.5 mM), and NmCSS (5 μ M). Other conditions were the same as described above.

4.6. MALDI-TOF MS Analyses of RNase B samples and the Released N-Glycans

Fresh solutions of 2,5-DHB (15 mg/mL) dissolved in ddH₂O and sinapinic acid (SA) (20 mg/mL) dissolved in ACN/0.1%TFA (7:3) were prepared. A mixed solution of these two with a 1:1 (*v/v*) ratio was used as the matrix for MALDI-TOF MS analysis of RNase B samples (~1 mg/mL) dialyzed against ddH₂O using Slide-A-Lyzer™ MINI Dialysis (10 k MWCO) devices.

To release N-glycans from RNase B samples, RNase B (1 mg) was denatured by adding 0.5% SDS and dithiothreitol (DTT) (40 mM) in 200 μ L followed by incubation at 98 °C for 10 min and then at room temperature for 5 min. PNGase F (100 ng) was then added and the glycans were released by incubation at 37 °C for 2 h. The deglycosylated proteins were precipitated by adding three volumes of pre-chilled ethanol followed by incubation on ice for 20 min. The mixtures were then centrifuged at 16,200 \times *g* for 5 min and the supernatants containing the glycans were purified with graphitized carbon cartridge and dried in a speed vacuum. They were dissolved in ddH₂O and used for MALDI-TOF MS analysis using 2,5-DHB dissolved in ACN/0.1%TFA (7:3) (25 mg/mL) as the matrix.

4.7. HRMS Analysis of Sialylated N-Glycans

RNase B (100 µg) samples were denatured and the N-glycans were released by treating with PNGase F (10 ng) similar to the conditions described above. The samples were cleaned using a homemade cotton tip via hydrophilic interaction liquid chromatography-solid phase extraction (HILIC-SPE) [60]. Briefly, samples were mixed with acetonitrile (ACN, 50% *v/v*) centrifuged (16,200 × *g*) at 4 °C for 5 min. The supernatant was transferred to a clean tube, pipetting up-and-down for a total of 20 times in a 10 µL-tip packed with a small volume of cotton. The cotton tip was washed 3 times with 20 µL of 85% ACN with 1% TFA, followed by 3 times with 20 µL of 85% ACN, and eluted with ddH₂O (10 µL). The eluant was used for HRMS analysis.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/molecules28062753/s1>. The DNA and amino acid sequences of MBP-Δ28hGnT-I-His₆, Δ24Bt3994-His₆, Δ18Bt1769-His₆, MBP-Δ27hGnT-II-His₆, MBP-Δ128Bβ4GalT1-His₆, MBP-Δ89hST6GAL-I-His₆, and MBP-CjCst-IΔ145-His₆.

Author Contributions: Conceptualization, L.Z., Y.L., R.L. and X.C.; data curation, L.Z., Y.L., R.L., X.Y., Z.Z., J.F. and H.Y.; formal analysis, L.Z., Y.L., R.L. and X.C.; funding acquisition, X.C.; investigation, L.Z., Y.L., R.L., X.Y., Z.Z., J.F., H.Y. and X.C.; methodology, L.Z., Y.L., R.L., X.Y., Z.Z., J.F., H.Y. and X.C.; project administration, X.C.; resources, X.C.; supervision, X.C.; writing—original draft, L.Z., Y.L., R.L., X.Y., Z.Z. and X.C.; writing—review and editing, L.Z., Y.L., R.L., X.Y., Z.Z., J.F., H.Y. and X.C. All authors have read and agreed to the published version of the manuscript.

Funding: This work was financially supported by the United States (US) Defense Threat Reduction Agency (HDTRA1-15-1-0054) and by the financial assistance award 70NANB22H017 from US Department of Commerce (DOC), National Institute of Standards and Technology (NIST). Bruker UltraFLEXtreme MALDI TOF/TOF was funded by NIH grant No. S10OD018913. Thermo Q-Exactive HF (High-field Orbitrap) was funded by NIH grant No. S10OD025271. The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official view of DTRA, DOC, NIST, or NIH. The funding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the wiring of the manuscript; and in the decision to publish the results.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors would like to acknowledge the collaboration of Karen A. McDonald and Somen Nandi on glycoprotein in vitro enzymatic glycoengineering projects. The authors would like to thank William Jewell for his assistance and advice on the MALDI-TOF MS and HRMS analyses.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Li, Y.; Li, X.; Chen, X. Glycoproteins: Chemical features and biological roles. In *Carbohydrates Chemistry: State-of-the-Art and Challenges for Drug Development*; Cipolla, L., Ed.; Imperial College Press: London, UK, 2015; pp. 3–33.
2. Stanley, P.; Moremen, K.W.; Lewis, N.E.; Taniguchi, N.; Aebi, M. N-Glycans. In *Essentials of Glycobiology*, 4th ed.; Varki, A., Cummings, R.D., Esko, J.D., Stanley, P., Hart, G.W., Aebi, M., Mohnen, D., Kinoshita, T., Packer, N.H., Prestegard, J.H., et al., Eds.; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, NY, USA, 2022; pp. 103–116.
3. Borza, B.; Hajba, L.; Guttman, A. N-glycan analysis in molecular medicine: Innovator and biosimilar protein therapeutics. *Curr. Mol. Med.* **2020**, *20*, 828–839. [[CrossRef](#)]
4. Wang, L.X.; Tong, X.; Li, C.; Giddens, J.P.; Li, T. Glycoengineering of antibodies for modulating functions. *Annu. Rev. Biochem.* **2019**, *88*, 433–459. [[CrossRef](#)] [[PubMed](#)]
5. Delobel, A. Glycosylation of therapeutic proteins: A critical quality attribute. *Methods Mol. Biol.* **2021**, *2271*, 1–21. [[PubMed](#)]
6. Majewska, N.I.; Tejada, M.L.; Betenbaugh, M.J.; Agarwal, N. N-Glycosylation of IgG and IgG-like recombinant therapeutic proteins: Why is it important and how can we control it? *Annu. Rev. Chem. Biomol. Eng.* **2020**, *11*, 311–338. [[CrossRef](#)]
7. Higel, F.; Seidl, A.; Sorgel, F.; Friess, W. N-Glycosylation heterogeneity and the influence on structure, function and pharmacokinetics of monoclonal antibodies and Fc fusion proteins. *Eur. J. Pharm. Biopharm.* **2016**, *100*, 94–100. [[CrossRef](#)] [[PubMed](#)]

8. Cymer, F.; Beck, H.; Rohde, A.; Reusch, D. Therapeutic monoclonal antibody N-glycosylation—Structure, function and therapeutic potential. *Biologicals* **2018**, *52*, 1–11. [[CrossRef](#)] [[PubMed](#)]
9. Zhou, Q.; Qiu, H. The mechanistic impact of N-glycosylation on stability, pharmacokinetics, and immunogenicity of therapeutic proteins. *J. Pharm. Sci.* **2019**, *108*, 1366–1377. [[CrossRef](#)] [[PubMed](#)]
10. Reusch, D.; Tejada, M.L. Fc glycans of therapeutic antibodies as critical quality attributes. *Glycobiology* **2015**, *25*, 1325–1334. [[CrossRef](#)]
11. Jones, J.; Krag, S.S.; Betenbaugh, M.J. Controlling N-linked glycan site occupancy. *Biochim. Biophys. Acta* **2005**, *1726*, 121–137. [[CrossRef](#)] [[PubMed](#)]
12. de Haas, P.; Hendriks, W.; Lefeber, D.J.; Cambi, A. Biological and technical challenges in unraveling the role of N-glycans in immune receptor regulation. *Front. Chem.* **2020**, *8*, 55. [[CrossRef](#)]
13. Caval, T.; Heck, A.J.R.; Reiding, K.R. Meta-heterogeneity: Evaluating and describing the diversity in glycosylation between sites on the same glycoprotein. *Mol. Cell. Proteom.* **2021**, *20*, 100010. [[CrossRef](#)] [[PubMed](#)]
14. Fisher, P.; Thomas-Oates, J.; Wood, A.J.; Ungar, D. The N-glycosylation processing potential of the mammalian golgi apparatus. *Front. Cell Dev. Biol.* **2019**, *7*, 157. [[CrossRef](#)] [[PubMed](#)]
15. Edwards, E.; Livanos, M.; Krueger, A.; Dell, A.; Haslam, S.M.; Mark Smales, C.; Bracewell, D.G. Strategies to control therapeutic antibody glycosylation during bioprocessing: Synthesis and separation. *Biotechnol. Bioeng.* **2022**, *119*, 1343–1358. [[CrossRef](#)]
16. Rexer, T.; Laaf, D.; Gottschalk, J.; Frohnmeier, H.; Rapp, E.; Elling, L. Enzymatic synthesis of glycans and glycoconjugates. *Adv. Biochem. Eng. Biotechnol.* **2021**, *175*, 231–280.
17. Nomura, K.; Liu, Y.; Kajihara, Y. Synthesis of homogeneous glycoproteins with diverse N-glycans. *Adv. Carbohydr. Chem. Biochem.* **2022**, *81*, 57–93. [[PubMed](#)]
18. Fairbanks, A.J. Chemoenzymatic synthesis of glycoproteins. *Curr. Opin. Chem. Biol.* **2019**, *53*, 9–15. [[CrossRef](#)]
19. Li, C.; Wang, L.X. Chemoenzymatic methods for the synthesis of glycoproteins. *Chem. Rev.* **2018**, *118*, 8359–8413. [[CrossRef](#)]
20. Mastrangeli, R.; Palinsky, W.; Bierau, H. Glycoengineered antibodies: Towards the next-generation of immunotherapeutics. *Glycobiology* **2018**, *29*, 199–210. [[CrossRef](#)]
21. Wang, T.; Liu, L.; Voglmeir, J. mAbs N-glycosylation: Implications for biotechnology and analytics. *Carbohydr. Res.* **2022**, *514*, 108541. [[CrossRef](#)]
22. Mastrangeli, R.; Audino, M.C.; Palinsky, W.; Broly, H.; Bierau, H. The formidable challenge of controlling high mannose-type N-glycans in therapeutic mAbs. *Trends Biotechnol.* **2020**, *38*, 1154–1168. [[CrossRef](#)]
23. Hsu, Y.P.; Verma, D.; Sun, S.; McGregor, C.; Mangion, I.; Mann, B.F. Successive remodeling of IgG glycans using a solid-phase enzymatic platform. *Commun. Biol.* **2022**, *5*, 328. [[CrossRef](#)] [[PubMed](#)]
24. Warnock, D.; Bai, X.; Autote, K.; Gonzales, J.; Kinealy, K.; Yan, B.; Qian, J.; Stevenson, T.; Zopf, D.; Bayer, R.J. In vitro galactosylation of human IgG at 1 kg scale using recombinant galactosyltransferase. *Biotechnol. Bioeng.* **2005**, *92*, 831–842. [[CrossRef](#)] [[PubMed](#)]
25. Hodoniczky, J.; Zheng, Y.Z.; James, D.C. Control of recombinant monoclonal antibody effector functions by Fc N-glycan remodeling in vitro. *Biotechnol. Prog.* **2005**, *21*, 1644–1652. [[CrossRef](#)] [[PubMed](#)]
26. Zhou, Q.; Stefano, J.E.; Manning, C.; Kyazike, J.; Chen, B.; Gianolio, D.A.; Park, A.; Busch, M.; Bird, J.; Zheng, X.; et al. Site-specific antibody-drug conjugation through glycoengineering. *Bioconjug. Chem.* **2014**, *25*, 510–520. [[CrossRef](#)] [[PubMed](#)]
27. Li, X.; Fang, T.; Boons, G.J. Preparation of well-defined antibody-drug conjugates through glycan remodeling and strain-promoted azide-alkyne cycloadditions. *Angew. Chem. Int. Ed. Engl.* **2014**, *53*, 7179–7182. [[CrossRef](#)]
28. Thomann, M.; Schlothauer, T.; Dashivets, T.; Malik, S.; Avenal, C.; Bulau, P.; Ruger, P.; Reusch, D. In vitro glycoengineering of IgG1 and its effect on Fc receptor binding and ADCC activity. *PLoS ONE* **2015**, *10*, e0134949. [[CrossRef](#)]
29. Jarontomeechai, T.; Kwon, Y.H.; Liu, Y.; Young, O.; Bhawal, R.; Wilson, J.D.; Li, M.; Chapla, D.G.; Moremen, K.W.; Jewett, M.C.; et al. A universal glycoenzyme biosynthesis pipeline that enables efficient cell-free remodeling of glycans. *Nat. Commun.* **2022**, *13*, 6325. [[CrossRef](#)]
30. Moremen, K.W.; Ramiah, A.; Stuart, M.; Steel, J.; Meng, L.; Forouhar, F.; Moniz, H.A.; Gahlay, G.; Gao, Z.; Chapla, D.; et al. Expression system for structural and functional studies of human glycosylation enzymes. *Nat. Chem. Biol.* **2018**, *14*, 156–162. [[CrossRef](#)]
31. Hidari, K.I.; Horie, N.; Murata, T.; Miyamoto, D.; Suzuki, T.; Usui, T.; Suzuki, Y. Purification and characterization of a soluble recombinant human ST6Gal I functionally expressed in *Escherichia coli*. *Glycoconj. J.* **2005**, *22*, 1–11. [[CrossRef](#)]
32. Williams, R.L.; Greene, S.M.; McPherson, A. The crystal structure of ribonuclease B at 2.5-Å resolution. *J. Biol. Chem.* **1987**, *262*, 16020–16031. [[CrossRef](#)] [[PubMed](#)]
33. Rudd, P.M.; Joao, H.C.; Coghill, E.; Fiten, P.; Saunders, M.R.; Opdenakker, G.; Dwek, R.A. Glycoforms modify the dynamic stability and functional activity of an enzyme. *Biochemistry* **1994**, *33*, 17–22. [[CrossRef](#)]
34. Li, Y.; Li, R.; Yu, H.; Sheng, X.; Wang, J.; Fisher, A.J.; Chen, X. *Enterococcus faecalis* alpha1-2-mannosidase (EfMan-I): An efficient catalyst for glycoprotein N-glycan modification. *FEBS Lett.* **2020**, *594*, 439–451. [[CrossRef](#)] [[PubMed](#)]
35. Henrissat, B. A classification of glycosyl hydrolases based on amino acid sequence similarities. *Biochem. J.* **1991**, *280 Pt 2*, 309–316. [[CrossRef](#)] [[PubMed](#)]
36. Henrissat, B.; Bairoch, A. Updating the sequence-based classification of glycosyl hydrolases. *Biochem. J.* **1996**, *316 Pt 2*, 695–696. [[CrossRef](#)]

37. Campbell, J.A.; Davies, G.J.; Bulone, V.; Henrissat, B. A classification of nucleotide-diphospho-sugar glycosyltransferases based on amino acid sequence similarities. *Biochem. J.* **1997**, *326 Pt 3*, 929–939. [[CrossRef](#)]
38. Coutinho, P.M.; Deleury, E.; Davies, G.J.; Henrissat, B. An evolving hierarchical family classification for glycosyltransferases. *J. Mol. Biol.* **2003**, *328*, 307–317. [[CrossRef](#)]
39. Fujiyama, K.; Ido, Y.; Misaki, R.; Moran, D.G.; Yanagihara, I.; Honda, T.; Nishimura, S.; Yoshida, T.; Seki, T. Human N-acetylglucosaminyltransferase I. Expression in *Escherichia coli* as a soluble enzyme, and application as an immobilized enzyme for the chemoenzymatic synthesis of N-linked oligosaccharides. *J. Biosci. Bioeng.* **2001**, *92*, 569–574. [[CrossRef](#)]
40. Zhu, Y.; Suits, M.D.; Thompson, A.J.; Chavan, S.; Dinev, Z.; Dumon, C.; Smith, N.; Moremen, K.W.; Xiang, Y.; Siriwardena, A.; et al. Mechanistic insights into a Ca²⁺-dependent family of alpha-mannosidases in a human gut symbiont. *Nat. Chem. Biol.* **2010**, *6*, 125–132. [[CrossRef](#)]
41. Kadirvelraj, R.; Yang, J.Y.; Sanders, J.H.; Liu, L.; Ramiah, A.; Prabhakar, P.K.; Boons, G.J.; Wood, Z.A.; Moremen, K.W. Human N-acetylglucosaminyltransferase II substrate recognition uses a modular architecture that includes a convergent exosite. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, 4637–4642. [[CrossRef](#)]
42. Boeggeman, E.E.; Balaji, P.V.; Sethi, N.; Masibay, A.S.; Qasba, P.K. Expression of deletion constructs of bovine beta-1,4-galactosyltransferase in *Escherichia coli*: Importance of Cys134 for its activity. *Protein Eng.* **1993**, *6*, 779–785. [[CrossRef](#)] [[PubMed](#)]
43. Gastinel, L.N.; Cambillau, C.; Bourne, Y. Crystal structures of the bovine beta4galactosyltransferase catalytic domain and its complex with uridine diphosphogalactose. *EMBO J.* **1999**, *18*, 3546–3557. [[CrossRef](#)] [[PubMed](#)]
44. Ramakrishnan, B.; Boeggeman, E.; Ramasamy, V.; Qasba, P.K. Structure and catalytic cycle of beta-1,4-galactosyltransferase. *Curr. Opin. Struct. Biol.* **2004**, *14*, 593–600. [[CrossRef](#)]
45. Li, Y.; Xue, M.; Sheng, X.; Yu, H.; Zeng, J.; Thon, V.; Chen, Y.; Muthana, M.M.; Wang, P.G.; Chen, X. Donor substrate promiscuity of bacterial beta1-3-N-acetylglucosaminyltransferases and acceptor substrate flexibility of beta1-4-galactosyltransferases. *Bioorg. Med. Chem.* **2016**, *24*, 1696–1705. [[CrossRef](#)] [[PubMed](#)]
46. Chen, X.; Varki, A. Advances in the biology and chemistry of sialic acids. *ACS Chem. Biol.* **2010**, *5*, 163–176. [[CrossRef](#)]
47. Yang, X.; Yu, H.; Yang, X.; Kooner, A.S.; Yuan, Y.; Luu, B.; Chen, X. One-pot multienzyme (OPME) chemoenzymatic synthesis of brain ganglioside glycans with human ST3GAL II expressed in *E. coli*. *ChemCatChem* **2022**, *14*, e202101498. [[CrossRef](#)]
48. Zhang, Y.; Malinovskii, V.A.; Fiedler, T.J.; Brew, K. Role of a conserved acidic cluster in bovine beta1,4 galactosyltransferase-1 probed by mutagenesis of a bacterially expressed recombinant enzyme. *Glycobiology* **1999**, *9*, 815–822. [[CrossRef](#)]
49. Yu, H.; Chokhawala, H.; Karpel, R.; Yu, H.; Wu, B.; Zhang, J.; Zhang, Y.; Jia, Q.; Chen, X. A multifunctional *Pasteurella multocida* sialyltransferase: A powerful tool for the synthesis of sialoside libraries. *J. Am. Chem. Soc.* **2005**, *127*, 17618–17619. [[CrossRef](#)]
50. Sugiarto, G.; Lau, K.; Qu, J.; Li, Y.; Lim, S.; Mu, S.; Ames, J.B.; Fisher, A.J.; Chen, X. A sialyltransferase mutant with decreased donor hydrolysis and reduced sialidase activities for directly sialylating LewisX. *ACS Chem. Biol.* **2012**, *7*, 1232–1240. [[CrossRef](#)]
51. Thon, V.; Li, Y.; Yu, H.; Lau, K.; Chen, X. PmST3 from *Pasteurella multocida* encoded by Pm1174 gene is a monofunctional alpha2-3-sialyltransferase. *Appl. Microbiol. Biotechnol.* **2012**, *94*, 977–985. [[CrossRef](#)] [[PubMed](#)]
52. Yu, H.; Huang, S.; Chokhawala, H.; Sun, M.; Zheng, H.; Chen, X. Highly efficient chemoenzymatic synthesis of naturally occurring and non-natural alpha-2,6-linked sialosides: A *P. damsela* alpha-2,6-sialyltransferase with extremely flexible donor-substrate specificity. *Angew. Chem. Int. Ed. Engl.* **2006**, *45*, 3938–3944. [[CrossRef](#)]
53. Bai, Y.; Yang, X.; Yu, H.; Chen, X. Substrate and process engineering for biocatalytic synthesis and facile purification of human milk oligosaccharides. *ChemSusChem* **2022**, *15*, e202102539. [[CrossRef](#)] [[PubMed](#)]
54. Ding, L.; Yu, H.; Lau, K.; Li, Y.; Muthana, S.; Wang, J.; Chen, X. Efficient chemoenzymatic synthesis of sialyl Tn-antigens and derivatives. *Chem. Commun.* **2011**, *47*, 8691–8693. [[CrossRef](#)]
55. Ding, L.; Zhao, C.; Qu, J.; Li, Y.; Sugiarto, G.; Yu, H.; Wang, J.; Chen, X. A *Photobacterium sp.* alpha2-6-sialyltransferase (Psp2,6ST) mutant with an increased expression level and improved activities in sialylating Tn antigens. *Carbohydr. Res.* **2015**, *408*, 127–133. [[CrossRef](#)] [[PubMed](#)]
56. Li, Y.; Chen, X. Sialic acid metabolism and sialyltransferases: Natural functions and applications. *Appl. Microbiol. Biotechnol.* **2012**, *94*, 887–905. [[CrossRef](#)]
57. Mbua, N.E.; Li, X.; Flanagan-Steet, H.R.; Meng, L.; Aoki, K.; Moremen, K.W.; Wolfert, M.A.; Steet, R.; Boons, G.J. Selective exo-enzymatic labeling of N-glycans on the surface of living cells by recombinant ST6Gal I. *Angew. Chem. Int. Ed. Engl.* **2013**, *52*, 13012–13015. [[CrossRef](#)] [[PubMed](#)]
58. Chiu, C.P.; Lairson, L.L.; Gilbert, M.; Wakarchuk, W.W.; Withers, S.G.; Strynadka, N.C. Structural analysis of the alpha-2,3-sialyltransferase Cst-I from *Campylobacter jejuni* in apo and substrate-analogue bound forms. *Biochemistry* **2007**, *46*, 7196–7204. [[CrossRef](#)]
59. Geissner, A.; Baumann, L.; Morley, T.J.; Wong, A.K.O.; Sim, L.; Rich, J.R.; So, P.P.L.; Dullaghan, E.M.; Lessard, E.; Iqbal, U.; et al. 7-Fluorosialyl glycosides are hydrolysis resistant but readily assembled by sialyltransferases providing easy access to more metabolically stable glycoproteins. *ACS Cent. Sci.* **2021**, *7*, 345–354. [[CrossRef](#)] [[PubMed](#)]
60. Selman, M.H.; Hemayatkar, M.; Deelder, A.M.; Wuhler, M. Cotton HILIC SPE microtips for microscale purification and enrichment of glycans and glycopeptides. *Anal. Chem.* **2011**, *83*, 2492–2499. [[CrossRef](#)]
61. Yu, H.; Yu, H.; Karpel, R.; Chen, X. Chemoenzymatic synthesis of CMP-sialic acid derivatives by a one-pot two-enzyme system: Comparison of substrate flexibility of three microbial CMP-sialic acid synthetases. *Bioorg. Med. Chem.* **2004**, *12*, 6427–6435. [[CrossRef](#)]

62. Yu, H.; Chen, X. One-pot multienzyme (OPME) systems for chemoenzymatic synthesis of carbohydrates. *Org. Biomol. Chem.* **2016**, *14*, 2809–2818. [[CrossRef](#)]
63. Chen, Y.; Thon, V.; Li, Y.; Yu, H.; Ding, L.; Lau, K.; Qu, J.; Hie, L.; Chen, X. One-pot three-enzyme synthesis of UDP-GlcNAc derivatives. *Chem. Commun.* **2011**, *47*, 10815–10817. [[CrossRef](#)] [[PubMed](#)]
64. Muthana, M.M.; Qu, J.; Li, Y.; Zhang, L.; Yu, H.; Ding, L.; Malekan, H.; Chen, X. Efficient one-pot multienzyme synthesis of UDP-sugars using a promiscuous UDP-sugar pyrophosphorylase from *Bifidobacterium longum* (BLUSP). *Chem. Commun.* **2012**, *48*, 2728–2730. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.