UC Irvine

UC Irvine Previously Published Works

Title

Soil warming and organic carbon content.

Permalink

<https://escholarship.org/uc/item/38m7b07m>

Journal

Nature, 408(6814)

ISSN

0028-0836

Authors

Davidson, EA
Trumbore, SE
Amundson, R

Publication Date

2000-12-01

DOI

10.1038/35048672

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Biogeochemistry

Soil warming and organic carbon content

Soils store two or three times more carbon than exists in the atmosphere as CO₂, and it is thought that the temperature sensitivity of decomposing organic matter in soil partly determines how much carbon will be transferred to the atmosphere as a result of global warming¹. Giardina and Ryan² have questioned whether turnover times of soil carbon depend on temperature, however, on the basis of experiments involving isotope analysis and laboratory incubation of soils. We believe that their conclusions are undermined by methodological factors and also by their turnover times being estimated on the assumption that soil carbon exists as a single homogeneous pool, which can mask the dynamics of a smaller, temperature-dependent soil-carbon fraction. The real issue about release of carbon from soils to the atmosphere, however, is how temperature, soil water content and other factors interact to influence decomposition of soil organic matter. And, contrary to one interpretation³ of Giardina and Ryan's results, we believe that positive feedback to global warming is still a concern.

Climate has long been believed to affect the accumulation of detrital carbon and nitrogen⁴ — cool conditions and poor aeration in wet soils can impede decomposition⁴, and soil carbon stocks are highest in cool and moist biomes and lowest in hot and dry biomes⁵; climate also affects plant productivity and the input of carbon to the soil. To determine decomposition (carbon output), a simplistic one-pool model (soil carbon stock/(field-measured soil respiration minus estimated root respiration)) can be used to calculate average turnover times (soil carbon stock/decomposition rate) of total soil carbon⁶. Although this one-pool model is not without problems (see later), a trend of increasing average turnover times with declining mean annual temperature and increasing latitude is apparent in these field data.

Temperature should not be viewed in isolation. For example, temperature affects evapotranspiration, which affects soil water content, which influences decomposition of soil organic matter (SOM)^{1,6}. Climate is only one of a hierarchy of factors controlling decomposition, including clay mineralogy, chemical properties of leaf and root litter, activity of macrofauna and micro-organisms, and disturbance⁷. Giardina and Ryan's proposal² that temperature has no effect on the turnover of soil carbon is based on their search for only a single factor in this complex matrix and also on, in our view, inappropriate tools and sampling

design (involving a type II statistical error).

Giardina and Ryan used changes in ¹³C content of SOM following conversion of C₃ vegetation to C₄ vegetation, which necessitates different degrees of site disturbance. Ploughing causes rapid, large changes in decomposition, exposing SOM previously protected inside soil aggregates^{8,9}. Conversion of forest to cattle pasture, however, causes only moderate soil disturbance and small changes in SOM decomposition¹⁰. The degree of disturbance may thus have a larger effect on calculated turnover times than either temperature or the time elapsed since disturbance.

Giardina and Ryan base their estimates of soil-carbon turnover times on the assumption that SOM can be represented as a single homogeneous pool, an approach that has yielded mixed results^{2,6} because it ignores the widely varying age (from months to millennia)¹¹ of carbon in SOM. The proportion of SOM cycling over millennia ranges from zero to over 50%, depending on mineral type, clay content and soil depth. This variation has a large effect on the calculated turnover times when soil-carbon fractions that cycle at different rates are averaged together¹¹.

Figure 1 illustrates this point, showing results from radiocarbon analysis of two soils sampled along an elevation transect in the Sierra Nevada mountains of California¹², where turnover times of fractionated carbon pools in each soil range from 25 to over 1,000 years. Although the total soil carbon is comparable in both soils, the soil at lower elevation (Musick soil), which experiences higher annual temperatures and slightly less precipitation (not higher²), contains less labile carbon with a faster turnover time than the higher-elevation (Shaver) soil. Decomposition of the SOM fractions during a one-year incubation would sum to similar amounts of CO₂ for the two soils, hence the single-pool model of Giardina and Ryan would ascribe similar turnover times to both soils, obscuring the faster cycling of the smaller labile fraction of carbon in the warmer, lower-elevation soil.

Figure 1 Radiocarbon estimates of turnover times of carbon fractions of two soils on an elevational gradient with similar parent material, vegetation and disturbance history¹². Fractions were separated by density and hydrolysis for each soil depth¹². The CO₂ that would be evolved during one-year incubations (98 and 92 g C m⁻² yr⁻¹ for Musick and Shaver soils, respectively) was calculated from carbon stocks and turnover times. Dividing respired CO₂ by total soil carbon, as Giardina and Ryan² do in their one-pool model, yields nearly identical turnover times estimates for the two soils (53 and 54 yr for Musick and Shaver soils, respectively). However, the cooler Shaver soil contains twice as much carbon with turnover times of about 50 yr and the warmer Musick soil has a small but important pool that cycles more rapidly.

Laboratory incubations, as used by Giardina and Ryan², involve procedures such as sieving and mixing, so are not comparable to decomposition *in situ*. In some soils, the small amount of easily decomposable humus can be spent before the end of a one-year incubation, so differences in calculated turnover times among samples may reflect different substrate limitations rather than temperature sensitivity¹³; they may also reflect differences in sampling depths, which affect the proportions of labile and non-labile carbon.

Rather than averaging turnover times of total extant SOM, it is the carbon that has already left the soil that may reveal the latitudinal importance of temperature on SOM dynamics — for example, slash-and-burn agriculture in the tropics must be abandoned after a few years as the small amount of labile soil humus releases nutrients for only a short time. By contrast, agricultural productivity in a northern temperate climate was maintained for decades without extra nutrients after ploughing soils with a similar sand content, but with more humus that decomposed gradually in the cooler climate⁹. Variations in mineralogy and aggregate stability also contribute to these differences.

The temperature sensitivity of soil carbon subject to climate change has been tested by fractionating SOM into pools that can be compared along climate gradients^{9,11,12}. *In situ* experimental soil warming also increases soil CO₂ emissions unless it provokes drought¹. Increased emission was limited to a few years after warming treatments began, indicating that only a fraction of the soil carbon is temperature-sensitive and supporting our assertion that carbon dynamics should be studied using multi-pool models.

In a field study along climate gradients¹⁴, plant biomass increased in all the EUROFLUX sites, even those with a carbon budget close to neutral and one losing carbon annually — the site losing carbon was drained, exposing anaerobic soil layers to decomposition. Soil carbon loss also happens where warming helps to dry wetlands

or thaw frozen soil¹⁵. The EUROFLUX sites with almost neutral carbon balance are at high latitudes where significant warming has occurred, which may provoke loss of soil carbon at a rate comparable to plant biomass gains¹⁵.

Unlike Grace and Rayment³, we consider that the key to climatic sensitivity of soil carbon is not total ecosystem respiration but decomposition rates (carbon loss per unit of fractionated carbon pool). The high respiration rates at the northern EUROFLUX sites¹⁴ are probably due to detritus accumulation under cool moist conditions, and not to temperature insensitivity of decomposition; even slow turnover of large stocks produces a lot of CO₂.

We agree with Giardina and Ryan that temperature effects on soil carbon dynamics may be overestimated in current biogeochemical models. However, feedback to global warming does not concern just temperature, but also includes its effect on soil water content and drainage, for example, and applies to all detrital carbon, whether on top of mineral soil or buried in peatlands and permafrost.

On the basis of the new results^{2,14}, Grace and Rayment³ suggest that the doomsday view of runaway global warming now seems unlikely. We believe that, on the contrary, the evidence remains in favour of a strong climatic control over storage of detrital carbon in terrestrial ecosystems.

Eric A. Davidson*, **Susan E. Trumbore†**,
‡Ronald Amundson

*Woods Hole Research Center, Woods Hole,
Massachusetts 02543, USA

e-mail: edavidson@whrc.org

†Department of Earth System Science, University of
California, Irvine, California 92697-3100, USA

‡Division of Ecosystem Sciences, University of
California, Berkeley, California 94720, USA

1. Kirchbaum, M. U. F. *Biogeochemistry* **48**, 21–51 (2000).
2. Giardina, C. P. & Ryan, M. G. *Nature* **404**, 858–861 (2000).
3. Grace, J. & Rayment, M. *Nature* **404**, 819–820 (2000).
4. Jenny, H. *Factors of Soil Formation* (McGraw-Hill, New York, 1941).
5. Post, W. M., Emanuel, R., Zinke, P. J. & Stangenberger, A. G. *Nature* **298**, 156–159 (1982).
6. Raich, J. W. & Schlesinger, W. H. *Tellus* **44B**, 81–99 (1992).
7. Lavelle, P. et al. *Biotropica* **25**, 130–150 (1993).
8. Davidson, E. A. & Ackerman, I. L. *Biogeochemistry* **20**, 161–193 (1993).
9. Tiessen, H., Cuevas, E. & Chacon, P. *Nature* **371**, 783–785 (1994).
10. Neill, C. & Davidson, E. A. in *Carbon Pools and Dynamics in Tropical Ecosystems* (eds Lal, R., Kimble, J. M. & Stewart, B. A.) 197–211 (CRC, Boca Raton, 2000).
11. Trumbore, S. E. *Ecol. Appl.* **10**, 399–411 (2000).
12. Trumbore, S., Chadwick, O. A. & Amundson, R. A. *Science* **272**, 393–396 (1996).
13. Holland, E. A., Townsend, A. R. & Vitousek, P. M. *Glob. Change Biol.* **1**, 115–123 (1995).
14. Valentini, R. et al. *Nature* **404**, 861–865 (2000).
15. Goulden, M. L. et al. *Science* **279**, 214–217 (1998).

Giardina and Ryan reply — Davidson et al. question the validity of our conclusions on the grounds that differences in disturbance, high variability among sites, and the use of a single-pool model to estimate turnover

time (TT) obscured the effect of temperature on the decomposition of soil carbon.

In our comparison of incubated soils, disturbance associated with sieving and mixing probably increased decomposition rates. However, soil processing was similar across sites, and the resulting increase in carbon availability should have accentuated an effect of temperature on TT. In our *in situ* comparison, most tropical sites were pastures and all of the temperate sites were cropped. But these differences did not obscure an effect of temperature on TT. First, a subset of tropical sites had been continuously cultivated with sugar cane, which involves biannual mechanized ploughing, burning and fertilization. For sites that had been similarly disturbed after conversion from forest — that is, under intensive cultivation — TT tended to increase with mean annual temperature (MAT; $R^2 = 0.14$, $P = 0.07$, $n = 25$). Second, tropical pastures are more severely disturbed than Davidson et al. suggest — bulldozers may be used to level a site and remove stumps, and mechanized disking is often used to control weeds^{1,2}. Further, the effects of pasture conversion on carbon decomposition rates remain poorly understood².

There are two ways to test whether we missed significance because of high variability. First, how important is site-to-site variability? And second, given our sample size, how much variation must temperature explain in order to be significant? For the incubation comparison, variation in TT was low (coefficient of variation, 37%), and TT actually increased with temperature ($TT = 0.09 \times MAT + 6.2$). For the *in situ* comparison, time since conversion explained 34% of the variance in per cent carbon mass loss, a measure that assumes no model or age distribution of carbon in soil. With a sample size of 44 sites for the *in situ* comparison, MAT need only explain 8% of the variance to be significant at the 0.05 level. Given that MAT explained less than 1% of this variance, more than 1,000 sites would be needed to establish that there is a global relationship between temperature and soil carbon turnover.

We examined whether a single-pool model could obscure a temperature effect, and concluded that this is unlikely because our methods were robust^{1,3}, the tests were independent and well replicated, and neither comparison showed the negative trends between TT and temperature that would be expected from modelled patterns of carbon mass loss across latitude⁴. We agree with Davidson et al. that only a small

fraction of total soil carbon may be sensitive to temperature — this was our point.

Although the ¹⁴C approach preferred by Davidson et al. is useful for examining carbon turnover in soil, it has some methodological problems. For example, the ¹⁴C decay of interest begins at photosynthesis, not after incorporation into soil, as is now assumed^{1,5}. In forests, this lag could affect estimates of soil-carbon TT that are based on ¹⁴C because carbon can reside in living biomass and the litter layer for decades before being released to mineral soil. The ¹⁴C approach is also sensitive to contamination³ and has yielded much faster estimates of soil-carbon turnover than more direct methods¹.

Modelling terrestrial carbon storage depends on an accurate understanding of how temperature affects carbon TT in soil, but this effect cannot be inferred from large-scale relationships between soil-carbon content and MAT. Along gradients similar to those used by Jenny⁶ and cited by Davidson et al., plant primary production declines with increasing MAT, and it is this decline, not an increase in soil-carbon decomposition rate, that explains decreasing soil-carbon content with increasing temperature⁷. Similar findings have been reported⁸ for MAT gradients in northern Europe, and the gradient study cited by Davidson et al. actually reports⁹ longer carbon TT for tropical forest than for temperate forest. Despite the evidence against an exponential effect of temperature on soil carbon turnover, future *in situ* warming studies are needed to settle this issue.

Christian P. Giardina*, **Michael G. Ryan†**

*Department of Natural Resources and
Environmental Management, University of Hawaii
at Manoa, 1910 East-West Road, Honolulu,
Hawaii 96822, USA

e-mail: giardina@hawaii.edu

†United States Department of Agriculture Forest
Service, Rocky Mountain Research Station, 240 West
Prospect Street, Fort Collins, Colorado 80526, USA
and Graduate Degree Program in Ecology, Colorado
State University, Fort Collins, Colorado 80523, USA

1. Trumbore, S. E. et al. *Glob. Biogeochem. Cycles* **9**, 515–528 (1995).
2. Neill, C. & Davidson, E. in *Carbon Pools and Dynamics in Tropical Ecosystems* (eds Lal, R., Kimble, J. & Stewart, B.) 197–211 (CRC, Boca Raton, 2000).
3. Townsend, A. R., Vitousek, P. M. & Trumbore, S. E. *Ecology* **76**, 721–733 (1995).
4. Schimel, D. S. et al. *Glob. Biogeochem. Cycles* **8**, 279–293 (1994).
5. Trumbore, S. E., Chadwick, O. A. & Amundson, R. A. *Science* **272**, 393–396 (1996).
6. Jenny, H. *Factors of Soil Formation* (McGraw-Hill, New York, 1941).
7. Epstein, H. E., Burke, I. C. & Lauenroth, W. K. *Ecology* (in the press).
8. Liski, J. et al. *Ambio* **28**, 171–174 (1999).
9. Raich, J. W. & Schlesinger, W. H. *Tellus* **44B**, 81–99 (1992).

brief communications is intended to provide a forum for short, topical reports of general scientific interest and for technical discussion of recently published material of particular interest to non-specialist readers. Priority will be given to contributions that have fewer than 500 words, 10 references and only one figure. Detailed guidelines are available on Nature's website (www.nature.com) or on request from nature@nature.com