

UC Davis

UC Davis Previously Published Works

Title

Eicosanoids The Overlooked Storm in Coronavirus Disease 2019 (COVID-19)?

Permalink

<https://escholarship.org/uc/item/38s0c5t5>

Journal

American Journal Of Pathology, 190(9)

ISSN

0002-9440

Authors

Hammock, Bruce D

Wang, Weicang

Gilligan, Molly M

et al.

Publication Date

2020-09-01

DOI

10.1016/j.ajpath.2020.06.010

Peer reviewed



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



MINI-REVIEW

Q3 Q1 Eicosanoids

The Overlooked Storm in Coronavirus Disease 2019 (COVID-19)?

Q20 Bruce D. Hammock,* Weicang Wang,* Molly M. Gilligan,^{†‡} and Dipak Panigrahy^{†‡}

Q4 Q5 From the Department of Entomology and Nematology and UCD Comprehensive Cancer Center,* University of California, Davis, California; and the Center for Vascular Biology Research[†] and the Department of Pathology,[‡] Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts

Accepted for publication
June 30, 2020.

Address correspondence to
Bruce D. Hammock, Ph.D.,
Department of Entomology,
University of California, Davis,
One Shields Ave., Davis,
CA 95616; or Dipak Panigrahy,
M.D., Department of Pathol-
ogy, Beth Israel Deaconess
Medical Center, 99 Brookline
Ave., Room 220, Boston,
MA 02215. E-mail:
bdhammock@ucdavis.edu or
dpanigra@bidmc.harvard.edu.

Severe coronavirus disease 2019 (COVID-19) symptoms, including systemic inflammatory response and multisystem organ failure, are now affecting thousands of infected patients and causing widespread mortality. Coronavirus infection causes tissue damage, which triggers the endoplasmic reticulum stress response and subsequent eicosanoid and cytokine storms. Although proinflammatory eicosanoids, including prostaglandins, thromboxanes, and leukotrienes, are critical mediators of physiological processes, such as inflammation, fever, allergy, and pain, their role in COVID-19 is poorly characterized. Arachidonic acid-derived epoxyeicosatrienoic acids could alleviate the systemic hyperinflammatory response in COVID-19 infection by modulating endoplasmic reticulum stress and stimulating the resolution of inflammation. Soluble epoxide hydrolase (sEH) inhibitors, which increase endogenous epoxyeicosatrienoic acid levels, exhibit potent anti-inflammatory activity and inhibit various pathologic processes in preclinical disease models, including pulmonary fibrosis, thrombosis, and acute respiratory distress syndrome. Therefore, targeting eicosanoids and sEH could be a novel therapeutic approach in combating COVID-19. In this review, we discuss the predominant role of eicosanoids in regulating the inflammatory cascade and propose the potential application of sEH inhibitors in alleviating COVID-19 symptoms. We also discuss the host-protective action of omega-3 fatty acid-derived epoxyeicosanoids and specialized proresolving mediators in regulating anti-inflammation and antiviral response. Future studies determining the eicosanoid profile in COVID-19 patient or preclinical model are pivotal in providing the novel insight of coronavirus-host interaction and inflammation modulation. (*Am J Pathol* 2020, ■: 1–7; <https://doi.org/10.1016/j.ajpath.2020.06.010>)

Q9 Early in the coronavirus disease 2019 (COVID-19) pandemic, it was observed that morbidity and mortality were associated with a drastic systemic inflammatory response caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection.^{1–3} This was initially observed in the lung, with many patients experiencing respiratory distress, pneumonia, fever, and multi-organ failure. Severe pathologies are now widely appreciated in many organs throughout the human body, including the lung, heart, spleen, lymph nodes, brain, liver, kidneys, eyes, and vasculature, because of widespread inflammatory responses.^{1,4,5} The proinflammatory cytokine storm is a critical driving force in severe COVID-19 infection and may lead to multisystem

organ failure.^{2,3,5,6} One of the first descriptions of the cytokine storm was detailed on graft-versus-host disease in 1993.⁷ Since then, cytokine storms and associated hyperinflammatory mechanisms have been characterized in studies of viral infection [influenza, SARS-associated coronavirus

Supported by NIH grants, including National Institute of Environmental Health Science Superfund Research Program P42 ES004699 (B.D.H. and D.P.); National Institute of Environmental Health Science River Award R35ES030443 (B.D.H.); the Credit Unions Kids at Heart Team; the CJ Buckley Pediatric Brain Tumor Fund; and the Joe Andruzzi Foundation (D.P.).

Disclosures: None declared.

(SARS-CoV), and SARS-CoV-2], macrophage activation syndrome, immunotherapy, chemotherapy, rheumatic diseases, traumatic injury, and acute respiratory disease and may thus be utilized as potential diagnostic markers of inflammatory disease progression.^{8–12}

Both SARS-CoV and the influenza virus induce apoptosis and necrosis of various cell types, including T cells, endothelial cells, and epithelial cells, through proinflammatory cytokine- and chemokine-mediated vascular leakage and suboptimal T-cell responses.^{6,13–16} Moreover, viral clearance may be impaired by a macrophage-derived proinflammatory cytokine storm and thus targeting viral replication may not be sufficient to promote host survival.¹⁶ Locally and systemically elevated proinflammatory cytokines, including IL-1 β , interferon- γ , IP-10, IL-1, IL-6, and monocyte chemoattractant protein 1, have been detected in select critically ill patients with COVID-19 and found to correlate with disease severity.^{2,6} Interestingly, this hyperinflammatory innate host response to SARS-CoV may additionally correlate with death in large animal models (eg, primates) more directly than virus titers.¹⁷ Thus, a therapeutic approach to stimulating the resolution of the host inflammatory response, including the cytokine storm, may be as critical as preventing viral replication for patient survival.³

Cell death, including apoptosis, is induced by host defense mechanisms in response to infections.¹⁸ However, cell death (debris) can trigger an eicosanoid and cytokine storm in inflammatory diseases.¹⁰ Virus-induced inflammatory proteins and cell debris trigger a cellular endoplasmic reticulum (ER) stress response, which mediates the virus-host interaction in infected cells.¹⁹ Normally, an ER stress response down-regulates global protein synthesis to diminish further viral replication in infected cells. However, prolonged responses initiate proinflammatory and later proapoptotic programs of ER stress in host cells, which cause elevated cytokine production during infection. The spike protein, which mediates SARS-CoV binding to angiotensin-converting enzyme 2 for cellular entry, has been shown to activate ER stress transducer X-box binding protein 1 and up-regulate ER-resident protein Herpud1 and chemokine Cxcl2 in infected murine fibroblast L cells.²⁰ In addition, overexpression of SARS-CoV ORF8b protein elevates ER stress effector CHOP, which leads to cell apoptosis in HeLa cells.²¹ Coronavirus-induced tissue damage and accumulation of cellular debris (ie, cell fragments and proteins) has further been implicated in stimulating cell apoptosis and boosting inflammation by regulating other cellular pathways (JNK/p38, Bcl2/Bax, inflammasome, and NF- κ B) with specific ER stress sensor arms.¹⁹ Manipulations of the coronavirus-induced ER stress response upstream of the devastating cytokine storm could be a powerful approach in combating COVID-19 pathogenesis in patients with severe pneumonia or multi-organ failure (Figure 1).

Arachidonic acid-derived lipid autacoids, including prostaglandins (PGs), thromboxanes, and leukotrienes, generated by cyclooxygenases and lipoxygenases are

collectively termed eicosanoids and are critical mediators of inflammation, resolution, and tissue homeostasis.²² Eicosanoids play pivotal roles in a broad range of physiological processes, such as inflammation, fever, allergy, and pain.^{22,23} On activation by cell debris or infection, the inositol-requiring enzyme 1 α -X-box binding protein 1 branch of the ER stress pathway up-regulates expression of microsomal prostaglandin E synthase-1 and prostaglandin-endoperoxide synthase 2 [cyclooxygenase 2 (COX-2)], and accelerates the biosynthesis of prostaglandins (PGE₂, PGD₂, and PGF_{2 α}) and thromboxane B₂ from arachidonic acid.^{24,25} This ensuing eicosanoid storm may be associated with production of a cytokine storm.^{10,26} Indeed, activated ER stress signaling alone only slightly increases the level of IL-6, whereas the presence of PGE₂ and/or other cytokines (interferon- γ) with an activated ER stress response greatly enhances IL-6 production in glial cells.²⁷ More important, this eicosanoid surge has not been adequately evaluated in the setting of COVID-19 infection.

Although elevated cytokine levels in COVID-19 have been identified as a major factor contributing to morbidity and mortality, the role of eicosanoids in COVID-19 as key mediators of both inflammation and its active resolution remains poorly characterized.^{3,28–30} More important, not all eicosanoids are proinflammatory as arachidonic acid and related fatty acids are also metabolized into anti-inflammatory and proresolution docosanoids in certain temporal situations.^{22,31–33} Infectious processes often activate inflammasome formation and the subsequent formation of an eicosanoid storm consisting of both proinflammatory and anti-inflammatory mediators, thereby disrupting the temporal progression of inflammation and its resolution.³⁴ Namely, phospholipase A₂ regulates eicosanoid class switching during inflammasome and caspase activation, triggering arachidonic acid to generate proresolution mediators, such as lipoxins.^{34,35} As increased proinflammatory cytokines may be a driving force in severe COVID-19, the balance of proinflammatory/anti-inflammatory eicosanoids and proresolution lipid mediators during the initiation and resolution of infection can regulate the cytokine storm.³⁴ Thus, we hypothesize that SARS-CoV-2 may trigger a temporal production of an eicosanoid storm, including an imbalance of both proinflammatory and proresolution mediators.^{3,22}

Although eicosanoids, such as prostaglandins and leukotrienes, are best known as products of arachidonic acid metabolism by cyclooxygenases and lipoxygenases, arachidonic acid is also a substrate for another enzymatic pathway, the cytochrome P450 system. This eicosanoid pathway consists of two main branches: ω -hydroxylases, which convert arachidonic acid to hydroxyeicosatetraenoic acids; and epoxygenases, which convert it to four regiosomeric epoxyeicosatrienoic acids (EETs; 5,6-EET, 8,9-EET, 11,12-EET, and 14,15-EET).³⁶ EETs regulate inflammation and vascular tone and are produced predominantly in the endothelium.³⁶ More important, the epoxides

187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248




Figure 1 Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection leads to severe tissue damage, which releases cell debris. Both primary infection and the accumulation of cell debris initiate the endoplasmic reticulum (ER) stress response and up-regulate inflammatory enzymes, including microsomal prostaglandin E synthase-1 (mPGES-1) and prostaglandin-endoperoxide synthase 2 [cyclooxygenase 2 (COX-2)], which subsequently produce eicosanoids, including prostaglandins (PGs), leukotrienes (LTs), and thromboxanes (TXs). These proinflammatory lipid autacoids induce cytokine storms that mediate widespread inflammatory responses and organ damage in severe coronavirus disease 2019 (COVID-19) patients. By contrast, epoxyeicosatrienoic acids (EETs), which are stabilized by inhibition of their metabolizing enzyme, soluble epoxide hydrolase (sEH), are anti-inflammatory and proresolving mediators that promote the termination (resolution) of inflammation by suppressing the ER stress response, inflammatory enzyme induction, and proinflammatory cytokine production. EETs also shift arachidonic acid metabolism to favor the production of specialized proresolving mediators (SPMs), which initiate downstream anti-inflammatory and proresolving programs. EETs and sEH inhibitors may counterregulate the unabated systemic inflammatory response and organ failure associated with COVID-19 infection. DHET, dihydroxyeicosatrienoic acid; TNF- α , tumor necrosis factor- α .

print & web 4C/FPO

of EETs are rapidly converted into dihydroxyeicosatrienoic acids by the soluble epoxide hydrolase (sEH) enzyme (Figure 1). Soluble epoxide hydrolase inhibitors, which raise endogenous EET levels, exhibit potent anti-inflammatory activity, including inhibiting proinflammatory cytokines in various pathologic diseases, including inflammatory bowel disease, atherosclerosis, pancreatitis, diabetes, hypertension, stroke, cerebral ischemia, dyslipidemia, pain, immunologic disorders, ocular diseases, neurologic diseases, renal disease (eg, acute kidney injury), organ damage, vascular remodeling, ischemia-reperfusion, lung disease (chronic obstructive pulmonary disease), fibrosis (eg, pulmonary and cardiac fibrosis), graft stenosis, and other medical conditions.^{37–39} Although EETs and sEH signaling play a key role in hyperinflammatory diseases, their role in COVID-19 should be explored.

EETs, which are generated from arachidonic acid by cytochrome P450 enzymes, promote the active termination (resolution) of inflammation through mediating a broad array of anti-inflammatory and proresolving mechanisms, including mitigation of the cytokine storm.^{3,40,41} In an endotoxemia mouse model, increased EET biosynthesis suppresses the endotoxin-induced surge of proinflammatory cytokines (IL-6 and IL-1 β), chemokines (monocyte chemoattractant protein 1 and ENA-78), adhesion molecules (E-selectin), and NF- κ B activation in lung.⁴² Similarly, treatment with 14,15-EET inhibits activation of ER stress effectors (p-eIF2 α , CHOP, and GRP78) by cigarette smoke and suppresses injury-induced oxidative stress and cell apoptosis in human bronchial epithelial cells.⁴³ Reduction of the ER stress response by EETs and promoting the activity of anti-inflammatory, proresolving mediators may

represent a promising new therapeutic avenue in COVID-19 treatment.

Although inflammation can induce sEH expression,^{36,37} COVID-19 infection may further stimulate sEH levels throughout the body in various tissues. Stabilizing the anti-inflammatory, proresolving EETs by using sEH inhibitors (sEHIs) has been shown to have a valuable therapeutic application in various preclinical models and human trials, including sepsis, cardiovascular disease, neuroinflammatory disease, and cancer.^{44–46} More important, administration of sEHIs suppresses pulmonary cytokine expression and neutrophil infiltration, thereby alleviating pulmonary inflammation and edema and decreasing mortality in murine models of endotoxin-induced acute respiratory distress syndrome.⁴⁷ Similar anti-inflammatory activity has been observed in other models of chronic obstructive pulmonary disease, asthma, and pulmonary fibrosis.⁴⁸ In addition, sEHIs dramatically reduce NF- κ B induction of inflammatory enzymes (ie, COX-2), as well as the downstream production of proinflammatory mediators, such as PGE₂.²⁴

These omega-3 fatty acid-derived epoxyeicosanoids exhibit anti-inflammatory activity in various inflammatory diseases, including in the lung, heart, ocular angiogenesis, and pain.⁴⁹ Omega-3 supplementation may also synergize with sEH inhibition to suppress inflammation.⁵⁰ Omega-3 epoxides, stabilized via inhibition of sEH, are important regulators of inflammation and autophagy in metabolic diseases.⁵¹ Dietary eicosapentaenoic acid/docosahexaenoic acid supplementation causes a profound shift of the endogenous cytochrome P450–eicosanoid profile from arachidonic acid– to eicosapentaenoic acid– and docosahexaenoic acid–derived metabolites, increasing, in particular, the

plasma and tissue levels of 17,18-epoxyeicosatetraenoic acid and 19,20-epoxydocosapentaenoic acid. COVID-19 is characterized by increased angiogenesis demonstrated in autopsy samples of lungs from patients who died from COVID-19 compared with influenza patients.⁵² Inhibition of sEH prevents angiogenic diseases, such as diabetic retinopathy.⁵³ Cytochrome P450–derived lipid metabolites epoxydocosapentaenoic acids and epoxyeicosatetraenoic acids also dampen choroidal angiogenesis.⁵⁴

Moreover, EETs shift arachidonic acid metabolism to stimulate the production of specialized proresolving mediators, such as lipoxins, which stimulate clearance of inflammatory cellular debris and counter proinflammatory cytokine production without being immunosuppressive.^{24,48} Pharmacological enhancement of resolution via resolvins at nanogram doses per day or omega-3 fatty acid supplementation restores endogenous specialized proresolving mediators.^{55–57} Stimulation of resolution of inflammation by immunoresolved agonists, including resolvins, is host protective during infection and sepsis.^{58–61} The anti-inflammatory agent dexamethasone may have activity in COVID-19 patients.⁶² Dexamethasone stimulates specialized proresolving mediators to stimulate resolution of airway inflammation.⁶³ The ER stress response can be reduced by dexamethasone by promoting protein folding and degradation of misfolded proteins from the ER.⁶⁴

Targeting eicosanoid metabolism could be a promising new approach in COVID-19 infection given the critical role that eicosanoids play in both the initiation and resolution of inflammation.²² Although current therapeutic strategies are aimed at inhibiting individual inflammatory cytokines (ie, IL-1 and/or IL-6) or viral cell entry and intracellular processing, these strategies neglect the critical upstream role that ER stress and eicosanoids play between generation of cell fragments and proteins released during viral-induced death (debris) and the downstream cytokine production in coronavirus infection (Figure 1). Nonsteroidal anti-inflammatory drugs (NSAIDs), the pan-COX inhibitors that block prostaglandin biosynthesis, are a routine and effective choice for the relief of influenza-caused fever and pain.^{23,37} There may be minimal to no benefit for severe SARS-CoV-2 infections with NSAIDs, which also increase the risk of gastrointestinal and cardiovascular complications.⁶⁵ Initially during infection, the classic mediators of inflammation, including prostaglandins and leukotrienes, are produced, leading to the initiation of inflammation.⁶⁶ Subsequently, prostaglandin E₂ and prostaglandin D₂ induce a lipid mediator class switching of eicosanoid production by neutrophils from leukotriene B₄ and 5-lipoxygenase pathways to lipoxins, resolvins, and protectins to promote resolution.^{33,66–69} Thus, NSAIDs should be used with caution as they may block subsequent activity of prostaglandins, which are essential for the resolution of inflammation.²² NSAIDs are currently recommended to be on an individual basis because of the insufficient evidence in clinical trials and their well-known gastrointestinal and cardiovascular toxicities.⁷⁰

Interestingly, sEHIs synergize with COX inhibitors in reducing inflammation and blocking the gastrointestinal erosion and cardiovascular events associated with these drugs.^{40,71,72} Shifting arachidonic acid metabolism via sEH inhibition also alters the ER stress response toward a more homeostatic role in cell maintenance and promotes the production of anti-inflammatory, proresolving lipids, thus representing an attractive new approach to controlling inflammation in COVID-19. Indeed, sEHIs have been demonstrated to suppress the inflammatory response and inflammation-driven diseases in lung, heart, liver, kidney, and vasculature in numerous preclinical models.^{37,40} Recently, sEHIs have proved nontoxic at high doses in clinical development (phase 2A trials) and synergistic with other well-established anti-inflammatory medications, such as NSAIDs, to promote anti-inflammatory programs and reduce the gastrointestinal adverse effects.⁷¹ Interestingly, risk factors for COVID-19 are similar to idiopathic pulmonary fibrosis, and previous coronavirus outbreaks have been characterized by fibrosis.⁷³ Pharmacologic or genetic inhibition of soluble epoxide hydrolase prevents inflammation and liver fibrosis in various tissues, including lung, heart, liver, and kidney.^{74–80} Thus, the antifibrotic activity of sEHIs may additionally be useful in speeding recovery after COVID-19 infection.⁴⁰ Temporal data on eicosanoid levels in patients are critical to further establish the role of a putative eicosanoid storm in COVID-19, and additional preclinical and clinical studies are needed to test the safety and efficacy of sEHIs in COVID-19 patients. Herein, we propose sEHIs alone or in combination with other agents, such as COX inhibitors or omega-3 lipids, are envisioned to be of benefit in blocking or moderating the inflammatory cascade when the patient's condition first appears to be deteriorating in COVID-19.

References

- Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, Liu S, Zhao P, Liu H, Zhu L, Tai Y, Bai C, Gao T, Song J, Xia P, Dong J, Zhao J, Wang FS: Pathological findings of COVID-19 associated with acute respiratory distress syndrome. Lancet Respir Med 2020; 8:420–422
- Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ; Hlh Across Speciality Collaboration UK: COVID-19: consider cytokine storm syndromes and immunosuppression. Lancet 2020; 395:1033–1034
- Panigrahy D, Gilligan MM, Huang S, Gartung A, Cortes-Puch I, Sime PJ, Phipps RP, Serhan CN, Hammock BD: Inflammation resolution: a dual-pronged approach to averting cytokine storms in COVID-19? Cancer Metastasis Rev 2020; 39:337–340
- Azkur AK, Akdis M, Azkur D, Sokolowska M, van de Veen W, Bruggen MC, O'Mahony L, Gao Y, Nadeau K, Akdis CA: Immune response to SARS-CoV-2 and mechanisms of immunopathological changes in COVID-19. Allergy 2020 [Epub ahead of print]. doi: 10.1111/all.14364
- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B: Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet 2020; 395:497–506
- Ye Q, Wang B, Mao J: The pathogenesis and treatment of the “cytokine storm” in COVID-19. J Infect 2020; 80:607–613

433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496



- 497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
7. Ferrara JL, Abhyankar S, Gilliland DG: Cytokine storm of graft-versus-host disease: a critical effector role for interleukin-1. *Transplant Proc* 1993, 25:1216–1217
8. Cron RQ, Chatham WW: The rheumatologist's role in COVID-19. *J Rheumatol* 2020, 47:639–642
9. Filippou PS, Karagiannis GS: Cytokine storm during chemotherapy: a new companion diagnostic emerges? *Oncotarget* 2020, 11:213–215
10. Gartung A, Yang J, Sukhatme VP, Bielenberg DR, Fernandes D, Chang J, Schmidt BA, Hwang SH, Zurakowski D, Huang S, Kieran MW, Hammock BD, Panigrahy D: Suppression of chemotherapy-induced cytokine/lipid mediator surge and ovarian cancer by a dual COX-2/sEH inhibitor. *Proc Natl Acad Sci U S A* 2019, 116:1698–1703
11. Sulciner ML, Serhan CN, Gilligan MM, Mudge DK, Chang J, Gartung A, Lehner KA, Bielenberg DR, Schmidt B, Dalli J, Greene ER, Gus-Brautbar Y, Piwowarski J, Mammoto T, Zurakowski D, Perretti M, Sukhatme VP, Kaipainen A, Kieran MW, Huang S, Panigrahy D: Resolvins suppress tumor growth and enhance cancer therapy. *J Exp Med* 2018, 215:115–140
12. Tisoncik JR, Korth MJ, Simmons CP, Farrar J, Martin TR, Katze MG: Into the eye of the cytokine storm. *Microbiol Mol Biol Rev* 2012, 76:16–32
13. Channappanavar R, Fehr AR, Vijay R, Mack M, Zhao J, Meyerholz DK, Perlman S: Dysregulated type I interferon and inflammatory monocyte-macrophage responses cause lethal pneumonia in SARS-CoV-infected mice. *Cell Host Microbe* 2016, 19:181–193
14. Herold S, Steinmueller M, von Wulffen W, Cakarova L, Pinto R, Pleschka S, Mack M, Kuziel WA, Corazza N, Brunner T, Seeger W, Lohmeyer J: Lung epithelial apoptosis in influenza virus pneumonia: the role of macrophage-expressed TNF-related apoptosis-inducing ligand. *J Exp Med* 2008, 205:3065–3077
15. Hogner K, Wolff T, Pleschka S, Plog S, Gruber AD, Kalinke U, Walmarth HD, Bodner J, Gattenlohner S, Lewe-Schlosser P, Matrosovich M, Seeger W, Lohmeyer J, Herold S: Macrophage-expressed IFN-beta contributes to apoptotic alveolar epithelial cell injury in severe influenza virus pneumonia. *PLoS Pathog* 2013, 9:e1003188
16. Rodrigue-Gervais IG, Labbe K, Dagenais M, Dupaul-Chicoine J, Champagne C, Morizot A, Skeldon A, Brincks EL, Vidal SM, Griffith TS, Saleh M: Cellular inhibitor of apoptosis protein cIAP2 protects against pulmonary tissue necrosis during influenza virus infection to promote host survival. *Cell Host Microbe* 2014, 15:23–35
17. Smits SL, de Lang A, van den Brand JM, Leijten LM, van IWF, Eijkemans MJ, van Amerongen G, Kuiken T, Andeweg AC, Osterhaus AD, Haagmans BL: Exacerbated innate host response to SARS-CoV in aged non-human primates. *PLoS Pathog* 2010, 6: e1000756
18. Jorgensen I, Rayamajhi M, Miao EA: Programmed cell death as a defence against infection. *Nat Rev Immunol* 2017, 17:151–164
19. Fung TS, Huang M, Liu DX: Coronavirus-induced ER stress response and its involvement in regulation of coronavirus-host interactions. *Virus Res* 2014, 194:110–123
20. Versteeg GA, van de Nes PS, Bredenbeek PJ, Spaan WJ: The coronavirus spike protein induces endoplasmic reticulum stress and upregulation of intracellular chemokine mRNA concentrations. *J Virol* 2007, 81:10981–10990
21. Shi CS, Nabar NR, Huang NN, Kehrl JH: SARS-coronavirus open reading frame-8b triggers intracellular stress pathways and activates NLRP3 inflammasomes. *Cell Death Discov* 2019, 5:101
22. Serhan CN: Pro-resolving lipid mediators are leads for resolution physiology. *Nature* 2014, 510:92–101
23. Wang D, Dubois RN: Eicosanoids and cancer. *Nat Rev Cancer* 2010, 10:181–193
24. Schmelzer KR, Kubala L, Newman JW, Kim IH, Eisrich JP, Hammock BD: Soluble epoxide hydrolase is a therapeutic target for acute inflammation. *Proc Natl Acad Sci U S A* 2005, 102:9772–9777
25. Chopra S, Giovanelli P, Alvarado-Vazquez PA, Alonso S, Song M, Sandoval TA, Chae CS, Tan C, Fonseca MM, Gutierrez S, Jimenez L, Subbaramaiah K, Iwawaki T, Kingsley PJ, Marnett LJ, Kossenkov AV, Crespo MS, Dannenberg AJ, Glimcher LH, Romero-Sandoval EA, Cubillos-Ruiz JR: IRE1alpha-XBPI signaling in leukocytes controls prostaglandin biosynthesis and pain. *Science* 2019, 365:eaau6499
26. von Moltke J, Trinidad NJ, Moayeri M, Kintzer AF, Wang SB, van Rooijen N, Brown CR, Krantz BA, Leppla SH, Gronert K, Vance RE: Rapid induction of inflammatory lipid mediators by the inflammasome in vivo. *Nature* 2012, 490:107–111
27. Hosoi T, Honda M, Oba T, Ozawa K: ER stress upregulated PGE(2)/IFNgamma-induced IL-6 expression and down-regulated iNOS expression in glial cells. *Sci Rep* 2013, 3:3388
28. Das UN: Can bioactive lipids inactivate coronavirus (COVID-19)? *Arch Med Res* 2020, 51:282–286
29. Regidor PA: Covid-19 management with inflammation resolving mediators? perspectives and potential. *Med Hypotheses* 2020, 142: 109813
30. Hoxha M: What about COVID-19 and arachidonic acid pathway? *Eur J Clin Pharmacol* 2020 [Epub ahead of print]. doi:10.1007/s00228-020-02941-w
31. Yang R, Chiang N, Oh SF, Serhan CN: Metabolomics-lipidomics of eicosanoids and docosanoids generated by phagocytes. *Curr Protoc Immunol* 2011. Chapter 14:Unit 14.26
32. Serhan CN: Novel eicosanoid and docosanoid mediators: resolvins, docosatrienes, and neuroprotectins. *Curr Opin Clin Nutr Metab Care* 2005, 8:115–121
33. Hong S, Gronert K, Devchand PR, Moussignac RL, Serhan CN: Novel docosatrienes and 17S-resolvins generated from docosahexaenoic acid in murine brain, human blood, and glial cells: autacoids in anti-inflammation. *J Biol Chem* 2003, 278:14677–14687
34. Dennis EA, Norris PC: Eicosanoid storm in infection and inflammation. *Nat Rev Immunol* 2015, 15:511–523
35. Norris PC, Gosselin D, Reichart D, Glass CK, Dennis EA: Phospholipase A2 regulates eicosanoid class switching during inflammasome activation. *Proc Natl Acad Sci U S A* 2014, 111:12746–12751
36. Zeldin DC: Epoxigenase pathways of arachidonic acid metabolism. *J Biol Chem* 2001, 276:36059–36062
37. Imig JD, Hammock BD: Soluble epoxide hydrolase as a therapeutic target for cardiovascular diseases. *Nat Rev Drug Discov* 2009, 8: 794–805
38. Shen HC: Soluble epoxide hydrolase inhibitors: a patent review. *Expert Opin Ther Pat* 2010, 20:941–956
39. Wang W, Zhu J, Lyu F, Panigrahy D, Ferrara KW, Hammock B, Zhang G: Omega-3 polyunsaturated fatty acids-derived lipid metabolites on angiogenesis, inflammation and cancer. *Prostaglandins Other Lipid Mediat* 2014, 113–115:13–20
40. Wagner KM, McReynolds CB, Schmidt WK, Hammock BD: Soluble epoxide hydrolase as a therapeutic target for pain, inflammatory and neurodegenerative diseases. *Pharmacol Ther* 2017, 180:62–76
41. Gilroy DW, Edin ML, De Maeyer RP, Bystrom J, Newson J, Lih FB, Stables M, Zeldin DC, Bishop-Bailey D: CYP450-derived oxylipins mediate inflammatory resolution. *Proc Natl Acad Sci U S A* 2016, 113:E3240–E3249
42. Deng Y, Edin ML, Theken KN, Schuck RN, Flake GP, Kannon MA, DeGraff LM, Lih FB, Foley J, Bradbury JA, Graves JP, Tomer KB, Falck JR, Zeldin DC, Lee CR: Endothelial CYP epoxigenase over-expression and soluble epoxide hydrolase disruption attenuate acute vascular inflammatory responses in mice. *FASEB J* 2011, 25: 703–713
43. Yu G, Zeng X, Wang H, Hou Q, Tan C, Xu Q, Wang H: 14,15-Epoxyeicosatrienoic acid suppresses cigarette smoke extract-induced apoptosis in lung epithelial cells by inhibiting endoplasmic reticulum stress. *Cell Physiol Biochem* 2015, 36:474–486
44. Kodani SD, Hammock BD: The 2014 Bernard B. Brodie award lecture-epoxide hydrolases: drug metabolism to therapeutics for chronic pain. *Drug Metab Dispos* 2015, 43:788–802
45. Lazaar AL, Yang L, Boardley RL, Goyal NS, Robertson J, Baldwin SJ, Newby DE, Wilkinson IB, Tal-Singer R, Mayer RJ, Cherian J: Pharmacokinetics, pharmacodynamics and adverse event
- 559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620

- profile of GSK2256294, a novel soluble epoxide hydrolase inhibitor. *Br J Clin Pharmacol* 2016; 81:971–979
46. Belyanskaia SL, Ding Y, Callahan JF, Lazaar AL, Israel DI: Discovering drugs with DNA-encoded library technology: from concept to clinic with an inhibitor of soluble epoxide hydrolase. *Chembiochem* 2017; 18:837–842
47. Zhou Y, Liu T, Duan JX, Li P, Sun GY, Liu YP, Zhang J, Dong L, Lee KSS, Hammock BD, Jiang JX, Guan CX: Soluble epoxide hydrolase inhibitor attenuates lipopolysaccharide-induced acute lung injury and improves survival in mice. *Shock* 2017; 47:638–645
48. Ono E, Dutile S, Kazani S, Wechsler ME, Yang J, Hammock BD, Douda DN, Tabet Y, Khaddaj-Mallat R, Sirois M, Sirois C, Rizcallah E, Rousseau E, Martin R, Sutherland ER, Castro M, Jarjour NN, Israel E, Levy BD; National Heart, Lung, and Blood Institute's Asthma Clinical Research Network: Lipoxin generation is related to soluble epoxide hydrolase activity in severe asthma. *Am J Respir Crit Care Med* 2014; 190:886–897
49. Schunck WH, Konkel A, Fischer R, Weylandt KH: Therapeutic potential of omega-3 fatty acid-derived epoxyeicosanoids in cardiovascular and inflammatory diseases. *Pharmacol Ther* 2018; 183:177–204
50. Xia R, Sun L, Liao J, Li H, You X, Xu D, Yang J, Hwang SH, Jones RD, Hammock B, Yang GY: Inhibition of pancreatic carcinoma growth through enhancing omega-3 epoxy polyunsaturated fatty acid profile by inhibition of soluble epoxide hydrolase. *Anticancer Res* 2019; 39:3651–3660
51. Lopez-Vicario C, Alcaraz-Quiles J, Garcia-Alonso V, Rius B, Hwang SH, Titos E, Lopategui A, Hammock BD, Arroyo V, Claria J: Inhibition of soluble epoxide hydrolase modulates inflammation and autophagy in obese adipose tissue and liver: role for omega-3 epoxides. *Proc Natl Acad Sci U S A* 2015; 112:536–541
52. Ackermann M, Verleden SE, Kuehnel M, Haverich A, Welte T, Laenger F, Vanstapel A, Werlein C, Stark H, Tzankov A, Li WW, Li VW, Mentzer SJ, Jonigk D: Pulmonary vascular endothelialitis, thrombosis, and angiogenesis in COVID-19. *N Engl J Med* 2020; 383:120–128
53. Hu J, Dziumbla S, Lin J, Bibli SI, Zukunft S, de Mos J, Awwad K, Fromel T, Jungmann A, Devraj K, Cheng Z, Wang L, Fauser S, Eberhart CG, Sodhi A, Hammock BD, Liebner S, Muller OJ, Glaubitz C, Hammes HP, Popp R, Fleming I: Inhibition of soluble epoxide hydrolase prevents diabetic retinopathy. *Nature* 2017; 552:248–252
54. Hasegawa E, Inafuku S, Mulki L, Okunuki Y, Yanai R, Smith KE, Kim CB, Klokman G, Bielenberg DR, Puli N, Falck JR, Husain D, Miller JW, Edin ML, Zeldin DC, Lee KSS, Hammock BD, Schunck WH, Connor KM: Cytochrome P450 monooxygenase lipid metabolites are significant second messengers in the resolution of choroidal neovascularization. *Proc Natl Acad Sci U S A* 2017; 114: E7545–E7553
55. Panigrahy D, Gartung A, Yang J, Yang H, Gilligan MM, Sulciner ML, Bhasin SS, Bielenberg DR, Chang J, Schmidt BA, Piwowarski J, Fishbein A, Soler-Ferran D, Sparks MA, Staffa SJ, Sukhatme V, Hammock BD, Kieran MW, Huang S, Bhasin M, Serhan CN, Sukhatme VP: Preoperative stimulation of resolution and inflammation blockade eradicates micrometastases. *J Clin Invest* 2019; 129:2964–2979
56. Barden AE, Mas E, Mori TA: n-3 Fatty acid supplementation and proresolving mediators of inflammation. *Curr Opin Lipidol* 2016; 27:26–32
57. Souza PR, Marques RM, Gomez EA, Colas RA, De Matteis R, Zak A, Patel M, Collier DJ, Dalli J: Enriched marine oil supplements increase peripheral blood specialized pro-resolving mediators concentrations and reprogram host immune responses: a randomized double-blind placebo-controlled study. *Circ Res* 2020; 126:75–90
58. Norris PC, Arnardottir H, Sanger JM, Fichtner D, Keyes GS, Serhan CN: Resolin D3 multi-level proresolving actions are host protective during infection. *Prostaglandins Leukot Essent Fatty Acids* 2018; 138:81–89
59. Spite M, Norling LV, Summers L, Yang R, Cooper D, Petasis NA, Flower RJ, Perretti M, Serhan CN: Resolvin D2 is a potent regulator of leukocytes and controls microbial sepsis. *Nature* 2009; 461: 1287–1291
60. Chiang N, Fredman G, Backhed F, Oh SF, Vickery T, Schmidt BA, Serhan CN: Infection regulates pro-resolving mediators that lower antibiotic requirements. *Nature* 2012; 484:524–528
61. Dalli J, Chiang N, Serhan CN: Elucidation of novel 13-series resolvins that increase with atorvastatin and clear infections. *Nat Med* 2015; 21:1071–1075
62. Ledford H: Coronavirus breakthrough: dexamethasone is first drug shown to save lives. *Nature* 2020; 582:469
63. Pyrillou K, Chairakaki AD, Tamvakopoulos C, Andreakos E: Dexamethasone induces omega3-derived immunoresolvents driving resolution of allergic airway inflammation. *J Allergy Clin Immunol* 2018; 142:691–695.e4
64. Das I, Png CW, Oancea I, Hasnain SZ, Lourie R, Proctor M, Eri RD, Sheng Y, Crane DI, Florin TH, McGuckin MA: Glucocorticoids alleviate intestinal ER stress by enhancing protein folding and degradation of misfolded proteins. *J Exp Med* 2013; 210:1201–1216
65. FitzGerald GA: Misguided drug advice for COVID-19. *Science* 2020; 367:1434
66. Serhan CN, Chiang N, Van Dyke TE: Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nat Rev Immunol* 2008; 8:349–361
67. Levy BD, Clish CB, Schmidt B, Gronert K, Serhan CN: Lipid mediator class switching during acute inflammation: signals in resolution. *Nat Immunol* 2001; 2:612–619
68. Serhan CN, Clish CB, Brannon J, Colgan SP, Chiang N, Gronert K: Novel functional sets of lipid-derived mediators with antiinflammatory actions generated from omega-3 fatty acids via cyclooxygenase 2-nonsteroidal antiinflammatory drugs and transcellular processing. *J Exp Med* 2000; 192:1197–1204
69. Serhan CN, Hong S, Gronert K, Colgan SP, Devchand PR, Mirick G, Moussignac RL: Resolvins: a family of bioactive products of omega-3 fatty acid transformation circuits initiated by aspirin treatment that counter proinflammation signals. *J Exp Med* 2002; 196:1025–1037
70. Yousefifard M, Zali A, Zarghi A, Madani Neishaboori A, Hosseini M, Safari S: Non-steroidal anti-inflammatory drugs in management of COVID-19: a systematic review on current evidence. *Int J Clin Pract* 2020; 00:e13557
71. Goswami SK, Wan D, Yang J, Trindade da Silva CA, Morisseau C, Kodani SD, Yang GY, Inceoglu B, Hammock BD: Anti-ulcer efficacy of soluble epoxide hydrolase inhibitor TPPU on diclofenac-induced intestinal ulcers. *J Pharmacol Exp Ther* 2016; 357:529–536
72. Ghosh R, Goswami SK, Feitoza L, Hammock B, Gomes AV: Diclofenac induces proteasome and mitochondrial dysfunction in murine cardiomyocytes and hearts. *Int J Cardiol* 2016; 223:923–935
73. George PM, Wells AU, Jenkins RG: Pulmonary fibrosis and COVID-19: the potential role for antifibrotic therapy. *Lancet Respir Med* 2020 [Epub ahead of print]. doi:10.1016/S2213-2600(20)30225-3
74. Kim J, Imig JD, Yang J, Hammock BD, Padanilam BJ: Inhibition of soluble epoxide hydrolase prevents renal interstitial fibrosis and inflammation. *Am J Physiol Renal Physiol* 2014; 307:F971–F980
75. Chiang CW, Lee HT, Targ NC, Kuo KL, Cheng LC, Lee TS: Genetic deletion of soluble epoxide hydrolase attenuates inflammation and fibrosis in experimental obstructive nephropathy. *Mediators Inflamm* 2015; 2015:693260
76. Zhang CH, Zheng L, Gui L, Lin JY, Zhu YM, Deng WS, Luo M: Soluble epoxide hydrolase inhibition with t-TUCB alleviates liver fibrosis and portal pressure in carbon tetrachloride-induced cirrhosis in rats. *Clin Res Hepatol Gastroenterol* 2018; 42:118–125
77. Deng W, Zhu Y, Lin J, Zheng L, Zhang C, Luo M: Inhibition of soluble epoxide hydrolase lowers portal hypertension in cirrhotic rats by ameliorating endothelial dysfunction and liver fibrosis. *Prostaglandins Other Lipid Mediat* 2017; 131:67–74

621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744



- 745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
78. Dong XW, Jia YL, Ge LT, Jiang B, Jiang JX, Shen J, Jin YC, Guan Y, Sun Y, Xie QM: Soluble epoxide hydrolase inhibitor AUDA decreases bleomycin-induced pulmonary toxicity in mice by inhibiting the p38/Smad3 pathways. *Toxicology* 2017, 389: 31–41
79. Zhou C, Huang J, Li Q, Zhan C, He Y, Liu J, Wen Z, Wang DW: Pharmacological inhibition of soluble epoxide hydrolase ameliorates chronic ethanol-induced cardiac fibrosis by restoring autophagic flux. *Alcohol Clin Exp Res* 2018, 42:1970–1978
80. Harris TR, Bettaieb A, Kodani S, Dong H, Myers R, Chiamvimonvat N, Haj FG, Hammock BD: Inhibition of soluble epoxide hydrolase attenuates hepatic fibrosis and endoplasmic reticulum stress induced by carbon tetrachloride in mice. *Toxicol Appl Pharmacol* 2015, 286:102–111

813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880