

Lawrence Berkeley National Laboratory

LBL Publications

Title

Gamma Ray Spectroscopy on 155-Day Lu177 Using Lithium-Drifted Germanium Counters

Permalink

<https://escholarship.org/uc/item/38s6x7ck>

Authors

Blok, J

Shirley, D A

Publication Date

1964-10-01

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

University of California
Ernest O. Lawrence
Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

GAMMA RAY SPECTROSCOPY ON 155-DAY Lu^{177} USING
LITHIUM-DRIFTED GERMANIUM COUNTERS

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.



UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

GAMMA RAY SPECTROSCOPY ON 155-DAY Lu¹⁷⁷ USING
LITHIUM-DRIFTED GERMANIUM COUNTERS

J. Blok and D. A. Shirley

October 1964

GAMMA RAY SPECTROSCOPY ON 155-DAY Lu^{177} USING
LITHIUM-DRIFTED GERMANIUM COUNTERS

J. Blok and D. A. Shirley

Department of Chemistry and Lawrence Radiation Laboratory
University of California
Berkeley, California

October 1964

We have used recently-developed lithium drifted germanium gamma-ray detectors¹⁾ to study the spectrum from the decay of the 155-day isomer of Lu^{177} . This isomer has been shown to be a three quasi-particle state of an odd proton and an uncoupled neutron pair which gives rise to the very high spin of $23/2^{2,3)}$.

Quite thorough studies have been made of the decay scheme of 155-day Lu^{177} $2,3)$. The isomer was shown to decay partly by isomeric transition to high-lying levels of the $K = 7/2^+$ [404] rotational band of Lu^{177} and partly by β^- decay to corresponding three quasi-particle states in Hf^{177} . Alexander, et al.²⁾ were able to confirm all the levels of the Lu^{177} rotational band up to spin $17/2$ and all the members of the two Hf^{177} rotational bands up to spin $21/2$ except for the spin $19/2$ level of the $K = 7/2^-$ [514] band. The decay scheme is shown in Fig. 1.

We feel that this decay is a particularly important one for nuclear structure theory, involving as it does up to eight members of a rotational band in Hf^{177} , and it seemed worthwhile to confirm the proposed level schemes independently by definitive coincidence measurements and high-sensitivity gamma-ray spectroscopy using the newly-developed gamma-ray counters. In particular we were interested in filling the gap in the $7/2^-$ [514] band, in which there was no experimental evidence for the $19/2$ state.

A sample of 155-day Lu^{177} was prepared by irradiating natural lutetium metal for one 41-day cycle with a neutron flux of 5×10^{14} in the ETR reactor at Arco, Idaho. After a waiting period of several months to permit the 7-day isomer to decay, the sample was dissolved and separated from long-lived activities of all other elements by means of an ion-exchange procedure.

The Ge(Li) counters which were used to obtain the spectra measured 1 cm. \times 2 cm. in sensitive area and 6 mm. in depth. They were produced at this laboratory, as were the associated preamplifiers, amplifiers, and coincidence circuitry. The gamma-ray absorption efficiency of the counters was such that sources of less than one millicurie gave reasonable counting rates, enabling us to obtain the spectrum shown in Fig. 2 in about two days.

In Fig. 2 we show a gamma-ray spectrum of 155-day Lu^{177} obtained with the Ge(Li) counter. Comparison with the bent-crystal data²⁾ shows that, although not as good for low energies, the resolution of the Ge(Li) counter exceeds that of the bent-crystal spectrometer for energies above about 300 keV. In two other respects the Ge(Li) counters are far superior to a bent-crystal spectrometer for these reasonably high energies. First, the efficiency is far higher; our experiments were done with sources of less than one millicurie. Secondly, the signal-to-noise ratio is very high for Ge(Li) counters, since there is little background due to incoherent scattering.

In Fig. 3 we show a spectrum of the region around 420 keV on an expanded scale. One observes a peak at about 426 keV which was not seen by Alexander, et al. This peak fits energetically as the cross-over transition between the missing $19/2$ level and the $15/2$ level in the $K = 7/2^-$ band of Hf^{177} . To establish that this 426 keV gamma ray actually was the $19/2$ to $15/2$ transition; we used two Ge(Li) counters and obtained a spectrum which was in coincidence with this gamma ray. Due to the low efficiency of the Ge(Li) counters, the coincidence counting rate was very low; hence only the more prominent

gamma rays could be seen. Peaks were observed in the coincidence spectrum at 113, 136, 249 and 341 keV and the other prominent peaks of the singles spectrum were either sharply reduced in intensity or else were completely missing, as shown in Fig. 4. Thus the 426 keV gamma ray must certainly come from the $19/2$ to $15/2$ transition in the $K = 7/2^-$ band of Hf^{177} . A careful search was made for the transitions which must feed the $19/2$ level, namely the $21/2$ to $19/2$ cascade of about 243 keV or the $21/2$ to $19/2$ interband transition of about 284 keV. Neither peak could be found; the former because of the high Compton background in the low energy region, and the latter because resolution from the intense 281 keV peak was impossible. Even without this final proof, however, the position of the $19/2$ level appears to be well established.

As an additional independent check of the decay scheme of Alexander, et al. we measured the spectra in coincidence with the 105, 413, and 418 keV transitions. Prominent peaks were again present or absent in agreement with the above decay scheme.

We conclude that the proposed decay scheme is confirmed in detail by the gamma-ray spectra and the coincidence work with Ge(Li) counters, and in particular that the 426 keV gamma ray is present, thus establishing the $19/2$ member of the $7/2^-$ [514] band. These measurements indicate the power of the Ge(Li) counters for gamma ray spectroscopy. Accurate intensity determinations and nuclear orientation measurements with 155-day Lu^{177} are underway.

REFERENCES

*This work was done under the auspices of the U. S. Atomic Energy Commission.

- 1) A. J. Tavendale, in Proc. Int. Symp. Nuclear Electronics, Paris (1963) 235
- 2) P. Alexander, F. Boehm and E. Kankeleit, Phys. Rev. 133, (1964) B284
- 3) L. Kristensen, M. Jorgensen, O. B. Nielsen and G. Sidenius, Phys. Letters 8 (1964) 57


FIGURE CAPTIONS


Fig. 1. Decay scheme of Lu^{177} taken from Ref. 2. We have added the $I = 19/2$ level of the $K = 7/2^-$ [514] band and its associated gamma ray. Energies of gamma rays above 318 keV are ours. Gamma rays observed in this work are underlined.


Fig. 2. Gamma ray spectrum of 155-day Lu^{177} with Ge(Li) detectors. The peaks are identified by their energy in keV.


Fig. 3. Detail of 155-day Lu^{177} gamma ray spectrum showing the 426 keV transition. The experimental points are indicated as filled circles.

Fig. 4. Gamma ray spectrum of 155-day Lu^{177} in coincidence with the 426 keV transition. Some of the peaks in coincidence with the 418 keV peak are also present (weakly), since a small amount of the 418 keV peak was accepted through the gate. Coincidence of the 113 with the 426 keV peak was determined in a separate run.









This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

[The page contains extremely faint and illegible text, likely bleed-through from the reverse side of the document. No specific words or phrases can be discerned.]

