

Lawrence Berkeley National Laboratory

Lawrence Berkeley National Laboratory

Title

Canonical Representations $sp(2n, R)$

Permalink

<https://escholarship.org/uc/item/39c1h1qw>

Author

Rangarajan, G.

Publication Date

1991-08-01



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA


Accelerator & Fusion Research Division

Submitted to the Journal of Mathematical Physics

Canonical Representations $sp(2n, \mathbb{R})$

G. Rangarajan

August 1991



DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. Neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California and shall not be used for advertising or product endorsement purposes.

Lawrence Berkeley Laboratory is an equal opportunity employer.

Canonical Representations $sp(2n, R)^*$

G. Rangarajan

Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

Submitted to Journal of Mathematical Physics

* This work was supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, High Energy Physics Division, of the U. S. Department of Energy, under Contract No. DE-AC03-76SF00098.

CANONICAL REPRESENTATIONS OF $\text{sp}(2n, \mathbb{R})$

Govindan Rangarajan

*Lawrence Berkeley Laboratory, University of California,
Berkeley, CA 94720*

Filippo Neri

*AT Division, Los Alamos National Laboratory,
Los Alamos, NM 87545*

In this paper, we study a rather unconventional real basis for the real symplectic algebra $\text{sp}(2n, \mathbb{R})$. We demonstrate the utility of this basis for practical computations by giving a simple derivation of the second and fourth order indices of irreducible representations of $\text{sp}(2n, \mathbb{R})$.

PACS numbers: 02.20.+b, 03.20.+i

I. INTRODUCTION

In this paper, we study irreducible representations (irreps) of the real symplectic Lie algebra $\mathfrak{sp}(2n, \mathbb{R})$ [1] carried by homogeneous polynomials of the $2n$ phase-space variables $\{q_i, p_i\}$ ($i = 1, 2, \dots, n$). These representations of $\mathfrak{sp}(2n, \mathbb{R})$ are the physically interesting ones in classical mechanics[2].

First, we define a real basis for $\mathfrak{sp}(2n, \mathbb{R})$. Using this basis, irreps of $\mathfrak{sp}(2n, \mathbb{R})$ carried by homogeneous polynomials of phase-space variables are obtained. Weight vectors corresponding to these irreps are also computed. Using these results, we finally give a simple derivation of the second and fourth order indices corresponding to these irreps.

II. A REAL BASIS FOR $\mathfrak{SP}(2N, \mathbb{R})$

In this section, we obtain a real basis for $\mathfrak{sp}(2n, \mathbb{R})$. Since we are interested in representations carried by functions of phase-space variables, it is convenient to define the operators constituting a basis for $\mathfrak{sp}(2n, \mathbb{R})$ using these variables. This can be done using the concept of a Lie operator[2]. Let us denote the collection of $2n$ phase-space variables q_i, p_i ($i = 1, 2, \dots, n$) by the symbol z . The Lie operator corresponding to a phase-space function $f(z)$ is denoted by $:f(z):$. It is defined by its action on a phase-space function $g(z)$ as shown below

$$:f(z): g(z) = [f(z), g(z)]. \quad (2.1)$$

Here $[f(z), g(z)]$ denotes the usual Poisson bracket of the functions $f(z)$ and $g(z)$:

$$[f(z), g(z)] \equiv \sum_{i=1}^n \left(\frac{\partial f(z)}{\partial q_i} \frac{\partial g(z)}{\partial p_i} - \frac{\partial f(z)}{\partial p_i} \frac{\partial g(z)}{\partial q_i} \right). \quad (2.2)$$

We are now in a position to give a basis for $\mathfrak{sp}(2n, \mathbb{R})$. Consider the following set of real operators:

$$\begin{aligned}
:C_{jk}: &= :q_j p_k:, \\
:L_{jk}: &= :q_j q_k:, \quad j \leq k, \\
:R_{jk}: &= :p_j p_k:, \quad j \leq k,
\end{aligned} \tag{2.3}$$

where the indices j and k range from 1 to n . We note that these are nothing but Lie operators corresponding to the set of all quadratic monomials in variables q_j and p_k . It can be shown that these operators constitute a real basis for the Lie algebra $\mathfrak{sp}(2n, \mathbb{R})$ [3]. As expected, there are $n(2n + 1)$ elements in this basis. We will denote a general basis element by the symbol w_l .

The commutator of two Lie operators $:f:$ and $:g:$ can be shown[2] to satisfy the following relation

$$\{ :f:, :g: \} \equiv :f::g: - :g::f: = :[f, g]: \tag{2.4}$$

Using this property, the basis elements are seen to satisfy the following commutation relations

$$\begin{aligned}
\{ :C_{jk} :, :C_{rs}: \} &= :C_{rk}: \delta_{js} - :C_{js}: \delta_{kr}, \\
\{ :C_{jk} :, :L_{rs}: \} &= - :L_{js}: \delta_{kr} - :L_{jr}: \delta_{ks}, \\
\{ :C_{jk} :, :R_{rs}: \} &= :R_{ks}: \delta_{jr} + :R_{kr}: \delta_{js}, \\
\{ :L_{jk} :, :L_{rs}: \} &= 0, \\
\{ :R_{jk} :, :R_{rs}: \} &= 0, \\
\{ :L_{jk} :, :R_{rs}: \} &= :C_{ks}: \delta_{jr} + :C_{kr}: \delta_{js} + :C_{js}: \delta_{kr} + :C_{jr}: \delta_{ks}.
\end{aligned} \tag{2.5}$$

Here the indices j, k, r , and s range from 1 to n .

The basis given in Eq. (2.3) is not the conventional basis used for $\mathfrak{sp}(2n, \mathbb{R})$. It does not contain a basis for the unitary algebra $\mathfrak{u}(3)$ as a subset. However, this shortcoming is of no consequence when one studies only the symplectic algebra without any reference to its unitary subalgebras. In such cases, the real basis defined in Eq. (2.3) offers several advantages due to its one-to-one correspondence with the set of quadratic monomials. These advantages will become apparent in the discussions that follow.

III. REPRESENTATION OF $\mathfrak{SP}(2N, \mathbb{R})$

In this section, we first obtain the irreps of $\mathfrak{sp}(2n, \mathbb{R})$ carried by homogeneous polynomials of phase-space variables. Next, we compute the weight vectors corresponding to these irreps.

From the previous section, it is clear that Lie operators corresponding to the set of all quadratic polynomials in z constitute the symplectic Lie algebra $\mathfrak{sp}(2n, \mathbb{R})$. Therefore, an N dimensional representation of $\mathfrak{sp}(2n, \mathbb{R})$ is obtained by mapping each element (denoted in general by $:f_2:$) onto a $N \times N$ matrix $d(f_2)$ such that the following conditions are satisfied for all $:f_2:, :g_2:$ belonging to $\mathfrak{sp}(2n, \mathbb{R})$:

$$d(af_2 + bg_2) = a d(f_2) + b d(g_2), \quad a, b \in \mathcal{R} \quad (3.1)$$

$$d([f_2, g_2]) = \{d(g_2), d(f_2)\}. \quad (3.2)$$

A representation d is said to be (completely) reducible if it can be converted into a block diagonal form using a similarity transformation. Otherwise, it is said to be irreducible.

Irreducible representations of $\mathfrak{sp}(2n, \mathbb{R})$ carried by homogeneous polynomials in z can be obtained in a straightforward fashion. Let $\mathcal{P}^{(m)}(z)$ denote the set of all homogeneous polynomials in z of degree m and $g_m(z)$ denote a general element belonging

to this set. Consider the following property of the Lie operator $:f_2:$ [cf. Eq. (2.2)]

$$:f_2:g_m = [f_2, g_m] = h_m, \quad (3.3)$$

where h_m is an element of $\mathcal{P}^{(m)}(z)$. That is, the elements $:f_2:$ of $\mathfrak{sp}(2n, \mathbb{R})$ preserve the degree of the polynomial on which they act. We therefore get the following relation

$$:f_2:g_m = [f_2, g_m] \in \mathcal{P}^{(m)} \quad \forall g_m \in \mathcal{P}^{(m)}. \quad (3.4)$$

In other words, the set of all elements belonging to $\mathfrak{sp}(2n, \mathbb{R})$ leaves $\mathcal{P}^{(m)}$ invariant.

Let $\{P_\alpha^{(m)}\}$ be a basis for the set $\mathcal{P}^{(m)}$. Typically, we choose these basis elements to be the monomials of degree m in the $2n$ phase-space variables. The number of basis monomials $N(m)$ of degree m in the $2n$ phase-space variables is given by the relation[4, 3]

$$N(m) = \binom{2n + m - 1}{m}. \quad (3.5)$$

From Eq. (3.4) and the completeness of the set $\{P_\alpha^{(m)}\}$, we get the following relation

$$:f_2:P_\alpha^{(m)}(z) = \sum_{\beta=1}^{N(m)} d^{(m)}(f_2)_\alpha^\beta P_\beta^{(m)}(z) \quad \alpha = 1, 2, \dots, N(m). \quad (3.6)$$

where $d^{(m)}(f_2)_\alpha^\beta$ are coefficients multiplying the basis elements.

As we vary α from 1 to $N(m)$ in Eq. (3.6), the set of coefficients $d^{(m)}(f_2)_\alpha^\beta$ gives rise to an $N(m) \times N(m)$ matrix, $d^{(m)}(f_2)$, for each $:f_2:$ belonging to $\mathfrak{sp}(2n, \mathbb{R})$. It is easily verified that the set of such matrices (obtained by letting $:f_2:$ range over the entire Lie algebra) gives an $N(m)$ -dimensional representation of $\mathfrak{sp}(2n, \mathbb{R})$. It can be shown[3] that these representations (for each m) are in fact irreducible.

Next, we obtain the weight vectors for irreps of $\mathfrak{sp}(2n, \mathbb{R})$ defined above. First, we need the Cartan subalgebra of $\mathfrak{sp}(2n, \mathbb{R})$. It is easily verified that the Cartan subalgebra of $\mathfrak{sp}(2n, \mathbb{R})$ is spanned by the following n elements:

$$H_1 = :q_1 p_1:, \quad H_2 = :q_2 p_2:, \quad \dots, \quad H_n = :q_n p_n:. \quad (3.7)$$

As expected, the rank of $\mathfrak{sp}(2n, \mathbb{R})$ is equal to n . The $N(m)$ -dimensional irrep of H_i is given by Eq. (3.6).

To proceed further, we need to choose the $N(m)$ basis elements $P_\alpha^{(m)}(z)$. We make the simplest choice. We choose $\{P_\alpha^{(m)}(z)\}$ to be the set of all m th degree monomials in the $2n$ phase-space variables. Therefore, the general element $P_\alpha^{(m)}(z)$ is given by the following relation

$$P_\alpha^{(m)}(z) = q_1^{r_1} p_1^{r_2} \dots q_n^{r_{2n-1}} p_n^{r_{2n}}, \quad r_1 + r_2 + \dots + r_{2n} = m. \quad (3.8)$$

Now, the advantage of using the rather unconventional basis given in Eq. (2.3) becomes apparent. In this basis, the $N(m)$ -dimensional irreps of the Cartan basis elements H_i are automatically diagonal! This is true for all the irreps defined by Eq. (3.6). Moreover, it is a simple matter to calculate the general weight vector. Using Eq. (3.8), we obtain the following results

$$:q_i p_i: P_\alpha^{(m)}(z) = (r_{2i} - r_{2i-1}) P_\alpha^{(m)}(z), \quad i = 1, 2, \dots, n. \quad (3.9)$$

The weight vector $\lambda^{(\alpha)}$ is defined by the relation

$$:q_i p_i: P_\alpha^{(m)}(z) = \lambda_i^{(\alpha)} P_\alpha^{(m)}(z), \quad i = 1, 2, \dots, n. \quad (3.10)$$

Comparing Eqs. (3.9) and (3.10), we obtain the result

$$\lambda^{(\alpha)} = (r_2 - r_1, r_4 - r_3, \dots, r_{2n} - r_{2n-1}). \quad (3.11)$$

IV. INDICES OF IRREDUCIBLE REPRESENTATIONS

Indices of irreducible representations[5] play an important part in representation theory. For example, they can be used to decompose direct products of representations and compute branching rules. In this section, we illustrate the utility of the basis given in Eq. (2.3) by explicitly computing the second and fourth order indices of the irreps carried by homogeneous polynomials.

A. Second order index

The second order index is defined as follows[5]

$$I_m^{(2)} = \sum_{\alpha=1}^{N(m)} [\lambda^{(\alpha)}]^2, \quad (4.1)$$

where

$$[\lambda^{(\alpha)}]^2 = \sum_{i=1}^n [\lambda_i^{(\alpha)}]^2. \quad (4.2)$$

Here n is the rank of $\mathfrak{sp}(2n, \mathbb{R})$, $\lambda_i^{(\alpha)}$ is the i th component of the α th weight vector and $N(m)$ is the dimension of the irrep carried by the m th degree homogeneous polynomials [cf. Eq. (3.5)]. The expression for $I_m^{(2)}$ in Eq. (4.1) can be simplified as follows. From symmetry considerations, we have the following relation

$$\sum_{\alpha=1}^{N(m)} [\lambda_i^{(\alpha)}]^2 = \sum_{\alpha=1}^{N(m)} [\lambda_j^{(\alpha)}]^2 \quad \forall i, j. \quad (4.3)$$

Hence, Eq. (4.1) can be rewritten as follows:

$$I_m^{(2)} = n \sum_{\alpha=1}^{N(m)} [\lambda_1^{(\alpha)}]^2. \quad (4.4)$$

Before proceeding further, we first establish the relation between the second order index and the more familiar Dynkin index[5]. Using the fact that $d^{(m)}(q_i p_i)$ is diagonal, we obtain the relation [cf. Eq. (3.6)]

$$:q_1 p_1: P_{\alpha}^{(m)}(z) = d^{(m)}(q_1 p_1)_{\alpha}^{\alpha} P_{\alpha}^{(m)}(z) \quad \alpha = 1, 2, \dots, N(m) \quad (4.5)$$

Comparing this with Eq. (3.10), we discover that Eq. (4.4) can be rewritten as follows

$$I_m^{(2)} = n \text{Tr}[d^{(m)}(q_1 p_1) d^{(m)}(q_1 p_1)]. \quad (4.6)$$

However, the Dynkin index $\nu(m)$ is defined by the relation[6]

$$\nu(m) = \text{Tr}[d^{(m)}(w_{\alpha}) d^{(m)}(w_{\beta})] / \text{Tr}[d^{(2)}(w_{\alpha}) d^{(2)}(w_{\beta})]. \quad (4.7)$$

Since $\nu(m)$ is independent of the the choice for w_α and w_β , we can take both to be equal to $q_1 p_1$. Then, we get the relation

$$I_m^{(2)} = \nu(m) I_2^{(2)}. \quad (4.8)$$

We now turn to the task of computing $I_m^{(2)}$. From Eq. (4.4), we see that we need to sum over squares of the first components of the weight vectors corresponding to all the monomials $P_\alpha^{(m)}(z)$ ($\alpha = 1, 2, \dots, N(m)$). Since $:q_1 p_1:$ acts only on the q_1 and p_1 variables, we need to know only the q_1 and p_1 content of the basis monomials $P_\alpha^{(m)}(z)$. Hence, we write the general monomial as follows

$$P_\alpha^{(m)}(z) = q_1^{m-k-l} p_1^l h_k(q_2, p_2, \dots, q_n, p_n), \quad (4.9)$$

where h_k stands for a monomial of degree k in the $2n - 2$ variables q_2, p_2, \dots, p_n . The index k varies from 0 to m and the index l from 0 to $m - k$. The total number of monomials h_k of degree k is given by the relation [cf. Eq. (3.5)]

$$N'(k) = \binom{2n + k - 3}{k}. \quad (4.10)$$

From Eqs. (3.11) and (4.9), we find the following expression for the first component $\lambda_1^{(\alpha)}$ of the weight vector corresponding to the monomial $P_\alpha^{(m)}(z)$

$$\lambda_1^{(\alpha)} = -m + k + 2l. \quad (4.11)$$

Since all $N'(k)$ monomials with the same value of k have the same weight, we obtain the result:

$$I_m^{(2)} = n \sum_{k=0}^m \sum_{l=0}^{m-k} \binom{2n + k - 3}{k} (-m + k + 2l)^2. \quad (4.12)$$

However, the following identity can be proved

$$\sum_{l=0}^{m-k} (-m + k + 2l)^2 = 2 \binom{m - k + 2}{m - k - 1}. \quad (4.13)$$

Therefore, we obtain the relation

$$I_m^{(2)} = 2n \sum_{k=0}^{m-1} \binom{2n+k-3}{k} \binom{m-k+2}{m-k-1}. \quad (4.14)$$

After some manipulation, this can be shown to equal the following result

$$I_m^{(2)} = 2n \binom{2n+m}{2n+1}. \quad (4.15)$$

It is easily verified that this expression gives the correct second order indices[7] for the irreps being studied (apart from an overall numerical factor).

From Eq. (4.8), the Dynkin index $\nu(m)$ is given by the relation

$$\nu(m) = \binom{2n+m}{2n+1} / (2n+2). \quad (4.16)$$

Further, the eigenvalue $C_2(\Lambda)$ of the second order Casimir operator can be calculated using the following relation[6]

$$C_2(\Lambda) = N(2)\nu(m)/N(m), \quad (4.17)$$

where $N(2)$ is equal to the dimension of $\text{sp}(2n, \mathbb{R})$. We obtain the following result

$$C_2(\Lambda) = (m^2 + 2nm)/4(n+1). \quad (4.18)$$

B. Fourth order index

The fourth order index is defined as follows[5]

$$I_m^{(4)} = \sum_{\alpha=1}^{N(m)} \left[\sum_{i=1}^n (\lambda_i^{(\alpha)})^2 \right]^2. \quad (4.19)$$

Using symmetry considerations, this can be rewritten as follows

$$I_m^{(4)} = n \sum_{\alpha=1}^{N(m)} (\lambda_1^{(\alpha)})^4 + n(n-1) \sum_{\alpha=1}^{N(m)} (\lambda_1^{(\alpha)})^2 (\lambda_2^{(\alpha)})^2. \quad (4.20)$$

Proceeding as before, we find after some manipulation the following relation

$$\sum_{\alpha=1}^{N(m)} (\lambda_1^{(\alpha)})^4 = \sum_{k=0}^{m-1} \binom{2n+k-3}{k} \binom{m-k+2}{3} [-8 + 12(m-k) + 6(m-k)^2]/5. \quad (4.21)$$

Evaluating the sums, we obtain the result

$$\sum_{\alpha=1}^{N(m)} (\lambda_1^{(\alpha)})^4 = 2 \binom{2n+m}{2n+1} (2n^2 + 6m^2 + 12mn - 7n - 3)/(2n+3)(n+1). \quad (4.22)$$

Similarly, we can derive the relation

$$\sum_{\alpha=1}^{N(m)} (\lambda_1^{(\alpha)})^2 (\lambda_2^{(\alpha)})^2 = \sum_{k=0}^{m-1} \sum_{l=0}^{m-k} (-m+k+2l)^2 \sum_{k'=0}^{k-1} \sum_{l'=0}^{k-k'} \binom{2n+k'-5}{k'} (-k+k'+2l')^2. \quad (4.23)$$

This can be simplified to the following expression

$$\sum_{\alpha=1}^{N(m)} (\lambda_1^{(\alpha)})^2 (\lambda_2^{(\alpha)})^2 = 2 \sum_{k=1}^{m-1} \binom{2n+k-2}{2n-1} \sum_{l=0}^{m-k} (-m+k+2l)^2. \quad (4.24)$$

Evaluating this double sum, we obtain the result

$$\sum_{\alpha=1}^{N(m)} (\lambda_1^{(\alpha)})^2 (\lambda_2^{(\alpha)})^2 = 4 \binom{2n+m+1}{2n+3}. \quad (4.25)$$

Substituting Eqs. (4.22) and (4.25) in Eq. (4.20), we finally get the following relation

$$I_m^{(4)} = 2n \binom{2n+m}{2n+1} (2mn^2 + m^2n + 10mn + 5m^2 - 6n - 2)/(2n+3)(n+1). \quad (4.26)$$

Again, it is easily verified that the above expression for $I_m^{(4)}$ gives correct results (upto an overall numerical factor). Comparing Eqs. (4.15) and (4.26) we also obtain the relation

$$I_m^{(4)} = I_m^{(2)} (2mn^2 + m^2n + 10mn + 5m^2 - 6n - 2)/(2n+3)(n+1). \quad (4.27)$$

V. SUMMARY

In this paper, we discussed a real basis for $sp(2n, \mathbb{R})$. In this basis, the Cartan basis elements are automatically diagonal for all irreps carried by homogeneous polynomials of phase-space variables. This fact facilitates the computation of various quantities characterizing these irreps. As an illustration, we calculated the second and fourth order indices of these irreps.

ACKNOWLEDGMENTS

One of us (GR) was supported by the United States Department of Energy under contract number DE-AC03-76SF00098. We would like to thank Dr. Alex Dragt for useful discussions.

REFERENCES

- [1] The reader should be warned that there are several conflicting notations in use for denoting the various symplectic algebras. The notation $\text{sp}(2n)$ is often used to denote the real symplectic algebra $\text{sp}(2n, \mathbb{R})$. Sometimes, $\text{sp}(n, \mathbb{R})$ is used to denote this algebra. In this paper, we have adopted the least ambiguous set of notations found in the literature. The real symplectic algebra is denoted by $\text{sp}(2n, \mathbb{R})$; the complex symplectic algebra is denoted by $\text{sp}(2n, \mathbb{C})$; the so-called unitary symplectic algebra is denoted by $\text{sp}(2n)$.
- [2] A. J. Dragt, in *Physics of High Energy Particle Accelerators*, edited by R. A. Carrigan, F. R. Huson, and M. Month, AIP Conference Proceedings No. 87 (American Institute of Physics, New York, 1982), p. 147; see also A. J. Dragt *et. al.*, *Annu. Rev. Nucl. Part. Sci.* **38**, 455 (1988).
- [3] G. Rangarajan, Ph. D. thesis, University of Maryland, 1990.
- [4] A. Giorgilli, *Comp. Phys. Comm.* **16**, 331 (1979).
- [5] J. Patera, R. T. Sharp, and P. Winternitz, *J. Math. Phys.* **17**, 1972 (1976); J. Patera, R. T. Sharp, and P. Winternitz, *J. Math. Phys.* **18**, 1519 (1977).
- [6] J. F. Cornwell, *Group Theory in Physics*, volumes 1 and 2 (Academic Press, London, 1984).
- [7] W. G. McKay and J. Patera, *Tables of Dimensions, Indices, and Branching Rules for Representations of Simple Lie Groups* (Marcel Dekker, New York, 1981).