UCLA

UCLA Previously Published Works

Title

Burkholderia caballeronis sp. nov., a nitrogen fixing species isolated from tomato (Lycopersicon esculentum) with the ability to effectively nodulate Phaseolus vulgaris.

Permalink

<https://escholarship.org/uc/item/39h5h24n>

Journal

Antonie van Leeuwenhoek, 104(6)

ISSN

0003-6072

Authors

Martínez-Aguilar, Lourdes
Salazar-Salazar, Corelly
Méndez, Rafael Díaz
et al.

Publication Date

2013-12-01

DOI

10.1007/s10482-013-0028-9

Peer reviewed

Phylogenetic Analysis of *Burkholderia* Species by Multilocus Sequence Analysis

Paulina Estrada-de los Santos · Pablo Vinuesa ·
Lourdes Martínez-Aguilar · Ann M. Hirsch ·
Jesús Caballero-Mellado

Received: 11 August 2012 / Accepted: 21 January 2013
© Springer Science+Business Media New York 2013

Abstract *Burkholderia* comprises more than 60 species of environmental, clinical, and agro-biotechnological relevance. Previous phylogenetic analyses of 16S rRNA, *recA*, *gyrB*, *rpoB*, and *acdS* gene sequences as well as genome sequence comparisons of different *Burkholderia* species have revealed two major species clusters. In this study, we undertook a multilocus sequence analysis of 77 type and reference strains of *Burkholderia* using *atpD*, *gltB*, *lepA*, and *recA* genes in combination with the 16S rRNA gene sequence and employed maximum likelihood and neighbor-joining criteria to test this further. The phylogenetic analysis revealed, with high supporting values, distinct lineages within the genus *Burkholderia*. The two large groups were named A and B, whereas the *B. rhizoxinica*/*B. endofungorum*, and *B. andropogonis* groups consisted of two and one species, respectively. The group A encompasses several plant-associated and saprophytic

bacterial species. The group B comprises the *B. cepacia* complex (opportunistic human pathogens), the *B. pseudomallei* subgroup, which includes both human and animal pathogens, and an assemblage of plant pathogenic species. The distinct lineages present in *Burkholderia* suggest that each group might represent a different genus. However, it will be necessary to analyze the full set of *Burkholderia* species and explore whether enough phenotypic features exist among the different clusters to propose that these groups should be considered separate genera.

Abbreviations

MLSA	Multilocus sequence analysis
BCC	<i>Burkholderia cepacia</i> complex
ML	Maximum likelihood
NJ	Neighbor-joining

Introduction

The genus *Burkholderia*, a β -proteobacteria group, was created to accommodate seven species from the *Pseudomonas* ribosomal RNA group II [51, 89]. Eventually, many more species were described and included in this new genus. Others were removed or reclassified, such as *Burkholderia pickettii* (now *Ralstonia pickettii*) and *B. solanacearum* (now *Ralstonia solanacearum*) [90], or *B. cocovenenans*, which was a synonym of *B. gladioli* [17, 41] and *B. vandii*, a synonym of *B. plantarii* [17]. Currently, *Burkholderia* comprises more than 60 species, which are distributed in diverse habitats. For example, several species are important components of the rhizosphere [22]. Others have been found in water, plant roots, or legume nodules, or can be opportunistic pathogens on plants or humans [22, 77]. Among the pathogenic

Electronic supplementary material The online version of this article (doi:10.1007/s00284-013-0330-9) contains supplementary material, which is available to authorized users.

P. Estrada-de los Santos (✉) · P. Vinuesa · L. Martínez-Aguilar ·
J. Caballero-Mellado
Centro de Ciencias Genómicas, Universidad Nacional Autónoma de
México, Ap. Postal 565-A, Cuernavaca, Morelos, Mexico
e-mail: pestradadelossantos@gmail.com

P. Estrada-de los Santos
Departamento de Microbiología, Escuela Nacional de Ciencias
Biológicas, I.P.N., Prolongación de Carpio y Plan de Ayala,
Mexico, D.F., Mexico

A. M. Hirsch
Department of Molecular, Cell and Developmental Biology,
Molecular Biology Institute, University of California,
Los Angeles, 621 Charles E. Young Drive South,
Los Angeles, CA 90095-1606, USA

Burkholderia, *B. mallei* and *B. pseudomallei* are extremely important because they are the etiologic agents of known diseases such as glanders and melioidosis, respectively [10, 87]. The *Burkholderia cepacia* complex (BCC; 17 species up to now) is another important group that includes many opportunistic pathogens, which may be found in patients with cystic fibrosis [38].

In 2001, two reports presented a totally different view of *Burkholderia*. First, the genus was rich in nitrogen-fixing species [28], and second, several nitrogen-fixing species were found to nodulate legume plants [48]. This last feature was striking because previously legumes were thought to be nodulated only by α -proteobacteria. Today, eight *Burkholderia* species that elicit effective nodule formation on legume roots have been reported and more are in the pipeline awaiting description [37, 46]. Recently, strains from *B. fungorum* were found to nodulate *Phaseolus vulgaris*, although ineffectively [30].

As the number of described *Burkholderia* species has increased, 16S rRNA sequence analyses have revealed two sub-lineages within the genus [8, 37, 53, 54, 58]. These two groups have been recovered with different phylogenetic reconstruction methods using 16S rRNA sequences [52], Multilocus sequence analysis (MLSA) with *gyrB* and *rpoB* genes sequences [2] or seven housekeeping genes [63], or *recA* gene sequences [73]. A phylogenetic analysis of the *acdS* gene in *Burkholderia* species also revealed the same two clusters [49]. One of these sub-lineages contains the BCC opportunistic human pathogens, some environmental species, the *B. pseudomallei* group, as well as plant pathogenic species, which have been also detected in human infections [43, 60, 86]. The other sub-lineage comprises soil, water, and plant-associated species, which thrive in rhizospheres, live as endophytes, or nodulate legumes. Many species are diazotrophs and so far, evidence for pathogenicity is lacking, although a few reports exist of single strains of *B. xenovorans*, *B. tropica*, and *B. fungorum* that were isolated from animal or human clinical samples [18, 24, 35]. Additionally, the phylogenetic analysis of 617 genes, from the *Burkholderia* core genome, demonstrated that the BCC–*B. pseudomallei*–plant pathogen group was clearly separated from the cluster of environmental species [69].

Not only MLSA but also Multilocus sequence typing (MLST) have been used for epidemiologic and population genetics studies, delineation of species, and assignment of strains to defined bacterial species in *Burkholderia* [2, 40, 80, 82]. Different housekeeping genes have been useful for this purpose and an MLST database exists for BCC members (<http://pubmlst.org/bcc/>).

In view of these findings, we performed MLSA of a set of *Burkholderia* species (77 *Burkholderia* type and reference species) using four housekeeping genes, *atpD*, *gltB*,

lepA, and *recA*, combined with the 16S rRNA gene sequence, to explore the positioning of the *Burkholderia* species in a phylogenetic analysis. In this report, we present evidence that the genus *Burkholderia* is composed of distinctly different phylogenetic lineages.

Materials and Methods

Bacterial Strains

A set of *Burkholderia* species was analyzed by MLSA. Several species with sequenced genomes were also chosen for the analysis. The set of *Burkholderia* species analyzed included a total of 77 *Burkholderia* species, 51 type species and 26 reference species. Additionally, type and reference strains from *Cupriavidus* (3) and *Ralstonia* (4) species were included. A list of strains used in this study is presented in Additional file 1.

atpD, *gltB*, *lepA*, and *recA* Gene Sequencing

The housekeeping genes were randomly selected based on the BCC MLST Database (<http://pubmlst.org/bcc/>). The following set of primers was designed to obtain DNA fragments in the range of 850–1,100 bp. These DNA fragments were more than twice as long as the ones used in MLST for the BCC because we wished to retrieve more information for the phylogenetic analysis. For the ATP synthase beta chain (*atpD*), the primers atpD-F2 (5'-CCACCAGCACAAGCCGCT-3') and atpD-R (5'-ATCCGCTCGTCGTCGGCG-3') were used. For the glutamate synthase large subunit (*gltB*), the primers gltB-F (5'-CTGCGCTCGAAGATCAAGCAGGG-3') and gltB-R (5'-TGCGCACCGGCTGGATGAACG-3') were utilized. For the GTP binding protein (*lepA*), the primers lepA-F2 (5'-TGGTTGCACAACACTACGTCGG-3') and lepA-R (5'-ATCAGCATGTGACCTTCAC-3') were employed, and for recombinase A (*recA*), BUR1 and BUR2 primers were used [52]. The PCR was performed as described previously [28]. The DNA fragments on each strand were sequenced with the primers used in the initial PCR amplification by Macrogen (www.macrogen.com). The housekeeping genes from some *Burkholderia* species were retrieved from the NCBI Genome Database (<http://www.ncbi.nlm.nih.gov/sites/entrez?db=genome>). The accession numbers are provided in Additional file 1 and displayed on the phylogenetic trees.

Phylogenetic Inference

Data processing and phylogenetic analyses were performed as described elsewhere [77]. In brief, nucleotide sequences

of the housekeeping genes were translated and aligned using Muscle 3.57 [25]. The resulting multiple-sequence alignments of proteins were used as masks to generate the corresponding nucleotide codon alignments using custom Perl scripts. Individual alignments were concatenated using *ad hoc* Perl scripts. Models of nucleotide substitution were selected by the Akaike information criterion, using MODELTEST3.7 [56, 57]. Among-site rate variation was modeled by a gamma distribution, approximated with four rate categories [91], with each category being represented by its mean. ML trees were inferred for each dataset under the models of nucleotide substitution selected by the Akaike information criterion [57], using PhyML v3.0 [36]. Tree searches were initiated from a BioNJ seed tree, retaining the best tree among those found with NNI and TBR branch swapping. The robustness of the ML topologies was evaluated using a recently developed Shimodaira–Hasegawa (SH)-like test [6] for branches implemented in PhyML v3.0. In brief, the test assesses whether the branch being studied provides a significant likelihood gain, in comparison with the null hypothesis that involves collapsing that particular branch, but leaving the rest of the tree topology the same. The SH-like procedure was chosen for assessing bipartition significance because the test is nonparametric and much less liberal than the diverse (parametric) approximate-likelihood ratio tests that are also implemented in that program. The resulting SH-like *P* values, therefore, indicate the probability that the corresponding split is significant. The ML phylogenetic trees were visualized with the program MEGA version 5 [66]. The alignment dataset was also used for NJ analysis. The NJ trees were constructed based on Tamura–Nei distances, using 1,000 bootstrap replicates with the program MEGA version 5.

Diazotrophy Test

The capacity to fix nitrogen by most *Burkholderia* species was tested by acetylene reduction activity [28], by *nifH* gene amplification [53, 55, 70], or by collecting information from the literature.

Results and Discussion

Phylogenetic Inference of 16S rRNA Gene Sequences

All *Burkholderia* species were included in the analysis of the 16S rRNA gene sequence by ML (Fig. 1). The phylogenetic analyses resulted in two large clusters; one (designated as group A) comprised the plant-associated and saprophytic species. The second cluster (group B) contained the BCC opportunistic human pathogens, the

B. pseudomallei group, and the plant pathogenic species. *Burkholderia andropogonis*, *B. rhizoxinica*, and *B. endofungorum* were placed at the edge of group A. A distance analysis (using MEGA version 5) of the 16S rRNA sequences from group A and B, *B. rhizoxinica*/*B. endofungorum* and *B. andropogonis* was carried out. The analysis showed a larger dissimilarity of 4 % between group A and B. The intra-group 16S rRNA similarity values were 98.7 % among the species of the group B and 96.0 % for the group A. Also, the 16S rRNA similarity value of *B. andropogonis* with the different groups was <96: 95.5 % with group B and 93.8 % with group A. Similarly, the *B. endofungorum*/*B. rhizoxinica* group was 96.1 % identical to the group B and 95.1 % to group A. The *B. endofungorum*/*B. rhizoxinica* group was found to be 95.1 % similar to *B. andropogonis*. This evidence clearly indicates that the genus *Burkholderia* consists of different bacterial lineages. Generally, values below ~95 % 16S rRNA gene sequence similarity indicate different genera [67].

Furthermore, the incongruent position of *B. andropogonis* was previously reported by Viallard et al. [84] based on analysis of 16S rRNA gene. To investigate this species in greater detail, fifteen 16S rRNA gene sequences from the NCBI database belonging to different strains of *B. andropogonis* were included in an NJ tree with the rest of the *Burkholderia* species (data not shown). The analysis showed that the *B. andropogonis* strains clustered outside of groups A, B and *B. endofungorum*/*B. rhizoxinica* groups, and distantly from *Cupriavidus*/*Ralstonia* species. The phylogenetic position of this species and the 16S similarity percentages among *B. andropogonis* and the other groups suggest that this species might represent a new genus. This may also be the case for *B. endofungorum* and *B. rhizoxinica*, but these species were not analyzed further because both species were described with only single strains.

The 16S sequence genes were also analyzed by NJ. The analysis showed that the genus split in different lineages, interestingly group A was further divided in two groups and *B. andropogonis* was placed outside from all clusters (Additional file 2).

Phylogenetic Inference of Individual and Concatenated Housekeeping Genes

The set of *Burkholderia* species analyzed in this study was selected from our in-house *Burkholderia* collection and some of the other species were gathered from the LMG culture collection (BCCM/LMG, Belgium). Whenever possible, more than one strain from each *Burkholderia* species was selected to perform robust species phylogeny estimation by MLSA. A phylogenetic analysis of the individual *atpD*, *gltB*, *lepA*, and *recA* sequences under the ML criterion revealed two distinct lineages within the

Fig. 1 Maximum likelihood tree inferred from the 16S rRNA gene, showing the phylogenetic relationships among *Burkholderia* species. The bar represents the number of expected substitutions per site under the GTR + G model. One asterisk indicates sequences obtained from the Genome database at NCBI. Two asterisks indicate sequences obtained from an ongoing genomic project. In parenthesis are accession numbers at NCBI

genus *Burkholderia*, the same groups A and B found in the analysis of 16S rRNA gene. However, in the *gltB* and *lepA* phylogenetic trees, group A was further split in two distinct lineages (Additional file 3–6). The dataset was also analyzed under the NJ criterion (Additional files 7–10), again showing two main groups in the *atpD* and *lepA* trees and three clusters in the *gltB* and *recA* trees. Moreover, the position of certain *Burkholderia* species was not clear. Generally, *B. andropogonis*, *B. soil*, and *B. rhizoxinica* were found to be not related to the two main groups.

A highly resolved ML species tree was estimated from the five gene concatenated dataset (Fig. 2). Only those strains having the full set of sequenced genes were included in this analysis. The results showed the two clusters, which are designated as group A and B. However, *B. soli*, *B. rhizoxinica*, *B. kururiensis*, and *B. glathei* were positioned outside these two main clusters, which indicates that more strains from these species must be analyzed to define their actual position in the phylogenetic trees. *Burkholderia andropogonis* was not included in the concatenated tree because the *recA* gene sequence was not successfully amplified and hence not examined. The concatenated dataset analyzed by NJ showed three groups with group A further split in two clusters (Additional file 11).

GC Content Analysis

Previously, Gyaneshwar et al. [37] reported that some of the species included in the group A have a lower GC content than the species in the pathogenic group B. We expanded upon this analysis, and surveyed the literature and various databases to learn the GC content of all the currently described *Burkholderia* species (Additional file 12, 13). The analysis of the GC content of the different groups showed a substantial divergence between the group B (67.1 ± 1.0 %) compared to the group A (62.9 ± 1.3 %), *B. andropogonis* (59.0 %), and the *B. rhizoxinica/B. endofungorum* (60.7 %) cluster. Evidently, the group B GC content is higher than any of the other *Burkholderia* groups. Tindall et al. [67] observed that, with some exceptions, GC content may be fairly constant in a bacterial group. Taken together, this observation strongly suggests that the groups observed in *Burkholderia* consist of distinct bacterial lineages.

Diazotrophy Test

The ability to fix nitrogen is well known in *B. vietnamiensis* [31], the only member of the BCC within the group B that has unequivocally been shown to perform this activity. Starting in 2001 and continuing, the genus *Burkholderia*, especially members of the group A, is rich in nitrogen-fixing species [8, 28, 48, 53, 54, 58, 74]. Previously, *B. ferrariae*, a member of the group A, was found to have a

nifH gene (GenBank:EF158799) [47] and the capacity of this species to fix nitrogen was confirmed by acetylene reduction activity (ARA) in the present study (Additional file 1). However, Aizawa et al. [5] did not detect ARA activity either in *B. ferrariae* or in *B. acidipaludis* or *B. bannensis*, although a *nifH* sequence was detected by PCR analysis. Thus it is still unknown whether the *nif* operon is complete or whether these strains actually fix nitrogen. One possibility to explain the difference in the reports about *B. ferrariae* diazotrophy is that the growth medium used to test ARA by Aizawa et al. [5] was not optimal. We found that using different medium [28] we could detect that *B. ferrariae* is a diazotroph. A similar situation could exist for both *B. acidipaludis* and *B. bannensis*. Other *Burkholderia* species not tested in this study by MLSA are also diazotrophs; such as *B. heleaia*, *B. symbiotica*, and *B. diazotrophica*, the latter two species are able to nodulate *Mimosa* spp. plants [3, 61, 62]. However, *B. oxyphila*, *B. rhizoxinica*, and *B. endofungorum* have not been reported to be diazotrophs. *Burkholderia andropogonis* was found to be ARA-minus in this study. In conclusion, approximately half of the species from the group A are able to fix nitrogen as either free-living bacteria or in symbiotic associations.

Conclusions

The results presented in this report show that the genus *Burkholderia* is composed of different lineages. A glimpse of this observation was showed by Ussery et al. [69] where 56 *Burkholderia* genomes were analyzed in a phylogenetic study. Vandamme and Dawyndt [79] came to similar conclusions using ML to examine 100,000 base positions randomly extracted from single-copy core genes of the available *Burkholderia* genomes released up to date.

As new *Burkholderia* species are continually being described, the presence of different lineages will become more obvious. However, to propose a split of the *Burkholderia* genus at this time may be premature. The presence of two and sometimes three main groups plus *B. andropogonis* and *B. endofungorum/B. rhizoxinica* shows that the genus is evolving over time such that different bacterial lineages are developing. To understand this process better, the entire assemblage of species must be thoroughly analyzed, which means that both phylogenetic and physiological/biochemical traits need to be examined as well.

One example is the pathogenicity shown by the species of the group B compared to the species of the group A. Preliminary experiments found no pathogenic activity toward *Caenorhabditis elegans* or HeLa cells after testing *B. unamae*, *B. phytofirmans*, *B. tuberum*, and two strains of

Fig. 2 Maximum likelihood species tree inferred from the concatenated alignment *atpD*–*gltB*–*lepA*–*recA*–16S rRNA genes, showing the phylogenetic relationships among *Burkholderia* species. The *bar* represents the number of expected substitutions per site under the GTR + G model

B. silvatlantica (A. Angus and A.M. Hirsch, ms. in prep.). In any case, differential phenotypic characteristics may not be as relevant at the genus level as they are for the description of new species because phenotypic traits may be inconsistent when large populations are studied [85]. Nevertheless, it is critical to analyze the full set of *Burkholderia* species, including more strains from each species as well as including extra housekeeping genes in the analysis.

Acknowledgments This work is dedicated to the memory of Dr. Jesus Caballero Mellado (1953–2010), for his many years of fruitful work, support, generosity, and friendship. We are grateful to Jorge Eduardo Buendia Buendia, Isaac Fernando Lopez Moyado, Mariana del Rosario Ruiz Velasco Leyva, Jorge Arturo Zepeda Martinez, and Marie Lisandra Zepeda Mendoza for technical support during training as students from the Undergraduate Program on Genomic Sciences (Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México). We also thank Annette A. Angus (University of California, Los Angeles) for sharing her unpublished data with us. The housekeeping gene sequences were provided from the ongoing sequencing project of *B. unamae* MTI-641^T, *B. tuberum* STM678^T, *B. silvatlantica* SRMrh-20^T, and *B. silvatlantica* PVA5 from a project, funded in part by the U.S. National Science Foundation (Grant IOB-0537497) to George Weinstock (Washington University, St. Louis, MO) and AMH.

References

- Achouak W, Christen R, Barakat M, Martel MH, Heulin T (1999) *Burkholderia caribensis* sp. nov., an exopolysaccharide producing bacterium isolated from vertisol microaggregates in Martinique. *Int J Syst Bacteriol* 49:787–794
- Ait-Tayeb L, Lefevre M, Passet V, Diancourt L, Brisse S, Grimont PAD (2008) Comparative phylogenies of *Burkholderia*, *Ralstonia*, *Comamonas*, *Brevundimonas* and related organism derived from *rpoB*, *gyrB* and *rrs* gene sequences. *Res Microbiol* 159:169–177
- Aizawa T, Bao Ve N, Nakajima M, Sunairi M (2010) *Burkholderia heleia* sp. nov., a nitrogen-fixing bacterium isolated from an aquatic plant, *Eleocharis dulcis*, that grows in highly acidic swamps in actual acid sulfate soil areas of Vietnam. *Int J Syst Evol Microbiol* 60:1152–1157
- Aizawa T, Nguyen BV, Vijarnsorn P, Nakajima M, Sunairi M (2010) *Burkholderia acidipaludis* sp. nov., aluminum-tolerant bacteria isolated from the Chinese water chestnut, *Eleocharis dulcis*, that grows in highly acidic swamps in Southeast Asia. *Int J Syst Evol Microbiol* 60:2036–2041
- Aizawa T, Vijarnsorn P, Nakajima M, Sunairi M (2011) *Burkholderia bannensis* sp. nov., and acidic pH-neutralizing bacterium isolated from torpedo grass (*Panicum repens*) that grows in highly acidic swamps in Thailand. *Int J Syst Evol Microbiol* 61:1645–1650
- Anisimova M, Gascuel O (2006) Approximate likelihood-ratio test for branches: a fast, accurate, and powerful alternative. *Syst Biol* 55:539–552
- Bramer CO, Vandamme P, da Silva LF, Gomez JGC, Steinbuchel A (2001) *Burkholderia sacchari* sp. nov., a polyhydroxyalkanoate-accumulating bacterium isolated from soil of a sugar-cane plantation in Brazil. *Int J Syst Evol Microbiol* 51:1709–1713
- Caballero-Mellado J, Martinez-Aguilar L, Paredes-Valdez G, Estrada-de los Santos P (2004) *Burkholderia unamae* sp. nov., a N₂-fixing rhizospheric and endophytic species. *Int J Syst Evol Microbiol* 54:1165–1172
- Caballero-Mellado J, Onofre-Lemus J, Estrada-de los Santos P, Martinez-Aguilar L (2007) The tomato rhizosphere, an environment rich in nitrogen-fixing *Burkholderia* species with capabilities of interest for agriculture and bioremediation. *Appl Environ Microbiol* 73:5308–5319
- Chen WM, Laevens S, Lee TM, Coenye T, De Vos P, Mergeay M, Vandamme P (2001) *Ralstonia taiwanensis* sp. nov., isolated from root nodules of *Mimosa* species and sputum of cystic fibrosis patient. *Int J Syst Evol Microbiol* 51:1729–1735
- Chen WM, Moulin L, Bontemps C, Vandamme P, Bena G, Boivin-Masson C (2003) Legume symbiotic nitrogen fixation by β -proteobacteria is widespread in nature. *J Bacteriol* 185:7266–7272
- Chen WM, de Faria SM, Stralioetto R, Pitard RM, Simoes-Araujo JL, Chou JH, Chou YJ, Barrios E, Prescott AR, Elliot GN, Sprent JI, Young JPW, James EK (2005) Proof that *Burkholderia* strains from effective symbioses with legumes: a study of novel *Mimosa*-nodulating strains from South America. *Appl Environ Microbiol* 71:7461–7471
- Chen WM, James EK, Chou JH, Sheu SY, Yang SZ, Sprent JI (2005) β -rhizobia from *Mimosa pigra*, a newly discovered invasive plant in Taiwan. *New Phytol* 168:661–675
- Chen WM, James EK, Coenye T, Chou JH, Barrios E, de Faria SM, Elliott GN, Sheu SY, Sprent JI, Vandamme P (2006) *Burkholderia mimosarum* sp. nov., isolated from root nodules of *Mimosa* spp. from Taiwan and South America. *Int J Syst Evol Microbiol* 56:1847–1851
- Chen WM, de Faria SM, James EK, Elliott GN, Lin KY, Chou JH, Sheu SY, Cnockaert M, Sprent JI, Vandamme P (2007) *Burkholderia nodosa* sp. nov., isolated from root nodules of the woody Brazilian legumes *Mimosa bimucronata* and *Mimosa scabrella*. *Int J Syst Evol Microbiol* 57:1055–1059
- Chen WM, de Faria SM, Chou JH, James EK, Elliott GN, Sprent JI, Bontemps C, Young JPW, Vandamme P (2008) *Burkholderia sabiae* sp. nov., isolated from root nodules of *Mimosa caesalpinifolia*. *Int J Syst Evol Microbiol* 58:2174–2179
- Coenye T, Holmes B, Kersters K, Govan JRW, Vandamme P (1999) *Burkholderia cocovenenans* (van Damme et al. 1960) Gillis et al. 1995 and *Burkholderia vandii* Urakami et al. 1994 are junior synonyms of *Burkholderia gladioli* (Severini 1913) Yabuuchi et al. 1993 and *Burkholderia plantarii* (Azegami et al. 1987) Urakami et al. 1994, respectively. *Int J Syst Bacteriol* 49:37–42
- Coenye T, Laevens S, Willems A, Ohlen M, Hannant W, Govan JRW, Gillis M, Falsen E, Vandamme P (2001) *Burkholderia fungorum* sp. nov. and *Burkholderia caledonica* sp. nov., two new species isolated from the environment, animals and human clinical samples. *Int J Syst Evol Microbiol* 51:1099–1107
- Coenye T, Mahenthiralingam E, Henry D, LiPuma JJ, Laevens S, Gillis M, Speert DP, Vandamme P (2001) *Burkholderia ambifaria* sp. nov., a novel member of the *Burkholderia cepacia* complex including biocontrol and cystic fibrosis-related isolates. *Int J Syst Evol Microbiol* 51:1481–1490
- Coenye T, Goris J, De Vos P, Vandamme P, LiPuma JJ (2003) Classification of *Ralstonia pickettii*-like isolates from the environment and clinical samples as *Ralstonia insidiosa* sp. nov. *Int J Syst Evol Microbiol* 53:1075–1080
- Coenye T, Henry D, Speert DP, Vandamme P (2004) *Burkholderia phenoliruptrix* sp. nov., to accommodate the 2,4,5-trichlorophenoxyacetic acid and halophenol-degrading strain AC1100. *Syst Appl Microbiol* 27:623–627
- Compant S, Nowak J, Coenye T, Clement C, Barka EA (2008) Diversity and occurrence of *Burkholderia* spp. in the natural environment. *FEMS Microbiol Rev* 32:607–626

23. De Baere T, Steyaert S, Wauters G, De Vos P, Goris J, Coenye T, Suyama T, Verschraegen G, Vaneechoutte M (2001) Classification of *Ralstonia pickettii* biovar 3/‘thomasi’ strains (Pickett 1994) and of new isolates related to nosocomial recurrent meningitis as *Ralstonia mannitolytica* sp. nov. *Int J Syst Evol Microbiol* 51:547–558
24. Deris ZZ, Van Rostenberghe H, Habsah H, Noraida R, Tan GC, Chan YY, Rosliza RN, Ravichandran M (2010) First isolation of *Burkholderia tropica* from neonatal patient successfully treated with imipenem. *Int J Infect Dis* 14:e73–e74
25. Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797
26. Elliot GN, Chen WM, Chou JH, Wang HC, Sheu SY, Perin L, Reis VM, Moulin L, Simon MF, Bontemps C, Sutherland JM, Bessi R, de Faria SM, Trinick MJ, Prescott AR, Sprent JJ, James EK (2007) *Burkholderia phymatum* is a highly effective nitrogen-fixing symbiont of *Mimosa* spp. and fixes nitrogen ex planta. *New Phytol* 173:168–180
27. Elliot GN, Chen WM, Bontemps C, Chou JH, Young JP, Sprent JJ, James EK (2007) Nodulation of *Cyclopia* spp. (Leguminosae, Papilionoidae) by *Burkholderia tuberum*. *Ann Bot* 100:1403–1411
28. Estrada-de los Santos P, Bustillos-Cristales R, Caballero-Mellado J (2001) *Burkholderia*, a genus rich in plant-associated nitrogen fixers with wide environmental and geographic distribution. *Appl Environ Microbiol* 67:2790–2798
29. Estrada-de los Santos P, Martínez-Aguilar L, López-Lara IM, Caballero-Mellado J (2012) *Cupriavidus alkaliphilus* sp. nov., a new species associated with agricultural plants that grow in alkaline soils. *Syst Appl Microbiol* 35:310–314
30. Ferreira PAA, Bomfeti CA, Soares BL, Moreira FMS (2012) Efficient nitrogen-fixing *Rhizobium* strains isolated from Amazonian soils are highly tolerant to acidity and aluminum. *World J Microbiol Biotechnol* 28:1947–1959
31. Gillis M, Van Van T, Bardin R, Goor M, Hebbard P, Willems A, Segers P, Kersters K, Heulin T, Fernandez MP (1995) Polyphasic taxonomy in the genus *Burkholderia* leading to an emended description of the genus and proposition of *Burkholderia vietnamiensis* sp. nov. for N₂-fixing isolates from rice in Vietnam. *Int J Syst Bacteriol* 45:274–289
32. Glass MB, Steigerwalt AG, Jordan JG, Wilkins PP, Gee JE (2006) *Burkholderia oklahomensis* sp. nov., a *Burkholderia pseudomallei*-like species formerly known as the Oklahoma strain of *Pseudomonas pseudomallei*. *Int J Syst Evol Microbiol* 56:2171–2176
33. Goris J, De Vos P, Coenye T, Hoste B, Janssens D, Brim H, Diels L, Mergeay M, Kersters K, Vandamme P (2001) Classification of metal-resistant bacteria from industrial biotopes as *Ralstonia campinensis* sp. nov., *Ralstonia metallidurans* sp. nov. and *Ralstonia basilensis* Steinle et al. 1998 emend. *Int J Syst Evol Microbiol* 51:1773–1782
34. Goris J, Dejonghe W, Falsen E, De Clerck E, Geeraerts B, Willems A, Top EM, Vandamme P, De Vos P (2002) Diversity of transconjugants that acquired plasmid pJP4 or pEMT1 after inoculation of a donor strain in the A- and B-horizon of an agricultural soil and description of *Burkholderia hospita* sp. nov. and *Burkholderia terricola* sp. nov. *Syst Appl Microbiol* 25:340–352
35. Goris J, De Vos P, Caballero-Mellado J, Park JH, Falsen E, Quensen JF III, Tiedje JM, Vandamme P (2004) Classification of the PCB- and biphenyl-degrading strain LB400 and relatives as *Burkholderia xenovorans* sp. nov. *Int J Syst Evol Microbiol* 54:1677–1681
36. Guindon S, Delsuc F, Dufayard JF, Gascuel O (2009) Estimating maximum likelihood phylogenies with PhyML. *Methods Mol Biol* 537:113–137
37. Gyaneshwar P, Hirsch AM, Chen WM, Elliott GN, Bontemps C, Gross E, dos Reis Junior FB, Sprent JJ, Young JPW, James EK (2011) Legume nodulating β -proteobacteria: diversity, host range and future prospects. *Mol Plant Microbe Interact* 24:1276–1288
38. Hauser AR, Jain M, Bar-Meir M, McColley SA (2011) Clinical significance of microbial infection and adaptation in cystic fibrosis. *Clin Microbiol Rev* 24:29–70
39. Hee-Chan Y, Wan-Taek I, Kwang KK, Dong-Shan A, Sung-Taik L (2006) *Burkholderia terrae* sp. nov., isolated from a forest soil. *Int J Syst Evol Microbiol* 56:453–457
40. Ho-Bin K, Min-Ju P, Hee-Chan Y, Dong-Shan A, Hai-Zhu J, Deok-Chun Y (2006) *Burkholderia ginsengisoli* sp. nov. a β -glucosidase-producing bacterium isolated from soil of a ginseng field. *Int J Syst Evol Microbiol* 56:2529–2533
41. Jiao Z, Kawamura Y, Mishima N, Yang R, Li N, Liu X, Ezaki T (2003) Need to differentiate lethal toxin-producing strains of *Burkholderia gladioli*, which cause severe food poisoning: description of *B. gladioli* pathovar *cocovenenans* and emended description of *B. gladioli*. *Microbiol Immunol* 47:915–925
42. Lackner G, Moebius N, Partida-Martinez L, Hertweck C (2011) Complete genome sequence of *Burkholderia rhizoxinica*, an endosymbiont of *Rhizopus microsporus*. *J Bacteriol* 193:783–784
43. Lestin F, Kraak R, Podbielski A (2008) Two cases of keratitis and corneal ulcers caused by *Burkholderia gladioli*. *J Clin Microbiol* 46:2445–2449
44. Lim YW, Baik KS, Han SK, Kim SB, Bae KS (2003) *Burkholderia sordidicola* sp. nov., isolated from the white-rot fungus *Phanerochaete sordida*. *Int J Syst Evol Microbiol* 53:1631–1636
45. Lim JH, Baek SH, Lee ST (2008) *Burkholderia sediminicola* sp. nov., isolated from freshwater sediment. *Int J Syst Evol Microbiol* 58:565–569
46. Lu P, Zheng LQ, Sun JJ, Liu HM, Li SP, Li WJ, Hong Q (2012) *Burkholderia zhejiangensis* sp. nov., a novel methyl parathion-degrading bacterium isolated from a wastewater-treating system. *Int J Syst Evol Microbiol* 62:1337–1341
47. Martínez-Aguilar L, Díaz R, Peña-Cabriaes JJ, Estrada-de los Santos P, Dunn MF, Caballero-Mellado J (2008) Multichromosomal genome structure and confirmation of diazotrophy in novel plant-associated *Burkholderia* species. *Appl Environ Microbiol* 74:4574–4579
48. Moulin L, Munive A, Dreyfus B, Boivin-Masson C (2001) Nodulation of legumes by members of the beta-subclass of Proteobacteria. *Nature* 411:948–950
49. Onofre-Lemus J, Hernandez-Lucas I, Girard L, Caballero-Mellado J (2009) ACC (1-aminocyclopropane-1-carboxylate) deaminase activity, a widespread trait in *Burkholderia* species, and its growth-promoting effect on tomato plants. *Appl Environ Microbiol* 75:6581–6590
50. Otsuka Y, Muramatsu Y, Nakagawa Y, Matsuda M, Nakamura M, Murata H (2010) *Burkholderia oxyphila* sp. nov., isolated from acidic forest soil that catabolizes (+)-catechin and its putative aromatic derivatives. *Int J Syst Evol Microbiol* 61:249–254
51. Palleroni NJ, Kunisawa R, Contopoulou R, Doudoroff M (1973) Nucleic acid homologies in the genus *Pseudomonas*. *Int J Syst Bacteriol* 23:333–339
52. Payne GW, Vandamme P, Morgan SH, LiPuma JJ, Coenye T, Weightman AJ, Jones TH, Mahenthalingam E (2005) Development of a *recA* gene-based identification approach for the entire *Burkholderia* genus. *Appl Environ Microbiol* 71:3917–3927
53. Perin L, Martínez-Aguilar L, Castro-Gonzalez R, Estrada-de los Santos P, Cabellos-Avelar T, Guedes HV, Reis VM, Caballero-Mellado J (2006) Diazotrophic *Burkholderia* species associated with field-grown maize and sugarcane. *Int J Syst Evol Microbiol* 72:3103–3110
54. Perin L, Martínez-Aguilar L, Paredes-Valdez G, Baldani JI, Estrada-de los Santos P, Reis VM, Caballero-Mellado J (2006) *Burkholderia silvatlantica* sp. nov., a diazotrophic bacterium associated with sugarcane and maize. *Int J Syst Evol Microbiol* 56:1931–1937

55. Poly F, Monrozier LJ, Bally R (2001) Improvement in the RFLP procedure for studying the diversity of *nifH* genes in communities of nitrogen fixers in soil. *Res Microbiol* 152:95–103
56. Posada D, Crandall KA (1998) MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14:817–818
57. Posada D, Buckley TR (2004) Model selection and model averaging in phylogenetics: advantages of Akaike information criterion and bayesian approaches over likelihood ratio tests. *Syst Biol* 53:793–808
58. Reis VM, Estrada-de los Santos P, Tenorio-Salgado S, Vogel J, Stoffels M, Guyon S, Mavingui P, Baldani VLD, Schmid M, Baldani JJ, Balandreau J, Hartmann A, Caballero-Mellado J (2004) *Burkholderia tropica* sp. nov., a novel nitrogen-fixing, plant-associated bacterium. *Int J Syst Evol Microbiol* 54:2155–2162
59. Sato Y, Nishihara H, Yoshida M, Watanabe M, Rondal JD, Concepcion RN, Ohta H (2006) *Cupriavidus pinatubonensis* sp. nov. and *Cupriavidus laharis* sp. nov., novel hydrogen oxidizing, facultatively chemolithotrophic bacteria isolated from volcanic mudflow deposits from Mt. Pinatubo in the Philippines. *Int J Syst Evol Microbiol* 56:973–978
60. Segonds C, Clavel-Batut P, Thouverez M, Grenet D, Le Coustumier L, Plesiat P, Chabanon G (2009) Microbiological and epidemiological features of clinical respiratory isolates of *Burkholderia gladioli*. *J Clin Microbiol* 47:1510–1516
61. Sheu SY, Chou JH, Bontemps C, Elliott GN, Gross E, James EK, Sprent JJ, Young JPW, Chen WM (2012) *Burkholderia symbiotica* sp. nov., isolated from root nodules of *Mimosa* spp. native to north east Brazil. *Int J Syst Evol Microbiol* 62:2272–2278
62. Sheu SY, Chou JH, Bontemps C, Elliott GN, Gross E, dos Reis FB Jr., Melkonian R, Moulin L, James EK, Sprent JJ, Young JPW, Chen WM (2012) *Burkholderia diazotrophica* sp. nov., isolated from root nodules of *Mimosa* spp. *Int J Syst Evol Microbiol*. doi:10.1099/ijs.0.039859-0
63. Spilker T, Baldwin A, Bumford A, Dowson CG, Mahenthalingam E, LiPuma J (2009) Expanded multilocus sequence typing for *Burkholderia* species. *J Clin Microbiol* 47:2607–2610
64. Storms V, Van Den Vracken N, Coenye T, Mahenthalingam E, LiPuma JJ, Gillis M, Vandamme P (2004) Polyphasic characterisation of *Burkholderia cepacia*-like isolates leading to the emended description of *Burkholderia pyrocinia*. *Syst Appl Microbiol* 27:517–526
65. Talbi C, Delgado MJ, Girard L, Ramírez-Trujillo A, Caballero-Mellado J, Bedmar EJ (2008) *Burkholderia phymatum* strains capable of nodulating *Phaseolus vulgaris* are present in Moroccan soils. *Appl Environ Microbiol* 76:4587–4591
66. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distances, and maximum parsimony methods. *Mol Biol Evol* 28:2731–2739
67. Tindall BJ, Rossello-Mora R, Busse HJ, Ludwig W, Kampfer P (2010) Notes on the characterization of prokaryote strains for taxonomic purposes. *Int J Syst Evol Microbiol* 60:249–266
68. Urakami T, Ito-Yoshida C, Araki H, Kijima T, Suzuki KI, Komagata K (1994) Transfer of *Pseudomonas plantarii* and *Pseudomonas glumae* to *Burkholderia* as *Burkholderia* spp. and description of *Burkholderia vandii* sp. nov. *Int J Syst Bacteriol* 44:235–245
69. Ussery DW, Kiil K, Lagesen K, Sicheritz-Ponten T, Bohlin J, Wassenaar TM (2009) The genus *Burkholderia*: analysis of 56 genomic sequences. *Genome Dyn* 6:140–157
70. Valdes M, Perez NO, Estrada-de los Santos P, Caballero-Mellado J, Peña-Cabriaes JJ, Normand P, Hirsch AM (2005) Non-Frankia actinomycetes isolated from surface-sterilized roots of *Casuarina equisetifolia* fix nitrogen. *Appl Environ Microbiol* 71:460–466
71. Valverde A, Delvasto P, Peix A, Velazquez E, Santa-Regina I, Ballester A, Rodriguez-Barrueco C, Garcia-Balboa C, Igual JM (2006) *Burkholderia ferrariae* sp. nov., isolated from an iron ore in Brazil. *Int J Syst Evol Microbiol* 56:2421–2425
72. Vandamme P, Holmes B, Vancannet M, Coenye T, Hoste B, Coopman R, Revets H, Lauwers S, Gillis M, Kersters K, Govan JRW (1997) Occurrence of multiple genomovars of *Burkholderia cepacia* in cystic fibrosis patients and proposal of *Burkholderia multivorans* sp. nov. *Int J Syst Bacteriol* 47:1188–1200
73. Vandamme P, Mahenthalingam E, Holmes B, Coenye T, Hoste B, De Vos P, Henry D, Speert DP (2000) Identification and population structure of *Burkholderia stabilis* sp. nov. (formerly *Burkholderia cepacia* genomovar IV). *J Clin Microbiol* 38:1042–1047
74. Vandamme P, Goris J, Chen WM, de Vos P, Willems A (2002) *Burkholderia tuberum* sp. nov. and *Burkholderia phymatum* sp. nov. nodulate the roots of tropical legumes. *Syst Appl Microbiol* 25:507–512
75. Vandamme P, Henry D, Coenye T, Nzula S, Vancannet M, LiPuma JJ, Speert DP, Govan JRW, Mahenthalingam E (2002) *Burkholderia anthina* sp. nov. and *Burkholderia pyrocinia*, two additional *Burkholderia cepacia* complex bacteria, may confound results of new molecular diagnostics tools. *FEMS Immunol Med Microbiol* 33:143–149
76. Vandamme P, Holmes B, Coenye T, Goris J, Mahenthalingam E, LiPuma J, Govan JRW (2003) *Burkholderia cenocepacia* sp. nov., a new twist to an old story. *Res Microbiol* 154:91–96
77. Vandamme P, Govan J, LiPuma J (2007) Diversity and role of *Burkholderia* spp. In: Coenye T, Vandamme P (eds) *Burkholderia* molecular microbiology and genomics. Horizon Bioscience, Norfolk, pp 1–28
78. Vandamme P, Opelt K, Knochel N, Berg C, Schonmann S, De Brandt E, Eberl L, Falsen E, Berg G (2007) *Burkholderia bryophila* sp. nov. and *Burkholderia megapolitana* sp. nov., moss associated species with antifungal and plant-growth-promoting properties. *Int J Syst Evol Microbiol* 57:2228–2235
79. Vandamme P, Dawyndt P (2011) Classification and identification of the *Burkholderia cepacia* complex: past, present and future. *Syst Appl Microbiol* 34:87–95
80. Vanlaere E, LiPuma JJ, Baldwin A, Henry D, De Brandt E, Speert D, Mahenthalingam E, Dowson C, Vandamme P (2008) *Burkholderia latens* sp. nov., *Burkholderia diffusa* sp. nov., *Burkholderia arboris* sp. nov., *Burkholderia seminalis* sp. nov. and *Burkholderia metallica* sp. nov., novel species within the *Burkholderia cepacia* complex. *Int J Syst Evol Microbiol* 58:1580–1590
81. Vanlaere E, van der Meer JR, Falsen E, Salles JF, de Brandt E, Vandamme P (2008) *Burkholderia sartisoli* sp. nov., isolated from a polycyclic aromatic hydrocarbon-contaminated soil. *Int J Syst Evol Microbiol* 58:420–423
82. Vanlaere E, Baldwin A, Gevers D, Henry D, De Brandt E, LiPuma JJ, Mahenthalingam E, Speert DP, Dowson C, Vandamme P, Taxon K (2009) a complex within the *Burkholderia cepacia* complex, comprises at least two novel species, *Burkholderia contaminans* sp. nov. and *Burkholderia lata* sp. nov. *Int J Syst Evol Microbiol* 59:102–111
83. Vermis K, Coenye T, LiPuma JJ, Mahenthalingam E, Nelis HJ, Vandamme P (2004) Proposal to accommodate *Burkholderia cepacia* genomovars VI as *Burkholderia dolosa* sp. nov. *Int J Syst Evol Microbiol* 54:689–691
84. Viallard V, Poirier I, Cournoyer B, Haurat J, Wiebkin S, Ophel-Keller K, Balandreau J (1998) *Burkholderia graminis* sp. nov., a rhizospheric *Burkholderia* species, and reassessment of [*Pseudomonas*] *phenazinum*, [*Pseudomonas*] *pyrocinia* and [*Pseudomonas*] *glathiei* as *Burkholderia*. *Int J Syst Bacteriol* 48:549–563
85. Vinuesa P, Leon-Barrios M, Silva C, Willems A, Jarabo-Lorenzo A, Perez-Galdona R, Werner D, Martinez-Romero E (2005) *Bradyrhizobium canariense* sp. nov., and acidic-tolerant

- endosymbiont that nodulates endemic genistoid legumes (Papilionoideae: Genisteae) from the Canary Islands, along with *Bradyrhizobium japonicum* bv. *genistearum*, *Bradyrhizobium* geno-species alpha and *Bradyrhizobium* genospecies beta. *Int J Syst Evol Microbiol* 55:569–575
86. Weinberg JB, Alexander BD, Majure JM, Williams LW, Kim JY, Vandamme P, LiPuma JJ (2008) *Burkholderia glumae* infection in an infant with chronic granulomatous disease. *J Clin Microbiol* 45:662–665
 87. zWhitlock GC, Estes DM, Torres AG (2007) Glanders: off to the races with *Burkholderia mallei*. *FEMS Microbiol Lett* 277:115–122
 88. Wong-Villarreal A, Caballero-Mellado J (2010) Rapid identification of nitrogen-fixing and legume-nodulating *Burkholderia* species based on PCR 16S rRNA species-specific oligonucleotides. *Syst Appl Microbiol* 33:35–43
 89. Yabuuchi E, Yoshimasa K, Oyaizu H, Yano I, Hotta H, Hashimoto Y, Ezaki T, Arakawa M (1992) Proposal of *Burkholderia* gen. nov. and transfer of seven species of the genus *Pseudomonas* homology group II to the new genus, with the type species *Burkholderia cepacia* (Palleroni and Holmes 1981) comb. nov. *Microbiol Immunol* 36:1251–1275
 90. Yabuuchi E, Kosako Y, Yano I, Hotta H, Nishiuchi Y (1995) Transfers of two *Burkholderia* and an *Alcaligenes* species to *Ralstonia* gen. nov.: proposal of *Ralstonia pickettii* (Ralston, Palleroni and Doudoroff 1973) comb. nov., *Ralstonia solanacearum* (Smith 1896) comb. nov. and *Ralstonia eutropha* (Davis 1969) comb. nov. *Microbiol Immunol* 39:897–904
 91. Yang Z (1996) Among-site rate variation and its impact on phylogenetic analyses. *Trends Ecol Evol* 11:367–372
 92. Yoo SH, Kim BY, Weon HY, Kwon SW, Go SJ, Stackebrandt E (2007) *Burkholderia soli* sp. nov., isolated from soil cultivated with Korean ginseng. *Int J Syst Evol Microbiol* 57:122–125
 93. Zhang H, Hanada S, Shigematsu T, Shibuya K, Kamagata Y, Kanagawa T, Kurane R (2000) *Burkholderia kururiensis* sp. nov., a trichloroethylene (TCE)-degrading bacterium isolated from an aquifer polluted with TCE. *Int J Syst Evol Microbiol* 50:743–749