

Lawrence Berkeley National Laboratory

LBL Publications

Title

Small Angle Neutron-Proton Scattering at 90 Mev

Permalink

<https://escholarship.org/uc/item/39x2907d>

Authors

Chamberlain, Owen

Easley, James W

Publication Date

1954-02-01

UCRL 2475

UNCLASSIFIED

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-2475

UNIVERSITY OF CALIFORNIA

UNCLASSIFIED

Radiation Laboratory

Contract No. W-7405-eng-48

SMALL ANGLE NEUTRON-PROTON SCATTERING AT 90 MEV

Owen Chamberlain and James W. Easley

February 4, 1954

Berkeley, California

Small Angle Neutron-Proton Scattering at 90 Mev.

Owen Chamberlain and James W. Easley

Radiation Laboratory, Department of Physics, University of California,
Berkeley, California

February 4, 1954

The relative neutron-proton differential cross section has been measured for several angles near zero degrees. This range of angles has been difficult to investigate because of the very short range of the recoil protons. Previously published work¹⁻⁵, done by measurement of proton recoils, has not covered these angles. However, some cloud chamber experiments are currently in progress at this laboratory⁶ in which recoil protons are employed in this range.

The neutron beam was obtained by stripping 190 Mev deuterons on a 1/2-in. beryllium target in the 184-inch Berkeley cyclotron. A steel shielding block 6-ft long X 6-in. high X 3-in. thick was placed close to the beam on the same side as the counters to reduce background.

The target was of liquid hydrogen contained in a cylindrical vessel of 5.6-in. diameter with axis vertical. The beam dimensions at the target were 3-in. high X 1-in. wide. A bismuth fission counter was employed as a beam monitor.

Scattered neutrons were counted at small angles to the neutron beam. Fig. 1 shows the neutron counter, which consisted of the following components, listed in the order that they would be traversed by a scattered particle: (a) absorber number 1, 25 g cm^{-2} of copper to prevent protons from reaching the scintillation counters directly; (b) scintillation counter number 1, made of plastic scintillant 4.0 in. X 3.5 in. X 0.080-in. thick; (c) the converter, 1.74 g cm^{-2} of poly-

ethylene; (d) scintillation counter number 2, a liquid counter with active volume 2.2 in. x 1.4 in. x 0.18 in. thick; (e) absorber number 2, which was from 1.0 to 1.8 g cm⁻² of copper plus a shaped aluminum absorber of maximum thickness 1.8 g cm⁻²; (f) scintillation counter number 3, made of plastic scintillant 3.6 in. x 2 in. x 0.4 in. thick. The distance from the center of the target to the center of the converter was 48 in. for the range of the laboratory scattering angle θ from 18.5° to 5.0° and 70 in. for the range of 5.0° to 2.5°.

The neutrons actually counted were those that were "converted" in the polyethylene converter or in counter number 2. The term "converted neutron" refers to a neutron that yields a high energy proton by n-p scattering or nuclear disintegration. These recoil protons were counted in coincidence in counters 2 and 3. In order to reject charged particles originating in other processes, counter 1 was connected in anticoincidence. Only events detected in counters 2 and 3 but not 1 were counted.

The thickness of absorber 2 determined the minimum energy which a scattered neutron could have and still be counted. In the present work the minimum energy was adjusted at each scattering angle to be $[(80 \text{ Mev}) \cos^2 \theta]$. Therefore, a neutron in the beam must have had at least 80 Mev in order to have been counted after scattering, since otherwise the proton yielded at the converter would have had insufficient range to reach counter 3. The shaped portion of absorber number 2 was thickest in the center and was shaped so that to good approximation the minimum neutron energy to count would be independent of the angle with which the recoil proton emerged from the converter. To determine the shape of the absorber we assumed that most of the recoil protons were

from n-p scattering processes in the converter. In order to ascertain whether the relative differential cross sections were particularly sensitive to the value of the minimum energy for detection of the scattered neutrons a measurement was made with this minimum energy increased to 66 Mev. The results were in agreement with those obtained for the 60 Mev minimum within the uncertainty of the counting statistics.

The results are tabulated in Table I. Fig. 2 shows the data in their relation to prior work, indicating a marked similarity of the differential cross section in the region of 0° to that in the region of 180° .

Thanks are due to Dr. Martin O. Stern for initial studies of background and design of the neutron collimator.

This work was done under the auspices of the Atomic Energy Commission.

¹Hadley, Kelly, Leith, Segre, Wiegand, and York, Phys. Rev. 75, 351 (1949)

²Brueckner, Hartsough, Hayward, and Powell, Phys. Rev. 75, 555 (1949)

³R. H. Fox, Atomic Energy Commission Report UCRL-387, 1950 (unpublished).

⁴R. Wallace, Phys. Rev. 81, 493 (1951)

⁵Selove, Strauch, and Titus, Phys. Rev. 92, 724 (1953).


⁶C. Y. Chih and W. M. Powell (private communication).

TABLE I. Experimental data for neutron-proton scattering at 30 Kev. θ is the scattering angle in the center-of-mass system. Column I gives the ratio of the differential cross section in the center-of-mass system at the angle θ to that at 30° . Column II gives the absolute differential cross section obtained by normalizing to the results of Hadley et al. (see reference 1). The errors listed are the standard deviations of the counting statistics.

θ	I relative cross section	II absolute cross section (10^{-27} cm ² /steradian)
5.1	$1.70 \pm .14$	12.9 ± 1.2
10.3	$1.53 \pm .06$	12.0 ± 0.7
20.6	$1.35 \pm .05$	10.5 ± 0.6
33.0	1.00	7.6 ± 0.4

FIG. 1. Arrangement of components of neutron counter.

FIG. 2. Differential neutron-proton cross section in the center-of-mass system in 10^{-27} cm²/steradian.



NEUTRON COUNTER.

