

UCSF

UC San Francisco Previously Published Works

Title

Attribution of cause of end-stage renal disease among patients with systemic lupus erythematosus: the Georgia Lupus Registry.

Permalink

<https://escholarship.org/uc/item/3b70n731>

Journal

Lupus Science and Medicine, 3(1)

ISSN

2053-8790

Authors

Drenkard, Cristina

Pastan, Stephen

Lim, S

et al.

Publication Date

2016

DOI

10.1136/lupus-2015-000132

Peer reviewed

Attribution of cause of end-stage renal disease among patients with systemic lupus erythematosus: the Georgia Lupus Registry

Laura C Plantinga,¹ Cristina Drenkard,² Stephen O Pastan,¹ S Sam Lim²

To cite: Plantinga LC, Drenkard C, Pastan SO, *et al.* Attribution of cause of end-stage renal disease among patients with systemic lupus erythematosus: the Georgia Lupus Registry. *Lupus Science & Medicine* 2016;**3**: e000132. doi:10.1136/lupus-2015-000132

Received 2 October 2015
Revised 21 November 2015
Accepted 23 December 2015



CrossMark

¹Division of Renal Medicine, Department of Medicine, Emory University, Atlanta, Georgia, USA

²Division of Rheumatology, Department of Medicine, Emory University, Atlanta, Georgia, USA

Correspondence to
Dr Laura C Plantinga;
laura.plantinga@emory.edu

ABSTRACT

Objective: Whether using provider-attributed end-stage renal disease (ESRD) cause of systemic lupus erythematosus (SLE) in national surveillance data captures the entire population of patients with SLE and ESRD remains uncertain. Our goal was to examine attributed cause of ESRD in US surveillance data among patients with SLE who have developed ESRD.

Methods: Data from a national registry of treated ESRD (United States Renal Data System (USRDS)) were linked to the population-based Georgia Lupus Registry (GLR). The provider-attributed cause of ESRD was extracted from the USRDS for each validated patient with SLE in the GLR (diagnosed through 2004) who initiated treatment for ESRD through 2012. The percentage of these patients with SLE whose ESRD was subsequently attributed to SLE in the USRDS was calculated, overall and by patient characteristics.

Results: Among 251 patients with SLE who progressed to ESRD, 78.9% had SLE as their attributed cause of ESRD. Of the remaining 53 patients, 43.4%, 18.9% and 15.6% had ESRD attributed to hypertension, diabetes mellitus type II and non-SLE-related glomerulonephritis, respectively. Attribution of ESRD to SLE was higher among patients aged ≤ 30 (87.9–93.9%) vs >30 (52.6%; $p < 0.001$) but did not differ by sex or race. Having Medicaid (86.2%) or no insurance (93.5%) was associated with greater attribution of ESRD to SLE than having private insurance (72.5%; $p = 0.02$), as was having two or more providers state a diagnosis of SLE (89.0% vs 73.5% with a rheumatologist diagnosis alone; $p = 0.008$).

Conclusions: These estimates indicate that USRDS-based studies may underreport ESRD among US patients with SLE. However, observed patterns of differential attribution of ESRD cause, particularly by age, suggest that providers may be correctly attributing ESRD to causes other than SLE among some patients with SLE.

INTRODUCTION

The study of US patients with end-stage renal disease (ESRD) attributed to systemic lupus

KEY MESSAGES

- ▶ Among 251 systemic lupus erythematosus (SLE) patients from a U.S. population-based registry, who subsequently progressed to end-stage renal disease (ESRD), 79% had SLE as their attributed cause of ESRD in national administrative data.
- ▶ Attribution of ESRD to SLE was higher among younger patients (88–94% and 53% among patients aged <30 and >30 years, respectively) but it did not differ by sex or race.
- ▶ These estimates indicate that national studies that rely on administrative data may underreport ESRD among SLE patients, particularly among older SLE patients.

erythematosus (SLE) is possible through the United States Renal Data System (USRDS).¹ The USRDS captures all treated ESRD in the USA, due to the universal Centers for Medicare & Medicaid Services (CMS) coverage of ESRD treatment for Social Security-eligible patients, regardless of patient age or disability status. The USRDS includes data on the provider-attributed cause of ESRD from the CMS ESRD Medical Eligibility Form 2728 (CMS-2728), which is required for all treated patients with ESRD. Several investigators have leveraged these data to estimate the population-based incidence of SLE-attributed ESRD^{2–5} and to demonstrate disparities in this incidence by age,^{4–6} sex,⁴ race,^{2 4–6} socioeconomic status,^{7 8} access to care⁹ and geography.^{4 6} Our group also recently examined wait time to transplant and graft failure¹⁰ and quality of ESRD care^{11 12} in patients with ESRD attributed to SLE using the USRDS.

However, the degree to which this method, which relies upon provider identification of SLE as the cause of ESRD at the start of treatment, captures the entire population of patients with SLE and ESRD remains

uncertain. One prior validation study¹³ conducted using biopsy samples prior to 2001 suggested potentially low sensitivity of this attributed cause. Recently, we linked the population-based Georgia Lupus Registry (GLR)^{14–16} to the USRDS to estimate ESRD incidence among patients with SLE in metropolitan Atlanta.¹⁷ Using these linked data, we aimed to examine the attribution of ESRD cause among patients with known SLE who had progressed to ESRD.

PATIENTS AND METHODS

Data sources and study population

GLR. The primary aim of the GLR was to more accurately estimate the prevalence and incidence of SLE in 2002–2004 in Atlanta, Georgia (Fulton and DeKalb Counties).^{14–16} Investigators partnered with the state health department under a public health surveillance exemption to the Health Insurance Portability and Accountability Act to comprehensively search for and review cases. Potential SLE cases were identified from multiple sources, including hospitals; providers in rheumatology, dermatology and nephrology; commercial and hospital-based laboratories; regional pathology laboratories; lupus research databases and population databases, including the USRDS, Veterans Affairs data, Medicaid claims data, and state mortality and hospital discharge data. The presence of a suggestive pathology report, laboratory test and/or diagnostic code (International Classification of Diseases, 9th revision (ICD-9)) for SLE (710.0) or related conditions that might evolve into SLE—including discoid lupus (695.4), other specified connective tissue disease (710.8) and other unspecified connective tissue disease (710.9)—in any of these sources flagged patients as potential SLE cases.¹⁶ Medical records for all potential cases with residence in Fulton or DeKalb County in 2002–2004 were fully abstracted by trained abstractors. Patient date of SLE diagnosis, fulfilment of classification criteria, age, sex, race and other abstracted medical information were available in the GLR. The project was reviewed and approved by the Emory University and Georgia Department of Public Health Institutional Review Boards.

USRDS. The USRDS data¹ originate primarily from the CMS and the United Networks for Organ Sharing. Date of initiation of ESRD treatment (through 30 September 2012), attributed cause of ESRD, age and insurance at ESRD start, and pre-ESRD nephrology care (2005+ only) were obtained from the USRDS. The USRDS also provided inpatient diagnostic codes for any hospitalisations ≥ 90 days after ESRD start (when Medicare coverage is established for all patients with ESRD, regardless of initial insurance). Up to ten discharge codes for each hospitalisation event were available.

US Census. Publicly available data on characteristics of US residential neighbourhoods, as defined by census tracts, were obtained from the 2000 US Census ([\[www.census.gov\]\(http://www.census.gov\)\) via the Minnesota Population Center \(<http://www.nghis.org>\).¹⁸ Aggregate census tract-level data on racial composition, education and poverty were extracted.](http://</p></div><div data-bbox=)

Data linkage. Identifiers (Social Security number, date of birth, sex, first name and surname) on incident patients with SLE from the GLR were sent to the USRDS for a probabilistic match to identify those who progressed to ESRD from diagnosis of SLE (2002–2004) to the last date of follow-up currently available from the USRDS (30 September 2012). First patient residential addresses from the GLR were geocoded, which identified the census tract for spatial linkage with Census data. The GLR-USRDS data linkage was approved by the Emory Institutional Review Board.

Study population. SLE cases were defined by a combined case definition previously described:¹⁴ (1) the presence of ≥ 4 American College of Rheumatology (ACR) criteria¹⁹ in the medical record, or (2) the presence of three ACR criteria plus a treating, board-certified rheumatologist's final diagnosis of SLE; or (3) < 4 ACR criteria plus a biopsy consistent with class II–VI lupus nephritis. Among the 1666 SLE cases by this definition in the GLR, 284 (17.0%) were identified in the USRDS as having started ESRD treatment in the USA between 1 January 1979 and 30 September 2012. Patients whose ESRD treatment start date preceded their SLE diagnosis date ($n=31$) or who lacked an attributed cause of ESRD ($n=2$) were excluded, leaving 251 patients with SLE for the analysis of attributed ESRD cause.

Study variables and definitions

Attributed cause of ESRD. The attributed cause of ESRD was defined by the single ICD-9 code listed on the CMS-2728, which is completed by the dialysis provider (usually the nephrologist) within 45 days of dialysis start. Attribution to SLE was defined by CMS-2728 ICD-9 code 710.0.

Patient characteristics. Patient demographics, including age at SLE diagnosis, sex, race and ethnicity, were obtained from the GLR. Age at ESRD start was available from the USRDS. Census tract-level sociodemographic variables, including percentage black residents, percentage residents living below poverty and percentage residents not completing high school, were available from the US Census and were assigned to individuals based on patient residence at the time of GLR data collection (2002–2004). Insurance at the start of ESRD (private, Medicaid, other and none) and reported pre-ESRD nephrology care (collected 2005+ only) were available from USRDS data, via the CMS-2728.

Duration of SLE prior to ESRD start was calculated as the time from date of SLE diagnosis (from GLR) to the date of ESRD start (from USRDS). Information on whether patients had renal biopsies prior to 2004, physician-stated SLE-related diagnoses and ACR criteria were obtained from the GLR. The 11 ACR criteria¹⁹

were defined by medical record-extracted data, available through 2004.

Statistical analysis

Overall patient characteristics were summarised by means, medians or percentages, as appropriate. Attribution of ESRD to SLE was defined as the percentage of patients with SLE who progressed to ESRD whose providers recorded SLE as the cause of ESRD on their CMS-2728. All attributed causes were tabulated, and attribution of ESRD cause was examined by patient characteristics. The effects of (1) using inpatient ICD-9 diagnostic discharge codes for hospitalisations subsequent to ESRD treatment initiation, in addition to the provider-attributed cause on the CMS-2728 and (2) restricting the population to those with biopsy-proven lupus nephritis on capture of SLE cases were also examined. Stata V. 14 (StataCorp, College Station, Texas, USA) was used for all analyses.

RESULTS

Characteristics of patients with SLE who developed ESRD

Overall, the mean age at SLE diagnosis among 251 Atlanta-area patients with SLE who had ESRD was 25 years, with a third being <18 years (table 1); on average, patients were 39 years old when they started ESRD treatment. Table 1 shows that the majority were female (85%) and black (92%) and that these patients lived in neighbourhoods that were predominantly black (median, 89% of residents in census tract). Patients' residential neighbourhoods also had median values of 11% and 13% of residents living below poverty and not finishing high school, respectively. Most patients had private insurance (40%) or Medicaid (28%) at the start of ESRD; 14% were not insured (table 1). Most were diagnosed with SLE by a rheumatologist, either alone or in combination with a nephrologist or dermatologist, and 69% had had at least one renal biopsy (table 1). Among patients with at least one biopsy, 149 (86.1%) had a biopsy consistent with lupus nephritis. The median duration of SLE prior to the start of ESRD treatment was 8 years, and 32% and 43% had had SLE <5 and >10 years, respectively, prior to ESRD start (table 1). Most had a physician-stated diagnosis of SLE (88%) or SLE in combination with connective tissue disease (9%) (table 1). The most common ACR criteria met by these patients were renal (96%), ANA (95%), haematological (92%) and immunological (73%) (table 1).

Patients with ESRD attributed to SLE were younger at SLE diagnosis and more likely to have Medicaid than patients with ESRD attributed to other causes; no differences by race, sex or neighbourhood sociodemographics were noted (table 1). Having an SLE diagnosis, having that diagnosis from more than specialist and having greater numbers of ACR criteria were all associated with greater likelihood of attribution of ESRD to SLE versus other causes (table 1). Among the individual criteria,

malar rash, arthritis, immunologic criteria (positive anti-DNA, anti-Smith or anti-cardiolipin antibodies; positive Russell viper venom time/mixing test or false-positive syphilis), haematological criteria and photosensitivity (not statistically significant) were more common among patients with ESRD attributed to SLE versus other causes, whereas prevalence of renal, discoid rash and ANA criteria did not differ by attributed cause (table 1). Of patients with at least one biopsy, 90.1% vs 68.8% of those with ESRD attributed to SLE versus other causes had biopsies consistent with lupus nephritis.

Attribution of ESRD cause in patients with SLE

Of the 251 ESRD cases, 198 (78.9%) were attributed to SLE on the CMS-2728 (table 2 and figure 1A). Most of the 53 of 251 ESRD cases that were not attributed to SLE were attributed to hypertension (HTN) (23/53, 43%) and diabetes mellitus type II (10/53, 19%), with 8 (16%) patients having ESRD attributed to non-SLE glomerulonephritis (figure 1A). Among patients with SLE who also had CMS hospitalisation data subsequent to ESRD start (n=196), capture of SLE increased from 77% by CMS-2728 alone to 91% and 96% by CMS-2728 or the presence of inpatient diagnostic discharge codes for SLE (in two or more hospitalisations or at least one hospitalisation, respectively; table 2). Among the 149 cases with biopsy-defined lupus nephritis in the GLR through 2004 who subsequently developed ESRD, 85.2% had their ESRD attributed to SLE in national surveillance data. In contrast, among the 24 patients with biopsy data that were not consistent with lupus nephritis and 78 patients without biopsy data, 58.3% and 73.1%, respectively, had their ESRD attributed to SLE (p=0.004).

Differential attribution of ESRD cause by patient characteristics

Attribution of ESRD to SLE was higher among patients aged ≤ 30 (88–94%) compared with patients aged >30 (53%; p<0.001; table 3). Among those aged >50 (n=15), only three (20.0%) had their ESRD attributed to SLE. Attribution of ESRD to SLE did not differ by sex, race or neighbourhood sociodemographics (table 3). Among the 149 patients with biopsy-proven lupus nephritis prior to 2004, black patients with SLE-ESRD did have higher attribution of their ESRD to SLE than their white counterparts (86.5% vs 62.5%), but the difference was not statistically significant (p=0.10). Those with Medicaid (86%) and no insurance (94%) were more likely than those with private (73%) or other (71%) insurance to have their ESRD attributed to SLE (p=0.02; table 3). There was a non-statistically significant trend towards higher attribution of ESRD to SLE among patients with renal biopsies (table 3). Attribution of ESRD to SLE was highest among those diagnosed with SLE by a nephrologist (82%) or by a combination of providers (89%; table 3). Diagnosis of SLE without other connective tissue

Table 1 Selected characteristics of 251 Georgia Lupus Registry patients who initiated treatment for ESRD (through 2012), on or after the date of SLE diagnosis, overall and by attributed cause of ESRD in the United States Renal Data System

Characteristic	Overall	ESRD attributed to		p Value*
		SLE	Other	
Sociodemographics				
Age in years at SLE diagnosis, mean (SD)	25.3 (13.6)	21.9 (11.2)	38.2 (14.3)	<0.001
Age (years) at SLE diagnosis, no (%)				<0.001
<18	82 (32.7%)	77 (38.9%)	4 (9.4%)	
18–30	91 (36.3%)	80 (40.4%)	11 (20.8%)	
>30	78 (31.1%)	41 (20.7%)	37 (69.8%)	
Sex, no (%)				0.66
Male	38 (15.1%)	31 (15.7%)	7 (13.2%)	
Female	213 (84.9%)	167 (84.3%)	46 (86.8%)	
Race, no (%)				0.64
Black	232 (92.4%)	184 (92.9%)	48 (90.6%)	
White	18 (7.2%)	13 (6.6%)	5 (9.4%)	
Asian	1 (0.4%)	1 (0.5%)	0 (0.0%)	
Hispanic, no (%)				>0.9
Yes	3 (1.2%)	3 (1.5%)	0 (0.0%)	
No	247 (98.8%)	194 (98.5%)	53 (100.0%)	
% black residents in residential census tract, median (IQR)	89.1 (73.7–95.5)	88.8 (73.7–95.3)	89.6 (77.7–96.3)	0.46
% living below poverty in residential census tract, median (IQR)	11.2 (7.0–22.7)	10.8 (6.5–22.5)	12.0 (7.9–24.5)	0.27
% high-school dropouts in residential census tract, median (IQR)	13.3 (8.5–25.8)	13.2 (8.4–24.4)	14.8 (9.6–26.6)	0.27
Access to care				
Insurance† at start of ESRD, no (%)				0.02
Private	91 (39.7%)	66 (36.5%)	25 (52.1%)	
Medicaid	65 (28.4%)	56 (30.9%)	9 (18.8%)	
Other	42 (18.3%)	30 (16.6%)	12 (25.0%)	
None	31 (13.5%)	29 (16.0%)	2 (4.2%)	
Pre-ESRD nephrology care, no (%)				0.09
Yes	77 (77.0%)	56 (72.7%)	21 (91.3%)	
No	23 (23.0%)	21 (27.3%)	2 (8.7%)	
Physician stating final diagnosis, no (%)				0.005
No rheumatologist, dermatologist or nephrologist	33 (13.2%)	20 (10.2%)	13 (24.5%)	
Rheumatologist	83 (33.2%)	61 (31.0%)	22 (41.5%)	
Dermatologist	4 (1.6%)	3 (1.5%)	1 (1.9%)	
Nephrologist	39 (15.6%)	32 (16.2%)	7 (13.2%)	
Combination of rheumatologist, dermatologist or nephrologist	91 (36.4%)	8 (41.1%)	10 (18.9%)	
At least one renal biopsy, no (%)				0.13
Yes	173 (68.9%)	141 (71.2%)	32 (60.4%)	
No	78 (31.1%)	57 (28.8%)	21 (39.6%)	
Multiple renal biopsies, no (%)				0.16
Yes	82 (32.7%)	69 (34.9%)	13 (24.5%)	
No	169 (67.3%)	129 (65.1%)	40 (75.5%)	
Clinical				
Duration of SLE prior to ESRD start, years	8.0 (4.0–14.1)	8.0 (4.1–13.6)	9.0 (4.0–16.3)	0.47
Duration of SLE prior to ESRD start (years), no (%)				0.74
<5	80 (31.9%)	64 (32.3%)	16 (30.2%)	
5–10	64 (25.5%)	52 (26.3%)	12 (22.6%)	
>10	107 (42.6%)	82 (41.4%)	25 (47.2%)	
Physician-stated diagnosis, no (%)				<0.001
SLE	219 (88.3%)	184 (93.4%)	35 (68.6%)	
SLE and connective tissue disease	22 (8.9%)	10 (5.1%)	12 (23.5%)	
Drug-induced lupus	1 (0.4%)	0 (0.0%)	1 (2.0%)	
Specified connective tissue disease	5 (2.0%)	3 (5.9%)	2 (1.0%)	
Unspecified diffuse connective tissue disease	1 (0.4%)	1 (0.5%)	0 (0.0%)	

Continued

Table 1 Continued

Characteristic	Overall	ESRD attributed to		p Value*
		SLE	Other	
No of ACR criteria, median (IQR)	5 (4–7)	6 (4–7)	5 (3–5)	<0.001
ACR criteria, %				0.08
≥4 criteria	193 (76.9%)	157 (79.3%)	36 (67.9%)	
<3 criteria	58 (23.1%)	41 (20.7%)	17 (32.1%)	
ACR criterion present, no (%)				
Malar rash	72 (28.7%)	63 (31.8%)	9 (17.0%)	0.03
Discoid rash	35 (13.9%)	29 (14.7%)	6 (11.3%)	0.54
Photosensitivity	47 (18.7%)	42 (21.2%)	5 (9.4%)	0.07
Mucocutaneous ulcers	43 (17.1%)	37 (18.7%)	6 (11.3%)	0.21
Arthritis	161 (64.1%)	136 (68.7%)	24 (47.2%)	0.004
Serositis	119 (47.4%)	99 (50.0%)	20 (37.7%)	0.11
Renal	241 (96.0%)	190 (96.0%)	51 (96.2%)	>0.9
Neurological	59 (23.9%)	51 (25.8%)	8 (15.1%)	0.10
Haematological	231 (92.0%)	188 (95.0%)	43 (81.1%)	0.001
Immunological	184 (73.3%)	158 (79.8%)	26 (49.1%)	<0.001
ANA	206 (94.9%)	162 (95.3%)	44 (93.6%)	0.71
Year of ESRD start, no (%)				0.80
Before 1995	27 (10.8%)	22 (11.1%)	5 (9.4%)	
1995–2004	120 (47.9%)	96 (48.5%)	24 (45.3%)	
2005 or later	104 (41.4%)	80 (40.4%)	24 (45.3%)	

n=251 for all characteristics except: Hispanic ethnicity, n=250; census tract characteristics, n=230; insurance at start of ESRD, n=229; pre-ESRD nephrology care, n=100 (collected 2005+ only); final physician stating diagnosis, n=250; physician-stated diagnosis, n=248.

*By t, rank-sum, χ^2 or Fisher's exact test, as appropriate.

†Other insurance includes Medicare, VA and other coverage not listed.

ACR, American College of Rheumatology; CMS-2728, Centers for Medicare & Medicaid Services ESRD Medical Eligibility Form 2728; ESRD, end-stage renal disease; SLE, systemic lupus erythematosus.

disease was also associated with high attribution (84% vs 46% among those with SLE and other connective tissue disease; table 3). There was a trend towards higher attribution of ESRD to SLE among patients with more severe disease by ACR criteria, but the differences were not statistically significant; however, attribution of ESRD to SLE was particularly high among patients with malar rash (88%) and photosensitivity (89%; table 3). There were no trends over time by year of ESRD start (table 3).

The majority of the attributions of ESRD to HTN (18/23, 78.3%) and diabetes mellitus type II (8/10, 80.0%) were among patients aged >30 years (figure 1B). Among those aged >50 years, 13% and 60% of ESRD cases were

attributed to diabetes mellitus type II and HTN, respectively. ESRD was more commonly attributed to diabetes mellitus II among females than males and to HTN among blacks than whites (figure 1B).


Among those 53 patients with SLE whose ESRD was not attributed to SLE, 46 had hospitalisations subsequent to ESRD; of these, 21 (45.6%) did not have diagnostic codes for their non-SLE attributed ESRD cause for any of their hospitalisations. However, this varied by attributed cause: only 5.0% of patients with HTN as cause did not have matching codes in their hospitalisation data, vs 33.3%, 62.5% and 100% of patients with ESRD attributed to diabetes mellitus type II, focal

Table 2 Identification of SLE by attributed cause of ESRD in the USRDS combined with inpatient diagnostic codes, among Georgia Lupus Registry (GLR) patients who initiated treatment for ESRD (through 2012) and were subsequently hospitalised

Identification of patients with SLE by	No of patients with SLE identified in USRDS	No of patients with SLE progressing to ESRD, with subsequent hospitalisation data	Attribution to SLE (95% CI)
CMS-2728 only	150	196	76.5% (70.0% to 82.3%)
CMS-2728 or ICD-9 code (710.0) in two or more hospitalisations ≥30 days apart	178	196	90.8% (85.9% to 94.5%)
CMS-2728 or ICD-9 code (710.0) in at least one hospitalisation	188	196	95.9% (92.1% to 98.2%)

CMS-2728, Centers for Medicare & Medicaid Services ESRD Medical Eligibility Form 2728; ESRD, end-stage renal disease; ICD-9, International Classification of Diseases V.9; SLE, systemic lupus erythematosus; USRDS, United States Renal Data System.

Figure 1 Attributed cause of end-stage renal disease (ESRD) among 251 patients with SLE in the Georgia Lupus Registry who progressed to ESRD (1979–2012), overall (A) and by age, sex and race (B). SLE, systemic lupus erythematosus; HTN, hypertension; DMII, diabetes mellitus type II; FSGS, focal segmental glomerulosclerosis; other GN, glomerulonephritis not attributed to SLE. Other includes tubular necrosis, IgA nephropathy, scleroderma, diabetes mellitus type I, multiple myeloma, other renal disorder and uncertain aetiology.



segmental glomerulosclerosis (FSGS) and glomerulonephritis, respectively (figure 2). Overall, 97.8% of patients with non-SLE attributed causes of ESRD had at least one hospitalisation with a code for SLE.

DISCUSSION

We found that, among a population of patients diagnosed with SLE in Atlanta through 2004 who subsequently started treatment for ESRD, attribution of ESRD cause to SLE using the CMS-2728 was 79%. This means that approximately one in five patients with SLE who progress to ESRD may be missed when the USRDS is used to identify the population of US patients with SLE who have developed ESRD. A prior validation study that compared biopsy results from a Southeastern registry

(with data on glomerular disease biopsies performed in 1979–2000) and CMS-2728-attributed causes showed that only 27% of patients with a biopsy indicating SLE had SLE as their attributed ESRD cause¹³—much lower than both our overall estimate and the more directly comparable estimate of attribution of ESRD to SLE among patients with biopsy-proven lupus nephritis (85%). The small sample size (n=30 biopsies indicating SLE) and more complete data on attributed cause of ESRD likely explains some of these differences, since the prior study considered missing attributed cause to be a mismatch and the attributed cause was missing in more than half of the cases.¹³

We have provided an updated estimate of the sensitivity of USRDS data in capturing the US population of patients with SLE and all-cause ESRD. However, without

Table 3 Attribution of ESRD to SLE in the USRDS, among GLR patients who initiated ESRD treatment through 2012, overall and by patient characteristics

Patient subgroup	No of GLR patients with SLE with ESRD attributed to SLE in USRDS	No of GLR patients with SLE progressing to ESRD	Attribution of ESRD to SLE (95% CI)
Overall	198	251	78.9% (73.3% to 83.8%)
Demographics			
Age at SLE diagnosis, years			
<18	77	82	93.9% (86.3% to 98.0%)
18–30	80	91	87.9% (79.4% to 93.8%)
>30	41	78	52.6% (40.9% to 64.0%)
p Value*			<0.001
Sex			
Male	31	38	81.6% (65.7% to 92.3%)
Female	167	213	78.4% (72.3% to 83.7%)
p Value*			0.66
Race†			
Black	184	232	79.3% (73.5% to 84.3%)
White	13	18	72.2% (46.5% to 90.3%)
p Value*			0.48
% black residents in residential census tract			
Above median	93	116	80.2% (71.7% to 87.0%)
Below median	88	114	77.2% (68.4% to 84.5%)
p Value*			0.58
% living below poverty in residential census tract			
Above median	87	115	75.7% (66.8% to 83.2%)
Below median	94	115	81.7% (73.5% to 88.3%)
p Value*			0.26
% high-school dropouts in residential census tract			
Above median	89	115	77.4% (68.7% to 84.7%)
Below median	92	115	80.0% (71.5% to 86.9%)
p Value*			0.63
Access to care			
Insurance at start of ESRD			
Private	66	91	72.5% (62.2% to 81.4%)
Medicaid	56	65	86.2% (75.3% to 93.5%)
Other	30	42	71.4% (55.4% to 84.3%)
None	29	31	93.5% (78.6% to 99.2%)
p Value*			0.02
Pre-ESRD nephrology care‡			
Yes	56	77	72.7% (61.4% to 82.3%)
No	21	23	91.3% (72.0% to 98.9%)
p Value*			0.06
Physician stating final SLE diagnosis			
No rheumatologist, dermatologist or nephrologists	20	33	60.6% (42.1% to 77.1%)
Rheumatologist	61	83	73.5% (62.7% to 82.6%)
Dermatologist	3	4	75.0% (19.4% to 99.4%)
Nephrologist	32	39	82.1% (66.5% to 92.5%)
Combination of rheumatologist, dermatologist or nephrologist	81	91	89.0% (80.7% to 94.6%)
p Value*			0.008
At least one renal biopsy			
Yes	141	173	81.5% (74.9% to 87.0%)
No	57	78	73.1% (61.8% to 82.5%)
p Value*			0.13
Multiple renal biopsies			
Yes	129	169	76.3% (69.2% to 82.5%)
No	69	82	84.2% (74.4% to 91.3%)
p Value*			0.16

Continued

Table 3 Continued

Patient subgroup	No of GLR patients with SLE with ESRD attributed to SLE in USRDS	No of GLR patients with SLE progressing to ESRD	Attribution of ESRD to SLE (95% CI)
Clinical			
Duration of SLE prior to ESRD start, years			
<5	64	80	80.0% (69.6% to 88.1%)
5–10	52	64	81.3% (69.5% to 89.9%)
>10	82	107	76.6% (67.5% to 84.3%)
p Value*			0.74
Physician-stated SLE-related diagnosis			
SLE	184	219	84.0% (78.5% to 88.6%)
SLE and connective tissue disease	10	22	45.5% (24.4% to 67.8%)
Drug-induced lupus	0	1	0%
Specified connective tissue disease	2	5	40.0% (5.3% to 85.3%)
Unspecified diffuse connective tissue disease	1	1	100%
p Value*			<0.001
ACR criteria, %			
≥4 criteria	157	193	81.4% (75.1% to 86.6%)
<3 criteria	41	58	70.7% (57.3% to 81.9%)
p Value*			0.08
ACR criterion present			
Malar rash	63	72	87.5% (77.6% to 94.1%)
Discoid rash	29	35	82.9% (66.4% to 93.4%)
Photosensitivity	42	47	89.4% (76.9% to 96.5%)
Mucocutaneous ulcers	37	43	86.1% (72.1% to 94.7%)
Arthritis	136	161	84.5% (77.9% to 89.7%)
Serositis	99	119	83.2% (75.2% to 89.4%)
Renal	190	241	78.8% (73.1% to 83.8%)
Neurological	51	59	86.4% (75.0% to 94.0%)
Haematological	188	231	81.4% (75.8% to 86.2%)
Immunological	158	184	85.9% (80.0% to 90.6%)
ANA	162	206	78.6% (72.4% to 84.0%)
Year of ESRD start			
Before 1995	22	27	81.5% (61.9% to 93.7%)
1995–2004	96	120	80.0% (71.7% to 86.7%)
2005 or later	80	104	76.9% (67.6% to 84.6%)
p Value*			0.80

*By χ^2 or Fisher's exact test.

†Restricted to white and black patients only.

‡Pre-ESRD nephrology care collected 2005+ only.


ACR, American College of Rheumatology; ESRD, end-stage renal disease; ESRD Medical Eligibility Form 2728; SLE, systemic lupus erythematosus; USRDS, United States Renal Data System.

a gold standard that provides a definitive diagnosis for each GLR patient (eg, renal biopsies) prior to ESRD start, we cannot estimate the sensitivity of this method to capture ESRD that is truly due to SLE. Further, we are unable to determine the population of patients with ESRD living in Atlanta during the same time at risk for ESRD (rather than just at ESRD start) as the GLR patients, due to unknown residential mobility in and out of the area over time. We also cannot identify individuals who are not in the GLR who have ESRD attributed to SLE in the USRDS (potential false-positive cases), among the same source population from which the GLR was drawn. Thus, we cannot properly perform a true

validation study and estimate other important aspects of validity of CMS-2728-attributed cause, including the specificity (probability that patients with ESRD without SLE will have their ESRD attributed to non-SLE causes) and the positive predictive value (probability that a patient with SLE-attributed ESRD actually has SLE).

The addition of diagnostic codes from CMS-covered hospitalisations subsequent to ESRD start to the CMS-2728 in identification of patients with SLE with ESRD increased capture of patients with SLE from 79% to >90%. However, diagnostic codes may be unreliable, and this method of identifying patients with SLE with ESRD to increase capture may be especially problematic

Figure 2 Presence of inpatient diagnostic codes for attributed cause of end-stage renal disease (ESRD) among 46 patients with systemic lupus erythematosus (SLE) in the Georgia Lupus Registry who progressed to ESRD (1979–2012) and had subsequent hospitalisations but whose ESRD was not attributed to SLE. HTN, hypertension; DMII, diabetes mellitus type II; FSGS, focal segmental glomerulosclerosis; other GN, glomerulonephritis not attributed to SLE. Other includes tubular necrosis, IgA nephropathy, scleroderma, diabetes mellitus type I, multiple myeloma, other renal disorder and uncertain aetiology.



in patients with SLE, who are not likely to have Medicare coverage prior to the start of ESRD (based on age ≥ 65 years or disability). Thus, hospitalisations that take place prior to ESRD start and before Medicare eligibility is established may be captured differentially. Additionally, these patients are likely to have secondary insurance that may cover hospitalisation costs, may lose ESRD coverage 3 years after kidney transplantation,²⁰ or may die soon after ESRD start. Further, not all patients will be hospitalised after ESRD start. We found that 21% of these patients did not have any claim data for Medicare-covered hospitalisations subsequent to ESRD.

Attribution of ESRD to SLE was greatly reduced among older patients (53% among those aged >30 and only 20% among those aged ≥ 50). This is likely to reflect increasing comorbidity in older patients with SLE, particularly HTN and diabetes, which are, by far, the most common causes of ESRD in the general population¹ and which are common in SLE, due to metabolic effects of chronic inflammation, steroid treatment or both.^{21–25} In fact, we found that many patients with non-SLE attributed ESRD causes such as HTN had subsequent inpatient codes for these conditions. This observation would suggest that these patients were likely to have the conditions listed as their attributed ESRD cause and that the non-SLE ESRD cause attributed by the provider may not have always been incorrect, even if it did not capture the SLE status of the patient. Studies of older patients with SLE-ESRD should recognise the limitation that patients with SLE with other comorbidities that are associated with ESRD may not be captured using only the dialysis provider-attributed cause of ESRD.

Age was associated with the most striking differences in attribution of SLE cause, but we generally did find differential attribution by other patient characteristics. While we did find that black patients with SLE who

progressed to ESRD were more likely than their white counterparts to have their ESRD attributed to SLE (79% vs 72%), and that this difference was more pronounced among those with biopsy-proven lupus nephritis (87% vs 63%)—suggesting provider perceptions may play a role in recognition of SLE at ESRD start—neither difference was statistically significant. This is likely at least partially due to the small sample size of white patients progressing to ESRD in the GLR ($n=18$). Blacks were also more likely than whites to have their ESRD attributed to HTN rather than SLE. While attribution of ESRD cause to SLE was not differential by sex overall, females were more likely than males to have their ESRD attributed to diabetes mellitus type II. Having Medicaid or no insurance was associated with greater attribution to SLE than private insurance, which may partially reflect the ages of the patients with each type of insurance.

Interestingly, among those patients with pre-ESRD nephrology care information, those without nephrology care were more likely to have their ESRD attributed to SLE, although this association was not statistically significant. Nephrologists may be more likely to be aware of renal biopsy results that indicate non-SLE-related kidney disease and also to provide care pre-ESRD and post-ESRD start and thus attribute ESRD to other comorbid causes. Having at least two providers stating the diagnosis was associated with ESRD being attributed to SLE nearly 90% of the time, vs 73% with a rheumatologist alone, indicating the potential importance of communication between providers in recognition of SLE in ESRD. More severe SLE and biopsy history were associated with higher attribution of ESRD to SLE, relative to milder SLE and no biopsies performed, but these associations were not statistically significant. Longer duration of SLE prior to ESRD was not associated with greater attribution. More visible symptoms of SLE,

including malar rash and photosensitivity, were associated with slightly higher attribution than other symptoms, but whether these symptoms persisted at ESRD start is unknown. Despite substantial changes to the CMS-2728 in 1995 and 2005, there was no increase in attribution over time.

Additional limitations deserve mention. First, these estimates may not be generalisable to other US SLE populations. We had limited power to detect differential attribution of ESRD cause by patient characteristics and particularly to stratify on or adjust for multiple covariates. Given the number of exposures assessed, it is possible that some statistically significant associations are due to chance alone. Non-CMS-eligible patients (eg, undocumented residents) and patients who moved out of the USA between SLE diagnosis and start of ESRD treatment would not be captured by USRDS. However, these estimates apply to studies with the same limitations. We also did not have data on individual socioeconomic status. Data on medications, including prescribed immunosuppressant medications and patient adherence to these regimens, were not available in the GLR. It is possible that younger patients were more often prescribed fertility-preserving immunosuppressant regimens and/or had lower adherence to their prescribed regimens, leading to more clinically apparent lupus nephritis at ESRD presentation and, thus, greater attribution of ESRD to SLE among this population. Only one cause is allowed per patient on the CMS-2728 form, which may decrease attribution to SLE: for example, FSGS could be secondary to lupus nephritis but still listed as the attributed cause of ESRD; these patients would remain uncaptured by attempts to identify the US SLE-ESRD population using data on the attributed cause, even if they are recognised by providers as patients with SLE. Finally, although renal biopsy information was collected, when available, in the GLR, attempts to validate non-SLE-attributed ESRD causes using biopsy data would be confounded by indication and subject to selection bias as well. Furthermore, since GLR case ascertainment ended in 2004, no clinical data, including biopsy results, are available after this time; even among patients with biopsy-proven lupus nephritis at a single time point, other conditions related to treatment or complications of SLE and lupus nephritis could still develop and lead to ESRD.

Despite these limitations, this study provides an updated estimate of sensitivity of USRDS data to capture US patients with SLE who have progressed to ESRD, regardless of cause. Additionally, our findings of differential attribution of ESRD cause by age and severity of SLE suggest that older patients or patients with milder cases of SLE, whose ESRD may be attributed to comorbid conditions—perhaps even resulting from prolonged SLE treatment—could have been excluded from previous studies based on attributed ESRD cause alone incidence of SLE-attributed ESRD.^{2–12} These results contribute to the appropriate interpretation of past and

future US population-based studies of incidence and quality of care in ESRD patients with SLE.

Acknowledgements Some of the data reported here have been supplied by the United States Renal Data System (USRDS). The interpretation and reporting of these data are the responsibility of the authors and in no way should be seen as an official policy or interpretation of the US government.

Contributors All contributors meet the criteria for authorship and approved the submitted version of the manuscript.

Funding This research was supported by the Lupus Foundation of America. The Georgia Lupus Registry was supported in part by the CDC, and by cooperative agreement CDC-RFA-DP08-806 and earlier by cooperative agreement PA03022 from the CDC. This publication was supported by the Grant or Cooperative Agreement Number, DP005119, funded by the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services.

Competing interests None declared.

Ethics approval The Emory University institutional review board and the Georgia Department of Public Health provided ethics approval for this study.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement USRDS data are publicly available by request (<http://www.usrds.org>). Census data, including the American Community Survey, are publicly available (<http://www.census.gov> or <http://www.nhgis.org>.)

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. United States Renal Data System. 2014 Annual Data Report: Atlas of Chronic Kidney Disease and End-Stage Renal Disease in the United States. Bethesda, MD: National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, 2014.
2. Ward MM. Changes in the incidence of end-stage renal disease due to lupus nephritis, 1982–1995. *Arch Intern Med* 2000;160:3136–40.
3. Ward MM. Changes in the incidence of endstage renal disease due to lupus nephritis in the United States, 1996–2004. *J Rheumatol* 2009;36:63–7.
4. Costenbader KH, Desai A, Alarcón GS, *et al*. Trends in the incidence, demographics, and outcomes of end-stage renal disease due to lupus nephritis in the US from 1995 to 2006. *Arthritis Rheum* 2011;63:1681–8.
5. Sexton DJ, Reule S, Solid C, *et al*. ESRD from lupus nephritis in the United States, 1995–2010. *Clin J Am Soc Nephrol* 2015;10:251–9.
6. Hiraki LT, Lu B, Alexander SR, *et al*. End-stage renal disease due to lupus nephritis among children in the US, 1995–2006. *Arthritis Rheum* 2011;63:1988–97.
7. Ward MM. Medical insurance, socioeconomic status, and age of onset of endstage renal disease in patients with lupus nephritis. *J Rheumatol* 2007;34:2024–7.
8. Ward MM. Socioeconomic status and the incidence of ESRD. *Am J Kidney Dis* 2008;51:563–72.
9. Ward MM. Access to care and the incidence of endstage renal disease due to systemic lupus erythematosus. *J Rheumatol* 2010;37:1158–63.
10. Plantinga LC, Patzer RE, Drenkard C, *et al*. Association of time to kidney transplantation with graft failure among US patients with end-stage renal disease due to lupus nephritis. *Arthritis Care Res* 2015;67:571–81.
11. Plantinga LC, Drenkard C, Patzer RE, *et al*. Sociodemographic and geographic predictors of quality of care in United States patients with end-stage renal disease due to lupus nephritis. *J Rheumatol* 2015;67:761–72.
12. Plantinga LC, Patzer RE, Drenkard C, *et al*. Comparison of quality-of-care measures in US patients with end-stage renal

- disease secondary to lupus nephritis vs. other causes. *BMC Nephrol* 2015;16:39.
13. Layton JB, Hogan SL, Jennette CE, *et al.* Discrepancy between Medical Evidence Form 2728 and renal biopsy for glomerular diseases. *Clin J Am Soc Nephrol* 2010;5:2046–52.
 14. Lim SS, Bayakly AR, Helmick CG, *et al.* The incidence and prevalence of systemic lupus erythematosus, 2002–2004: the Georgia Lupus Registry. *Clin J Am Soc Nephrol* 2014;66:357–68.
 15. Lim SS, Drenkard C. Epidemiology of systemic lupus erythematosus: capturing the butterfly. *Curr Rheumatol Rep* 2008;10:265–72.
 16. Lim SS, Drenkard C, McCune WJ, *et al.* Population-based lupus registries: advancing our epidemiologic understanding. *Arthritis Rheum* 2009;61:1462–6.
 17. Plantinga LC, Lim SS, Patzer RE, *et al.* Incidence of end-stage renal disease among newly diagnosed systemic lupus erythematosus patients: the Georgia Lupus Registry. *Arthritis Care Res* 2016; doi:10.1002/acr.22685
 18. Minnesota Population Center. *National Historical Geographic Information System: Version 2.0*. Minneapolis, MN: University of Minnesota, 2011. <http://www.nhgis.org>
 19. Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1997;40:1725. doi:10.1002/1529-0131(199709)40:9<1725::AID-ART29>2-Y
 20. Grubbs V, Plantinga LC, Vittinghoff E, *et al.* Medicare immunosuppressant coverage and access to kidney transplantation: a retrospective national cohort study. *BMC Health Serv Res* 2012;12:254.
 21. Shaharir SS, Gafor AH, Said MS, *et al.* Steroid-induced diabetes mellitus in systemic lupus erythematosus patients: analysis from a Malaysian multi-racial lupus cohort. *Int J Rheum Dis* 2015;18:541–7.
 22. Parker B, Bruce I. SLE and metabolic syndrome. *Lupus* 2013;22:1259–66.
 23. Parker B, Urowitz MB, Gladman DD, *et al.* Clinical associations of the metabolic syndrome in systemic lupus erythematosus: data from an international inception cohort. *Ann Rheum Dis* 2013;72:1308–14.
 24. Bultink IE, Turkstra F, Diamant M, *et al.* Prevalence of and risk factors for the metabolic syndrome in women with systemic lupus erythematosus. *Clin Exp Rheumatol* 2008;26:32–8.
 25. Molina MJ, Mayor AM, Franco AE, *et al.* Prevalence of systemic lupus erythematosus and associated comorbidities in Puerto Rico. *J Clin Rheumatol* 2007;13:202–4.