UC Irvine

UC Irvine Previously Published Works

Title

POSSIBLE VARIATIONS IN ATMOSPHERIC METHANE.

Permalink

<https://escholarship.org/uc/item/3bb6023n>

Journal

J Geophys Res, 82(12)

ISSN

0148-0227

Authors

Chameides, WL

Liu, SC

Cicerone, RJ

Publication Date

1977

DOI

10.1029/JC082i012p01795

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Possible Variations in Atmospheric Methane

W. L. CHAMEIDES

Department of Physics and Astronomy, University of Florida
Gainesville, Florida 32611

S. C. LIU AND R. J. CICERONE

Department of Atmospheric and Oceanic Sciences, University of Michigan
Ann Arbor, Michigan 48109

A model coupling photochemistry and vertical transport in the troposphere has been used to investigate the magnitude of possible future perturbations to atmospheric CH_4 . We have studied the response of atmospheric CH_4 to an increase in the concentration of CO or to a variation in stratospheric O_3 . Both of the above mechanisms, which could be caused by man's activities, arise because a change in the concentration of tropospheric OH will lead to a change in the CH_4 abundance. Our calculations imply that a perturbation of ~30–40% in atmospheric CH_4 could occur if stratospheric O_3 were perturbed by 10% or if man-made CO continued to increase. The possible consequences of a CH_4 perturbation may entail a perturbation in stratospheric photochemistry or in the thermal balance of the atmosphere.

INTRODUCTION

Atmospheric CH_4 is produced at the earth's surface primarily by biological fermentation in anaerobic environments such as swamps, tropical rain forests, and paddy fields [Koyama, 1963]. Tropospheric CH_4 is characterized by a well-mixed profile with a vertically constant mixing ratio of 1.4 ppm [Ehhalt *et al.*, 1974]. Studies of the flux of CH_4 from various soil types imply a global production rate of the order of 10^{11} CH_4 molecules $\text{cm}^{-2} \text{ s}^{-1}$ (420 Mt yr^{-1} (420 million metric tons per year)) [Koyama, 1963; Baker-Blocker *et al.*, 1977] and thus a CH_4 atmospheric residence time of about 10 years.

The major sink of CH_4 is photochemical destruction via oxidation by OH [Levy, 1971]:

Calculations indicate that owing to destruction by reaction (1) in the troposphere, less than 10% of the CH_4 produced at the earth's surface enters the stratosphere. Note also that the oxidation of methane initiates a long series of reactions, known as the methane oxidation chain, which is believed to be a large source of tropospheric H_2 and CO and which may also play a major role in controlling the abundance of tropospheric ozone [Levy, 1972; Wofsy *et al.*, 1972; Crutzen, 1973, 1974a; Chameides and Walker, 1973, 1976]. There is, however, a possibility that the methane oxidation chain does not lead to CO exclusively; instead, the direct evolution of CO_2 may result, depending upon several formyl radical reaction paths (H. Niki, private communication, 1976).

The tropospheric CH_4 abundance is determined by a balance between biological production at the ground, transport upward, and destruction by reaction (1). In the steady state the CH_4 continuity equation may be written as

$$0 = -d/dz F(z) - k_1 n(\text{OH}) n(\text{CH}_4) \quad (2)$$

where F is the vertical methane flux, n is the number density, and k_1 is the rate coefficient for reaction (1), and we assume no net horizontal transport. Integrating (2) from $z = 0$ to $z = z_{tp}$, the tropopause, we obtain

$$F(0) - F(z_{tp}) = \int_0^{z_{tp}} [n(\text{OH}) n(\text{CH}_4) k_1] dz \quad (3)$$

where $F(z_{tp}) < 0.1F(0)$. Since CH_4 has a constant mixing ratio, it follows from (3) that

$$X_{\text{CH}_4} = [F(0) - F(z_{tp})]/D \sim F(0)/D \quad (4)$$

where X_{CH_4} is the equilibrium CH_4 mixing ratio, and that

$$D = \int_0^{z_{tp}} n(\text{OH}) n_M k_1 dz$$

where n_M is the background number density. Thus the tropospheric abundance of CH_4 is inversely proportional to the OH abundance.

As Chameides and Walker [1975] have pointed out, (4) implies that a perturbation in tropospheric OH will result in a change in the abundance of atmospheric CH_4 . Since CH_4 takes part in several key stratospheric reactions [Crutzen, 1974b] and also contributes significantly to the atmospheric 'greenhouse effect' [Wang *et al.*, 1976], a perturbation in atmospheric CH_4 could result in perturbations to stratospheric O_3 and the thermal balance of the atmosphere. In this work, two possible causes of a perturbation to atmospheric CH_4 are investigated: an increase in the anthropogenic production rate of CO and a variation in stratospheric ozone. Both processes cause a change in tropospheric OH, which then results in a variation in CH_4 . To determine the dependence of CH_4 upon these parameters, numerical calculations were made by using a steady state one-dimensional tropospheric model which couples photochemistry and transport, as described by Chameides and Stedman [1977] (their 'standard model'), with input data appropriate for 30°N latitude at equinox. In the calculations an iterative scheme with a 1% convergence test on all species was used to determine the mixing ratios of the long-lived gases, such as CH_4 , as a function of stratospheric O_3 or anthropogenic CO, given a fixed flux condition at the earth's surface. Note that in these steady state calculations we calculated the equilibrium CH_4 abundance. Since CH_4 has a residence time of about 10 years, it would probably take a decade or so for CH_4 to adjust to its equilibrium value after a perturbation. We have also assumed that CH_4 is presently in equilibrium and that the CH_4 production rate is constant at $1 \times 10^{11} \text{ cm}^{-2} \text{ s}^{-1}$. Note also that in our model, for every CH_4 molecule oxidized, a CO molecule is produced.

ANTHROPOGENIC CO AND ATMOSPHERIC CH₄

According to (4) a decrease in the concentration of OH would lead to an increase in atmospheric CH₄. A major sink of tropospheric OH is believed to be reaction with atmospheric CO [Weinstock, 1969; Crutzen, 1974a; Wofsy, 1976]:

Thus an increase in CO could lead to an increase in tropospheric CH₄ by causing a decrease in the OH abundance. Atmospheric CO is produced in nature via the CH₄ oxidation chain [McConnell *et al.*, 1971] at a rate of about 1×10^{11} CO molecules cm⁻² s⁻¹ according to our one-dimensional model calculations. While isotope data have been interpreted to indicate that at least 80% of the CO present in the atmosphere is produced by natural degradation of CH₄ [Stevens *et al.*, 1972], other evidence implies a much larger role for anthropogenic CO sources, as was discussed by Wofsy [1976]. Measurements indicate that approximately 3 times as much CO is present in the northern hemisphere, where anthropogenic CO production is largest, as is present in the southern hemisphere [Seiler, 1974]. Our model calculations, similar to the two-dimensional calculations of Wofsy [1976], indicate that in the southern hemisphere, with an average CO mixing ratio of 0.06 ppm, an approximate balance exists between CO production via CH₄ oxidation and CO destruction by reaction (1). In the northern hemisphere, however, with an average CO abundance of 0.15 ppm, a large imbalance was calculated between production and loss, possibly indicating the importance of anthropogenic CO production. For an average global CO mixing ratio of about 0.11 ppm we calculated a CO column destruction rate of 2.1×10^{11} cm⁻² s⁻¹ (1530 Mt yr⁻¹) and a CO production rate via CH₄ oxidation of 1×10^{11} cm⁻² s⁻¹ (730 Mt yr⁻¹). Since it is estimated that other natural CO sources produce about 3×10^{10} cm⁻² s⁻¹ (220 Mt yr⁻¹) [Seiler, 1974], this result implies that anthropogenic activities could produce as much as 8×10^{10} cm⁻² s⁻¹ (580 Mt yr⁻¹), or about 40% of the total CO production. This conclusion is supported by Seiler's [1974] estimate that anthropogenic CO sources produce about 9×10^{10} cm⁻² s⁻¹ (640 Mt yr⁻¹). If we included a CO soil sink of about 6×10^{10} cm⁻² s⁻¹ (450 Mt yr⁻¹), as extrapolated from laboratory experiments and reported by Seiler [1974], an additional CO source would be needed, although in the southern hemisphere a soil sink of 2×10^{10} cm⁻² s⁻¹, or 30% of the total sink, would not cause a significant imbalance in the southern hemispheric CO budget. However, as Seiler [1974] points out, since the uptake of CO by soils depends upon the CO density above the soil, the soil temperature, and the type of soil, the global uptake rate extrapolated from laboratory data is highly uncertain.

While further research is needed to determine more accurately the role of anthropogenic activities in the CO budget, models based on our present understanding of tropospheric photochemistry indicate that anthropogenic CO has already begun to perturb the atmosphere, and its effect will likely increase if anthropogenic CO production continues to grow. For instance, we calculated about 20% less OH in the northern hemisphere than in the southern hemisphere under present-day conditions owing to the enhanced CO concentration in the northern hemisphere, results similar to those of Wofsy [1976]. Since OH acts as a scavenger of many trace gases (e.g., NO_x, SO_x, NH₃, CH₃Cl), this diminution could be indicative of a serious problem.

As was discussed above, one likely effect of a decreasing OH density is an increase in atmospheric CH₄. The results of

Fig. 1. The dependence of X_{CH_4} , the equilibrium CH₄ abundance, upon FCO, the non-CH₄ CO source strength, and upon time, where we assumed that $\text{FCO} = 3 \times 10^{10} + 8 \times 10^{10}(1.045)^{t-1976}$ cm⁻² s⁻¹; i.e., the anthropogenic production rate is presently 8×10^{10} cm⁻² s⁻¹ and is increasing at an annual rate of 4.5%.

model calculations which determined the dependence of X_{CH_4} upon FCO, the non-CH₄ CO source strength, are presented in Figure 1, where $\text{FCO} = 1.1 \times 10^{11}$ cm⁻² s⁻¹ and $X_{\text{CH}_4} = 1.4$ ppm represent present-day conditions. These results confirm that as FCO increases, thereby increasing CO and decreasing OH, the CH₄ abundance increases. Also illustrated in Figure 1 is the variation of X_{CH_4} with time, where we assumed an anthropogenic production rate of 8×10^{10} cm⁻² s⁻¹ in 1976 and an annual increase of 4.5%, similar to that of CO₂ [Broecker *et al.*, 1971]. Given these assumptions, the photochemical model predicts that by the year 2000 the steady state CH₄ abundance will have increased by almost 40%. Of course, improved pollution control devices and/or a switch from fossil fuel burning to alternative power sources could slow the anthropogenic production of CO and thereby lessen a perturbation to atmospheric CH₄. In fact, pollution control in the United States has probably already slowed the global anthropogenic production of CO [Jaffe, 1973]. It is also possible that as the CO abundance increases, the magnitude of the soil sink for CO will also increase, thereby moderating the effect of anthropogenic activities. Finally, it is possible that future increases in anthropogenic NO_x emissions may also perturb OH levels and thus affect the CH₄ abundance. However, if CO production continues to grow, a significant increase in atmospheric CH₄ may result in the coming decades. Further measurements, especially of CO, CH₄, and OH, are needed to understand further the relationship between anthropogenic CO and atmospheric CH₄.

STRATOSPHERIC O₃ AND ATMOSPHERIC CH₄

Chameides and Walker [1975] have pointed out that a change in stratospheric O₃ can lead to significant perturbations in CH₄ by causing a change in the tropospheric OH density. The primary source of tropospheric OH radicals is believed to be the photodissociation of O₃ by radiation of wavelengths shorter than about 3100 Å, producing excited oxygen atoms:

TABLE 1. The Equilibrium CH₄ Mixing Ratio, X_{CH_4} (ppm vol/vol), as a Function of $N(O_3)$ and $f(O_3)$ (11 km)

O ₃ Mixing Ratio at 11 km, $f(O_3)$ (11 km)	Value of the O ₃ Column Above 10 km, $N(O_3)$, cm ⁻²				
	7×10^{18}	8×10^{18}	9×10^{18}	10×10^{18}	11×10^{18}
6.5×10^{-8}	1.2	1.5	1.8	2.1	2.5
9.5×10^{-8}	1.1	1.3	1.6	1.9	2.3
1.3×10^{-7}	0.96	1.2	1.4*	1.7	2.0
1.65×10^{-7}	0.84	1.0	1.2	1.5	1.7
2.0×10^{-7}	0.74	0.90	1.1	1.3	1.5

*Present-day conditions are given by $N(O_3) = 9 \times 10^{18}$ cm⁻², $f(O_3)$ (11 km) = 1.3×10^{-7} , and $X_{CH_4} = 1.4$ ppm.

followed by

[Levy, 1971].

Since the O(¹D) density can be altered either by changing the flux of near-UV radiation or by changing the local O₃ density, a perturbation to stratospheric O₃ can affect the OH and therefore also the CH₄ tropospheric abundance in two ways: by changing the flux of near-UV radiation penetrating the tropopause and by changing the O₃ density in the lower stratosphere and thus the injection rate of stratospheric O₃ into the troposphere. (Note that in our model, approximately 60% of the ozone found in the troposphere originates in the stratosphere, the remainder being produced locally by photochemical processes [Chameides and Stedman, 1977].) For instance, for equinoctial conditions at 30°N near the ground we calculated that a change in $N(O_3)$, the ozone column above 10 km, from 9×10^{18} to 8×10^{18} cm⁻² resulted in a change in the production of O(¹D) from 4.2×10^6 to 4.8×10^6 cm⁻³ s⁻¹. Similarly, by changing $f(O_3)$ (11 km), the O₃ mixing ratio at 11 km, from 1.30×10^{-7} to 1.65×10^{-7} and keeping $N(O_3)$ constant we obtained an increase in the O(¹D) production from 4.2×10^6 to 4.6×10^6 cm⁻³ s⁻¹. Table 1 illustrates the calculated dependence of X_{CH_4} upon both $N(O_3)$ and $f(O_3)$ (11 km). These results, which could be incorporated into stratospheric model calculations as a lower boundary condition upon CH₄, indicate that a significant CH₄ perturbation could result from a change in stratospheric O₃. For example, our calculations predict an approximate 30% decrease in X_{CH_4} if a 10% decrease in $N(O_3)$ is accompanied by a 25% increase in $f(O_3)$ (11 km). Note that stratospheric model calculations predict that a decrease in $N(O_3)$ is generally accompanied by an increase in the ozone density near the tropopause due to photolytic healing. However, Liu *et al.* [1976] found the opposite effect when they used a combination of rate coefficients which maximized the odd hydrogen density in the lower stratosphere.

Since stratospheric CH₄, which originates in the troposphere, takes part in several key photochemical reactions, the possibility exists for feedback effects whereby an initial O₃ perturbation is reduced or enhanced by a subsequent variation in atmospheric CH₄. CH₄, as a source of stratospheric odd hydrogen via reactions (1) and (7), may act to increase or decrease stratospheric O₃ [Crutzen, 1971; McElroy *et al.*, 1974; Rao-Vupputuri, 1974; Liu *et al.*, 1976]. While reactions

and

act as an O₃ sink,

tends to enhance O₃ by removing an odd nitrogen molecule from the catalytic chain which destroys O₃. Similarly, in a stratosphere with chlorine, CH₄ can enhance O₃ by removing Cl from its catalytic O₃-destroying chain

Whether CH₄ acts to decrease or increase stratospheric O₃ and thus whether the system described above will stabilize or destabilize stratospheric O₃ depend upon the importance of reactions (8) and (9) relative to reactions (10) and (11). Further study, both in the laboratory and with numerical models, is needed to help clarify this point.

CONCLUSION

Our calculations indicate that a 30–40% perturbation in atmospheric CH₄ may occur in the coming decades. The effects of a CH₄ perturbation are not clear, and further research is needed to understand fully the consequences of such a perturbation. Model calculations of the effects of CH₄ upon stratospheric O₃ are conflicting. The findings of Liu *et al.* [1976] and Rao-Vupputuri [1974] imply that an increase in CH₄ will result in a decrease in stratospheric O₃, while Crutzen [1971] and McElroy *et al.* [1974] apparently find the opposite effect. A careful reexamination of the importance of stratospheric CH₄ as an odd hydrogen source and of the photochemical role of odd hydrogen in the lower stratosphere is necessary to clarify this discrepancy.

Another possible consequence of a CH₄ variation is a perturbation to the thermal equilibrium of the atmosphere. Wang *et al.* [1976] have found that the infrared absorption bands of CH₄ in the 7- to 14-μm spectral region contribute to the atmospheric greenhouse effect. Their calculations using a one-dimensional radiative-convective model imply that a 0.3°K increase in surface temperature will result from doubling the CH₄ concentration. Thus a 30–40% perturbation in atmospheric CH₄ could have a significant climatic impact. Further calculations are necessary to establish more accurately the magnitude and nature of this climatic effect.

It is interesting to note that other long-lived species, such as H₂, CH₃Cl, CHFCl₂, and CHF₂Cl, which are attacked by OH in the troposphere, should undergo variations similar to those discussed for CH₄. These variations are particularly significant in the case of the Cl-containing molecules, since they may affect stratospheric O₃ [cf. Cicerone *et al.*, 1975].

In view of the remaining gaps in our understanding of the tropospheric photochemical system, large uncertainties are associated with our results, and our predictions should therefore be taken as rough estimates of possible future trends. In order to better understand tropospheric photochemistry and therefore the magnitude of future perturbations to atmospheric CH₄, further observational and theoretical research is necessary. A global program for measurement of OH, odd nitrogen, and CO levels is of primary importance.

Acknowledgment. This research was supported in part by the National Aeronautics and Space Administration under grant NSG-1363.

REFERENCES

- Baker-Blocker, A., T. M. Donahue, and K. H. Mancy, Methane flux from wetlands areas, *Tellus*, in press, 1977.
- Broecker, W. S., Y.-H. Li, and T.-H. Peng, Carbon dioxide—Man's unseen artifact, in *Impingement of Man on the Oceans*, edited by D. W. Hood, pp. 287–324, John Wiley, New York, 1971.

- Chameides, W. L., and D. H. Stedman, Tropospheric ozone: Coupling transport and photochemistry, *J. Geophys. Res.*, **82**, in press, 1977.
- Chameides, W. L., and J. C. G. Walker, A photochemical theory of tropospheric ozone, *J. Geophys. Res.*, **78**, 8751-8760, 1973.
- Chameides, W. L., and J. C. G. Walker, Stratospheric ozone: The possible effects of tropospheric-stratospheric feedback, *Science*, **190**, 1294-1295, 1975.
- Chameides, W. L., and J. C. G. Walker, A time-dependent photochemical model for ozone near the ground, *J. Geophys. Res.*, **81**, 413-420, 1976.
- Cicerone, R. J., D. H. Stedman, and R. S. Stolarski, Estimate of late 1974 stratospheric concentration of gaseous chlorine compounds (ClX), *Geophys. Res. Lett.*, **2**, 219-222, 1975.
- Crutzen, P. J., Ozone production in an oxygen-hydrogen-nitrogen oxide atmosphere, *J. Geophys. Res.*, **76**, 7311-7333, 1971.
- Crutzen, P. J., Gas-phase nitrogen and methane chemistry in the atmosphere, in *Physics and Chemistry of the Upper Atmosphere*, edited by B. M. McCormac, D. Reidel, Dordrecht, Netherlands, 1973.
- Crutzen, P. J., Photochemical reactions initiated by and influencing ozone in unpolluted tropospheric air, *Tellus*, **26**, 47-57, 1974a.
- Crutzen, P. J., A review of upper atmospheric photochemistry, *Can. J. Chem.*, **52**, 1569-1581, 1974b.
- Ehhalt, D. H., L. E. Heidt, R. H. Lueb, and N. Roper, Vertical profiles of CH₄, H₂, CO, N₂O, and CO₂ in the stratosphere, Proceedings of the Third Conference on GIAP, *Rep. DOT-TSC-OST-74-15*, pp. 390-402, U.S. Dep. of Transp., Washington, D. C., 1974.
- Jaffe, L. S., Carbon monoxide in the biosphere: Sources, distribution, and concentration, *J. Geophys. Res.*, **78**, 5293-5305, 1973.
- Koyama, T., Gaseous metabolism in lake sediments and paddy soils and the production of atmospheric methane and hydrogen, *J. Geophys. Res.*, **68**, 3971-3973, 1963.
- Levy, H., Normal atmosphere: Large radical and formaldehyde concentrations predicted, *Science*, **173**, 141-143, 1971.
- Levy, H., Photochemistry of the lower troposphere, *Planet. Space Sci.*, **20**, 919-935, 1972.
- Liu, S. C., T. M. Donahue, R. J. Cicerone, and W. L. Chameides, Effect of water vapor on the destruction of ozone in the stratosphere perturbed by ClX or NO_x pollutants, *J. Geophys. Res.*, **81**, 3111-3118, 1976.
- McConnell, J. C., M. B. McElroy, and S. C. Wofsy, Natural sources of atmospheric CO, *Nature*, **233**, 187-188, 1971.
- McElroy, M. B., S. C. Wofsy, J. E. Penner, and J. C. McConnell, Atmospheric ozone: Possible impact of stratospheric aviation, *J. Atmos. Sci.*, **31**, 287-303, 1974.
- Rao-Vupputuri, R. K., A zonally averaged circulation model of the stratosphere incorporating radiative heating and ozone photochemistry in an oxygen-hydrogen-nitrogen atmosphere, Proceedings of the Third Conference on GIAP, *Rep. DOT-TSC-OST-74-15*, pp. 390-402, U.S. Dep. of Transp., Washington, D. C., 1974.
- Seiler, W., The cycle of atmospheric CO, *Tellus*, **26**, 116-135, 1974.
- Stevens, C. M., L. Krout, D. Walling, and A. Ventes, The isotopic composition of atmospheric carbon monoxide, *Earth Planet. Sci. Lett.*, **16**, 147-165, 1972.
- Wang, W. C., Y. L. Yung, A. A. Lacis, T. Mo, and J. E. Hansen, Greenhouse effects due to anthropogenic perturbations, *Science*, **194**, 685-690, 1976.
- Weinstock, B., Carbon monoxide: Residence time in the atmosphere, *Science*, **166**, 224-225, 1969.
- Wofsy, S. C., Interactions of CH₄ and CO in the earth's atmosphere, *Annu. Rev. Earth Planet. Sci.*, **4**, 1976.
- Wofsy, S. C., J. C. McConnell, and M. B. McElroy, Atmospheric CH₄, CO, and CO₂, *J. Geophys. Res.*, **77**, 4477-4493, 1972.

(Received June 1, 1976;
revised December 6, 1976;
accepted December 6, 1976.)