

UCLA

UCLA Previously Published Works

Title

Individual differences in pain sensitivity are associated with cognitive network functional connectivity following one night of experimental sleep disruption

Permalink

<https://escholarship.org/uc/item/3bd2m7rk>

Journal

Human Brain Mapping, 41(3)

ISSN

1065-9471

Authors

Letzen, Janelle E
Remeniuk, Bethany
Smith, Michael T
[et al.](#)

Publication Date

2020-02-15

DOI

10.1002/hbm.24824

Peer reviewed



HHS Public Access

Author manuscript

Hum Brain Mapp. Author manuscript; available in PMC 2020 February 15.

Published in final edited form as:

Hum Brain Mapp. 2020 February 15; 41(3): 581–593. doi:10.1002/hbm.24824.

Individual differences in pain sensitivity are associated with cognitive network functional connectivity following one night of experimental sleep disruption

Janelle E. Letzen¹, Bethany Remeniuk¹, Michael T. Smith¹, Michael R. Irwin², Patrick H. Finan^{1,†}, David A. Seminowicz^{3,†}

¹Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, Maryland ²Cousins Center for Psychoneuroimmunology, UCLA Semel Institute for Neuroscience and Human Behavior, Los Angeles, California ³Department of Neural and Pain Sciences, School of Dentistry, and Center to Advance Chronic Pain Research, University of Maryland Baltimore, Baltimore, Maryland

Abstract

Previous work suggests that sleep disruption can contribute to poor pain modulation. Here, we used experimental sleep disruption to examine the relationship between sleep disruption-induced pain sensitivity and functional connectivity (FC) of cognitive networks contributing to pain modulation. Nineteen healthy individuals underwent two counterbalanced experimental sleep conditions for one night each: uninterrupted sleep versus sleep disruption. Following each condition, participants completed functional MRI including a simple motor task and a noxious thermal stimulation task. Pain ratings and stimulus temperatures from the latter task were combined to calculate a pain sensitivity change score following sleep disruption. This change score was used as a predictor of simple motor task FC changes using bilateral executive control networks (RECN, LECN) and the default mode network (DMN) masks as seed regions of interest (ROIs). Increased pain sensitivity after sleep disruption was positively associated with increased RECN FC to ROIs within the DMN and LECN ($F_{(4,14)} = 25.28$, $pFDR = 0.05$). However, this pain sensitivity change score did not predict FC changes using LECN and DMN masks as seeds ($pFDR > 0.05$). Given that only RECN FC was associated with sleep loss-induced hyperalgesia, findings suggest that cognitive networks only partially contribute to the sleep-pain dyad.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Correspondence: Janelle E. Letzen, Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, 5510 Nathan Shock Dr., Suite 100, Baltimore, MD 21224. jletzen1@jhmi.edu.

[†]Shared senior authorship.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

CONFLICT OF INTEREST

The authors declare no competing financial interests.

Keywords

default mode network; executive control network; neuroimaging; pain sensitivity; sleep disruption

1 | INTRODUCTION

Evidence consistently shows that sleep disturbance leads to a variety of neurobiological and behavioral adaptations (Bosch et al., 2013; Short & Banks, 2014; Verweij et al., 2014). Even one night of total sleep deprivation is associated with brain changes, especially in neural networks underlying cognitive control, affect, and memory (Krause et al., 2017). However, these neurobiological differences are not always linked with detrimental behavioral responses. Some individuals actually benefit from acute sleep disturbance (Chuah, Venkatraman, Dinges, & Chee, 2006; Sletten, Segal, Flynn-Evans, Lockley, & Rajaratnam, 2015), begging the question: what mechanisms drive diverging responses to various types of sleep loss?

Among aberrant sequelae associated with sleep loss, maladaptive changes in pain perception represent one of the largest public health concerns. First, emerging evidence suggests that chronic sleep loss predicts new cases of chronic pain and exacerbations of existing pain symptoms (see Finan, Goodin, & Smith, 2013 for a review). Second, there is a high prevalence and rate of comorbidity between chronic pain and chronic sleep disturbance (Alsaadi et al., 2014; Karaman et al., 2014; Koffel et al., 2016; Tang, Wright, & Salkovskis, 2007). There are many protocols to experimentally alter sleep in order to explore mechanisms of the sleep-pain dyad, such as total sleep deprivation (i.e., preventing individuals from sleeping at all through >24-hr period), sleep restriction (i.e., limiting sleep under 8 hr), and forced awakenings (i.e., interrupting sleep at various intervals throughout the night). Because chronic pain patients tend to experience difficulty staying asleep—characterized by multiple night time awakenings (i.e., sleep maintenance insomnia)—the forced awakenings paradigm is a particularly relevant tool for understanding the relationship between sleep disruption and pain sensitivity (Rosseland, Pallesen, Nordhus, Matre, & Blågestad, 2018).

Pain itself is a complex experience that is influenced by sensory, affective, and cognitive processes (Apkarian, Bushnell, Treede, & Zubieta, 2005; Bushnell, eko, & Low, 2013; Price, 2002). While various networks contribute to this perception, brain regions associated with cognitive control are critical for our ability to modulate pain. Notably, cognitive control exerts top-down influence over sensory and affective pain components (Seminowicz & Ceko, 2015). For this reason, the present study focused on the relationship between sleep disruption-induced hyperalgesia and function across networks associated with cognitive control.

Two key networks involved in cognitive pain control are the executive control network (ECN) and default mode network (DMN). The ECN predominately includes the dorsolateral prefrontal cortex (dlPFC) and posterior parietal areas, and the DMN is composed of the medial prefrontal cortex (mPFC), posterior cingulate cortex/precuneus, and inferior parietal lobules. Functionally, the ECN is linked with attention to relevant information and

suppression of irrelevant information (Awh, Vogel, & Oh, 2006; Gazzaley & Nobre, 2012; Smith et al., 2018), and the DMN is associated with self-referential thought and focus on internal milieu (Andrews-Hanna, Reidler, Sepulcre, Poulin, & Buckner, 2010; Lorenzi et al., 2011; Raichle, 2015).

Previous work suggests that the functional dynamics of these networks can contribute to the detection and inhibition of nociceptive inputs. For example, greater ECN-mPFC functional connectivity (FC) was associated with better cognitive modulation of pain among healthy individuals (Kong et al., 2013). Commensurate with this finding, another study demonstrated greater ECN-DMN FC in individuals who learned and practiced cognitive modulation strategies over 21 days compared to participants who completed a treatment control (Kucyi, Salomons, & Davis, 2016). These results suggest that ECN-DMN FC is positively associated with pain modulation and is responsive to intervention.

Interactions between these networks are susceptible to experimental sleep manipulation. In this regard, greater ECN-DMN FC has been documented in healthy individuals after one night of experimental sleep deprivation (Bosch et al., 2013). Additional measures reflecting increased functional coupling among DMN and ECN nodes, such as a reduction in anticorrelation and loss of functional segregation, following sleep deprivation have also been reported (De Havas, Parimal, Soon, & Chee, 2012; Ben Simon, Lahav, Shamir, Hendler, & Maron-Katz, 2018). Per se, sleep deprivation is associated with the separate phenomena of hyperalgesia and greater ECN-DMN coupling; however, it is unclear how sleep disruption via forced awakenings might influence the relationship between ECN and DMN.

The present study used one night of forced awakenings to determine the effects on pain sensitivity and cognitive networks associated with pain modulation. The limited timeframe of the sleep manipulation used in this study may help to identify early brain responses to sleep disruption that contribute to sleep loss-related hyperalgesia. Based on the complex literature described above, we hypothesize that changes in ECN-DMN FC following sleep disruption will be associated with changes in pain sensitivity. However, given that previous studies suggest that the association between changes in ECN-DMN FC and pain sensitivity could be either positive or negative, the potential direction of this association remains unclear.

2 | METHODS

The present analyses were completed on data from an fMRI task that was part of a larger neuroimaging protocol used to examine the effects of one night of experimental sleep disruption via forced awakenings on the neural correlates of pain sensitivity and affective pain modulation in healthy individuals (Seminowicz et al., 2019). The primary data examined in the present study include (a) pain ratings and stimuli temperatures from a noxious thermal stimulation task (i.e., we did not use fMRI data from this task) and (b) a simple motor task (SMT) designed to elicit intrinsic network connectivity without the risk of participants falling asleep. Pain ratings and stimuli temperatures were computed in a change score (described below) and used as a predictor of individual differences in pain sensitivity

following sleep loss. More information about the task used to derive the pain ratings and stimuli temperatures is provided in Supporting Information.

The neuroimaging protocol was further nested within a larger study (R01DA032922) that included additional nights for each sleep conditions and separate daytime procedures conducted on the day following the MRI scan; these data are not examined for the present study. All procedures were approved by Institutional Review Boards (IRB) at Johns Hopkins University and University of Maryland, Baltimore in conformation to the ethical guidelines of the 1975 Declaration of Helsinki. All participants provided written informed consent to complete study procedures and were notified of their right to withdraw from study procedures at any time without penalty. Figure 1 demonstrates the methods used in the present study.

2.1 | Participants

SMT fMRI data were collected from 22 healthy individuals after uninterrupted and experimentally disrupted sleep. Participants were eligible if they met the following criteria: (a) met Research Diagnostic Criteria for Normal Sleepers, (b) scored <5 on the Pittsburgh Sleep Quality Index and <10 on the Epworth Sleepiness scale, (c) had average total sleep time between 6.5 and 8.5 hr/night and sleep efficiency >85% as reported on 2 weeks of sleep diaries and captured via actigraphy monitoring, (d) showed a stable sleep phase within 21:00 and 10:00, (e) polysomnography-confirmed apnea-hypopnea index <10, (f) denied a lifetime history of pain persisting >6 months or acute pain as measured on the McGill Pain Questionnaire and baseline sleep diaries, (g) reported no significant medical or psychiatric morbidity within 6 months of study participation and obtained *T*-scores <64 on the Brief Symptom Inventory global scales, (h) no lifetime history of bipolar disorder, psychotic disorder, recurrent major depression, posttraumatic stress disorder, substance abuse, reported history of traumatic brain injury, or seizures, (i) reported no tobacco/nicotine use and low caffeine use (<2 cups per day), (j) had a BMI <35, and (k) no MRI contraindications.

The present analyses were completed on data from 19 individuals (14 women; mean age = 23.95 years ($SD = 4.17$)) after two participants were excluded for incomplete fMRI data, and one individual was found to be an outlier with influential, extreme values for pain ratings. Seven participants identified as White Non-Hispanic, two individuals identified as White Hispanic, five participants identified as Black Non-Hispanic, one individual identified as Black Hispanic, two participants identified as mixed race, and one individual identified as Other Hispanic. Participant demographics and psychosocial function are listed in Table 1.

2.2 | Experimental design and statistical analyses

2.2.1 | Sleep monitoring and manipulation—Participants completed two nights of sleep monitoring at the Johns Hopkins Clinical Research Unit (CRU). Neuroimaging data analyzed for this study were collected following a night of 8-hr sleep opportunity (uninterrupted sleep, US) and following a night of sleep fragmentation via forced awakenings (FA). The order of these study visits was counterbalanced among participants to control for temporal confounds.

Uninterrupted sleep: For the US condition, participants were given the opportunity for 8 hours of sleep at the CRU without experimental disruption.

Sleep disruption procedure: The present study used a previously described sleep disruption protocol (see Finan et al., 2017, for a complete description). We chose sleep disruption (i.e., FA) based on previous work suggesting that this type of paradigm results in greater impairment of endogenous pain modulation compared to sleep restriction (Smith, Edwards, Mccann, & Haythornthwaite, 2007).

Briefly, the night was divided into eight 1-hr intervals, with one randomly chosen interval in which no sleep was allowed. The remaining seven 1-hr intervals were further divided into 20-min tertiles, so that one tertile per hour interval was randomly selected for a forced awakening. CRU staffs were responsible for waking up participants at these randomly chosen times and ensuring that participants remained awake throughout the interval. The maximum sleep allowed for the FA night was 280 min.

2.2.2 | MRI session—All structural and functional MRI scanning was completed at the University of Maryland, Baltimore Magnetic Resonance Research Center between 1 p.m. and 5 p.m. Data were acquired using a Siemens 3T Tim Trio scanner with a 32-channel head coil. Structural images were collected using a high-resolution T1-weighted MPRAGE anatomical scan with the following parameters: 240 slices, TR 2,300 ms, TE 2.98 ms, flip angle 9°, FOV 256 mm, matrix 256 × 256 mm, resolution 1.0 × 1.0 mm, slice thickness 1 mm, no gap. Participants also completed 3 fMRI runs of an experimental pain task, during which a noxious thermal stimulus was applied to the left medial forearm (fully described in Seminowicz et al. (2019) and Supporting Information). Neuroimaging data from this task were not used in the present study, but pain ratings aggregated over 15 stimulation periods were used to derive the predictor variable for FC analyses. Please see Supporting Information for complete methods related to the noxious thermal stimulation task.

The present study aimed to examine changes in FC as they related to individual differences in pain sensitivity before and after sleep disruption. Due to concerns that participants might fall asleep in the scanner using a traditional resting-state task, we instead collected data using a 10-min SMT, which required participants to respond to the direction of a projected arrow (left or right). Arrows were sequentially displayed for 500 ms seconds at a time, with a total of 150 trials. Left and right arrows were shown in equal proportion. Interstimulus intervals were jittered and ranged from 1s to 4 s each. Task stimuli were programmed and run using Presentation (Neurobehavioral Systems, Berkeley, CA) on a desktop computer in the scanner suite. Participants used an angled mirror mounted on the MRI head coil to view task images via a translucent screen with back-project images. The following instructions were provided: *During this time, we will help you stay awake by having you engage in a simple task that involves using your response box. You will look up and see a screen, and on the screen an arrow pointing either left or right will flash periodically. When the right facing arrow flashes, you should press 3 (the button farthest to the right). When the left facing arrow flashes you should press 1 (the button farthest to the left).*

Previous work has shown minimal FC differences between traditional resting-state tasks (i.e., acquired during a steady-state period of non-goal-directed activity) and simple motor tasks, such as the one we employed (Baliki, Geha, Apkarian, & Chialvo, 2008; Jurkiewicz, Crawley, & Mikulis, 2018). At each visit, the SMT scanning sequence included an interleaved T2*-weighted, echo planar imaging (EPI) sequence with the following parameters: spin-echo, 658 volumes/run, 66 slices, TR: 907 ms, TE: 30 ms, flip angle 52°, FOV 224 mm, matrix size 110 × 102 mm, resolution 2.0 × 2.0 mm, slice thickness 2.0 mm, multiband factor 6.

2.2.3 | Pain sensitivity change score—The present study aimed to examine changes in FC among cognitive networks as it related to sleep loss-induced hyperalgesia. In order to examine this aim, we computed a pain sensitivity change score as a predictor variable. For the parent study, participants underwent a quantitative sensory testing session before scanning at both study visits to derive participant-tailored temperatures for the fMRI noxious stimulation task. Temperatures rated between 5 and 8 out of 10 were selected. During fMRI scanning, these tailored temperatures were presented concomitant with music of neutral valence. Pain ratings were made following the delivery of each noxious thermal stimulus, which ended at the same time the music clip ended. The same procedure was employed at each study visit (see Supporting Information for additional details). Due to the aims of the parent study, music stimuli were paired with each noxious thermal stimulus. The parent study also overlaid positive-valence song clips on noxious thermal stimuli in separate blocks, but pain ratings associated with those blocks were not used in the present study because they were associated with pain inhibition (Seminowicz et al., 2019).

A pain sensitivity change score was calculated using pain ratings and tailored temperatures. Temperatures ranged from 40°C to 50°C, so we subtracted 40 from each temperature denominator to make the scales of pain ratings and temperatures equal (0–10). Our approach of subtracting 40 from temperature values was intended for scaling purposes only and does not assume an association between the temperature value and pain rating value.

Some individuals experienced changes in both temperature and pain ratings (e.g., lowered temperatures with higher pain ratings after FA sleep would indicate increased pain sensitivity). As such, we calculated a pain sensitivity change score using the following equation, with higher values indicating greater pain sensitivity following FA:

$$\text{pain sensitivity change score} = 1 - \frac{[\text{US pain rating} \div (\text{US temperature}-40)]}{[\text{FA pain rating} \div (\text{FA temperature}-40)]}$$

2.2.4 | Behavioral data analyses—Using SPSS 25 (IBM), we ran paired-samples *t* tests to examine condition-based differences in the following variables: total sleep time (TST), stage N1, stage N2, slow wave sleep (SWS), rapid eye movement sleep (REM), Stanford Sleepiness Scale (SSS), SMT percent accuracy, pain ratings collected during the noxious thermal stimulation task, and thermal stimuli temperatures used during the noxious thermal stimulation task. No transformations were applied to these data.

2.2.5 | Neuroimaging data processing

Functional MRI data preprocessing: SMT data were preprocessed in SPM12 v6906. The preprocessing pipeline included realignment (motion correction), coregistration of the T1 anatomical to the mean functional image, segmentation of the anatomical to six standard tissue probability maps in SPM12, normalization of functional images using forward deformation fields from anatomical segmentation, and spatial smoothing using a 6 mm full-width at half maximum (FWHM) kernel. Data were visually inspected for quality control to determine motion artifact. Further, the Artifact Detection Tool (ART; www.nitrc.org/projects/artifact_detect/) was also used to mark potentially confounding outliers in the SMT time series for exclusion (i.e., rotation $>.02$ rad from prior volume, or translation $>.4$ mm from the prior volume (Chai, Ofen, Gabrieli, & Whitfield-Gabrieli, 2014). No condition-based differences in outlier detection ($p = .4$) or motion correction ($p = .38$) were identified.

We used the CONN toolbox (Whitfield-Gabrieli & Nieto-Castanon, 2012) to complete denoising procedures and FC analyses. Gray matter, white matter, and CSF masks were created via anatomical segmentation and were eroded by one voxel to minimize partial volume effects (Whitfield-Gabrieli & Nieto-Castanon, 2012). Physiological confounds, including respiratory and cardiac activity, were accounted for by using the white matter and CSF masks as nuisance regressors during denoising (Behzadi, Restom, Liau, & Liu, 2007). Denoising including scrubbing of excessive motion and outliers detected via ART and regression of white matter and CSF masks (% volumes removed: mean = 1.9%, range = 0.53–5.34%). A stepwise, rather than simultaneous, regression approach was applied to maintain adequate degrees of freedom of the resting-state data (Patel & Bullmore, 2016). Data were bandpass filtered between .008 and .09 (Aurich, Alves Filho, Marques da Silva, & Franco, 2015).

Hybrid independent component analysis-seed-based functional connectivity: We conducted hybrid independent component analysis (ICA)-seed-based FC analyses on denoised fMRI data in the CONN toolbox. Previous work suggests that hybrid ICA-seed-based FC analyses result in better reproducibility of fMRI results than either ICA or seed-based analyses alone (Kelly et al., 2010). First, group ICA is a data-driven technique that identifies orthogonal sources of variance within the data (Calhoun, Adali, Pearlson, & Pekar, 2001), yielding independent components (ICs). Each IC contains brain regions with activity co-occurring across a unique time course, thought to represent a neural network (Calhoun, Adali, & Pekar, 2004).

Consistent with steps described by Calhoun et al. (Calhoun et al., 2001), denoised data first underwent participant- and condition-level BOLD signal concatenation. We estimated the optimum number of ICs within the GIFT toolbox using minimum description length criterion (Li, Adali, & Calhoun, 2007), yielding 20 potential ICs. Dimensionality reduction was then conducted at the group-level to identify these 20 dimensions/ICs. To estimate IC maps at the group-level, the fastICA algorithm was used. Group-level IC maps are then back reconstructed on individual-level data via GICA1 backprojection to create individual-level IC spatial maps that can be pooled for second-level GLM statistics; however, only the group-level ICs were used to create seeds for subsequent analyses. Resultant group-level ICs were

sorted to freely available spatial templates of RECN, LECN, and the DMN (Shirer, Ryali, Rykhlevskaia, Menon, & Greicius, 2012). The IC with the strongest match to each template (based on the Dice coefficient) was converted into a binarized map used in subsequent seed-based analyses (Table 2; Figure 2).

Binarized IC masks were added to the CONN toolbox as a priori regions of interest (ROIs). Further, the Power 264-region atlas was used to define 10 mm ROI spheres across the whole brain (Power et al., 2011). This whole-brain atlas describes subgraphs of functionally integrated nodes as defined by graph theory analyses on resting-state fMRI data, rather than structurally defined. First-level analyses were completed in CONN to extract time series data within each ROI and compute Fisher's r-to-Z transformed bivariate correlations among these ROIs.

GLM analyses: Contrast images were entered into a second-level seed-to-ROI analysis to determine FC differences between US and FA across the sample. The pain sensitivity change score described above was inserted as a regressor of interest, and RECN, LECN, and DMN masks were defined as seeds. In total, there were six primary sets of GLM tests, including three seed-to-ROI tests (i.e., RECN, LECN, and DMN as seeds) and three tests examining associations among RECN, LECN, and DMN masks (i.e., RECN-LECN, LECN-DMN, RECN-DMN). ROIs in the seed-to-ROI analyses included 264 spheres from the Power 264 atlas. A conservative analysis-level threshold across seeds was applied as a correction for multiple comparisons ($pFDR < .05$). We used a two-tailed test given the nature of our competing directional hypotheses.

2.2.6 | Data availability statement—We have shared the resultant ROI maps from the present study in neurovault.org (Gorgolewski et al., 2015), which is a public repository for neuroimaging statistical maps. Data are located at the following link: <https://neurovault.org/collections/ALMIXGCQ/>

2.2.7 | Validation analyses—Given the small sample size, we attempted to replicate findings using a distinct dataset. These analyses were exploratory in nature and not part of the primary aim. However, the complex and unique experimental approach used in our study limited our ability to test replication in an experimentally matched dataset. Instead, we examined whether clinical pain ratings moderated the relationship between sleep quality (i.e., Pittsburgh Sleep Quality Inventory scores) and FC values that resulted from our primary analyses. This validation sample included 30 individuals with migraine who reported poor sleep quality (i.e., PSQI >5). Methods from the study associated with our validation sample are reported at the following link: <https://clinicaltrials.gov/ct2/show/NCT02133209>.

3 | RESULTS

3.1 | Sleep manipulation check

To examine the efficacy of our sleep disruption manipulation we examined condition-based differences in several sleep parameters. As expected, there was a significant decrease in minutes of TST, stage N2, SWS, and REM sleep phases, and a significant increase in stage

N1 sleep. Participants reported significantly greater sleepiness during the FA compared to US condition. Table 3 provides means, *SD* values, and paired-samples *t* test results for these variables. Combined, these results indicate the efficacy of the experimental manipulation.

3.2 | Pain ratings, thermal stimuli temperatures, and SMT performance

Table 3 lists statistical information about condition-based differences in pain ratings, thermal stimuli temperatures, and SMT performance. Neither pain ratings nor thermal stimuli temperatures significantly differed between the US and FA conditions. On the SMT, participants' accuracy in identifying arrow directions was better during US compared to FA (74.8% and 63.3% accuracy, respectively). The low accuracy scores were due to missed trials or incorrect responses.

3.3 | Left and right ECN and DMN functional connectivity

3.3.1 | RECN as a seed—Several ROIs from the Power atlas demonstrated significant FC changes with the RECN seed associated with pain sensitivity change scores (omnibus model: $F_{(4,14)} = 25.28$, $pFDR = .05$; individual ROI statistics reported in Table 4; Figure 4). Specifically, there was a positive association between increased pain sensitivity and increased RECN FC to left dlPFC (part of the LECN), left medial and inferior frontal gyri (MFG, IFG, respectively; part of the DMN), and the right paracingulate gyrus (part of the DMN). There was also a positive association between increased pain sensitivity and decreased RECN FC to the right lateral occipital cortex (LOC; part of the DMN). Table 4 provides additional information about these ROIs, and Table 5 provides descriptive statistics for FC values based on condition.

3.3.2 | Null results—Although the RECN seed demonstrated significant associations between six RECN-ROI pairs and pain sensitivity change scores, the additional five primary analyses did not pass our significance threshold corrected for multiple comparisons. Specifically, neither the LECN nor DMN seed demonstrated significant seed-to-ROI FC associations with pain sensitivity changes ($pFDR > .05$). Further, FC between ECN and DMN seeds did not show significant pairwise associations between FC and pain sensitivity change scores ($pFDR$: RECN-LECN = .06, RECN-DMN = .16, LECN-DMN = .63).

3.4 | Validation analyses

We tested whether clinical pain ratings moderated the relationship between PSQI scores and FC values from the six significant RECN-ROI pairs on a separate sample of migraine patients who reported poor sleep quality. Among these six ROIs, there was only one significant moderation effect of pain on the relationship between RECN-right paracingulate gyrus ($F_{(1,29)} = 6.4$, $p = .02$). The other RECN-ROI moderation analyses were not significant (p s range = .37–.91). Because this was not an ideal validation sample, however, we do not further interpret results.

4 | DISCUSSION

The present study examined the association between cognitive network FC and individual differences in sleep loss-induced hyperalgesia following one night of sleep disruption. In a

healthy sample of normal sleepers, we used an experimental manipulation to produce multiple, prolonged FA during one night of sleep. We broadly hypothesized that neural systems associated with cognitive pain control would be affected by sleep loss and that these changes would be related to changes in pain sensitivity. Overall, we saw variability in the magnitude and direction of changes in participants' pain sensitivity following one night of acute sleep loss, with an overall positive association between pain sensitivity changes and changes in RECN FC to several brain regions (i.e., dIPFC, MFG, IFG, paracingulate gyrus, and LOC). These results, however, are qualified by null findings across five additional a priori tests. Taken together, results only partially support our hypothesis of greater ECN-DMN FC in individuals with higher pain sensitivity following sleep disruption, suggesting that cognitive network alterations only partially contribute to sleep loss-induced hyperalgesia.

4.1 | Pain sensitivity following forced awakenings

When combining pain ratings and thermal stimuli temperatures, some individuals displayed hyperalgesia while others showed decreased pain sensitivity. Even across pain ratings, which were obtained following tonic noxious thermal stimuli that were paired with neutral-valenced music, there were no overall mean changes in pain sensitivity across the sample after a single night of FA sleep. Although these findings are somewhat inconsistent with previous research demonstrating hyperalgesia following sleep loss, it should be noted that most of those studies were conducted over multiple nights of accumulated sleep loss. In fact, this study's parent project recently found that two nights of sleep disruption via FA-induced hyperalgesia in a larger sample ($n = 79$), reducing both heat pain threshold and cold pain tolerance. The present study's design using one night of FA did not induce a ubiquitous hyperalgesic response; however, results cautiously show the early changes in RECN FC contributes to sleep disruption-induced hyperalgesia.

Further, differences in quantitative sensory testing across studies make comparisons across studies challenging. In the present study, we analyzed pain ratings obtained following the application of tonic noxious thermal stimuli that were paired with neutral-valenced music, as the parent study design did not include noxious thermal stimuli presented alone (see Seminowicz et al., 2019). As with many quantitative sensory testing studies, the uniqueness of our study design may limit the generalizability of the present results. We encourage future replication attempts. Cautiously, these results suggest varied responses to acute sleep loss in healthy adults, with some individuals evidencing greater pain vulnerability to sleep loss, which may have more significant clinical implications (Lautenbacher, Kundermann, & Krieg, 2006).

4.2 | RECN functional connectivity changes following forced awakenings

Our study specifically examined FC of bilateral ECNs and DMN to functionally defined ROIs throughout the brain (Power et al., 2011), and FC among bilateral ECN and DMN masks. The ECN is a group of brain regions associated with top-down modulation of sensory stimuli via goal-directed attentional amplification of relevant information, as well as goal-directed suppression of irrelevant information (Awh et al., 2006; Gazzaley & Nobre, 2012; Smith et al., 2018). Importantly, the ECN is engaged during cognitive pain control in

individuals with and without chronic pain (Ceko et al., 2015; Ceko et al., 2015; Kong et al., 2013; Rainville, 2002; Sevel, Letzen, Staud, & Robinson, 2016; Villemure & Bushnell, 2002). The DMN, on the other hand, is a group of brain regions associated with self-referential thought and focus on internal milieu (Andrews-Hanna et al., 2010; Lorenzi et al., 2011; Raichle, 2015). Its activity is typically anticorrelated with ECN activity during cognitive tasks so that better suppression of the DMN is associated with better cognitive performance (Fox et al., 2005; Seminowicz & Davis, 2007; Smith, Sip, & Delgado, 2015). Because evidence links ECN and DMN interactions with cognitive control of sensory inputs that are susceptible to perturbations from sleep disruption, the combined previous findings were suggestive of ECN-DMN FC as a putative mechanism for sleep loss-induced hyperalgesia.

Among tested networks in the present study, however, changes in pain sensitivity were only associated with changes in RECN FC, and we did not observe significant changes in the association between pain sensitivity and ECN-DMN FC. Instead, there was a positive association between increased pain sensitivity following sleep deprivation and increased RECN FC to specific regions within the DMN and LECN as well as decreased RECN FC to a primary visual network region as a function of changes in pain sensitivity. These regions included the left dlPFC (LECN), left MFG/IFG and right paracingulate gyrus (DMN), and right LOC (primary visual network).

The dlPFC is among the brain regions commonly associated with pain vigilance, awareness, and attention (Lorenz, Minoshima, & Casey, 2003; Seminowicz et al., 2013; Seminowicz & Davis, 2007; Seminowicz & Moayedi, 2017; Valet et al., 2004). Findings from experimental sleep research suggest that individual differences in emotional responses following sleep loss are associated with changes in attention (Alfarra, Fins, Chayo, & Tartar, 2015). Attentional processes are well documented in the context of experimental pain so that pain vigilance contributes to exacerbated pain intensity and distraction is associated with lower pain report (for a review, see Wiech, Ploner, & Tracey, 2008). Individuals sleeping less than 6.5 hr per night have also been found to have impaired ability to derive analgesic benefits from distraction techniques (playing video games) during an experimental pain task. Further, because stronger anticorrelation between DMN-ECN activity is associated with better segregation of internally and externally directed awareness (Buckner, 2008; Horowitz et al., 2009; Larson-Prior et al., 2009), increased RECN-DMN FC—as identified in this study—potentially suggests poorer ability to modulate attention. The combined findings of increased intrinsic, inter-hemispheric dlPFC FC and RECN FC to regions within the DMN might represent heightened attention to sensory stimuli, contributing to greater pain sensitivity.

It is not fully clear why we observed lateralization effects of significantly altered RECN, but not LECN, FC. Although debated, there is some evidence suggesting a greater propensity of right hemisphere activity in relation to pain processing (Coghill, Gilron, & Iadarola, 2001; Ji & Neugebauer, 2009; Symonds, Gordon, Bixby, & Mande, 2006). Additionally, a previous study examining resting-state attentional networks in individuals with insomnia found altered right, but not left, hemisphere FC (Li et al., 2018). Although it is possible that RECN FC is one mechanism contributing to sleep loss-induced hyperalgesia, the fact that this was the only seed to demonstrate significant findings across all six a priori tests tempers the

strength of inference. Future research is encouraged to measure whether other neural networks contribute to sleep loss-induced hyperalgesia.

4.3 | Limitations

The present results should be interpreted in the context of several important limitations. First, the small sample size used in the present study, coupled with the high number of a priori tests, weakens the strength of inference. Second, previous studies using the FA protocol over two nights showed it maximally induced hyperalgesia (Iacovides, George, Kamerman, & Baker, 2017) and diminished pain inhibition (Finan et al., 2017; Smith et al., 2007). In the present study, we used a smaller time window (one night) than previously reported, which might account for our lack of robust hyperalgesia findings. Our previous data suggest that more than one night of FA sleep is beneficial for examining sleep-pain responses. Third, although the FA paradigm is thought to have face validity for the type of sleep disturbance profile commonly observed in patients with chronic pain (i.e., multiple prolonged awakenings with significant concomitant reduction in total sleep time) (Rosseland et al., 2018), it may not generalize fully to sleep disruption linked with chronic insomnia, which may be due to a variety of clinical factors that influence pain in addition to the multiple awakenings themselves. Our results should be extended to include patient populations that suffer from sleep loss. Fourth, as mentioned earlier, pain ratings were collected from a paradigm using neutral music overlay, so we could not rule out the possible influence of distraction on imaging findings. The primary findings should be replicated in a design that includes thermal stimuli presented alone. Fifth, we did not assess individuals' cognitive strategies used to control pain. Given that there are individual differences in cognitive pain control strategies (Seminowicz, Mikulis, & Davis, 2004), future studies should measure how ECN-DMN FC patterns relate to strategy use following sleep disturbance.

5 | CONCLUSION

The present findings suggest that healthy individuals have idiographic pain responses to one night of sleep disruption and that these responses are partially associated with changes in RECN FC to regions within the LECN and DMN. Future prospective studies are needed to examine whether additional neural networks further contribute to the sleep-pain dyad and whether RECN FC changes might act as one mechanism potentially involved in the chronification of pain via sustained sleep disruption.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

This research was supported with funds provided by NIH F32 HL143941 (JEL); NIH K23 DA035915 (PHF); Blaustein Pain Research Award (PHF); NIH R01 DA0329922 (MTS, MRI); NIH P30 NR014131 (MTS and PHF); and NIH T32 NS7020110 (BR, MTS). We would also like to thank Samuel Krimmel and Shana Burrowes for their time and guidance in preparing the present analyses.

Funding information

National Institutes of Health, Grant/Award Numbers: NIH F32 HL143941, NIH K23 DA035915, NIH P30 NR014131, NIH R01 DA0329922, NIH T32 NS7020110; Blaustein Pain Research Award

REFERENCES

- Alfarra R, Fins AI, Chayo I, & Tartar JL (2015). Changes in attention to an emotional task after sleep deprivation: Neurophysiological and behavioral findings. *Biological Psychology*, 104, 1–7. [PubMed: 25451382]
- Alsaadi SM, McAuley JH, Hush JM, Lo S, Bartlett DJ, Grunstein RR, & Maher CG (2014). The bidirectional relationship between pain intensity and sleep disturbance/quality in patients with low back pain. *The Clinical Journal of Pain*, 30, 755–765. [PubMed: 24451630]
- Andrews-Hanna JR, Reidler JS, Sepulcre J, Poulin R, & Buckner RL (2010). Functional-anatomic fractionation of the brain's default network. *Neuron*, 65, 550–562. [PubMed: 20188659]
- Apkarian AV, Bushnell MC, Treede R, & Zubieta J (2005). Human brain mechanisms of pain perception and regulation in health and disease. *European Journal of Pain*, 9, 463. [PubMed: 15979027]
- Aurich NK, Alves Filho JO, Marques da Silva AM, & Franco AR (2015). Evaluating the reliability of different preprocessing steps to estimate graph theoretical measures in resting state fMRI data. *Frontiers in Neuroscience*, 9, 48 <https://www.frontiersin.org/article/10.3389/fnins.2015.00048> [PubMed: 25745384]
- Awh E, Vogel EK, & Oh S-H (2006). Interactions between attention and working memory. *Neuroscience*, 139, 201–208. <http://www.sciencedirect.com/science/article/pii/S0306452205008985> [PubMed: 16324792]
- Baliki MN, Geha PY, Apkarian AV, & Chialvo DR (2008). Beyond feeling: Chronic pain hurts the brain, disrupting the default-mode network dynamics. *The Journal of Neuroscience*, 28, 1398–1403. [PubMed: 18256259]
- Behzadi Y, Restom K, Liao J, & Liu TT (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage*, 37, 90–101. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2214855/> [PubMed: 17560126]
- Ben Simon E, Lahav N, Shamir R, Hendler T, & Maron-Katz A (2018). 792. Tired and MISCONNECTED: A breakdown of brain modularity following sleep deprivation. *Biological Psychiatry*, 81, S322.
- Bosch OG, Rihm JS, Scheidegger M, Landolt H-P, Stämpfli P, Brakowski J, ... Seifritz E (2013). Sleep deprivation increases dorsal nexus connectivity to the dorsolateral prefrontal cortex in humans. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 19597–19602. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3845164/> [PubMed: 24218598]
- Buckner RL (2008). The brain's default network. *Annals of the New York of Sciences*, 1124, 1–38. <http://onlinelibrary.wiley.com/doi/10.1196/annals.1440.011/full>
- Bushnell MC, eko M, & Low LA (2013). Cognitive and emotional control of pain and its disruption in chronic pain. *Nature Reviews. Neuroscience*, 14, 502–511. [PubMed: 23719569]
- Calhoun VD, Adali T, Pearlson GD, & Pekar JJ (2001). A method for making group inferences from functional MRI data using independent component analysis. *Human Brain Mapping*, 14, 140–151. [PubMed: 11559959]
- Calhoun VD, Adali T, & Pekar JJ (2004). A method for comparing group fMRI data using independent component analysis: Application to visual, motor and visuomotor tasks. *Magnetic Resonance Imaging*, 22, 1181–1191. [PubMed: 15607089]
- Ceko M, Gracely JL, Fitzcharles M-A, Seminowicz DA, Schweinhardt P, & Bushnell MC (2015). Is a responsive default mode network required for successful working memory task performance? *The Journal of Neuroscience*, 35, 11595–11605. [PubMed: 26290236]
- Ceko M, Shir Y, Ouellet JA, Ware MA, Stone LS, & Seminowicz DA (2015). Partial recovery of abnormal insula and dorsolateral prefrontal connectivity to cognitive networks in chronic low back pain after treatment. *Human Brain Mapping*, 36, 2075–2092. [PubMed: 25648842]

- Chai XJ, Ofen N, Gabrieli JDE, & Whitfield-Gabrieli S (2014). Selective development of anticorrelated networks in the intrinsic functional organization of the human brain. *Journal of Cognitive Neuroscience*, 26, 501–513. [PubMed: 24188367]
- Chuah YML, Venkatraman V, Dinges DF, & Chee MWL (2006). The neural basis of interindividual variability in inhibitory efficiency after sleep deprivation. *The Journal of Neuroscience*, 26, 7156–7162. [PubMed: 16822972]
- Coghill RC, Gilron I, & Iadarola MJ (2001). Hemispheric lateralization of somatosensory processing. *Journal of Neurophysiology*, 85, 2602–2612. [PubMed: 11387404]
- De Havas JA, Parimal S, Soon CS, & Chee MWL (2012). Sleep deprivation reduces default mode network connectivity and anti-correlation during rest and task performance. *NeuroImage*, 59, 1745–1751. [PubMed: 21872664]
- Finan PH, Goodin BR, & Smith MT (2013). The association of sleep and pain: An update and a path forward. *The Journal of Pain*, 14, 1539–1552. [PubMed: 24290442]
- Finan PH, Quartana PJ, Remeniuk B, Garland EL, Rhudy JL, Hand M, ... Smith MT (2017). Partial sleep deprivation attenuates the positive affective system: Effects across multiple measurement modalities. *Sleep*, 40, 1–9.
- Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, & Raichle ME (2005). The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 9673–9678. [PubMed: 15976020]
- Gazzaley A, & Nobre AC (2012). Top-down modulation: Bridging selective attention and working memory. *Trends in Cognitive Sciences*, 16, 129–135. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3510782/> [PubMed: 22209601]
- Gorgolewski KJ, Varoquaux G, Rivera G, Schwarz Y, Ghosh SS, Maumet C, ... Margulies DS (2015). NeuroVault.Org: A web-based repository for collecting and sharing unthresholded statistical maps of the human brain. *Frontiers in Neuroinformatics*, 9, 8. [PubMed: 25914639]
- Horowitz SG, Braun AR, Carr WS, Picchioni D, Balkin TJ, Fukunaga M, & Dwyer JH (2009). Decoupling of the brain's default mode network during deep sleep. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 11376–11381. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2708777/> [PubMed: 19549821]
- Iacovides S, George K, Kamenman P, & Baker FC (2017). Sleep fragmentation hypersensitizes healthy young women to deep and superficial experimental pain. *The Journal of Pain*, 18, 844–854. [PubMed: 28300651]
- Ji G, & Neugebauer V (2009). Hemispheric lateralization of pain processing by amygdala neurons. *Journal of Neurophysiology*, 102, 2253–2264. <https://www.ncbi.nlm.nih.gov/pubmed/19625541> [PubMed: 19625541]
- Jurkiewicz MT, Crawley AP, & Mikulis DJ (2018). Is rest really rest? Resting state functional connectivity during rest and motor task paradigms. *Brain Connectivity*, 8, 268–275. [PubMed: 29665711]
- Karaman S, Karaman T, Dogru S, Onder Y, Citil R, Bulut YE, ... Suren M (2014). Prevalence of sleep disturbance in chronic pain. *European Review for Medical and Pharmacological Sciences*, 18, 2475–2481. [PubMed: 25268092]
- Kelly R, Wang Z, Alexopoulos G, Gunning-Dixon F, Murphy C, Morimoto S, ... Hoptman M (2010). Hybrid ICA-seed-based methods for fMRI functional connectivity. *International Journal of Biomedical Imaging*, 2010, 1–24.
- Koffel E, Kroenke K, Bair MJ, Leverty D, Polusny MA, & Krebs EE (2016). The bidirectional relationship between sleep complaints and pain: Analysis of data from a randomized trial. *Health Psychology*, 35, 41. [PubMed: 26076002]
- Kong J, Jensen K, Loiotile R, Cheetham A, Wey H-Y, Tan Y, ... Gollub RL (2013). Functional connectivity of the frontoparietal network predicts cognitive modulation of pain. *Pain*, 154, 459–467. [PubMed: 23352757]
- Krause AJ, Ben SE, Mander BA, Greer SM, Saletin JM, Goldstein-Piekarski AN, & Walker MP (2017). The sleep-deprived human brain. *Nature Reviews. Neuroscience*, 18, 404 10.1038/nrn.2017.55 [PubMed: 28515433]

- Kucyi A, Salomons TV, & Davis KD (2016). Cognitive behavioral training reverses the effect of pain exposure on brain network activity. *Pain*, 157, 1895–1904. [PubMed: 27101426]
- Larson-Prior LJ, Zempel JM, Nolan TS, Prior FW, Snyder AZ, & Raichle ME (2009). Cortical network functional connectivity in the descent to sleep. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 4489–4494. [PubMed: 19255447]
- Lautenbacher S, Kundermann B, & Krieg J-C (2006). Sleep deprivation and pain perception. *Sleep Medicine Reviews*, 10, 357–369. [PubMed: 16386930]
- Li S, Tian J, Li M, Wang T, Lin C, Yin Y, ... Jiang G (2018). Altered resting state connectivity in right side frontoparietal network in primary insomnia patients. *European Radiology*, 28, 664–672. [PubMed: 28828546]
- Li Y-O, Adali T, & Calhoun VD (2007). Estimating the number of independent components for functional magnetic resonance imaging data. *Human Brain Mapping*, 28, 1251–1266. [PubMed: 17274023]
- Lorenz J, Minoshima S, & Casey KL (2003). Keeping pain out of mind: The role of the dorsolateral prefrontal cortex in pain modulation. *Brain*, 126, 1079–1091. 10.1093/brain/awg102 [PubMed: 12690048]
- Lorenzi M, Beltramello A, Mercuri NB, Canu E, Zoccatelli G, Pizzini FB, ... Costardi D (2011). Effect of Memantine on resting state default mode network activity in Alzheimers disease. *Drugs & Aging*, 28, 205–217. [PubMed: 21250762]
- Patel AX, & Bullmore ET (2016). A wavelet-based estimator of the degrees of freedom in denoised fMRI time series for probabilistic testing of functional connectivity and brain graphs. *NeuroImage*, 142, 14–26. <http://www.sciencedirect.com/science/article/pii/S1053811915003523> [PubMed: 25944610]
- Power JD, Cohen AL, Nelson SM, Wig GS, Barnes KA, Church JA, ... Petersen SE (2011). Functional network organization of the human brain. *Neuron*, 72, 665–678. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3222858/> [PubMed: 22099467]
- Price DD (2002). Central neural mechanisms that interrelate sensory and affective dimensions of pain. *Molecular Interventions*, 2, 392. [PubMed: 14993415]
- Raichle ME (2015). The brain's default mode network. *Annual Review of Neuroscience*, 38, 433–447.
- Rainville P (2002). Brain mechanisms of pain affect and pain modulation. *Current Opinion in Neurobiology*, 12, 195–204. [PubMed: 12015237]
- Rosseland R, Pallesen S, Nordhus IH, Matre D, & Blågestad T (2018). Effects of sleep fragmentation and induced mood on pain tolerance and pain sensitivity in young healthy adults. *Frontiers in Psychology*, 9, 2089 <https://www.frontiersin.org/article/10.3389/fpsyg.2018.02089> [PubMed: 30429815]
- Seminowicz DA, & Ceko M (2015). Can we exploit cognitive brain networks to treat chronic pain? *Pain Management*, 5, 399–402. [PubMed: 26399153]
- Seminowicz DA, & Davis KD (2007). Pain enhances functional connectivity of a brain network evoked by performance of a cognitive task. *Journal of Neurophysiology*, 97, 3651–3659. [PubMed: 17314240]
- Seminowicz DA, Mikulis DJ, & Davis KD (2004). Cognitive modulation of pain-related brain responses depends on behavioral strategy. *Pain*, 112, 48–58. [PubMed: 15494184]
- Seminowicz DA, & Moayed M (2017). The dorsolateral prefrontal cortex in acute and chronic pain. *The Journal of Pain*, 18, 1027–1035. [PubMed: 28400293]
- Seminowicz DA, Remeniuk B, Krimmel SR, Smith MT, Barrett FS, Wulff AB, ... Finan PH (2019). Pain-related nucleus accumbens function: Modulation by reward and sleep disruption. *Pain*, 160, 1196–1207. [PubMed: 30753171]
- Seminowicz DA, Shpaner M, Keaser ML, Krauthamer GM, Mantegna J, Dumas JA, ... Naylor MR (2013). Cognitive-behavioral therapy increases prefrontal cortex gray matter in patients with chronic pain. *The Journal of Pain*, 14, 1573–1584. [PubMed: 24135432]
- Sevel LS, Letzen JE, Staud R, & Robinson ME (2016). Inter-hemispheric dorsolateral prefrontal cortex connectivity is associated with individual differences in pain sensitivity in healthy controls. *Brain Connectivity*, 6, 357–364. [PubMed: 26916416]

- Shirer WR, Ryali S, Rykhlevskaia E, Menon V, & Greicius MD (2012). Decoding subject-driven cognitive states with whole-brain connectivity patterns. *Cerebral Cortex*, 22, 158–165. [PubMed: 21616982]
- Short MA, & Banks S (2014). The functional impact of sleep deprivation, sleep restriction, and sleep fragmentation In *Sleep deprivation and disease* (pp. 13–26). New York, NY: Springer.
- Sletten TL, Segal AY, Flynn-Evans EE, Lockley SW, & Rajaratnam SMW (2015). Inter-individual differences in neurobehavioural impairment following sleep restriction are associated with circadian rhythm phase. *PLoS One*, 10, e0128273. [PubMed: 26043207]
- Smith DV, Sip KE, & Delgado MR (2015). Functional connectivity with distinct neural networks tracks fluctuations in gain/loss framing susceptibility. *Human Brain Mapping*, 36, 2743–2755. [PubMed: 25858445]
- Smith M, Edwards RR, Mccann U, & Haythornthwaite JA (2007). The effects of sleep deprivation on pain inhibition and spontaneous pain in women. *Sleep*, 30, 494–505. [PubMed: 17520794]
- Smith R, Lane RD, Alkozei A, Bao J, Smith C, Sanova A, ... Killgore WDS (2018). The role of medial prefrontal cortex in the working memory maintenance of one's own emotional responses. *Scientific Reports*, 8, 3460. [PubMed: 29472625]
- Symonds LL, Gordon NS, Bixby JC, & Mande MM (2006). Right-lateralized pain processing in the human cortex: An fMRI study. *Journal of Neurophysiology*, 95, 3823–3830. [PubMed: 16554508]
- Tang NKY, Wright KJ, & Salkovskis PM (2007). Prevalence and correlates of clinical insomnia co-occurring with chronic back pain. *Journal of Sleep Research*, 16, 85–95. [PubMed: 17309767]
- Valet M, Sprenger T, Boecker H, Willoch F, Rummeny E, Conrad B, ... Tolle TR (2004). Distraction modulates connectivity of the cingulo-frontal cortex and the midbrain during pain—An fMRI analysis. *Pain*, 109, 399–408. [PubMed: 15157701]
- Verweij IM, Romeijn N, Smit DJA, Piantoni G, Van Someren EJW, & van der Werf YD (2014). Sleep deprivation leads to a loss of functional connectivity in frontal brain regions. *BMC Neuroscience*, 15, 88. [PubMed: 25038817]
- Villemure C, & Bushnell CM (2002). Cognitive modulation of pain: How do attention and emotion influence pain processing? *Pain*, 95, 195–199. [PubMed: 11839418]
- Whitfield-Gabrieli S, & Nieto-Castanon A (2012). A functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain Connectivity*, 2, 125–141. [PubMed: 22642651]
- Wiech K, Ploner M, & Tracey I (2008). Neurocognitive aspects of pain perception. *Trends in Cognitive Sciences*, 12, 306–313. [PubMed: 18606561]




FIGURE 1.

This study is an analysis of a simple motor task collected during fMRI scanning as part of a larger protocol. Procedures used in the present study are outlined above. Participants underwent two nights of monitored sleep. One night included an 8-hr opportunity for uninterrupted sleep. The other night included a sleep disruption protocol, wherein participants were awakened during one randomly chosen 1-hr interval and seven 20-min tertiles for the remaining hours. Structural and functional MRI scanning was completed following each night. Structural MRI and simple motor task fMRI data were used to address the study's aims




FIGURE 2. Independent component analysis (ICA) was conducted on simple motor task fMRI data. Resultant independent components (ICs) were spatially sorted to standardized templates to identify bilateral executive control networks (blue and green) and the default mode network (red). Dice coefficients ranged from 0.26 to 0.39 for goodness of fit. These top fitting ICs were masked for use as a region of interest (ROI) in subsequent seed-to-ROI functional connectivity analyses




FIGURE 3.

Mean pain ratings and temperatures are shown in white (uninterrupted sleep, US) and gray (forced awakenings, FA) bars, whereas individual differences in these measures are shown in uniquely colored lines. The present study aimed to examine cognitive network FC changes as a function of sleep induced-hyperalgesia. To derive a measure of change in pain sensitivity following FA during the sleep fragmentation condition compared to US, we calculated a pain sensitivity change score accounting for changes in individually tailored stimulus temperatures used for a noxious thermal stimulation task and concomitant pain ratings. In the “Thermal Stimuli Temperature” panel, some lines overlap and are eclipsed for individuals who had identical changes in temperature




FIGURE 4.

To examine the relationship between change in pain sensitivity and change in cognitive network functional connectivity (FC) following forced awakenings, we conducted seed-to-region of interest (ROI) FC analyses using the right executive control network (RECN, red), left executive control network (LECN), and default mode network (DMN) as a priori seeds and a pain sensitivity change score as a predictor of interest. We did not identify significant associations between the LECN and DMN as seeds for seed-to-ROI FC analyses. However, there was a positive association between pain sensitivity change scores and change in FC between the RECN seed and several ROIs located within the LECN, dDMN, and the primary visual network (shown as colored spheres; as defined by the Power 264 atlas). Numbers represent the associated ROI within the Power 264 atlas (see Table 4 for corresponding region names). Scatterplots demonstrate the relationship between FC values and pain sensitivity change scores

TABLE 1Participant demographic characteristics ($n = 19$)

Sex	Male	6
	Female	13
Age		24 (4.3)
Race	Black	5
	White	10
	Asian	2
	Other/mixed	2
Ethnicity	Hispanic	5
	Non-Hispanic	14
Education	Current student	5
	Some college	5
	College graduate	1
	Master's degree	3
	Doctoral degree	5

Note: Sex, race, ethnicity, and education are represented as frequencies. Age is represented as mean (*SD*).

TABLE 2

Clusters identified within each independent component that was used as a seed for seed-to-ROI functional connectivity analyses. Independent components represented the right executive control network (RECN), left executive control network (LECN), and default mode network (DMN)

Independent component	Cluster hemisphere	Cluster coordinates	Cluster size	p-value (<.05 FWE)	Region
RECN	Right	34 18 54	5,746	<.001	dIPFC/MFG
	Right	44 -44 54	3,287	<.001	LOC
	Left	-34 -56 52	1,726	<.001	SPL
	Right	64 -32 -14	810	<.001	MTG
	Left	-30 -62 -32	681	<.001	Cerebellum
	Right	6 -26 32	544	<.001	PCC
	Left	-42 50 -10	333	<.001	Frontal pole
	Left	-10 -78 -26	315	<.001	Cerebellum
	Left	-64 -44 -22	309	<.001	ITG
	Right	32 33 -6	243	<.001	Insula cortex
LECN	Right	10 -58 46	233	<.001	Precuneus
	Left	-52 16 40	53	<.001	MFG
	Left	-34 4 30	45	<.001	Ventral premotor
	Left	-38 -28 8	19	.005	Planum temporale
	Left	-46 12 36	7,986	<.001	MFG
	Left	-34 -60 52	3,536	<.001	LOC
	Left	-56 -50 -10	1,461	<.001	ITG
	Right	30 -62 46	284	<.001	LOC
	Right	44 34 8	230	<.001	IFG
	Right	12 -76 -24	157	<.001	Cerebellum
DMN	Right	28 -70 -46	143	<.001	Cerebellum
	Right	28 -62 -30	136	<.001	Cerebellum
	Right	4 8 28	75	<.001	MCC
	Left	-6 -34 42	46	<.001	PCC
	Right	54 -48 -18	41	<.001	ITG
	Right	50 -12 60	31	<.001	Precentral gyrus
	Left	-10 -20 72	25	<.001	Precentral gyrus
	Midline	6 -46 20	8,750	<.001	Precuneus/PCC

Independent component	Cluster hemisphere	Cluster coordinates	Cluster size	p-value (<.05 FWE)	Region
	Left	-44 -62 32	2,529	<.001	LOC
	Right	40 -64 44	1,941	<.001	LOC
	Midline	0 52 -12	1,872	<.001	vMPFC/ACC
	Left	-22 30 42	436	<.001	SFG
	Right	22 32 38	357	<.001	SFG
	Right	34 -80 -6	204	<.001	LOC
	Left	-58 -22 -18	69	<.001	MTG
	Left	-8 50 -42	54	<.001	Cerebellum
	Right	54 -10 -18	34	<.001	MTG
	Left	-24 -88 -8	23	.003	Fusiform gyrus
	Left	-18 -96 4	20	.005	Occipital pole
	Right	10 -46 -46	19	.01	Cerebellum

Abbreviations: ACC, anterior cingulate cortex; IFG, inferior frontal gyrus; ITG, inferior temporal gyrus; LOC, lateral occipital cortex; MFG, middle frontal gyrus; MTG, middle temporal gyrus; OFC, orbitofrontal cortex; PCC, posterior cingulate cortex; SFG, superior frontal gyrus; SPL, superior parietal lobule.

Comparison of polysomnography findings detailing participants' total sleep time (TST) and times in stages N1, N2, slow wave sleep (SWS) and rapid eye movement (REM) sleep at the uninterrupted sleep (US) and forced awakenings (FA) conditions. Additional data reflect condition-based differences in self-reported sleepiness (Stanford Sleepiness Scale; SSS), performance on the fMRI simple motor task (SMT), and pain ratings/stimulus temperatures from the noxious thermal stimulation task

TABLE 3

Variable	Mean (SD)		Paired-samples <i>t</i> test
	US	FA	
TST (min)	415.9 (81.9)	269.3 (53.9)	$t_{17} = 5.2, p < .001$
N1 (min)	17.5 (9.3)	25.1 (11.3)	$t_{17} = -2.9, p = .009$
N2 (min)	191.6 (77.8)	121.3 (30.3)	$t_{17} = 3.6, p = .002$
SWS (min)	113.6 (24.6)	81.0 (24.6)	$t_{17} = 2.2, p = .045$
REM (min)	88.6 (28.4)	42.8 (29.3)	$t_{17} = 4.6, p < .001$
SSS	1.9 (.6)	2.98 (.9)	$t_{13} = -5.1, p < .001$
SMT % accuracy	74.8% (25.3)	63.3% (22.7)	$t_{18} = 2.5, p = .021$
Pain ratings	5.4 (1.8)	5.4 (1.4)	$t_{18} = 0.05, p = .96$
Thermal stimulus temperatures	46.4 (1.4)	46.1 (1.6)	$t_{18} = 1.1, p = .29$
Pain sensitivity change score	.04 (.31); range = -.81 – .55		

Note: Thermal stimuli temperatures were derived during a quantitative sensory testing session that took place outside of the scanner. Pain ratings were collected during a noxious thermal stimulation fMRI task that used the derived stimulus temperatures. The pain sensitivity change score accounts for changes in both thermal stimulus temperatures and pain ratings between US and FA.

Significant power atlas ROIs yielded from seed-to-ROI analyses examining differences in cognitive network FC between the uninterrupted sleep and forced awakenings sleep conditions associated with pain sensitivity change scores

TABLE 4

Seed	Power	264 atlas ROI	Associated network	Region	Coordinates			t-stat	pFDR
					x	y	z		
RECN	188		LECN	I-dIPFC	-42	38	21	4.21	.03
	201		LECN	I-dIPFC	-42	25	30	4.24	.03
	100		DMN	I-MFG	-35	20	51	4.35	.03
	137		DMN	I-IFG	-46	31	-13	4.23	.03
	108		DMN	r-paracingulate gyrus	9	54	3	3.97	.04
	161		Visual	r-LOC	42	-66	-8	4.28	.03

Abbreviations: dIPFC, dorsolateral prefrontal cortex; DMN, default mode network; FDR, false discovery rate; l-, left; LECN, left executive control network; IFG, inferior frontal gyrus; MFG, medial frontal gyrus; LOC, lateral occipital cortex; r-, right; RECN, right executive control network; ROI, region of interest.

TABLE 5

Descriptive information for functional connectivity values based on sleep condition

Seed	Power 264 atlas ROI	Associated network	Region	FC mean (SD)		Correlation with pain sensitivity changes (r, p)
				US	FA	
RECN	188	LECN	l-dIPFC	0.54 (0.27)	0.58 (0.26)	.72, .001
	201	LECN	l-dIPFC	0.60 (0.23)	0.64 (0.28)	.72, .001
	100	DMN	l-MFG	0.42 (0.18)	0.45 (0.22)	.73, .000
	137	DMN	l-IFG	0.14 (0.16)	0.26 (0.23)	.69, .001
	108	DMN	r-paracingulate gyrus	0.03 (0.21)	0.05 (0.24)	.72, .001
	161	Visual	r-LOC	0.28 (0.2)	0.27 (0.28)	.51, .03

Abbreviations: dIPFC, dorsolateral prefrontal cortex; DMN, default mode network; FDR, false discovery rate; l-, left; LECN, left executive control network; IFG, inferior frontal gyrus; MFG, medial frontal gyrus; LOC, lateral occipital cortex; r-, right; RECN, right executive control network; ROI, region of interest.