

UC Davis

UC Davis Previously Published Works

Title

Associations between Aircraft Noise Exposure and Self-Reported Sleep Duration and Quality in the United States-Based Prospective Nurses Health Study Cohort.

Permalink

<https://escholarship.org/uc/item/3cd2j6n9>

Journal

Environmental Health Perspectives, 131(4)

Authors

Bozigar, Matthew

Huang, Tianyi

Redline, Susan

et al.

Publication Date

2023-04-01

DOI

10.1289/EHP10959

Peer reviewed

Associations between Aircraft Noise Exposure and Self-Reported Sleep Duration and Quality in the United States-Based Prospective Nurses' Health Study Cohort

Matthew Bozigar,^{1*} Tianyi Huang,² Susan Redline,^{2,3,4} Jaime E. Hart,^{2,5} Stephanie T. Grady,¹ Daniel D. Nguyen,¹ Peter James,^{5,6} Bradley Nicholas,⁷ Jonathan I. Levy,¹ Francine Laden,^{2,3,5} and Junenette L. Peters¹

¹Department of Environmental Health, Boston University School of Public Health, Boston, Massachusetts, USA

²Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA

³Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

⁴Beth Israel Deaconess Medical Center, Boston, Massachusetts, USA

⁵Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

⁶Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, Massachusetts, USA

⁷Volpe National Transportation Systems Center, U.S. Department of Transportation, Cambridge, Massachusetts, USA

BACKGROUND: Sleep disruption is linked with chronic disease, and aircraft noise can disrupt sleep. However, there are few investigations of aircraft noise and sleep in large cohorts.

OBJECTIVES: We examined associations between aircraft noise and self-reported sleep duration and quality in the Nurses' Health Study, a large prospective cohort.

METHODS: Aircraft nighttime equivalent sound levels (L_{night}) and day–night average sound levels (DNL) were modeled around 90 U.S. airports from 1995 to 2015 in 5-y intervals using the Aviation Environmental Design Tool and linked to geocoded participant residential addresses. L_{night} exposure was dichotomized at the lowest modeled level of 45 A-weighted decibels [dB(A)] and at multiple cut points for DNL. Multiple categories of both metrics were compared with <45 dB(A). Self-reported short sleep duration (<7 h/24-h day) was ascertained in 2000, 2002, 2008, 2012, and 2014, and poor sleep quality (frequent trouble falling/staying asleep) was ascertained in 2000. We analyzed repeated sleep duration measures using generalized estimating equations and sleep quality by conditional logistic regression. We adjusted for participant-level demographics, behaviors, comorbidities, and environmental exposures (greenness and light at night) and examined effect modification.

RESULTS: In 35,226 female nurses averaging 66.1 years of age at baseline, prevalence of short sleep duration and poor sleep quality were 29.6% and 13.1%, respectively. In multivariable models, exposure to L_{night} ≥45 dB(A) was associated with 23% [95% confidence interval (CI): 7%, 40%] greater odds of short sleep duration but was not associated with poor sleep quality (9% lower odds; 95% CI: –30%, 19%). Increasing categories of L_{night} and DNL ≥45 dB(A) suggested an exposure–response relationship for short sleep duration. We observed higher magnitude associations among participants living in the West, near major cargo airports, and near water-adjacent airports and among those reporting no hearing loss.

DISCUSSION: Aircraft noise was associated with short sleep duration in female nurses, modified by individual and airport characteristics. <https://doi.org/10.1289/EHP10959>

Introduction

Sleep is an essential, natural process needed for healthy brain and general body functioning.¹ Disruption of sleep can cause drowsiness and poor concentration and adversely affect the metabolic, endocrine, and immune systems.^{2–5} Poor sleep quality and short sleep duration (alternatively referred to as insufficient sleep or reduced total sleep time) have been associated with many adverse health outcomes,⁶ including depression,⁷ metabolic disorders (e.g., obesity, type 2 diabetes),^{8–10} cardiovascular disease,¹¹ incident ulcerative colitis,¹² coronary events,¹³ cancer,¹⁴ kidney function decline,¹⁵ mental and physical functional decline,^{16–18} and mortality.^{19–21}

Noise is unwanted or harmful sound,²² where sound is defined as repetitive variations in air pressure (i.e., vibrations) sensed by the human ear. Humans recognize, evaluate, and react to environmental

sounds even when asleep.²³ Noise can disrupt sleep architecture via arousals, sleep-stage changes, and awakenings.^{24,25} These, in turn, increase cortical excitations, indicative of an elevated stress response to noise, which may activate the sympathetic nervous system.²⁴

Aircraft noise is unique for its multispectral acoustical properties that impact the human auditory system²⁶ and has been shown to disrupt sleep.^{24,25,27} However, most studies of aircraft noise and sleep have taken place in a small collection of homes,²⁸ human sleep laboratories,^{25,29,30} or around one or a few airports.^{28,31–37} Despite calls by researchers for more large-scale field studies,²⁴ only two previous studies to our knowledge have used large-scale methods across many airports.^{38,39} In addition, most studies have been cross-sectional in design and therefore unable to assess associations over time. Sleep patterns may be disrupted by lifestyle factors such as shift work,⁴⁰ as well as environmental exposures that include air pollution, greenness, inopportune light at night (LAN), and noise.^{41–45} But to our knowledge, there are no large-scale studies of aircraft noise that integrate and adjust for multiple environmental exposures such as greenness and LAN together, particularly in the United States,³⁸ where the health effects of aircraft noise have been understudied. Although a legal framework for noise abatement was established for high aircraft noise levels [i.e., ≥65 dB(A)] in the United States,^{46,47} few studies have assessed potential health effects at lower thresholds in the country. Finally, there has been limited assessment of effect measure modification in previous studies to identify potential vulnerable and susceptible subpopulations.

We seek to add to existing knowledge by investigating associations between aircraft noise and sleep repeatedly self-reported in a U.S.-based prospective cohort living near 90 U.S. airports,

Address correspondence to Matthew Bozigar, 160 SW 26th St., Corvallis, OR 97331 USA. Email: bozigarm@oregonstate.edu

Supplemental Material is available online (<https://doi.org/10.1289/EHP10959>).

*Current address: Matthew Bozigar, College of Public Health and Human Sciences, Oregon State University, Corvallis, Oregon, USA.

S.R. reports consulting fees and grant support from Jazz Pharma, unrelated to this article. The other authors declare they have no actual or potential competing financial interests.

Received 17 January 2022; Revised 21 February 2023; Accepted 3 March 2023; Published 14 April 2023.

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehpsubmissions@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

controlling for individual and area-level factors. We also examine whether any relationships found are modified by individual, airport, or area characteristics.

Methods

Study Population and Airports

We used the Nurses' Health Study (NHS), an ongoing nationwide U.S. prospective cohort started in 1976, in this research. The cohort was comprised of 121,701 female registered nurses who were 30–55 years of age at initial enrollment. Although recruitment originally focused on the 11 states with the most registered nurses, participants have moved during follow-up and now live throughout the United States. Questionnaires were self-administered biennially and contained questions about incident disease, medical history, and lifestyle factors. Response rates have been $\geq 90\%$.^{48,49} Home addresses indicated by the participants were geocoded every 2 y (corresponding with the biennial sequence of the NHS).

We used responses to questionnaires in survey years within the interval with aircraft noise assessment (1995–2015) that included questions about sleep—2000, 2002, 2008, 2012, and 2014—to define the study years. The year 2000, the first year for which both aircraft noise and sleep data were available, was the study baseline. To be included in this study, participants needed to have reported on sleep and have a valid residential address in the United States that could be used to successfully locate them geographically (i.e., geocoded). Residential moves among the participants during the study years were captured at the biennial assessment and aircraft noise exposure and environmental exposure estimates were updated, but participant-years were skipped when new addresses could not be geocoded. Participants were further excluded from the study if they did not live within a 22.2-mi (35.7-km) radius buffer around 1 of the 90 study airports at baseline, which represented the maximal empirical extent of aircraft noise above a day–night average sound level (DNL) of 45 decibels [A-weighted, dB(A)] surrounding any of the airports in this study. This exclusion criterion was used to limit the people unexposed to aircraft noise [here referring to those below our lowest modeled level of 45 dB(A)] to those who lived in places similar to where the exposed group lived and, therefore, reduce potential bias from unmeasured confounding factors associated with living close to an airport. Use of the 22.2-mi (35.7-km) buffers additionally helped limit exposure misclassification of participants living near airports that were not included in this study. Table S1 summarizes the number of participants excluded at each survey year for each successive criterion.

The 90 study airports were selected based on availability of model input (aircraft operations) data. They were located in 40 of 50 states plus the District of Columbia and captured 87% of all enplanements in the United States in 2010.⁵⁰

The study protocol was approved by the institutional review board of Brigham and Women's Hospital, Boston, Massachusetts. Consent was implied through the return of the questionnaires.

Outcome Assessment

Participants self-reported the average number of hours they slept in a 24-h day in 5 survey years [2000 (baseline), 2002, 2008, 2012, and 2014]. Response options included ≤ 5 , 6, 7, 8, 9, and ≥ 10 h/24-h day. Self-reported sleep duration on the 2002 questionnaire was previously shown to correlate strongly ($r_{\text{Spearman}} = 0.79$) with sleep assessed by a 1-wk sleep diary in a validation study among a subset of NHS participants.²¹ The American Academy of Sleep Medicine and the Sleep Research Society reports that physiological and neurobehavioral deficits can occur and worsen over time

with < 7 h of sleep every 24 h.^{14,51} Therefore, we defined short sleep duration as < 7 h/24-h day. For the sleep quality outcome, participants were asked a question at baseline (2000) about their frequency of having difficulty falling or staying asleep during the previous 4 wk. We defined poor sleep quality as participant answers to this question on a six-point Likert scale of: “all of the time,” “most of the time,” or “a good bit of the time.”

Exposure Assessment

Details about the generation of aircraft noise estimates are described elsewhere.^{52,53} Briefly, aircraft noise estimates around all 90 study airports were modeled comprehensively using analogous input data and modeling assumptions by the U.S. Department of Transportation's John A. Volpe National Transportation Systems Center (Volpe) using the U.S. Federal Aviation Administration's (FAA's) Aviation Environmental Design Tool (AEDT; <https://aedt.faa.gov/>). Source data for the AEDT were aircraft operations from the Enhanced Traffic Management System, excluding helicopter operations. All annualized operations (e.g., commercial, cargo, military) were grouped by Aircraft Noise and Performance aircraft type, day or nighttime, operation airport, and stage length.

Aircraft noise level contours were modeled every 5 y from 1995 to 2015 (1995, 2000, 2005, 2010, and 2015) surrounding the 90 U.S. airports. Two noise metrics were estimated: aircraft nighttime equivalent sound level (L_{night}, A-weighted) and DNL (A-weighted). Noise levels ranged from 45 to 71 dB(A). L_{night} was the primary aircraft noise metric in this analysis because it captures aircraft noise occurring when people typically sleep. However, DNL incorporates nighttime estimates as well, and it is the primary metric the FAA uses to inform decision-making about aircraft noise.⁵⁴ Where L_{night} assesses aircraft noise from 2200 to 0700 hours, DNL is a 24-h annualized average, capturing the average day of a year's operations with a 10 dB(A) penalty for nighttime aircraft noise from 2200 to 0700 hours, when levels of background noise tend to be lower compared with daytime. We therefore included DNL as a secondary aircraft noise exposure metric.

Spatially, each geocoded participant home address point was linked to the aircraft noise contour polygon in which it resided and assigned the respective aircraft noise level. If an address point was not within any contour, it was assigned an arbitrary value of 44 dB(A), just below the lowest modeled value of 45 dB(A), which indicated exposures estimated to be < 45 dB(A), for both L_{night} and DNL. DNL < 45 dB(A) aligns with the recommendation and guideline for protecting health by limiting aircraft noise from the World Health Organization (WHO) Regional Office for Europe using the day–evening–night average sound level (L_{den}), an aircraft noise metric comparable to DNL that includes an extra penalty for evening-time aircraft noise. However, the 45 dB(A) threshold we used for L_{night} is higher than the corresponding WHO guideline of limiting L_{night} to < 40 dB(A).⁵⁵

Temporally, the aircraft exposure estimates for a given year were matched to current home addresses of participants of the same survey year when they coincided. When the 5-y aircraft noise estimates were not temporally coincident to a survey year, the most recent previous aircraft noise estimates were temporally matched.

Covariates and Potential Confounders

A directed acyclic graph (DAG) proposed by Billings et al. was used to guide our theoretical DAG (Figure S1) and an operational DAG (Figure S2) on the adjustment for covariates and potential confounding factors.^{42,56} Based on variables identified in the DAG, we additionally determined inclusion of environmental variables in final models after evaluating correlation magnitudes

among these variables and whether they changed the effect estimates by at least 10%.

Demographic factors included age (continuous), age² (continuous), U.S. Census region of residence (Northeast, Midwest, South, West), race (White, Black, American Indian, Asian, Hawaiian), and individual socioeconomic status (SES). Race was derived from an NHS algorithm using a 1992 question about major ancestry in combination with a 2004 question about best fitting race category and was non-time-varying. Race is a social construct, and we include it as proxy metric to aid in adjusting for socio-historical and multigenerational effects of discrimination and racism, which have been shown to affect sleep.⁵⁷ The categories we analytically used for race mirror the survey response options. Age and census region of residence were reported every 2 y. Including age² in statistical models adjusted for quadratic effects of age on our sleep disturbance-related outcomes. Individual SES was captured by two variables, currently living alone (yes, no) and spouse's education (<high school, high school, >high school, missing/not married). In the NHS cohort, typical measures of individual SES are deemed less applicable given that participants are or were all nurses of moderate-to-high SES; instead, living alone and lower spousal education levels, indicating less or no joint financial resources, are used as markers of lower relative individual SES among this population.⁵⁸ Spouse's education was reported in 1992 and carried forward, whereas living alone was reported in 2000, 2008, and 2012 and was carried forward as necessary. A question asked about shift work in the 1988 survey year indicated that although 49.1% of participants reporting previously working night shifts at some point in their careers, in 1996 only 2.9% of NHS participants reported having rotating night shift work in the previous 6 months. Given the low frequency of shift work 4 y prior to this study's baseline year (2000) and no further data points on shift work after 1996, shift work was not included as a potential confounder. Biannually varying postmenopausal status until 2002 (yes, no, missing) and hormone replacement therapy (never, current, former, missing) were included in a sensitivity analysis, but these individual metrics did not vary significantly, likely because 98.8% of the participants were already post menopausal at baseline (Table 1). Behaviors included smoking status (never, former, current, missing) and alcohol consumption (none, >0–4 g/d, 5–9 g/d, 10–14 g/d, 15–29 g/d, ≥30 g/d, missing) and were reported every 4 y and carried forward 2 y as necessary. Comorbidities included diabetes and hypertension, determined by a self-report of a current clinician diagnosis, and were reported every 2 y. Potential environmental confounders included air pollution [particulate matter with an aerodynamic diameter of ≤2.5 μm (PM_{2.5}) per cubic meter], greenness, LAN, population density, and neighborhood SES (nSES).

Air pollution, particularly PM_{2.5} and nitrogen oxides, has been associated with sleep apnea,^{59,60} likely through inflammatory effects on upper airway function.⁴² The annual average of monthly ambient PM_{2.5} (i.e., each 1-month average in the calendar year of survey year, averaged) was estimated at participant residential addresses using generalized additive mixed models developed for spatiotemporal prediction.⁶¹ The prediction models integrate monitored data from the U.S. Environmental Protection Agency's Air Quality Monitoring System and other publicly available networks with geospatial predictors (e.g., road network data, land use, elevation). Predictive accuracy for PM_{2.5} was high ($R^2 = 0.77$). PM_{2.5} estimates were made through 2007; estimates from 2007 were carried forward to the remaining study years for each participant.

Greenness from natural vegetation has been linked with longer sleep duration, improved cardiovascular biomarkers, and decreased sympathetic activation and may be associated with improved mental

health and healthy behaviors promoting quality sleep, such as walking.^{42,62,63} However, greener areas have also been linked with more aircraft noise annoyance,⁴⁵ potentially resulting from greener areas being quieter and/or pleasing and thus modifying the experience of noise from aircraft. Greenness was estimated via the Normalized Difference Vegetation Index (NDVI), or the ratio of the difference between the near-infrared region and red reflectance (isolating visible light absorbed by chlorophyll in plants) to the sum of the two measures⁶⁴ from 30-m resolution Landsat data. We calculated annual average NDVI at a spatial resolution of 270 m annually using R (version 4.0.3; R Development Core Team) and Google Earth Engine, matching the corresponding grid cell to the home address of each study participant. We chose to use a 270-m buffer, which is roughly the size of an average city block in Manhattan, New York, because this size is more representative of the immediate greenness environment that could affect sleep.

Light is a major input into the timing of the circadian system. LAN has been documented as having adverse effects on sleep.^{65–67} Assessment of outdoor LAN was described by James et al.⁶⁸ Briefly, annual average LAN was estimated using nighttime radiance units (in nanowatts per centimeter squared per steradian) at a spatial resolution of 30 arc-seconds, or ~1 km². LAN estimates were developed from satellite imagery data from the U.S. Defense Meteorological Satellite Program's Operational Linescan System, managed by the National Oceanic and Atmospheric Administration's (NOAA's) Earth Observation Group.⁶⁹ LAN values were estimated for each survey year until 2010, after which estimates were carried forward.

Dense urban built environments have been linked to positive influences on health by promoting walking and other physical activity,^{70,71} but urban density may come at the cost of reduced and poorer quality sleep.⁷² Population density and neighborhood metrics of room crowding have previously been associated with worse sleep outcomes.^{72–74} Biannually varying census tract population density (in number of people per kilometer squared) metrics were derived from the 2000 U.S. Census for the 2000 and 2002 survey years and from the 2010 U.S. Census for the 2008, 2012, and 2014 survey years.

nSES is another potential confounder of aircraft noise affecting sleep, but previous findings have been mixed. Neighborhood disadvantage has been associated with adverse sleep outcomes in some studies^{75–77} but not in others.^{78–80} The social environment may be more highly associated with sleep disruption than nSES, directly.⁷⁹ However, previous research has suggested that nSES can act as an upstream factor affecting downstream mediating environmental exposures influencing sleep, including aircraft noise.^{42,81} The nSES metric comprised a summed z-score of SES-related census variables (e.g., race, education, income, home value, nativity, unemployment) at the level of the census tract.⁸² Annual environmental/area exposures considered in statistical models (PM_{2.5}, NDVI, LAN, population density, and nSES) were assigned to participant home addresses every 2 y to match the current home addresses of participants given that some of them moved residences throughout the study period.

Potential Effect Modifiers

Potential effect modifiers were determined *a priori* from hypothesized influences of individual, airport, and geographic characteristics on the aircraft noise and sleep relationship. We hypothesized that the association between aircraft noise and sleep would vary by census region (Northeast, Midwest, South, West) in the United States,⁸³ based on potential underlying acoustical differences resulting from seasonal and weather (e.g., temperature, humidity) factors,^{84,85} geographical development patterns,^{86,87} housing types,^{88,89} and intensity and daily temporal distribution of air services and peak hours of operation,⁹⁰

Table 1. Age-standardized characteristics of women in the Nurses' Health Study (NHS) at baseline (2000) overall, by nighttime aircraft sound (L_{night}) exposure group, and by aircraft day-night sound level (DNL) exposure group.

Characteristic	Overall N = 35,381	L _{night} <45 dB(A) N = 34,838	L _{night} ≥45 dB(A) N = 543	DNL <45 dB(A) N = 28,544	DNL 45–54 dB(A) N = 5,882	DNL 55–64 dB(A) N = 916	DNL ≥65 dB(A) N = 39
Demographics							
Age ly (mean ± SD)	66.1 ± 7.2 y	66.1 ± 7.2 y	67.0 ± 7.1	66.1 ± 7.2	66.1 ± 7.1	66.6 ± 7.1	68.3 ± 6.5
Region of residence (%)							
Northeast	48.7	48.5	58.7	46.7	56.8	57.4	59.1
Midwest	14.8	14.8	10.9	14.8	15.5	10.3	8.7
South	19.3	19.4	16.2	19.9	16.9	18.2	9.1
West	17.2	17.3	14.2	18.6	10.9	14.1	23.0
Race (%)							
White	96.1	96.2	89.1	96.6	94.5	90.7	86.6
Black	2.4	2.3	9.8	1.9	3.8	7.4	13.4
American Indian	0.2	0.2	0.0	0.2	0.3	0.2	0.0
Asian	1.3	1.3	1.1	1.2	1.3	1.6	0.0
Hawaiian	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Currently live alone	21.6	21.6	25.6	21.4	22.4	24.0	33.0
Spouse's education (%)							
<High school	4.0	4.0	2.9	3.8	4.6	4.5	0.0
High school	26.7	26.6	32.6	26.4	27.0	33.7	28.6
>High school	44.9	45.0	36.0	45.7	42.3	36.5	47.9
Missing	24.4	24.4	28.4	24.0	26.1	25.3	23.6
Postmenopausal (%)							
Yes	98.8	98.7	99.1	98.7	98.9	99.0	100.0
No	1.2	1.2	0.9	1.2	1.1	1.0	0.0
Missing	0.1	0.1	0.0	0.1	0.0	0.0	0.0
Hormone replacement therapy (%)							
Never	23.0	22.9	28.8	22.3	26.0	27.8	14.3
Former	27.2	27.2	26.9	27.2	27.5	27.4	39.5
Current	42.9	43.0	37.0	43.9	39.0	36.1	39.6
Missing	6.9	6.9	7.4	6.7	7.6	8.7	6.6
Behaviors							
Smoking status (%)							
Never	42.9	42.9	44.7	43.2	41.6	42.9	54.0
Former	47.8	47.8	47.5	47.7	48.0	47.1	43.1
Current	9.1	9.1	7.3	8.9	10.2	9.8	2.9
Missing	0.2	0.2	0.5	0.2	0.2	0.2	0.0
Alcohol consumption							
Nonmissing	91.1; 5.3 ± 9.1	91.2; 5.3 ± 9.2	89.4; 4.0 ± 7.5	91.3; 5.3 ± 9.2	90.4; 5.1 ± 9.0	90.0; 4.5 ± 8.1	85.8; 3.1 ± 5.0
Missing (%)	8.9	8.8	10.6	8.7	9.6	10.0	14.2
Comorbidities (%)							
Diabetes	6.1	6.1	7.8	6.1	6.1	8.1	5.6
Hypertension	35.1	35.0	40.6	35.0	35.1	39.2	47.7
Environmental (mean ± SD)							
NDVI greenness index	0.4 ± 0.1	0.4 ± 0.1	0.3 ± 0.1	0.4 ± 0.1	0.3 ± 0.1	0.3 ± 0.1	0.3 ± 0.1
LAN (nW/cm ² /sr)	38.7 ± 25.2	38.4 ± 24.9	61.5 ± 30.1	35.5 ± 22.9	51.2 ± 28.8	59.7 ± 30.3	63.1 ± 18.0
PM _{2.5} (µg/m ³)	13.0 ± 2.7	13.0 ± 2.7	13.7 ± 2.5	12.9 ± 2.7	13.4 ± 2.4	13.6 ± 2.5	14.2 ± 2.2
Population density (pp/km ²)	1,929.3 ± 3,724.5	1,888.8 ± 3,642.3	4,478.5 ± 6,728.3	1,611.1 ± 2,914.1	3,098.7 ± 5,848.2	4,173.4 ± 5,848.2	7,619.2 ± 9,457.4
nSES (z-score)	-0.4 ± 3.3	-0.4 ± 3.3	-1.0 ± 2.5	-0.4 ± 3.3	-0.4 ± 3.0	-0.9 ± 2.7	-0.8 ± 1.9
Sleep (%)							
Insufficient sleep	29.6	29.5	37.6	29.2	30.8	35.4	36.8
Poor sleep quality	13.1	13.1	12.2	13.2	12.5	13.5	21.6

Note: Environmental exposures considered included a 270-m buffer of greenness (NDVI), LAN measured in units of radiance (nanowatts per square centimeter per steradian, nW/cm²/sr), annual PM_{2.5} (µg/m³) at participant residential address, census tract population density (number of people per square kilometer, pp/km²), and census tract socioeconomic status z-score (nSES). Missing values are excluded from calculation of descriptive statistics. The study sample was age-standardized to reflect the age distribution of the NHS cohort. LAN, light at night; NDVI, Normalized Difference Vegetation Index; nSES, neighborhood socioeconomic status; PM_{2.5}, particulate matter with an aerodynamic diameter of ≤2.5 µm; SD, standard deviation.

which were covariates for which we did not have direct measurements. We also hypothesized that living near a major cargo airport could result in a stronger effect because cargo aircraft are typically larger and older planes with less aircraft noise reduction technology and frequently operate during the night.⁹¹ Heavier aircraft climb more slowly and subsequently generate higher noise exposure on the ground. Most of these components of cargo aircraft operations are captured in AEDT modeling. However, AEDT uses the stage length method that uses takeoff-to-landing distance as a proxy for aircraft weight,⁹² which may more accurately estimate the weight of passenger aircraft than the weight of cargo aircraft. We identified the 24 largest cargo airports of the 90 study airports by total landed weight of all-cargo operations.⁵⁰ The 24 cargo airports were consistently among the top 25 in landed weight over the study period.⁵⁰ We further hypothesized that the association would be stronger for participants living near airports adjacent to a large water body (i.e., water-adjacent airports), where water and local weather could acoustically alter or enhance the experience of noise from aircraft in such areas.^{93,94} Our aircraft noise metrics likely did not capture the within-year variation in weather conditions that can occur near large water bodies. There were 21 water-adjacent airports, which were determined by assessing whether any existing runway configuration at an airport allowed for an overwater approach or departure. In 2008 and 2012 most participants answered questions about hearing loss. Regardless of the cause of hearing loss, we posited that greater hearing loss would decrease the magnitude of the association between aircraft noise and short sleep duration. We were unable to investigate potential effect modification of hearing loss relative to sleep quality because data on hearing loss were not available for the relevant survey year. Owing to small numbers in some strata, hearing loss was categorized as none, mild, moderate/severe, and missing.

Statistical Analysis

We created a map of the locations of the 90 airports included in this study as points using ArcGIS (version 10.8.1; ESRI). Counts of NHS participants around each airport included at study baseline were categorized into quartiles. Regions were indicated as the four U.S. Census regions in the context of state outlines using shapefiles from the U.S. Census (<https://www.census.gov/geographies/mapping-files/time-series/geo/tiger-line-file.html>).

We examined associations between aircraft noise and short sleep duration longitudinally using generalized estimating equations for repeated measures, and poor sleep quality cross-sectionally at baseline using conditional logistic regression. For repeated measures, an unstructured covariance matrix was used to maximize model flexibility given that we were not limited by degrees of freedom. We used SAS (version 9.4; SAS Institute Inc.) for all statistical analyses.

A participant was defined as exposed if their respective Lnight or DNL was estimated at ≥ 45 dB(A). We also examined cut points of Lnight 50 dB(A) and DNL 55 and 65 dB(A). We were unable to examine Lnight cut points ≥ 50 dB(A) and DNL ≥ 65 dB(A) owing to low counts of exposed participants. To evaluate potential exposure–response with each sleep outcome, we also used mutually exclusive increasing aircraft noise exposure categories compared with < 45 dB(A). We used a three-category version of Lnight [< 45 (reference), 45–49, and ≥ 50 dB(A)] and a four-category version of DNL [< 45 (reference), 45–54, 55–64, and ≥ 65 dB(A)]. Significance of trends (p_{Trend}) were estimated by modeling categories of aircraft noise exposures (e.g., four-category DNL) as continuous variables. We also examined continuous aircraft noise exposure levels > 44 dB(A), but only as an ancillary analysis because 98.5% and

80.7% of participants were below the lowest available estimate of 45 dB(A) for Lnight and DNL, respectively.

We built five models for each outcome, starting with a crude model (model 0) adjusting only for age, age², and calendar period. Model 1 further adjusted for demographic factors, such as region of residence, race, living alone, and spouse's education. Models 2–4 then successively adjusted for behaviors (smoking status and alcohol consumption), comorbidities (diabetes and hypertension), and environmental factors (greenness and LAN), respectively, in which model 4 was the final, fully adjusted model. Given the high correlation between LAN and population density (Table S2), we chose to select only one of these variables for inclusion in the models. LAN was selected in favor of population density as a confounder in the final analyses owing to its possible direct effect on sleep in addition to its estimation on a relatively fine-scale, 1-km grid. Analyses included indicator variables for missing covariate data. We assessed effect measure modification hypotheses by including multiplicative interaction terms of potential effect modifiers and aircraft noise ($p_{Interaction}$) in models and stratifying them by the categories of each effect modifier.

Results

Figure 1 shows the location of the 90 airports included in this study. Study population characteristics at baseline for the 35,381 participants are listed overall, by Lnight exposure, and by DNL exposure in Table 1. At baseline, the age [mean \pm standard deviation (SD)] of participants was 66.1 ± 7.2 y, and only 13.0% were still working full time as nurses. Most participants lived in the Northeast (48.7%). Approximately 21.6% of the participants were living alone at baseline, and over half of the nurses' spouses had more than a high school education.

Characteristics were similar for aircraft noise exposure groups, with a few notable exceptions. Participants exposed to Lnight ≥ 45 dB(A) tended to live more in the Northeast (58.7%) and less in the remaining regions compared with unexposed participants [Lnight < 45 dB(A)]. Compared with unexposed participants, those exposed to Lnight ≥ 45 dB(A) were more likely to be Black (9.8% vs. 2.3%), more often lived alone (25.6% vs. 21.6%), less frequently had a spouse with more than a high school education (50.3% vs. 59.5%), and had higher rates of diabetes (7.8% vs. 6.1%) and hypertension (40.6% vs. 35.0%). Participants with high levels of exposure to either Lnight or DNL also experienced, on average, higher LAN and lived in more densely populated census tracts compared with participants exposed to low levels of aircraft noise.

Over time, participants provided an average of 3.48 survey-years of follow-up. Participant characteristics did not differ substantially for those who only provided data at baseline ($N = 2,901$) vs. those who provided 2–5 survey years of data ($N = 32,480$; see Table S3). The participants tended to live near the same airport throughout the study period (95.0%) and to have stable aircraft DNL exposure within 10 dB(A) (90.5%).

Table S4 describes levels of exposure to aircraft Lnight and DNL overall and by potentially effect-modifying airport characteristics for those exposed to ≥ 45 dB(A) of aircraft noise; 98.5% and 80.7% of participants were exposed to Lnight and DNL < 45 dB(A) of aircraft noise, respectively. Exposures did not vary much by region, living near a major cargo airport, nor living near a water-adjacent airport.

Table 2 shows results for the estimated association between aircraft noise and sleep, adjusted for individual factors, behaviors, comorbidities, and other environmental exposures. In longitudinal analysis, we found 34% greater odds [95% confidence interval (CI): 17%, 53%] of short sleep duration among those exposed to Lnight




Figure 1. Map of 90 study airports across the United States symbolized by U.S. Census region and quartiles of Nurses' Health Study (NHS) participants.

≥ 45 dB(A) compared with < 45 dB(A) in the crude, age-adjusted model. Adjusting for demographics, behaviors, comorbidities, and environmental factors, the magnitude of association attenuated to 23% greater odds (95% CI: 7%, 40%) of short sleep duration among the exposed compared with the unexposed (Table 2). Using DNL as the exposure, in the fully adjusted model, we found 3% greater odds of short sleep duration for DNL ≥ 45 dB(A) relative to < 45 dB(A) (95% CI: -1%, 8%). At the DNL 55-dB(A) cut point, there were stronger associations for all models, including the fully adjusted model (13% greater odds; 95% CI: 2%, 26%) (Table 2). The magnitude of the aircraft noise and short sleep duration association using a cut point of DNL 65 dB(A) was elevated (44% greater odds in the fully adjusted model) for those exposed compared with not exposed (Table 2), although the 95% CI was wide (-7%, 123%) given the small number of occurrences that any individual was an exposed case ($n = 39$; case counts by exposure category are included in Figures 2–5).

There was no evidence of a crude association between $L_{\text{night}} \geq 45$ dB(A) and poor sleep quality cross-sectionally (7% lower odds; 95% CI: -28%, 20%), and full adjustment for additional factors did not appreciably change the relationship (9% lower odds; 95% CI: -30%, 19%) (Table 2). Similarly, when using DNL as the exposure at 45 and 55 dB(A) cut points, we did not find evidence of associations between aircraft noise exposure and sleep quality. However, there were 101% greater odds of poor sleep quality from exposure to DNL ≥ 65 dB(A), although the 95% CI was wide and included the null (95% CI: -5%, 325%) (Table 2).

In sensitivity analyses, inclusion/exclusion of nSES, annual $PM_{2.5}$, postmenopausal status, and hormone replacement therapy, and swapping LAN for census tract population density did not change effect estimates by $> 10\%$ (Table S5); therefore, these variables were not included in final models.

Using continuous versions of aircraft noise in fully adjusted models, a 5-dB(A) increase in L_{night} was associated with 23% greater odds (95% CI: 6%, 44%) of short sleep duration, whereas

a 5-dB(A) increase in DNL was associated with 3% greater odds (95% CI: 0%, 6%). Figures 2 and 3 show the potential exposure–response relationships between aircraft noise exposure and short sleep duration from longitudinal modeling, whereas tabular results are included in Table S6. Similar to results for dichotomized and categorical aircraft noise exposures, controlling for covariates attenuated the estimated odds ratios within ordinal categories between either L_{night} or DNL and short sleep duration (Figures 2 and 3). For the crude models, results indicate the presence of an exposure–response relationship between increasing aircraft noise exposures and short sleep duration. For the fully adjusted models, results similarly suggest the presence of monotonic exposure–response relationships ($L_{\text{night}} p_{\text{Trend}} < 0.01$; DNL $p_{\text{Trend}} = 0.03$).

Using continuous versions of aircraft noise > 45 dB(A) in fully adjusted models, 5-dB(A) increases in L_{night} was associated (L_{night} : 2% greater odds; 95% CI: -23%, 34%; DNL: 2% lower odds; 95% CI: -7%, 4%) with poor sleep quality. Figures 4 and 5 (and Table S6) show that there were some indications of effects for the highest respectively exposed groups but there was no evidence of exposure–response relationships for lower exposure categories for poor sleep quality cross-sectionally ($L_{\text{night}} p_{\text{Trend}} = 0.77$; DNL $p_{\text{Trend}} = 0.37$).

Results from fully adjusted longitudinal models stratified by region, major cargo airport, water-adjacent airport, and hearing loss are shown in Table 3. They suggest that the association between nighttime aircraft noise and short sleep duration differed by region of residence ($p_{\text{Interaction}} = 0.06$). The association was strongest in the West, having 83% greater odds (95% CI: 32%, 152%). Participants living near a major cargo airport had 69% greater odds (95% CI: 21%, 136%) of short sleep duration associated with nighttime aircraft noise exposure, which was higher ($p_{\text{Interaction}} = 0.09$) than those not living near a major cargo airport (16% greater odds; 95% CI: 1%, 35%). For participants living near a water-adjacent airport, the association between L_{night} and short sleep duration resulted in 36% greater odds (95% CI: 12%,

Table 2. Estimated odds ratios (ORs) and 95% confidence intervals (CIs) for the relationship between aircraft nighttime sound level (L_{night}), day–night sound level exposure (DNL), and short sleep duration (<7 h/24-h d) and poor sleep quality (trouble falling/staying asleep \geq “a good bit of the time”) cases at specified aircraft noise metric cut points and levels of adjustment in the Nurses’ Health Study (NHS) (2000–2014).

Model	Short sleep duration	Poor sleep quality
	$N_{obs} = 123,023$, $N_{participants} = 35,381$, $N_{cases} = 35,497$	$N_{obs} = 35,226$, $N_{participants} = 35,226$, $N_{cases} = 4,617$
$L_{night} \geq 45$ vs. <45 dB(A)		
Crude: age-adjusted	1.34 (1.17, 1.53)	0.93 (0.72, 1.20)
1: Crude+other demographics	1.27 (1.11, 1.45)	0.94 (0.71, 1.21)
2: 1 + behaviors	1.26 (1.10, 1.44)	0.94 (0.72, 1.22)
3: 2 + comorbidities	1.26 (1.10, 1.44)	0.92 (0.71, 1.20)
4: 3 + environmental	1.23 (1.07, 1.40)	0.91 (0.70, 1.19)
DNL ≥ 45 vs. <45 dB(A)		
Crude: age-adjusted	1.10 (1.05, 1.15)	0.96 (0.89, 1.04)
1: Crude + other demographics	1.06 (1.01, 1.10)	0.96 (0.89, 1.04)
2: 1 + behaviors	1.06 (1.01, 1.10)	0.96 (0.88, 1.04)
3: 2 + comorbidities	1.05 (1.01, 1.10)	0.96 (0.88, 1.03)
4: 3 + environmental	1.03 (0.99, 1.08)	0.94 (0.86, 1.02)
DNL ≥ 55 vs. <55 dB(A)		
Crude: age-adjusted	1.23 (1.11, 1.37)	1.08 (0.89, 1.30)
1: Crude + other demographics	1.17 (1.05, 1.30)	1.08 (0.89, 1.30)
2: 1 + behaviors	1.16 (1.05, 1.29)	1.08 (0.90, 1.30)
3: 2 + comorbidities	1.16 (1.04, 1.29)	1.07 (0.89, 1.29)
4: 3 + environmental	1.13 (1.02, 1.26)	1.06 (0.87, 1.28)
DNL ≥ 65 vs. <65 dB(A)		
Crude: age-adjusted	1.53 (0.98, 2.39)	1.94 (0.92, 4.08)
1: Crude + other demographics	1.50 (0.97, 2.31)	2.00 (0.95, 4.22)
2: 1 + behaviors	1.49 (0.96, 2.31)	2.04 (0.97, 4.30)
3: 2 + comorbidities	1.48 (0.96, 2.29)	2.04 (0.97, 4.30)
4: 3 + environmental	1.44 (0.93, 2.23)	2.01 (0.95, 4.25)

Note: Age-adjusted (age, age²) models were sequentially further adjusted as indicated with other demographics, behaviors, comorbidities, and environmental factors. Other demographics: U.S. region of residence, race, living alone, spouse’s education. Behaviors: smoking status, alcohol consumption. Comorbidities: diabetes, hypertension. Environmental: greenness (NDVI), LAN. Models of short sleep duration used generalized estimating equations to estimate odds from repeated measures in survey years 2000 (study baseline), 2002, 2008, 2012, and 2014. Conditional logistic regression models of sleep quality were used to estimate odds only for the baseline study year. dB(A), A-weighted decibel; LAN, light at night; N_{cases} , number of cases; NDVI, Normalized Difference Vegetation Index; N_{obs} , number of observations; $N_{participants}$, number of participants.

65%) compared with 11% greater odds (95% CI: –9%, 34%; $p_{interaction} = 0.14$) for those living near a non–water-adjacent airport. For poor sleep quality in the cross-sectional analysis, there was little evidence of effect modification by region or by living near a water-adjacent airport (Table 3). However, for participants living near a major cargo airport, there were 45% greater odds (95% CI: –18%, 157%; $p_{interaction} = 0.09$) of poor sleep quality among those exposed to nighttime aircraft noise, vs. 18% reduced odds (95% CI: –39%, 10%) for participants not living near a major cargo airport, but both intervals contained the null. Finally, for participants who reported no hearing loss, there were an estimated 50% higher odds (95% CI: 11%, 103%) of short sleep duration from exposure to nighttime aircraft noise. Estimated odds of short sleep duration were in progressively lower but also more imprecise with higher reported hearing loss [mild: 31% higher odds (95% CI: –19%, 114%); moderate/severe: 15% lower odds (95% CI: –61%, 86%)]. However, we did not find definitive evidence in the two survey years with hearing loss data available in our cohort (2008 and 2012) of an interaction between hearing loss and aircraft noise ($p_{interaction} = 0.33$), although there was a suggested relationship.




Figure 2. Odds ratio (OR) point estimates and 95% confidence intervals (CIs) investigating exposure–response relationship ($p_{Trend} < 0.01$) between categorical aircraft nighttime sound level (L_{night}) exposure [<45 dB(A) (reference), 45–49 dB(A), and ≥ 50 dB(A)] and short sleep duration (<7 h/24-h day), using GEEs from repeated measures in survey years 2000 (study baseline), 2002, 2008, 2012, and 2014 in the Nurses’ Health Study (NHS). OR and CI estimates can be found in Table S6. Models adjusted for age (age, age²) were sequentially further adjusted for other demographics, behaviors, comorbidities, and environmental factors. Other demographics: U.S. region of residence, race, living alone, spouse’s education. Behaviors: smoking status, alcohol consumption. Comorbidities: diabetes, hypertension. Environmental: greenness (NDVI), LAN. Models of short sleep duration used GEEs to estimate odds from repeated measures in survey years 2000 (study baseline), 2002, 2008, 2012, and 2014. Note: dB(A), A-weighted decibel; GEE, generalized estimating equation; LAN, light at night; N_{cases} , number of cases; NDVI, Normalized Difference Vegetation Index; N_{obs} , number of observations.

Discussion

This study investigated the relationship between aircraft noise exposure and self-reported sleep in the NHS cohort participants living near 90 major U.S. airports. Adjusting for several individual and two environmental confounders (greenness and LAN), we found that exposure to aircraft noise was associated with short sleep duration in a repeated measures analysis, with an exposure–response relationship seemingly evident across all levels of aircraft noise included. Furthermore, we found evidence of potential effect modification by individual, area, and airport characteristics, with stronger associations with short sleep duration for participants living in the West, near major cargo airports, near water-adjacent airports, and among those reporting no hearing loss. Exposure to aircraft noise had a limited association with poor sleep quality cross-sectionally, with highly positive but uncertain associations seen only for the highest aircraft noise exposure category.

We found that short sleep duration was linked with two metrics of average annual aircraft noise exposure, L_{night} and DNL. This is only partly consistent with existing literature on aircraft noise and sleep quantity. Most laboratory and residential studies have documented that aircraft noise events can shorten sleep duration,^{34,95} but some found associations with longer sleep duration in conjunction with poorer sleep quality.^{31,96} A national study of aircraft noise and self-reported sleep from the large-scale U.S. Behavioral Risk Factor Surveillance System surveys did not find an association between aircraft noise and short sleep duration,³⁹ but the study differed in numerous ways from the




Figure 3. Odds ratio (OR) point estimates and 95% confidence intervals (CIs) investigating exposure–response relationship ($p_{\text{Trend}} = 0.03$) between categorical aircraft day–night average sound level (DNL) exposure [<45 (reference), 45–54, 55–64, and ≥ 65 dB(A)] and short sleep duration (<7 h/24-h day), using GEEs from repeated measures in survey years 2000 (study baseline), 2002, 2008, 2012, and 2014 in the Nurses’ Health Study (NHS). OR and CI estimates can be found in Table S6. Models adjusted for age (age, age²) were sequentially further adjusted for other demographics, behaviors, comorbidities, and environmental factors. Other demographics: U.S. region of residence, race, living alone, spouse’s education. Behaviors: smoking status, alcohol consumption. Comorbidities: diabetes, hypertension. Environmental: greenness (NDVI), LAN. Models of short sleep duration used GEEs to estimate odds from repeated measures in survey years 2000 (study baseline), 2002, 2008, 2012, and 2014. Note: dB(A), A-weighted decibel; GEE, generalized estimating equation; LAN, light at night; N_{cases} , number of cases; NDVI, Normalized Difference Vegetation Index; N_{obs} , number of observations.

present study. Although Holt et al. established feasibility of a large-scale study of self-reported sleep, the study differed in its cross-sectional design, aircraft noise exposure modeling (Integrated Noise Model), area-level (ZIP code) exposure assignment, absence of Lnight estimates, higher DNL cut points starting at 55 dB, dissimilar reference population that included participants far from study areas and possibly near non-study airports, and potential confounders.³⁹ Another study around two airports in France found that a Lden 10-dB(A) increase in aircraft noise was associated with 63% greater odds (95% CI: 15%, 132%) of short total sleep time, after adjusting for individual level potential confounders.³⁴ However, the study did not adjust for other environmental factors, potentially confounding the relationship between aircraft noise and sleep, nor did the researchers have data available to investigate potential effect measure modification in their sample of 1,244 adults. At the area level, elevated census block group environmental noise was not associated with total hours slept in a subsample of a national cross-sectional survey of urban individuals in the United States, although it was significantly associated with other adverse sleep outcomes.⁹⁷ Although Rudolph et al. applied a novel nationwide environmental noise model, the study could not isolate the association with aircraft noise and had to rely on area-level (census block group) exposure assignment.⁹⁷ The differences in study designs used in previous research may explain their mixed results in contrast to the longitudinal design used in this study that found a strong signal between aircraft noise and short sleep duration in the NHS.

For sleep assessments, most population-based studies used relatively high cut points of the DNL aircraft noise metric, such as 50^{34,62} or 55 dB,³⁹ although some smaller studies have assessed the influences on sleep of aircraft noise exposures as low as 30 or 35 dB.^{28,31} Large-scale sleep studies investigating aircraft noise have mostly been unable to assess low potential thresholds or to incorporate nighttime specific metrics such as Lnight, particularly in the United States, with a few exceptions.³⁸ We found limited evidence of thresholds across the range of exposures included. When considering exposed vs. unexposed people at a 45-dB(A) cut-point, we observed a stronger association between short sleep duration and the exposure metrics for the nighttime measure than the 24-h day–night metric. Given consistent associations between sleep and a wide range of health outcomes, this finding suggests value in including nighttime aircraft noise metrics in health assessments of airport noise. Furthermore, finding evidence of an exposure–response relationship using DNL, future studies should assess a range of exposures and consider the sensitivity of conclusions about exposure categorization (Table 1).

Our findings suggest a clear, monotonic exposure–response relationship between aircraft noise and short sleep duration that is consistent with the results of smaller studies analyzing a variety of sleep parameters.^{29,32,38,95} However, most of the studies documenting exposure–response relationships between transportation noise and sleep parameters often used laboratory- or home-based designs with limited adjustment for potential confounding factors that lacked generalizability to larger populations.




Figure 4. Odds ratio (OR) point estimates and 95% confidence intervals (CIs) investigating exposure–response relationship ($p_{\text{Trend}} = 0.37$) between categorical aircraft nighttime equivalent sound level (Lnight) exposure [<45 (reference), 45–49 dB(A), and ≥ 50 dB(A)] and poor sleep quality (trouble falling/staying asleep \geq “a good bit of the time”) using conditional logistic regression at study baseline (2000) in the Nurses’ Health Study (NHS). OR and CI estimates can be found in Table S6. Models were adjusted for age (age, age²) other demographics, behaviors, comorbidities, and environmental factors. Other demographics: U.S. region of residence (removed from the region-specific models), race, living alone, spouse’s education. Behaviors: smoking status, alcohol consumption. Comorbidities: diabetes, hypertension. Environmental: greenness (NDVI), LAN. Conditional logistic regression models of sleep quality were used to estimate odds only for the baseline study year. Note: dB(A), A-weighted decibel; LAN, light at night; N_{cases} , number of cases; NDVI, Normalized Difference Vegetation Index; N_{obs} , number of observations.




Figure 5. Odds ratio (OR) point estimates and 95% confidence intervals (CIs) investigating exposure–response relationship ($p_{\text{Trend}}=0.37$) between categorical aircraft day–night sound level (DNL) exposure [<45 (reference), 45–54, 55–64, and ≥ 65 dB(A)] and poor sleep quality (trouble falling/staying asleep \geq “a good bit of the time”) using conditional logistic regression at study baseline (2000) in the Nurses’ Health Study (NHS). OR and CI estimates can be found in Table S6. Models were adjusted for age (age, age²) other demographics, behaviors, comorbidities, and environmental factors. Other demographics: U.S. region of residence (removed from the region-specific models), race, living alone, spouse’s education. Behaviors: smoking status, alcohol consumption. Comorbidities: diabetes, hypertension. Environmental: greenness (NDVI), LAN. Conditional logistic regression models of sleep quality were used to estimate odds only for the baseline study year. Note: dB(A), A-weighted decibel; LAN, light at night; N_{cases} , number of cases; NDVI, Normalized Difference Vegetation Index; N_{obs} , number of observations.

An additional strength of this study was the examination of potential effect modification of the nighttime aircraft noise–sleep relationship by individual, area, and airport characteristics. Previous studies assessed potential effect modification, but in different ways. For example, one study used total sleep duration as an effect modifier and various other sleep parameters as outcomes.³⁸ Although greater risk for those near major cargo airports was expected, given that the aircraft noise metrics we used may not have sufficiently incorporated characteristics associated with nighttime aircraft noise dynamics, effect modification of the association with sleep had not been previously investigated. Our results suggest that participants living near major cargo airports had shorter and poorer quality sleep related to aircraft noise at night than those living near airports with less cargo activity. Evidence from Europe has shown that air cargo flights tend to use older, larger aircraft that commonly operate in the early morning hours when commercial flight activity is low.⁹¹ In the United States, the quantity of cargo shipped by air has increased from 56.4 M ton-miles in 2003 to 76.6 M ton-miles in 2020.⁹⁸ If the trend continues, further indicated by the rapid growth in e-commerce-related cargo flight activity during the COVID-19 pandemic,^{99–101} these results suggest that impacts on sleep may also increase in magnitude and for additional populations. Although our analyses incorporated noise exposure estimates that should, in theory, capture differential contributions by cargo flights, either the noise metrics we used were not well suited for characterizing sleep disturbance (e.g., differences in flight patterns or sound frequency distributions for cargo flights) or the AEDT modeling has differential error for cargo

flights relative to other flights. The AEDT does not directly incorporate cargo weight because aircraft takeoff weight is proprietary in the United States; instead, the AEDT uses stage length as a proxy.⁹² If true cargo weight is systematically greater than estimates from stage length formulas, such as the scenario when cargo flights have shorter stage lengths (short-haul operations) but are heavily laden, then we would expect to see differentially greater effects at airports with more cargo operations, as we indeed found.

Similarly, we found support for the hypothesis that the association between aircraft noise and short sleep duration was greater for participants living near a water-adjacent airport, but there was little evidence of a modifying effect on poor sleep quality. For water-adjacent airports, our sleep duration findings may indicate additional complexity in these acoustical environments whereby aircraft sound energy may propagate more easily over water,^{93,94} differentially influencing sleep. The AEDT assumes a soft ground surface (e.g., grass) near the sound receiver. Reflections off hard ground, such as pavement, or water generally cause higher sound levels. Under certain conditions, sound propagating over water can be channeled by the reflection off the surface and then refracted downward owing to cool air being immediately above the surface with warmer air above that.

It is not clear why aircraft noise seems to vary in its association with short sleep duration by region of the country, although it potentially relates to aspects of the surrounding environment and climate zones, as well as nationally and internationally influenced flight operations scheduling. Additional research is needed to disentangle potential place-based aircraft sound propagation mechanisms, such as varied surfaces (hard vs. grassy surfaces), local weather, seasonality, the influence of atmospheric conditions during overflights,¹⁰² and housing materials and types. Unique regional climates have been associated with patterns in heating/cooling system types and related behaviors (e.g., opening windows) to achieve thermal comfort in bedrooms while sleeping,¹⁰³ which may further modify the effects of aircraft noise on sleep. Daily flight scheduling can be a function of time zones, levels of business activity, airline hub and spoke networks, and markets served (e.g., geographic proximity to national and international destinations).¹⁰⁴

There are age-related changes in hearing as people age,¹⁰⁵ but we found a trend suggestive of a lower magnitude relationship between aircraft noise and short sleep duration with greater hearing loss independent of age. However, this study was underpowered to robustly investigate potential effect modification by hearing loss. In the literature, hearing loss is usually mentioned as a physiological outcome of aircraft noise,^{55,106} used as an exclusion criterion,^{30,96} or, as in one study, considered as a confounder.³⁵ However, we did not find sleep studies that assessed potential effect modification by hearing loss.

We did not find associations with sleep quality in most analyses. However, the NHS surveys did not capture longitudinal self-assessments of multiple dimensions of sleep quality. Thus, we were only able to use participant answers to one question in 2000, in which only 66 of the 541 participants who were exposed to $N_{\text{night}} \geq 45$ dB(A) reported poor sleep quality. Despite limited power, we cannot rule out a potential association with poor sleep quality at very high levels of DNL exposure. Other predominantly smaller-scale studies conducted in human sleep laboratories or participant homes have found deleterious effects of aircraft noise on sleep quality.^{31,35}

There were several limitations of our study. Our study population did not include males, younger individuals, or many individuals from underrepresented groups owing to the construction of the original cohort. Noise sensitivity and annoyance, which may influence the effects of aircraft noise on sleep, were not

Table 3. Estimated odds ratios (ORs) and 95% confidence intervals (CIs) for potential effect measure modifiers of the relationship between aircraft nighttime sound level [$L_{\text{night}} \geq 45$ dB(A)] and short sleep duration (< 7 h/24-h day) and poor sleep quality (trouble falling/staying asleep \geq “a good bit of the time”) cases from fully adjusted models (model 4) in the Nurses’ Health Study (NHS) (2000–2014).

Potential effect modifier	Short sleep duration $N_{\text{obs}} = 123,023, N_{\text{participants}} = 35,381, N_{\text{cases}} = 35,497$			Poor sleep quality $N_{\text{obs}} = 35,226, N_{\text{participants}} = 35,226, N_{\text{cases}} = 4,617$		
	N_{obs} (cases)	OR (95% CI)	p -Value	N_{obs} (cases)	OR (95% CI)	p -Value
Region			0.06			0.90
Northeast	59,358 (18,278)	1.15 (0.96, 1.37)		17,132 (2,251)	0.88 (0.62, 1.25)	
Midwest	18,720 (4,831)	1.33 (0.90, 1.96)		5,211 (641)	1.08 (0.51, 2.31)	
South	23,564 (6,533)	1.08 (0.77, 1.51)		6,803 (975)	1.05 (0.57, 1.90)	
West	21,381 (5,855)	1.83 (1.32, 2.52)		6,080 (750)	0.77 (0.37, 1.60)	
Water-adjacent airport			0.14			0.99
No	76,071 (20,781)	1.11 (0.91, 1.34)		21,401 (2,761)	0.91 (0.63, 1.31)	
Yes	46,952 (14,716)	1.36 (1.12, 1.65)		13,749 (1,856)	0.95 (0.65, 1.38)	
Major cargo airport			0.09			0.09
No	96,445 (27,760)	1.16 (1.01, 1.35)		27,431 (3,613)	0.82 (0.61, 1.10)	
Yes	26,578 (7,737)	1.69 (1.21, 2.36)		7,795 (1,004)	1.45 (0.82, 2.57)	
Hearing loss ^a			0.33		NA	
None	30,422 (8,974)	1.50 (1.11, 2.03)				
Mild	15,939 (4,899)	1.31 (0.81, 2.14)				
Moderate/severe	9,379 (2,711)	0.85 (0.39, 1.86)				

Note: Models were adjusted for age (age, age²), other demographics, behaviors, comorbidities, and environmental factors. Other demographics: U.S. region of residence (removed from the region-specific models), race, living alone, spouse’s education. Behaviors: smoking status, alcohol consumption. Comorbidities: diabetes, hypertension. Environmental: greenness (NDVI), LAN. Models of short sleep duration used generalized estimating equations to estimate odds from repeated measures in survey years 2000 (study baseline), 2002, 2008, 2012, and 2014. Conditional logistic regression models of sleep quality were used to estimate odds only for the baseline study year. dB(A), A-weighted decibel; LAN, light at night; N_{cases} , number of cases; NDVI, Normalized Difference Vegetation Index; N_{obs} , number of observations; $N_{\text{participants}}$, number of participants.

^aHearing loss was assessed and analyzed only for short sleep duration in 2008 and 2012. For potential effect modification by hearing loss: $N_{\text{obs}} = 55,740, N_{\text{participants}} = 25,627,$ and $N_{\text{cases}} = 16,584.$

included because they were not measured in the cohort. In addition, there may be residual confounding by weather conditions, such as temperature, which independently affects sleep.^{74,83} Furthermore, we could not adjust for sleep medication use, which might be a potential confounder. Sleep was subjectively self-reported, and it has been shown that subjective reporting of sleep duration may underestimate objective measures, although it can depend on how the question is asked. Only a single question about sleep quality was available on one biennial questionnaire. Although aircraft noise estimates used were average sound energies over time, single-event exposures of individual flights [e.g., maximum sound level (L_{max}) or sound exposure level (SEL)] may be more relevant to sleep, yet such metrics are infrequently used.⁵⁵ We could not capture within-5-y variations in noise exposure. Aircraft noise estimates were outside at home addresses, not in the actual bedrooms in which participants presumably slept, and they did not include noise from ground operations in and around airports. Although important housing and indoor home environments were not able to be captured, including sound insulation, window opening/closing or air conditioner use,¹⁰⁷ address-level exposure assignment for a large-scale study was a novel contribution to aircraft noise and sleep research. Other sources of environmental noise (e.g., natural, community, road, rail) were not captured directly, although they were likely partly captured by environmental confounders included. In the present study, LAN likely acted as a proxy for population density and sources of noise (e.g., road) and light happening more frequently in denser (e.g., urban) areas. The cohort and study population were not randomly distributed, given that participants were originally recruited from 11 U.S. states in 1976. For example, the Boston area, and the Northeast more broadly, was initially overrepresented. However, few nationwide populations have been followed with repeated sleep and aircraft noise measurements over time. Similarly, airports were not randomly selected into this study, but were included where operations data were available. However, the respective geographic coverage of both participants and airports were still wide over the study period, and the airports included in the study captured the vast majority of enplanements annually in the United States.⁵⁰ We did

not directly account for noise reduction policies of individual airports. However, this would be indirectly reflected in the aircraft noise estimates. In addition, although the Aviation Safety and Noise Abatement Act of 1976 and the Airport Noise and Capacity Act of 1990 established a legal framework for abatement corresponding to a threshold of DNL 65 dB,^{46,47} “airport sponsors have limited proprietary authority to restrict access as a means of reducing aircraft noise impacts” to local communities,¹⁰⁸ and, in practice, this authority is rarely exercised. There are currently no state or federal policies directly limiting aircraft noise in the United States.⁴⁶

Conclusions

The increasing recognition of the importance of adequate sleep for maintaining health and optimal daytime functioning has spurred research aimed at identifying modifiable factors for improving sleep duration and quality. Environmental risk factors—including noise pollution—represent targets for improving sleep health that have been underinvestigated. Estimated at participant’s home addresses, multiple metrics of aircraft noise were associated with self-reported short sleep duration even after adjustment for environmental characteristics, including greenness and LAN. Short sleep duration was associated with both L_{night} and DNL, and the L_{night} association varied by individual, area, and airport characteristics, including region, living near a major cargo airport, living near a water-adjacent airport, and by self-reported level of hearing loss. We found evidence for adverse effects on sleep at exposures as low as DNL 45 dB(A), the lowest modeled noise level, and evidence further showed an exposure–response relationship between aircraft noise and short sleep duration. There was little evidence that aircraft noise was associated with sleep quality as assessed by questionnaire at study baseline across most levels of aircraft noise exposure.

Acknowledgments

We thank the participants and staff of the Nurses’ Health Study; this study would not be possible without their valuable

contributions. E.J. Nelson helped prepare the numerous tables in this manuscript and its Supplemental Material.

This study was funded by the Federal Aviation Administration (FAA; 13-C-AJFE-BU-016, to J.L.P.), under the Aviation Sustainability Center. Any opinions, findings, conclusions, or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the FAA. S.T.G. and D.D.N. were supported by the National Institute of Environmental Health Sciences T32 training grant T32ES014562. J.L.P., J.I.L., and D.D.N. were additionally supported on R01ES025791-01A1, and J.E.H. and F.L. were additionally supported on P30 ES000002. S.R. was partially supported by National Heart, Lung, and Blood Institute grant R35 HL135818. We acknowledge the contribution made by the John A. Volpe National Transportation Systems Center in developing the aircraft noise contours. The Nurses' Health Studies are supported by UM1 CA186107, R01 HL150119, R01 HL034594, U01 CA176726, R01 HL35464, U01 HL145386.

References

1. Institute of Medicine Committee on Sleep Medicine and Research. 2006. *Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem*. Colten HR, Altevogt BM, eds. Washington, DC: National Academies Press.
2. Spiegel K, Tasali E, Leproult R, Van Cauter E. 2009. Effects of poor and short sleep on glucose metabolism and obesity. *Nat Rev Endocrinol* 5(5):253–261, PMID: 19444258, <https://doi.org/10.1038/nrendo.2009.23>.
3. Knutson KL, Spiegel K, Penev P, Van Cauter E. 2007. The metabolic consequences of sleep deprivation. *Sleep Med Rev* 11(3):163–178, PMID: 17442599, <https://doi.org/10.1016/j.smrv.2007.01.002>.
4. Besedovsky L, Lange T, Haack M. 2019. The sleep-immune crosstalk in health and disease. *Physiol Rev* 99(3):1325–1380, PMID: 30920354, <https://doi.org/10.1152/physrev.00010.2018>.
5. Spiegel K, Leproult R, Van Cauter E. 1999. Impact of sleep debt on metabolic and endocrine function. *Lancet* 354(9188):1435–1439, PMID: 10543671, [https://doi.org/10.1016/S0140-6736\(99\)01376-8](https://doi.org/10.1016/S0140-6736(99)01376-8).
6. Sivertsen B, Lallukka T, Salo P, Pallesen S, Hysing M, Krokstad S, et al. 2014. Insomnia as a risk factor for ill health: results from the large population-based prospective HUNT study in Norway. *J Sleep Res* 23(2):124–132, PMID: 24635564, <https://doi.org/10.1111/jsr.12102>.
7. Salo P, Sivertsen B, Oksanen T, Sjösten N, Pentti J, Virtanen M, et al. 2012. Insomnia symptoms as a predictor of incident treatment for depression: prospective cohort study of 40,791 men and women. *Sleep Med* 13(3):278–284, PMID: 22177343, <https://doi.org/10.1016/j.sleep.2011.06.022>.
8. Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. 2010. Quantity and quality of sleep and incidence of type 2 diabetes: a systematic review and meta-analysis. *Diabetes Care* 33(2):414–420, PMID: 19910503, <https://doi.org/10.2337/dc09-1124>.
9. Ayas NT, White DP, Al-Delaimy WK, Manson JE, Stampfer MJ, Speizer FE, et al. 2003. A prospective study of self-reported sleep duration and incident diabetes in women. *Diabetes Care* 26(2):380–384, PMID: 12547866, <https://doi.org/10.2337/diacare.26.2.380>.
10. Baden MY, Hu FB, Vetter C, Schernhammer E, Redline S, Huang T. 2020. Sleep duration patterns in early to middle adulthood and subsequent risk of type 2 diabetes in women. *Diabetes Care* 43(6):1219–1226, PMID: 32209646, <https://doi.org/10.2337/dc19-2371>.
11. Cappuccio FP, Cooper D, D'Elia L, Strazzullo P, Miller MA. 2011. Sleep duration predicts cardiovascular outcomes: a systematic review and meta-analysis of prospective studies. *Eur Heart J* 32(12):1484–1492, PMID: 21300732, <https://doi.org/10.1093/eurheartj/ehr007>.
12. Ananthakrishnan AN, Khalili H, Konijeti GG, Higuchi LM, de Silva P, Fuchs CS, et al. 2014. Sleep duration affects risk for ulcerative colitis: a prospective cohort study. *Clin Gastroenterol Hepatol* 12(11):1879–1886, PMID: 24780288, <https://doi.org/10.1016/j.cgh.2014.04.021>.
13. Ayas NT, White DP, Manson JE, Stampfer MJ, Speizer FE, Malhotra A, et al. 2003. A prospective study of sleep duration and coronary heart disease in women. *Arch Intern Med* 163(2):205–209, PMID: 12546611, <https://doi.org/10.1001/archinte.163.2.205>.
14. Luyster FS, Strollo PJ Jr, Zee PC, Walsh JK, Boards of Directors of the American Academy of Sleep Medicine and the Sleep Research Society. 2012. Sleep: a health imperative. *Sleep* 35(6):727–734, PMID: 22654183, <https://doi.org/10.5665/sleep.1846>.
15. McMullan CJ, Curhan GC, Forman JP. 2016. Association of short sleep duration and rapid decline in renal function. *Kidney Int* 89(6):1324–1330, PMID: 27165820, <https://doi.org/10.1016/j.kint.2015.12.048>.
16. Stenholm S, Kronholm E, Bandinelli S, Guralnik JM, Ferrucci L. 2011. Self-reported sleep duration and time in bed as predictors of physical function decline: results from the InCHIANTI study. *Sleep* 34(11):1583–1593, PMID: 22043129, <https://doi.org/10.5665/sleep.1402>.
17. Chen H, Schernhammer E, Schwarzschild MA, Ascherio A. 2006. A prospective study of night shift work, sleep duration, and risk of Parkinson's disease. *Am J Epidemiol* 163(8):726–730, PMID: 16495472, <https://doi.org/10.1093/aje/kwj096>.
18. Tworoger SS, Lee S, Schernhammer ES, Grodstein F. 2006. The association of self-reported sleep duration, difficulty sleeping, and snoring with cognitive function in older women. *Alzheimer Dis Assoc Disord* 20(1):41–48, PMID: 16493235, <https://doi.org/10.1097/01.wad.0000201850.52707.80>.
19. Rod NH, Vahtera J, Westerlund H, Kivimäki M, Zins M, Goldberg M, et al. 2011. Sleep disturbances and cause-specific mortality: results from the GAZEL cohort study. *Am J Epidemiol* 173(3):300–309, PMID: 21193534, <https://doi.org/10.1093/aje/kwq371>.
20. Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. 2010. Sleep duration and all-cause mortality: a systematic review and meta-analysis of prospective studies. *Sleep* 33(5):585–592, PMID: 20469800, <https://doi.org/10.1093/sleep/33.5.585>.
21. Patel SR, Ayas NT, Malhotra MR, White DP, Schernhammer ES, Speizer FE, et al. 2004. A prospective study of sleep duration and mortality risk in women. *Sleep* 27(3):440–444, PMID: 15164896, <https://doi.org/10.1093/sleep/27.3.440>.
22. Fink D. 2019. A new definition of noise: noise is unwanted and/or harmful sound. Noise is the new 'secondhand smoke.' *Proc Meet Acoust* 39:050002. <https://doi.org/10.1121/2.0001186>.
23. Oswald I, Taylor AM, Treisman M. 1960. Discriminative responses to stimulation during human sleep. *Brain* 83(3):440–453, PMID: 13731563, <https://doi.org/10.1093/brain/83.3.440>.
24. Basner M, Griefahn B, van den Berg M. 2010. Aircraft noise effects on sleep: mechanisms, mitigation and research needs. *Noise Health* 12(47):95–109, PMID: 20472955, <https://doi.org/10.4103/1463-1741.63210>.
25. Raschke F. 2004. Arousals and aircraft noise—environmental disorders of sleep and health in terms of sleep medicine. *Noise Health* 6(22):15–26, PMID: 15070525.
26. Kryter KD. 1966. Psychological reactions to aircraft noise. *Science* 151(3716):1346–1355, PMID: 5909784, <https://doi.org/10.1126/science.151.3716.1346>.
27. Scheuch K. 2004. Special assessment of aircraft noise effects during night by the Council of Experts for Environmental Questions of FRG. *Noise Health* 6(22):65–67, PMID: 15070531.
28. Smith MG, Rocha S, Witte M, Basner M. 2020. On the feasibility of measuring physiologic and self-reported sleep disturbance by aircraft noise on a national scale: a pilot study around Atlanta airport. *Sci Total Environ* 718:137368, PMID: 32092522, <https://doi.org/10.1016/j.scitotenv.2020.137368>.
29. Elmenhorst EM, Griefahn B, Rolny V, Basner M. 2019. Comparing the effects of road, railway, and aircraft noise on sleep: exposure–response relationships from pooled data of three laboratory studies. *Int J Environ Res Public Health* 16(6):1073, PMID: 30917492, <https://doi.org/10.3390/ijerph16061073>.
30. McGuire S, Müller U, Elmenhorst EM, Basner M. 2016. Inter-individual differences in the effects of aircraft noise on sleep fragmentation. *Sleep* 39(5):1107–1110, PMID: 26856901, <https://doi.org/10.5665/sleep.5764>.
31. Nassur AM, Léger D, Lefèvre M, Elbaz M, Miettlicki F, Nguyen P, et al. 2019. The impact of aircraft noise exposure on objective parameters of sleep quality: results of the DEBATS study in France. *Sleep Med* 54:70–77, PMID: 30529780, <https://doi.org/10.1016/j.sleep.2018.10.013>.
32. Basner M, Witte M, McGuire S. 2019. Aircraft noise effects on sleep—results of a pilot study near Philadelphia International Airport. *Int J Environ Res Public Health* 16(17):3178, PMID: 31480420, <https://doi.org/10.3390/ijerph16173178>.
33. Nassur AM, Léger D, Lefèvre M, Elbaz M, Miettlicki F, Nguyen P, et al. 2019. Effects of aircraft noise exposure on heart rate during sleep in the population living near airports. *Int J Environ Res Public Health* 16(2):269, PMID: 30669300, <https://doi.org/10.3390/ijerph16020269>.
34. Nassur AM, Lefèvre M, Laumon B, Léger D, Evrard AS. 2019. Aircraft noise exposure and subjective sleep quality: the results of the DEBATS study in France. *Behav Sleep Med* 17(4):502–513, PMID: 29172716, <https://doi.org/10.1080/15402002.2017.1409224>.
35. Rocha S, Smith MG, Witte M, Basner M. 2019. Survey results of a pilot sleep study near Atlanta International Airport. *Int J Environ Res Public Health* 16(22):4321, PMID: 31698800, <https://doi.org/10.3390/ijerph16224321>.
36. Perron S, Plante C, Ragettli MS, Kaiser DJ, Goudreau S, Smargiassi A. 2016. Sleep disturbance from road traffic, railways, airplanes and from total environmental noise levels in Montreal. *Int J Environ Res Public Health* 13(8):809, PMID: 27529260, <https://doi.org/10.3390/ijerph13080809>.
37. Franssen EAM, van Wiechen CMAG, Nagelkerke NJD, Lebreit E. 2004. Aircraft noise around a large international airport and its impact on general health and medication use. *Occup Environ Med* 61(5):405–413, PMID: 15090660, <https://doi.org/10.1136/oem.2002.005488>.

38. Brink M, Schäffer B, Vienneau D, Pieren R, Foraster M, Eze IC, et al. 2019. Self-reported sleep disturbance from road, rail and aircraft noise: exposure–response relationships and effect modifiers in the SIRENE study. *Int J Environ Res Public Health* 16(21):4186, PMID: 31671890, <https://doi.org/10.3390/ijerph16214186>.
39. Holt JB, Zhang X, Sizov N, Croft JB. 2015. Airport noise and self-reported sleep insufficiency, United States, 2008 and 2009. *Prev Chronic Dis* 12(4):140551, PMID: 25880768, <https://doi.org/10.5888/pcd12.140551>.
40. Knutsson A. 2003. Health disorders of shift workers. *Occup Med (Lond)* 53(2):103–108, PMID: 12637594, <https://doi.org/10.1093/occmed/kqg048>.
41. Muzet A. 2007. Environmental noise, sleep and health. *Sleep Med Rev* 11(2):135–142, PMID: 17317241, <https://doi.org/10.1016/j.smrv.2006.09.001>.
42. Billings ME, Hale L, Johnson DA. 2020. Physical and social environment relationship with sleep health and disorders. *Chest* 157(5):1304–1312, PMID: 31870910, <https://doi.org/10.1016/j.chest.2019.12.002>.
43. Grigsby-Toussaint DS, Turi KN, Krupa M, Williams NJ, Pandi-Perumal SR, Jean-Louis G. 2015. Sleep insufficiency and the natural environment: results from the US Behavioral Risk Factor Surveillance System survey. *Prev Med* 78:78–84, PMID: 26193624, <https://doi.org/10.1016/j.ypmed.2015.07.011>.
44. Astell-Burt T, Feng X, Kolt GS. 2013. Does access to neighbourhood green space promote a healthy duration of sleep? Novel findings from a cross-sectional study of 259 319 Australians. *BMJ Open* 3(8):e003094, PMID: 23943772, <https://doi.org/10.1136/bmjopen-2013-003094>.
45. Schäffer B, Brink M, Schlatter F, Vienneau D, Wunderli JM. 2020. Residential green is associated with reduced annoyance to road traffic and railway noise but increased annoyance to aircraft noise exposure. *Environ Int* 143:105885, PMID: 32619911, <https://doi.org/10.1016/j.envint.2020.105885>.
46. U.S. Congress. 1980. Public Law 96-193: Aviation Safety and Noise Abatement Act of 1979. <https://www.govtrack.us/congress/bills/96/hr2440/text> [accessed 25 July 2002].
47. U.S. Congress. 1990. Aircraft Noise and Capacity Act of 1990 (“ANCA”). https://www.lawa.org/-/media/lawa-web/environment/files/airport_noise_and_capacity_act_of_1990.ashx?la=en&hash=908B516999C1AE9F81A7852B0233081E707C1869 [accessed 25 July 2002].
48. Bao Y, Bertola ML, Lenart EB, Stampfer MJ, Willett WC, Speizer FE, et al. 2016. Origin, methods, and evolution of the three Nurses’ Health Studies. *Am J Public Health* 106(9):1573–1581, PMID: 27459450, <https://doi.org/10.2105/AJPH.2016.303338>.
49. Morabia A. 2016. 120 000 Nurses who shook public health. *Am J Public Health* 106(9):1528–1529, PMID: 27509274, <https://doi.org/10.2105/AJPH.2016.303345>.
50. FAA (Federal Aviation Administration). 2021. Passenger boarding (enplanement) and all-cargo data for U.S. airports. https://www.faa.gov/airports/planning_capacity/passenger_allcargo_stats/passenger/ [accessed 17 August 2021].
51. Consensus Conference Panel, Watson NF, Badr MS, Belenky G, Bliwise DL, Buxton OM, et al. 2015. Recommended amount of sleep for a healthy adult: a joint consensus statement of the American Academy of Sleep Medicine and Sleep Research Society. *J Clin Sleep Med* 11(6):591–592, PMID: 25979105, <https://doi.org/10.5664/jcs.m.4758>.
52. Kim CS, Grady ST, Hart JE, Laden F, VoPham T, Nguyen DD, et al. 2022. Long-term aircraft noise exposure and risk of hypertension in the Nurses’ Health Studies. *Environ Res* 207:112195, PMID: 34627796, <https://doi.org/10.1016/j.envres.2021.112195>.
53. Simon MC, Hart JE, Levy JI, VoPham T, Malwitz A, Nguyen D, et al. 2022. Sociodemographic patterns of exposure to civil aircraft noise in the United States. *Environ Health Perspect* 130(2):027009, PMID: 35167327, <https://doi.org/10.1289/EHP9307>.
54. FAA. 2020. Report to Congress: FAA Reauthorization Act of 2018 (Pub. L. 115-254): Section 188 and Sec 173. https://www.faa.gov/sites/faa.gov/files/2021-11/Day-Night_Average_Sound_Levels_COMPLETED_report_w_letters.pdf [accessed 6 October 2021].
55. World Health Organization (WHO) Regional Office for Europe. 2018. WHO Environmental Noise Guidelines for the European Region. <https://www.euro.who.int/en/publications/abstracts/environmental-noise-guidelines-for-the-european-region-2018>.
56. Textor J, van der Zander B, Gilthorpe MS, Liškiewicz M, Ellison GTH. 2016. Robust causal inference using directed acyclic graphs: the R package ‘dagitty.’ *Int J Epidemiol* 45(6):1887–1894, PMID: 28089956, <https://doi.org/10.1093/ije/dyw341>.
57. Bethae TN, Zhou ES, Schernhammer ES, Castro-Webb N, Cozier YC, Rosenberg L. 2020. Perceived racial discrimination and risk of insomnia among middle-aged and elderly Black women. *Sleep* 43(1):zsz208, PMID: 31555803, <https://doi.org/10.1093/sleep/zsz208>.
58. Lidfeldt J, Li TY, Hu FB, Manson JE, Kawachi I. 2007. A prospective study of childhood and adult socioeconomic status and incidence of type 2 diabetes in women. *Am J Epidemiol* 165(8):882–889, PMID: 17284723, <https://doi.org/10.1093/aje/kwk078>.
59. Billings ME, Gold D, Szpiro A, Aaron CP, Jorgensen N, Gasset A, et al. 2019. The association of ambient air pollution with sleep apnea: the Multi-Ethnic Study of Atherosclerosis. *Ann Am Thorac Soc* 16(3):363–370, PMID: 30571166, <https://doi.org/10.1513/AnnalsATS.201804-248OC>.
60. Shen YL, Liu WT, Lee KY, Chuang HC, Chen HW, Chuang KJ. 2018. Association of PM_{2.5} with sleep-disordered breathing from a population-based study in northern Taiwan urban areas. *Environ Pollut* 233:109–113, PMID: 29059625, <https://doi.org/10.1016/j.envpol.2017.10.052>.
61. Yanosky JD, Paciorek CJ, Laden F, Hart JE, Puett RC, Liao D, et al. 2014. Spatio-temporal modeling of particulate air pollution in the conterminous United States using geographic and meteorological predictors. *Environ Health* 13(1):1–15, PMID: 25097007, <https://doi.org/10.1186/1476-069X-13-63>.
62. Johnson BS, Malecki KM, Peppard PE, Beyer KMM. 2018. Exposure to neighborhood green space and sleep: evidence from the Survey of the Health of Wisconsin. *Sleep Health* 4(5):413–419, PMID: 30241655, <https://doi.org/10.1016/j.sleh.2018.08.001>.
63. Yeager R, Riggs DW, DeJarnett N, Tollerud DJ, Wilson J, Conklin DJ, et al. 2018. Association between residential greenness and cardiovascular disease risk. *J Am Heart Assoc* 7(24):e009117, PMID: 30561265, <https://doi.org/10.1161/JAHA.118.009117>.
64. Krieger FJ, Malila WA, Nalepka RF, Richardson W. 1969. Preprocessing transformations and their effects on multispectral recognition. In: *Proceedings of the Sixth International Symposium on Remote Sensing of Environment, Volume II*. 13–16 October 1969. Ann Arbor, MI: Infrared and Optics Laboratory, Willow Run Laboratories, Environmental Research Institute of Michigan, 97–131.
65. Ohayon MM, Milesi C. 2016. Artificial outdoor nighttime lights associate with altered sleep behavior in the American general population. *Sleep* 39(6):1311–1320, PMID: 27091523, <https://doi.org/10.5665/sleep.5860>.
66. Chepesiuk R. 2009. Missing the dark: health effects of light pollution. *Environ Health Perspect* 117(1):A20–A27, PMID: 19165374, <https://doi.org/10.1289/ehp.117-a20>.
67. Cho YM, Ryu SH, Lee BR, Kim KH, Lee E, Choi J. 2015. Effects of artificial light at night on human health: a literature review of observational and experimental studies applied to exposure assessment. *Chronobiol Int* 32(9):1294–1310, PMID: 26375320, <https://doi.org/10.3109/07420528.2015.1073158>.
68. James P, Bertrand KA, Hart JE, Schernhammer ES, Tamimi RM, Laden F. 2017. Outdoor light at night and breast cancer incidence in the Nurses’ Health Study II. *Environ Health Perspect* 125(8):087010, PMID: 28886600, <https://doi.org/10.1289/EHP935>.
69. NOAA (National Oceanic and Atmospheric Administration). 2014. Global radiance calibrated nighttime lights. https://ngdc.noaa.gov/eog/dmsp/download_radcal.html [accessed 8 August 2021].
70. Kredlow MA, Capozzoli MC, Hearon BA, Calkins AW, Otto MW. 2015. The effects of physical activity on sleep: a meta-analytic review. *J Behav Med* 38(3):427–449, PMID: 25596964, <https://doi.org/10.1007/s10865-015-9617-6>.
71. Hirsch JA, Diez Roux AV, Moore KA, Evenson KR, Rodriguez DA. 2014. Change in walking and body mass index following residential relocation: the Multi-Ethnic Study of Atherosclerosis. *Am J Public Health* 104(3):e49–e56, PMID: 24432935, <https://doi.org/10.2105/AJPH.2013.301773>.
72. Johnson DA, Hirsch JA, Moore KA, Redline S, Diez Roux AV. 2018. Associations between the built environment and objective measures of sleep. *Am J Epidemiol* 187(5):941–950, PMID: 29547912, <https://doi.org/10.1093/aje/kwx302>.
73. Johnson DA, Drake C, Joseph CLM, Krajaenta R, Hudgel DW, Cassidy-Bushrow AE. 2015. Influence of neighbourhood-level crowding on sleep-disordered breathing severity: mediation by body size. *J Sleep Res* 24(5):559–565, PMID: 25950087, <https://doi.org/10.1111/jsr.12305>.
74. Hale L, Do DP. 2007. Racial differences in self-reports of sleep duration in a population-based study. *Sleep* 30(9):1096–1103, PMID: 17910381, <https://doi.org/10.1093/sleep/30.9.1096>.
75. Watson NF, Horn E, Duncan GE, Buchwald D, Vitiello MV, Turkheimer E. 2016. Sleep duration and area-level deprivation in twins. *Sleep* 39(1):67–77, PMID: 26285009, <https://doi.org/10.5665/sleep.5320>.
76. Fuller-Rowell TE, Curtis DS, El-Sheikh M, Chae DH, Boylan JM, Ryff CD. 2016. Racial disparities in sleep: the role of neighborhood disadvantage. *Sleep Med* 27–28:1–8, PMID: 27938909, <https://doi.org/10.1016/j.sleep.2016.10.008>.
77. Bassett E, Moore S. 2014. Neighbourhood disadvantage, network capital and restless sleep: is the association moderated by gender in urban-dwelling adults? *Soc Sci Med* 108:185–193, PMID: 24650740, <https://doi.org/10.1016/j.socscimed.2014.02.029>.
78. Johnson DA, Brown DL, Morgenstern LB, Meurer WJ, Lisabeth LD. 2015. The association of neighborhood characteristics with sleep duration and daytime sleepiness. *Sleep Health* 1(3):148–155, PMID: 29073435, <https://doi.org/10.1016/j.sleh.2015.06.002>.
79. Johnson DA, Lisabeth L, Hickson D, Johnson-Lawrence V, Samdarshi T, Taylor H, et al. 2016. The social patterning of sleep in African Americans: associations of socioeconomic position and neighborhood characteristics with sleep in the Jackson Heart Study. *Sleep* 39(9):1749–1759, PMID: 27253767, <https://doi.org/10.5665/sleep.6106>.

80. DeSantis AS, Diez Roux AV, Moore K, Baron KG, Mujahid MS, Javier Nieto FJ. 2013. Associations of neighborhood characteristics with sleep timing and quality: the Multi-Ethnic Study of Atherosclerosis. *Sleep* 36(10):1543–1551, PMID: 24082314, <https://doi.org/10.5665/sleep.3054>.
81. Schisterman EF, Cole SR, Platt RW. 2009. Overadjustment bias and unnecessary adjustment in epidemiologic studies. *Epidemiology* 20(4):488–495, PMID: 19525685, <https://doi.org/10.1097/EDE.0b013e3181a819a1>.
82. DeVille NV, Iyer HS, Holland I, Bhupathiraju SN, Chai B, James P, et al. 2022. Neighborhood socioeconomic status and mortality in the Nurses' Health Study (NHS) and the Nurses' Health Study II (NHSII). *Environ Epidemiol* 7(1): e235, PMID: 36777531, <https://doi.org/10.1097/EE9.0000000000000235>.
83. Liu Y, Wheaton AG, Chapman DP, Cunningham TJ, Lu H, Croft JB. 2016. Prevalence of healthy sleep duration among adults—United States, 2014. *MMWR Morb Mortal Wkly Rep* 65(6):137–141, PMID: 26890214, <https://doi.org/10.15585/mmwr.mm6506a1>.
84. Obradovich N, Migliorini R, Mednick SC, Fowler JH. 2017. Nighttime temperature and human sleep loss in a changing climate. *Sci Adv* 3(5):e1601555, PMID: 28560320, <https://doi.org/10.1126/sciadv.1601555>.
85. Liao C, Akimoto M, Bivolarova MP, Sekhar C, Laverge J, Fan X, et al. 2021. A survey of bedroom ventilation types and the subjective sleep quality associated with them in Danish housing. *Sci Total Environ* 798:149209, PMID: 34323281, <https://doi.org/10.1016/j.scitotenv.2021.149209>.
86. Rasker R, Gude PH, Gude JA, van den Noort J. 2009. The economic importance of air travel in high-amenity rural areas. *J Rural Stud* 25(3):343–353, <https://doi.org/10.1016/j.jrurstud.2009.03.004>.
87. Grandner MA, Smith TE, Jackson N, Jackson T, Burgard S, Branas C. 2015. Geographic distribution of insufficient sleep across the United States: a county-level hotspot analysis. *Sleep Health* 1(3):158–165, PMID: 26989761, <https://doi.org/10.1016/j.sleh.2015.06.003>.
88. Johnson DA, Thorpe RJ, McGrath JA, Jackson WB, Jackson CL. 2018. Black–White differences in housing type and sleep duration as well as sleep difficulties in the United States. *Int J Environ Res Public Health* 15(4):564, PMID: 29561769, <https://doi.org/10.3390/ijerph15040564>.
89. U.S. Census Bureau. 1997. American housing brief: regional differences in housing. <https://www.census.gov/content/dam/Census/programs-surveys/ahs/publications/ahb-9501.pdf> [accessed 14 December 2022].
90. San Francisco International Airport. 2021. SFO flight patterns and operations. <https://www.flysfo.com/about/community-noise/noise-office/flight-patterns-operations> [accessed 9 June 2021].
91. Leleu C, Marsh D. 2009. Dependent on the dark: cargo and other night flights in European airspace. <https://www.eurocontrol.int/sites/default/files/publication/files/tat5-night-freight-report.pdf> [accessed 9 June 2021].
92. FAA (Federal Aviation Administration). 2020. Aviation Environmental Design Tool (AEDT). Version 3c. User Manual. https://aedt.faa.gov/Documents/AEDT3c_UserManual.pdf [accessed 18 December 2022].
93. Konishi K, Maekawa Z. 2001. Interpretation of long term data measured continuously on long range sound propagation over sea surfaces. *Appl Acoust* 62(10):1183–1210, [https://doi.org/10.1016/S0003-682X\(00\)00096-7](https://doi.org/10.1016/S0003-682X(00)00096-7).
94. Konishi K, Tanioku Y, Maekawa Z. 2000. Long time measurement of long range sound propagation over an ocean surface. *Appl Acoust* 61(2):149–172, [https://doi.org/10.1016/S0003-682X\(00\)00009-8](https://doi.org/10.1016/S0003-682X(00)00009-8).
95. Basner M, McGuire S. 2018. WHO Environmental Noise Guidelines for the European Region: a systematic review on environmental noise and effects on sleep. *Int J Environ Res Public Health* 15(3):519, PMID: 29538344, <https://doi.org/10.3390/ijerph15030519>.
96. Basner M, Samel A. 2005. Effects of nocturnal aircraft noise on sleep structure. *Somnologie* 9(2):84–95, <https://doi.org/10.1111/j.1439-054X.2005.00051.x>.
97. Rudolph KE, Shev A, Paksarian D, Merikangas KR, Mennitt DJ, James P, et al. 2019. Environmental noise and sleep and mental health outcomes in a nationally representative sample of urban US adolescents. *Environ Epidemiol* 3(4): e056, PMID: 31538137, <https://doi.org/10.1097/EE9.0000000000000056>.
98. Bureau of Transportation Statistics. 2021. Air cargo summary data (all). <https://www.transtats.bts.gov/freight.asp> [accessed 26 October 2021].
99. Airports Council International–North America Cargo Committee. 2019. *Air Cargo Guide*. <https://airportscouncil.org/wp-content/uploads/2020/03/Air-Cargo-Guide.pdf> [accessed 30 November 2022].
100. Dožić S, Babić D. 2015. Trends in airline cargo fleet. In: *2nd Logistics International Conference*. 21–23 May 2015. Belgrade, Serbia: LOGIC, 1–6.
101. IATA (International Air Transport Association). 2022. *Annual Review 2022*. <https://www.iata.org/contentassets/c81222d96c9a4e0bb4ff6ced0126f0bb/annual-review-2022.pdf> [accessed 15 December 2022].
102. Zaporozhets O. 2016. Aircraft noise models for assessment of noise around airports—improvements and limitations. https://www.icao.int/environmental-protection/Documents/EnvironmentalReports/2016/ENVReport2016_pg50-55.pdf [accessed 25 July 2022].
103. Liu J, Liu J, Lai D, Pei J, Wei S. 2021. A field investigation of the thermal environment and adaptive thermal behavior in bedrooms in different climate regions in China. *Indoor Air* 31(3):887–898, PMID: 33350529, <https://doi.org/10.1111/ina.12775>.
104. Wensveen JG. 2016. Managerial aspects of airlines. In: *Air Transportation: A Management Perspective*. 8th ed. New York, NY: Routledge, 423–452.
105. Lin FR, Niparko JK, Ferrucci L. 2011. Hearing loss prevalence in the United States. *Arch Intern Med* 171(20):1851–1852, PMID: 22083573, <https://doi.org/10.1001/archinternmed.2011.506>.
106. Basner M, Brink M, Bristow A, de Kluizenaar Y, Finegold L, Hong J, et al. 2015. ICBEN review of research on the biological effects of noise 2011–2014. *Noise Health* 17(75):57–82, PMID: 25774609, <https://doi.org/10.4103/1463-1741.153373>.
107. Hauptman M, Lockley SW, Landrigan CP. 2022. Invited commentary: there's no place like home—integrating a place-based approach to understanding sleep. *Am J Epidemiol* 191(9):1540–1543, PMID: 35524716, <https://doi.org/10.1093/aje/kwac086>.
108. FAA. 2021. Airport noise and access restrictions. In: *Airport Compliance Manual: Order 5190.6B Change 1*. Washington, DC: FAA. <https://www.federalregister.gov/documents/2021/12/03/2021-25936/airport-compliance-program> [accessed 2 April 2023].