

# UC Davis

## UC Davis Previously Published Works

### Title

A variety-specific analysis of climate change effects on California winegrapes

### Permalink

<https://escholarship.org/uc/item/3cd4358z>

### Journal

International Journal of Biometeorology, 68(8)

### ISSN

0020-7128

### Authors

Parker, Lauren E

Zhang, Ning

Abatzoglou, John T

et al.

### Publication Date

2024-08-01

### DOI

10.1007/s00484-024-02684-8

Peer reviewed



# A variety-specific analysis of climate change effects on California winegrapes

Lauren E. Parker<sup>1,2</sup> · Ning Zhang<sup>1,3</sup> · John T. Abatzoglou<sup>4</sup> · Isaya Kisekka<sup>3,5</sup> · Andrew J. McElrone<sup>6,7</sup> · Steven M. Ostoja<sup>1,2,8</sup>

Received: 9 November 2023 / Revised: 15 February 2024 / Accepted: 15 April 2024 / Published online: 23 April 2024  
© The Author(s) 2024

## Abstract

California contains a broad geography over which climate conditions can be suitable for cultivating multiple varieties of winegrapes. However, climate change is projected to make winegrape cultivation more challenging across many of California's winegrowing regions. In order to understand the potential effects of climate change on winegrapes, this study models variety-specific phenology for six winegrape varieties and quantifies the change in phenology and viticulturally-important agroclimate metrics over 12 of California's American Viticultural Areas (AVAs) by the mid-21st century. Results show more rapid development for winegrapes with earlier budburst, flowering, veraison, and maturation across all varieties and AVAs. Cabernet Sauvignon shows the greatest change in phenology timing, while Chardonnay shows the least change. Likewise, the West Sonoma Coast AVA shows the greatest average change in phenology timing across varieties and development stages and Lodi AVA shows the least. Projected changes in agroclimatic metrics include an additional month of potentially damaging heat days (above 35 °C) in some AVAs, and decreases in frost days. These results have implications for numerous factors related to viticultural production, including water resources management and crop yield and quality, and underscore the need for California winegrape growers to improve their resilience to climate change by adopting strategies such as increasing soil health and water use efficiency and selecting cultivars suited for future climate conditions. By conducting climate effects analyses at the variety-specific and AVA scale, important information is provided to the winegrowing industry at a resolution that can support decision-making towards resilience.

**Keywords** Agroclimatic metrics · California · Climate change · Phenology · Winegrapes

## Introduction

Winegrapes (*Vitis vinifera*) are thought to have been first cultivated from wild relatives in southwest Asia as early as 7000 BCE (Bouby et al. 2013). Winegrape cultivation in California began in earnest in the early 19th century (Anderson et al. 2003) and as of 2022, California grows winegrapes on more than 2,428 square kilometers, yielding approximately 3.4 million crushed tons (CDFA 2022; CDFA 2023). This scale of viticulture and the accompanying wine production contributes > 400,000 jobs and >\$70 billion to the state economy and makes California the fourth largest wine producing region globally (CWI, 2022).

California's Mediterranean macroclimate and diverse mesoclimates, ranging from cool and coastal to warm and inland, offer a broad geography over which climate conditions can be suitable for cultivating many varieties of winegrapes. However, in California, climate change is projected

✉ Lauren E. Parker  
leparker@ucdavis.edu

<sup>1</sup> USDA California Climate Hub, Davis, CA 95616, USA

<sup>2</sup> Institute of the Environment, University of California Davis, Davis, CA 95616, USA

<sup>3</sup> Department of Land, Air and Water Resources, University of California Davis, Davis, CA 95616, USA

<sup>4</sup> Department of Management of Complex Systems, University of California Merced, Merced, CA 95343, USA

<sup>5</sup> Department of Biological and Agricultural Engineering, University of California, Davis 95616, USA

<sup>6</sup> USDA-ARS Crops Pathology and Genetics Research Unit, Davis, CA 95616, USA

<sup>7</sup> Department of Viticulture and Enology, University of California Davis, Davis, CA 95616, USA

<sup>8</sup> USDA-ARS Sustainable Agricultural Water Systems Research Unit, Davis, CA 95616, USA

to increase temperature, interannual precipitation variability, and the frequency and intensity of extreme heat events, as well as exacerbate drought and alter pest and disease pressure (Pathak et al. 2018). Unsurprisingly, climate change is projected to affect winegrape yield and quality across many of California's winegrowing regions (Hannah et al. 2013). To this point, projected increases in extreme heat exposure could slow grapevine development, reduce berry weight, negatively impact berry quality, and alter the chemical composition of winegrapes and subsequently the characteristics of wine (Parker et al. 2020a). Similarly, multiple studies have shown that climate change, and extreme heat in particular, may reduce yields and suitable growing regions for wine in California (Diffenbaugh et al. 2011; Monteverde and De Sales 2020; White et al. 2006).

Despite viticulture's well-known sensitivity to climate, the diversity of winegrape varieties allows for grape production across a wide range of climates around the world (Jones 2015; Jones and Webb 2010). In fact, it has been argued that varietal variation could provide an adaptive opportunity to climate change and potentially reduce the projected contraction of suitable growing locations among today's wine regions; this argument is predicated on the variation in physiological tolerances and phenology across winegrape varieties (Morales-Castilla et al. 2020; Wolkovich et al. 2017). Climatic tolerances vary not only across varieties but also throughout the growing season. For winegrapes, changes in the timing of phenological development can influence berry size, color, chemistry and wine quality; however, by matching the climate conditions to variety-specific phenology

and climatic tolerances, growers can produce characteristic winegrapes (Parker et al. 2020b).

Previous studies have shown that the effects of climate change on winegrapes and grape phenology will vary by location and variety (e.g., Ausseil et al. 2021; Hannah et al. 2013; Webb et al. 2007), and understanding potential climate risks at local scales is useful for management planning (e.g., Babin et al., 2022). The objective of this study is to explore the potential effects of projected climatic change on winegrape production in California across multiple winegrape-growing regions known as American Viticultural Areas (AVAs) and across multiple winegrape varieties. To do this, we model potential shifts in variety-specific phenology and quantify the change in viticulturally-important climate metrics at the AVA scale. Given the economic importance of California viticulture and the climate changes expected, improving the understanding of how climate change may affect winegrape production across growing regions and varieties will assist winegrape growers and the broader wine industry in identifying and prioritizing adaptation actions to meet location- and variety-specific climate-mediated challenges.

## Data and methods

### Data

#### Climatological data

Daily maximum and minimum temperature ( $T_x$ ,  $T_n$ ), precipitation (Pr), and reference evapotranspiration (ET<sub>o</sub>) for the contemporary (1991–2020) period were obtained from the 4-km gridded dataset (gridMET, <https://www.climatologylab.org/gridmet.html>) of Abatzoglou (2013). gridMET is a spatially-continuous, daily dataset of surface meteorological conditions developed using the PRISM dataset of PRISM Climate Group, Oregon State University (<https://prism.oregonstate.edu>, see also Daly et al. 1997 and Daly et al. 2008) and regional reanalysis data (NLDAS-2, see Xia et al. 2012) covering the contiguous United States from 1979 to present (Abatzoglou 2013). Daily  $T_x$ ,  $T_n$ , Pr, and ET<sub>o</sub> were also acquired for 20 global climate models (GCMs, Table 1) participating in the fifth coupled model intercomparison project (CMIP5) for the mid-21st century (2040–2069) period for representative concentration pathway (RCP) 4.5. RCP 4.5 represents a future scenario in which greenhouse gas emissions peak in the early-mid 21st century and then decline, resulting in moderate warming. These GCM data were obtained from the 4-km Multivariate Adaptive Constructed Analogs (MACA, <https://www.climatologylab.org/maca.html>) dataset of Abatzoglou and Brown (2012). MACA is

**Table 1** Global Climate Models participating in the 5th Coupled Model Intercomparison Project (CMIP5) used in this study

GCM	Country of Origin
bcc-csm1-1	China
bcc-csm1-1-m	China
BNU-ESM	China
CanESM2	Canada
CCSM4	USA
CNRM-CM5	France
CSIRO-Mk3-6-0	Australia
GFDL-ESM2M	USA
GFDL-ESM2G	USA
HadGEM2-CC365	United Kingdom
HadGEM2-ES365	United Kingdom
inmcm4	Russia
IPSL-CM5A-LR	France
IPSL-CM5A-MR	France
IPSL-CM5B-LR	France
MIROC5	Japan
MIRCO-ESM	Japan
MIROC-ESM-CHEM	Japan
MRI-CGCM3	Japan
NorESM1-M	Norway

a statistical downscaling method that applies a constructed analog approach for mapping daily GCM data to observed data (Abatzoglou and Brown 2012). The approach further applies bias correction using an equidistant quantile mapping methodology (Li et al. 2010). Here, we use MACA trained using gridMET data which provides interoperability between the historical observed data and future projections. For both gridMET and MACA, ETo is calculated for a well-watered grass surface using the Penman-Monteith method (Allen et al. 1998; Walter et al., 2000). These datasets were chosen because they provide a spatially continuous and temporally complete record of climate conditions suitable for local and landscape-scale agricultural research. RCPs were used rather than the newer Shared Socioeconomic Pathways (SSPs) used in the current sixth coupled model intercomparison project (CMIP6) because downscaled SSP data are currently limited. Studies have generally shown similarities between climate models participating in the CMIP5 and CMIP6 over California (Krantz et al., 2021) and broadly similar hydroclimatic changes (Cook et al., 2021). While other RCP scenarios exist, we focus our results on RCP 4.5 because it provides a more conservative measure of potential change; we also note that the variability between models exceeds the variability between RCPs at mid-century time horizons (Kharin et al. 2013). Elevation data were acquired from the digital elevation model (DEM) associated with the gridMET dataset.

### Winegrape data

Six winegrape varieties – 3 red varieties and 3 white varieties – were selected for analysis based on a combination of economic importance, acreage, and variety-specific information available in the literature for developing phenology models and environmental tolerance thresholds. The red winegrape varieties selected include Cabernet Sauvignon, Pinot Noir, and Zinfandel, and the white winegrape varieties selected are Chardonnay, Pinot Gris, and Sauvignon Blanc. As of 2022, these six varieties comprise the top three red and top three white winegrape varieties by dollar value, and collectively these six varieties comprise more than 70% of the total dollar value of winegrapes sold in California (CDFA, 2023).

### American viticultural area data

Twelve American Viticultural Areas (AVAs) within California were chosen as representative of a range of climate conditions that can characterize climate change effects broadly. The selected AVAs provide a range of mesoclimates, are similar in size, represent both heterogeneous and homogenous topography, and cover both coastal and inland

locations as well as Northern and Southern California locations. The relative importance of the AVAs to statewide winegrape production was also considered; although production acreage, crushed tons, and value are calculated at the crush district scale, which is a larger spatial scale than the AVA, discussion with industry experts helped to identify key AVAs. The 12 AVAs are: El Dorado, Livermore Valley, Lodi, Madera, Mendocino, Monterey, Napa Valley, Paso Robles, Russian River Valley, Santa Ynez Valley, San Luis Obispo (SLO) Coast, and West Sonoma Coast (Fig. 1). AVA shapefiles were downloaded from the U.S. Department of Treasury Alcohol and Tobacco Tax Trade Bureau's AVA Map Explorer (<https://www.ttb.gov/images/AVA/>). A grid cell was considered as part of an AVA if its center was located within the AVA shapefile.

### Variety-specific phenology modeling


Winegrape development is a complex, multi-phasic cycle that has been simplified for our assessments of climate effects on variety-specific phenology and exposure. Following Parker and Abatzoglou (2018), we developed models for each variety using published climatic thresholds (Table 2); models assessed cold hardiness and changes in the timing of chill completion, budburst, flowering, veraison, and maturity.

### Chill accumulation

Sufficient time exposed to cool temperatures, known as “chill,” is necessary for fruit development and good yields in many perennial crops (Luedeling et al. 2009), including winegrapes. There are multiple models for calculating chill accumulation; here we calculate chilling degree days ( $DD_C$ ) using hourly temperature ( $T$ ) and we use a base temperature ( $T_{base}$ ) of 10 °C for all six varieties accumulating from November to February, after Ferguson et al. (2014).

### Budburst, flowering, and veraison

As vineyards are exposed to warmer temperatures in spring they begin to bloom. Budburst marks the first major stage in winegrape development and is reached when 50% of dormant buds show green tissue (Zapata et al. 2017). Flowering is the second major stage of winegrape development; it is reached when 50% of flower caps have dropped (Zapata et al. 2017). With further heat accumulation, fruit begins to develop and winegrapes will change color during a development phase known as veraison. Budburst, flowering, and veraison occur when growing degree day (GDD) accumulation reaches the defined threshold for the given stage. GDD are units of heat accumulation commonly used to track the



**Fig. 1** The 12 American Viticultural Areas (AVAs) explored in this analysis are outlined in red. Two AVAs selected to illustrate within-AVA spatial variability of results are shaded in pink. For reference, California counties are represented on the map by thin, grey lines

development of crops during the growing season. For each of these stages, GDD accumulation is calculated following Zapata et al. (2017) using variety- and stage-specific  $T_{base}$ . Although the base temperatures provided in Zapata et al. (2017) are based on a January 1 start date for GDD

accumulation, prior experiments showed this approach to result in poor model performance over our areas of interest, with particularly early development modeled at lower latitudes with warmer winter and spring temperatures. To address this, we do not begin GDD accumulation until

**Table 2** Variety-specific thresholds used to assess the changes in the timing of chill completion, budburst, flowering, veraison, and maturity. All DDc values are from Ferguson et al. (2014); GDD for budburst, flowering and veraison are from Zapata et al. (2017); and BEDD for maturity are from Gladstones (1992). For definitions of DDc and GDD, refer to Table 3; for a definition of BEDD, refer to the manuscript subsection on *Maturity*

Grape Variety	Chill			Budburst		Flowering		Veraison		Maturity
	DD <sub>c</sub>	GDD	T <sub>base</sub> °C	GDD	T <sub>base</sub> °C	GDD	T <sub>base</sub> °C	GDD	T <sub>base</sub> °C	BEDD
Cabernet Sauvignon	700	101	8.3	240	10.4	598	12.5	1300		
Pinot Noir	300	79	8.1	256	9.7	578	12.1	1150		
Zinfandel	500	132	7.2	294	9.1	639	11.2	1200		
Chardonnay	600	114	6.5	354	8.2	727	9.7	1150		
Pinot Gris	400	130	6.9	313	8.3	678	10.4	1100		
Sauvignon Blanc	300	101	7.4	327	8.8	607	11.1	1150		

daylength > 11-hours based on the growing season index (Jolly et al. 2005); this modification follows latitude or day-length adjustments in other heat summation models in viticulture (e.g., Gladstones 1992; Huglin 1978).

### Maturity

Fruit development continues through summer, and when sufficient heat accumulation has occurred the winegrapes reach maturity. Winegrape harvest dates are influenced by non-climatic conditions such as grape chemistry preferences (e.g., brix and acid levels within the fruit); here we estimate winegrape maturity by applying a threshold for the biologically effective degree days (BEDD) needed to reach maturation (Gladstones 1992). BEDD incorporates two factors: one for adjusting the diurnal temperature range and another for correcting day length. The diurnal temperature range (DTR) is the difference between daily maximum and minimum temperatures, and when calculating BEDD the DTR factor increases when the DTR exceeds 13 °C and decreases when it falls below 10 °C. The day length correction factor ranges from 1.00 at 40° latitude to 1.045 at 50° latitude. For detailed calculations, please refer to Hall and Jones (2010). Here we begin BEDD accumulation on April 1. BEDD were used to determine the timing of maturity due to a lack of threshold information for GDD or other indices for maturity (e.g., the Huglin Index) in the literature for all our selected varieties. Our maturity analysis is not limited by established minimum and maximum BEDD values for quality winegrape production (Jones et al. 2010) as experiments showed all of our selected AVAs maintain BEDD within the bounds for production under both contemporary and future climates.

### Quantifying changes in agroclimatic metrics

Quantifying agriculturally-relevant climate metrics offers a picture of physiologically-important environmental conditions, and monitoring these metrics for changes over time can be useful for identifying potential adaptation needs

within cropping systems. We selected 14 metrics of importance to viticulture (Table 3). For the majority of the metrics, we take a generalized approach to calculations and we do not use any variety- or stage-specific values for T<sub>base</sub>. However, for cold hardiness and frost damage, we use variety- and stage-specific thresholds, respectively.


For each metric, we calculated both the annual value and the 30-year average value over the contemporary period and the future period at the scale of the 4-km gridMET and MACA data. These calculations were done for each of 20 GCMs and the 20-model mean values were computed to represent the average condition of the future period. The grid cell values were then averaged across each AVA, providing a measure of the metric at the AVA scale and at the annual and climatological time step. The difference between 30-year average values for the future and contemporary periods were calculated (future [minus] contemporary); temporal trends in each metric were also assessed for each AVA over each 30-year period using a Theil-Sen estimator, and a Mann-Kendall test was applied to determine the significance of the trend ( $\alpha=0.05$ ) with a null hypothesis of no significant trend. Finally, a supplemental analysis explored correlations between both phenology and agroclimatic metrics and annual mean temperature and elevation.

## Results

Across all varieties and AVAs, we show that phenology shifts towards later chill completion and earlier budburst, flowering, veraison, and maturation. While the between-variety difference in phenology changes is not large, Cabernet Sauvignon consistently shows the greatest change in phenology timing between the contemporary and future periods, while Chardonnay shows the least change. The between-AVA difference in phenology change is more pronounced with the West Sonoma Coast showing the greatest change in phenology timing across development stages and Lodi showing the least. Beyond phenology, results also show that climate change increases the incidence of some

**Table 3** 14 Agroclimatic metrics selected for this analysis, their calculation, and their relevance to viticulture

Agroclimate Metric	Calculation	Relevance to Viticulture
Growing Degree Days (GDD)	GDD are calculated following Zapata et al. (2017) using $T_{\text{lower}} = 10\text{ }^{\circ}\text{C}$ and $T_{\text{upper}} = 32\text{ }^{\circ}\text{C}$ . GDD over the growing season is accumulated from April to October.	GDD are used in phenology models to estimate the timing of crop (and/or pest) development, which can be useful for in-season planning. Increased GDD can speed up crop development.
Cold Hardiness ( $H_{\text{ini}}$ )	Cold hardiness is calculated as the number of days between September 21 and December 21 with temperatures below variety-specific thresholds (Ferguson et al. 2014): <i>Cabernet Sauvignon</i> : $-10.3^{\circ}\text{C}$ <i>Pinot Noir</i> : $-11.5^{\circ}\text{C}$ <i>Zinfandel</i> : $-10.4^{\circ}\text{C}$ <i>Chardonnay</i> : $-11.8^{\circ}\text{C}$ <i>Pinot Gris</i> : $-12.0^{\circ}\text{C}$ <i>Sauvignon Blanc</i> : $-10.6^{\circ}\text{C}$	Cold hardiness is the minimum overwinter temperature that can be tolerated without damage. As winter progresses, vines become increasingly cold tolerant. Here we use the cold tolerance thresholds associated with the fall season to provide a conservative view of potential exposure.
Chilling Degree Days ( $DD_{\text{C}}$ )	Chill accumulation is calculated as $DD_{\text{C}}$ using hourly $T$ and $T_{\text{base}} = 10\text{ }^{\circ}\text{C}$ accumulating from November to February after Ferguson et al. (2014).	Chill is required in order for vines to fully break bud. Insufficient chill can lead to erratic budbreak and can impact fruit set and yields. Warmer winter and spring temperatures can reduce or delay chill accumulation.
Frost Damage Days (FDD)	FDD is calculated as the total number of days per year with $T_{\text{n}}$ below the following stage-specific frost damage thresholds (Jones 2003): <i>Sap Bleeding</i> : $-2.5\text{ }^{\circ}\text{C}$ ; <i>Budburst</i> : $-2.2\text{ }^{\circ}\text{C}$ ; <i>Flowering</i> : $-0.5\text{ }^{\circ}\text{C}$ ; <i>Veraison</i> and <i>Maturation</i> : $0\text{ }^{\circ}\text{C}$	Spring frost can damage buds and young shoots, resulting in uneven ripening and declines in yield. Fall frosts can cause defoliation of the vines prior to harvest, risking sun scald.
Last Spring Freeze (LSF)	The last day of the calendar year prior to June 30 with $T_{\text{n}} \leq 0\text{ }^{\circ}\text{C}$ .	LSF is an important consideration for early-blooming and frost-sensitive perennials. Earlier LSF can also increase pest pressure.
First Fall Freeze (FFF)	The first day of the calendar year commencing July 1 with $T_{\text{n}} \leq 0\text{ }^{\circ}\text{C}$ .	FFF is important for late-maturing varieties that may suffer damage from a fall freeze.
Freeze-Free Season (FFS)	The difference – in number of days – between the LSF and FFF (FFF [minus] LSF).	A certain length of FFS is required for vines to complete their annual development cycle. The FFS can inform the geography of cultivation. Longer FFS can also increase pest pressure.
Hot Days (HD)	The number of days with $T_{\text{x}} > 35\text{ }^{\circ}\text{C}$ (Jones 2015) during the growing season (April – October).	Hot days can negatively affect grape development and yield (White et al. 2006). Impacts are dependent on HD timing and vine heat tolerance.
Heatwaves (HW)	Heatwave events are defined as 3 + consecutive days (Gershunov et al. 2009; Sheridan and Lee 2018) with $T_{\text{x}} > 98$ th percentile of 1991–2020 summer (June–August) $T_{\text{x}}$ .	Consecutive days with high temperatures can impact grape yield and quality, depending on the timing and absolute temperatures (Martínez-Lüscher et al. 2020). Additionally, heatwaves have implications for vineyard worker safety and productivity.
Diurnal Temperature Range (DTR)	DTR is the difference between daily $T_{\text{x}}$ and $T_{\text{n}}$ . We calculate DTR over August – October.	Reduced DTR can speed berry development and alter berry chemistry (Cohen et al. 2012a, b), though effects can vary by cultivar (Jones et al. 2012)
Diurnal Temperature Range $> 20\text{ }^{\circ}\text{C}$ (DTR20)	The number of days with $DTR > 20\text{ }^{\circ}\text{C}$ (White et al. 2006) over August to October.	High DTR in cool climate regions keeps malic acid and acidity in grapes and wine, while low DTR in warm regions makes grapes and wine more fruity and less acidic (Gladstones 1992).
Excess Precipitation Days ( $Pr_{\text{ex}}$ )	The number of days with precipitation $> 5\text{ mm}$ (Mosedale et al. 2015) during May to October.	Dry weather from May to October is preferred for optimum photosynthesis, ripening and balance. Excess precipitation during this period can impact bloom, promote diseases, and dilute berries (Jones 2003).
Winter Accumulated Precipitation ( $Pr_{\text{acc}}$ )	Total accumulated precipitation in mm from November of the previous year to January of the current year.	Winter precipitation accumulation is necessary for soil moisture recharge (Jones 2003).
Crop Evapotranspiration ( $ET_{\text{c}}$ )	$ET_{\text{c}}$ is the adjusted reference ET ( $ET_{\text{o}}$ ) using the crop coefficient ( $K_{\text{c}}$ ) ( $ET_{\text{c}} = ET_{\text{o}} * K_{\text{c}}$ ). The total $ET_{\text{c}}$ is accumulated over the growing season (April – October) with units in mm.	$ET_{\text{c}}$ is commonly used in irrigation models and decision support systems (Zhang et al. 2021). Changes in $ET_{\text{c}}$ can indicate changes in irrigation demand.



**Fig. 2** Variety-specific phenological response to climate change across six varieties and 12 AVAs. The left y-axis illustrates the change in timing between the RCP4.5 and the contemporary period for five significant phenological stages (Chill, Budburst, Flowering, Veraison, and Maturity). The 12 American Viticultural Areas (AVAs) are shown

along the x-axis in geographical order from north to south. The red lines correspond to the right y-axis and indicate the annual mean temperature during the contemporary period for each AVA, with error bars signifying the standard deviation

**Table 4a** The 12-AVA average DOY of chill completion and the onset of four key growing stages (budbreak, flowering, veraison, and maturity) for six grape varieties under contemporary (1991–2020, observed) and future (2040–2069, RCP 4.5) climate conditions. The DOY is computed as the mean of means across 12 California AVAs

Variety	CHILL (DOY)		BUDBREAK (DOY)		FLOWERING (DOY)		VERAISON (DOY)		MATURITY (DOY)	
	1991–2020 Obs.	2040–2069 RCP4.5	1991–2020 Obs.	2040–2069 RCP4.5	1991–2020 Obs.	2040–2069 RCP4.5	1991–2020 Obs.	2040–2069 RCP4.5	1991–2020 Obs.	2040–2069 RCP4.5
<b>Cabernet Sauvignon</b>	Dec. 27	Jan. 09	Mar. 28	Mar. 21	May 22	May 08	Aug. 10	Jul. 22	Sep. 23	Sep. 13
<b>Pinot Noir</b>	Dec. 01	Dec. 10	Mar. 22	Mar. 16	May 16	May 03	Aug. 02	Jul. 15	Sep. 07	Aug. 28
<b>Zinfandel</b>	Dec. 14	Dec. 25	Mar. 28	Mar. 22	May 20	May 08	Aug. 03	Jul. 16	Sep. 13	Sep. 02
<b>Chardonnay</b>	Dec. 21	Jan. 02	Mar. 22	Mar. 17	May 19	May 07	Jul. 31	Jul. 14	Sep. 07	Aug. 28
<b>Pinot Gris</b>	Dec. 08	Dec. 18	Mar. 26	Mar. 20	May 17	May 05	Jul. 30	Jul. 13	Sep. 02	Aug. 22
<b>Sauvignon Blanc</b>	Dec. 01	Dec. 10	Mar. 23	Mar. 18	May 20	May 07	Jul. 31	Jul. 13	Sep. 07	Aug. 28

potentially damaging events, such as days above 35 °C and heatwaves, while decreasing the incidence of others, such as frost days. As with phenology, geography influences the degree of change projected for these and other agroclimatic metrics. Below we detail the results of our analysis, presenting relationships between phenology and variety, AVA, and geography, and the projected changes in viticulturally-important agroclimatic metrics at the AVA scale.

### Changes in phenology by variety

When exploring changes in phenology by variety (Fig. 2; Table 4a, Supplemental Fig. 1), which accounts for the phenology model output by variety across the 12 AVAs, those varieties with the lowest chilling degree days (DD<sub>C</sub>) requirement (see Table 2) complete chill accumulation earliest while the highest chill variety considered here, Cabernet Sauvignon, completes chill roughly one month later. Under future conditions, the 12 AVA average chill completion is delayed by approximately 9–13 days across the six varieties, with Cabernet Sauvignon showing the greatest



**Table 4b** The 6-variety average DOY of chill completion and the onset of four key growing stages (budburst, flowering, veraison, and maturity) for 12 California AVAs under contemporary (1991–2020, observed) and future (2040–2069, RCP 4.5) climate conditions. The DOY is computed as the mean of means across varieties

AVA	CHILL (DOY)		BUDBURST (DOY)		FLOWERING (DOY)		VERAISON (DOY)		MATURITY (DOY)	
	1991–2020 Obs.	2040–2069 RCP4.5	1991–2020 Obs.	2040–2069 RCP4.5	1991–2020 Obs.	2040–2069 RCP4.5	1991–2020 Obs.	2040–2069 RCP4.5	1991–2020 Obs.	2040–2069 RCP4.5
<b>El Dorado</b>	Dec. 05	Dec. 10	Apr. 04	Mar. 26	May 27	May 14	Jul. 24	Jul. 10	Sep. 12	Aug. 31
<b>Livermore Valley</b>	Dec. 15	Dec. 27	Mar. 24	Mar. 17	May 19	May 05	Jul. 29	Jul. 12	Sep. 11	Aug. 30
<b>Lodi</b>	Dec. 12	Dec. 22	Mar. 21	Mar. 14	May 07	Apr. 25	Jul. 07	Jun. 23	Aug. 26	Aug. 19
<b>Madera</b>	Dec. 11	Dec. 19	Mar. 20	Mar. 15	May 03	Apr. 23	Jun. 29	Jun. 18	Aug. 23	Aug. 17
<b>Mendocino</b>	Dec. 04	Dec. 12	Apr. 03	Mar. 25	May 30	May 18	Aug. 06	Jul. 21	Sep. 15	Sep. 05
<b>Monterey</b>	Dec. 14	Dec. 26	Mar. 22	Mar. 15	May 19	May 04	Aug. 18	Jul. 25	Sep. 17	Sep. 06
<b>Napa Valley</b>	Dec. 11	Dec. 20	Mar. 26	Mar. 19	May 17	May 05	Jul. 24	Jul. 08	Sep. 05	Aug. 27
<b>Paso Robles</b>	Dec. 07	Dec. 14	Mar. 24	Mar. 22	May 17	May 08	Jul. 24	Jul. 10	Sep. 01	Aug. 25
<b>Russian River Valley</b>	Dec. 10	Dec. 20	Mar. 25	Mar. 18	May 21	May 06	Aug. 10	Jul. 18	Sep. 09	Aug. 27
<b>Santa Ynez Valley</b>	Dec. 22	Jan. 06	Mar. 17	Mar. 17	May 09	Apr. 30	Jul. 31	Jul. 14	Sep. 06	Aug. 27
<b>SLO Coast</b>	Dec. 22	Jan. 06	Mar. 20	Mar. 20	May 23	May 11	Sep. 04	Aug. 10	Sep. 27	Sep. 13
<b>West Sonoma Coast</b>	Dec. 11	Dec. 24	Mar. 30	Mar. 20	Jun. 06	May 18	Sep. 09	Aug. 13	Sep. 29	Sep. 13

delay. Conversely, the 12 AVA average timing for budburst, flowering, veraison, and maturity advances by an average of 5–7 days, 12–14 days, 17–19 days, and 10–11 days, respectively. At budburst, Cabernet Sauvignon shows the greatest advancement of the red varieties and Pinot Gris of the white varieties. From flowering to maturity the differences between varieties is minimal. All varieties show average advancements in maturity of 10 or 11 days. While the advancement of phenology increases between budburst and veraison, the break in this pattern at maturity is likely attributed to the use of BEDD for determining the timing of maturation as opposed to the accumulation of GDD used for budburst, flowering, veraison stages.

### Changes in phenology by AVA

AVA-scale phenology analysis averages the phenology model output across the six varieties for each AVA, providing a view of how general winegrape phenology may shift under climate change at the AVA scale (Table 4b). Through this lens, results show that future phenology timing shifts towards a delay in chill accumulation and an advance in budburst, flowering, veraison and maturation across all AVAs. AVA chill completion is delayed by an average of 5–15 days, with SLO Coast and Santa Ynez Valley AVAs, which have the lowest winter  $DD_c$  under contemporary conditions, showing the greatest delay. Budburst advances by a week or more in 8 of the 12 AVAs analyzed, with only Madera, Paso Robles, SLO Coast, and Santa Ynez Valley AVAs showing less than seven days of advancement at this stage. Flowering advancements are two weeks or greater in 4 of 12 AVAs (West Sonoma Coast, Livermore Valley, Monterey, Russian River Valley), and four of the AVAs (West Sonoma Coast,

Monterey, Russian River Valley, SLO Coast) show veraison advancements of three weeks or greater. Advances in maturity range from less than one week (Madera) to more than two weeks (West Sonoma Coast), with maturity occurring an average of 11 days earlier.

### Geographic influences on phenology

At the AVA-scale, latitude is an important driver of phenology in earlier development phases (i.e., chill completion and budburst), illustrated by generally earlier chill accumulation and later budburst at more northerly AVAs (Fig. 2, Table 4b, Supplemental Table 1). In later phenology phases, climate drivers other than latitude (e.g., ocean proximity, local topography) influence temperature and therefore phenology. For example, moving from flowering to maturity, we observe a subtle contrast between inland and coastal regions with AVAs situated inland, such as El Dorado, Lodi, and Madera, exhibiting a smaller shift in timing compared to coastal AVAs during these later development stages. While these patterns are somewhat muted under future climate conditions as compared to the contemporary, they show one aspect of the influence of geography on phenology. Elevation can also influence temperatures and therefore phenology. At the AVA-scale, results show a positive correlation between elevation and the timing of budburst, suggesting that locations at higher elevations tend to experience a later budburst day; however, this relationship was not significant across all phenology stages (Fig. 2, Supplemental Table 2). Finally, although we focus on the AVA-scale, we note that local topography can also influence phenology. Consider our complementary analysis of projected changes in phenology across two AVAs at comparable latitudes: the

topographically complex Napa Valley AVA and the more homogeneous Lodi AVA (Supplemental Fig. 1). There is greater spatial variation in phenology timing over Napa, where across-AVA variations in phenology shifts can be as large as 31 days, as compared to Lodi where phenology shifts are more uniform across stages and varieties, further highlighting the importance of AVA geography on wine-grape phenology.

### Changes in agroclimatic metrics

Analysis of agroclimatic metrics across the 12 AVAs of interest reveal some notable changes between current and future climates (Fig. 3, Supplemental Tables 2 and 3). Under current climate conditions, expected geographic patterns appear across a number of agroclimatic metrics. For example, inland AVAs (e.g., El Dorado, Lodi, and Madera) exhibit higher Growing Degree Days (GDD) and a greater number of Hot Days above 35 °C (HD35) when compared to their coastal counterparts. Furthermore, metrics like Chilling Degree Days ( $DD_c$ ), excess precipitation days ( $Pr_{ex}$ ), and winter accumulated precipitation ( $Pr_{acc}$ ) exhibit a north-to-south gradient, with the northernmost AVA (Mendocino) and the highest elevation region (El Dorado) experiencing the greatest amount of winter chill and excess precipitation days. We also note that higher elevations correlate with

increased exposure to days below the threshold for cold hardiness ( $H_{ini}$ ) (Supplemental Table 2).

In regions with greater relative frost risk, such as El Dorado, Mendocino, and Paso Robles, the number of frost damage days (FDD) is projected to decline by approximately 1 to 2 days between the contemporary and future periods. Additionally, we anticipate higher GDD, HD35, and Evapotranspiration ( $ET_c$ ) in warmer locations, while lower values of  $Pr_{ex}$  and  $Pr_{acc}$  are expected in these areas. Analysis shows a positive correlation between annual mean temperature and the change in HD35 ( $\Delta HD35$ ), suggesting that warmer places are anticipated to experience a higher increase in the number of hot days. Moreover, elevation exhibits a positive correlation with the change in Diurnal Temperature Range ( $\Delta DTR$ ) and  $\Delta DTR20$ , indicating that areas with higher elevations are expected to experience a more pronounced increase in Diurnal Temperature Range and a greater reduction in FDD (Supplemental Table 2). Finally, we note that as with projected phenology changes, projected changes in agroclimatic metrics can be influenced by complex topography within AVAs (Supplemental Fig. 2).



**Fig. 3** The mean value of 14 general agroclimatic metrics over 12 AVAs for the contemporary period (blue bar) and the future (2040–2069) period under RCP 4.5 (yellow bar). The 12 AVAs are arranged from north to south. The full name of each metric is listed here: Growing Degree Days (GDD), Cold Hardiness ( $H_{ini}$ ), Chilling Degree Days

( $DD_c$ ), Frost Damage Days (FDD), Last Spring Freeze (LSF), First Fall Freeze (FFF), Freeze-Free Season (FFS), Hot Days (HD), Heatwaves (HW), Diurnal Temperature Range (DTR), Diurnal Temperature Range > 20 °C (DTR20), Excess Precipitation Days ( $Pr_{ex}$ ), Winter Accumulated Precipitation ( $Pr_{acc}$ ), Crop Evapotranspiration ( $ET_c$ ).

## Discussion

This study offers an analysis that reinforces the existing body of literature on climate effects on agricultural production. Our results corroborate other studies showing that agriculture in California will face the effects of warmer winters and subsequent reduced chill accumulation, longer frost-free seasons, increased evapotranspiration, and more heat extremes (e.g., Cayan et al. 2008; Gershunov and Guirguis 2012; Luedeling et al. 2009; Pathak et al. 2018). Moreover, our results are in line with recent observations of growing season shifts in Napa Valley vineyards (Cayan et al. 2023). Climate extremes have been associated with notable damages to California agriculture (Lobell et al. 2011), and our results show increased exposure to extreme heat under future climate. Heat extremes are a known problem for winegrape cultivation, decreasing berry size and influencing berry chemistry (Greer and Weston 2010; Parker et al. 2020a). In 2021 alone heat was cited as the cause of loss for more than \$25 M in crop indemnity claims in Napa and Sonoma counties, two of California's top wine-producing counties (AgRisk Viewer; Reyes and Elias 2019). Conversely, while other California crops may have to contend with increasingly warm winter temperatures in the form of lower chill accumulation (Luedeling et al. 2009), our results show that due to the low chilling requirements of winegrapes, warmer winters will not see similarly direct negative effects on winegrape cultivation. However, warmer winters – along with reduced frost exposure and longer growing seasons – have the potential to increase pest and disease pressure (Gross, 2021; Pathak et al. 2018).

Just as with non-cultivated plants (e.g., Gordo and Sanz 2010; Polgar and Primack 2011), shifts in crop phenology under climate change have been consistently reported in literature (e.g., Pathak and Stoddard 2018; Pope et al. 2013). In winegrapes, prior studies have shown that future warming will result in earlier development, but the degree of change varies by phenology phase, location, and variety (e.g., Ausseil et al. 2021; Fraga et al. 2016; Webb et al. 2007). While our results align with these studies broadly, there are some distinctions. For example, Fraga et al. (2016) showed that across Europe Pinot Noir harvest timing showed the greatest advancement under projected mid-century conditions, while the changes in timing of flowering were more modest. In contrast, our results show flowering and veraison to have greater advancements, though this is likely due to our use of BEDD to define maturity as compared to the use of a model-simulated estimation of alcohol content by Fraga et al. (2016). Similarly, Webb et al. (2007), investigating climate influence on the phenology of Cabernet Sauvignon and Chardonnay in Australia, suggested greater advancements in maturity than our results show. Although these distinctions

highlight the role of modeling approach in understanding phenology changes, our overall findings align with these prior studies and add to the body of literature underscoring the importance of geography and variety in climate-driven phenological shifts.

Although these results capture the overarching effects of climate change on winegrape production in California, there are some limitations to our study. The thresholds used in our phenology models come from trials conducted under different environmental conditions and outside of California and as such may not be precise relative to the thresholds (general or variety specific) that may emerge were these trials replicated in our AVAs of interest. For some metrics, there is a lack of variety-specific threshold data in the literature which limits our results. For example, while it has been suggested that cooler-climate winegrape varieties benefit from larger DTR than warmer-climate varieties (Jones 2015), variety-specific optimal DTR for our varieties is lacking. For metrics like BEDD, the limitations lie in their design; BEDD is not designed to model maturation as defined by a specific level of ripening (e.g., 24 °Brix). BEDD may limit the ability to capture warming-driven shifts in the timing of generalized maturity due to the  $T_{upper}$  placing a cap on heat accumulation, suggesting that BEDD may not be ideally suited for climate change modeling applications (Hall and Jones 2010). Even using an ideal metric for modeling thermally-driven development may still produce imprecise magnitudes of phenological change (e.g. Sadras and Moran 2013; Wolkovich et al. 2012). Beyond these thresholds and metrics limitations, it is important to acknowledge the spatial limitations of the climate data relative to the microclimatic factors that can influence winegrape phenology and alter climate exposure at the vineyard scale. It is also critical to acknowledge that these results do not account for the myriad farm management actions that growers can implement to mitigate exposure to or impacts of undesirable conditions.

The adoption of climate-smart adaptation practices can improve growers' resilience to climate change and help ameliorate the negative impacts of a warming world. For California winegrape growers faced with greater water demands by way of higher crop evapotranspiration and a need to alleviate the impacts of increasing exposure to heat extremes, climate-smart adaptation practices may include improving soil structure and incorporating soil amendments to increase soil water holding capacity; improving irrigation management to increase infiltration and improve water use efficiency; adopting minimum tillage or cover cropping practices to reduce soil water evaporation; planting new drought- and heat-tolerant rootstocks and varieties; and managing the vineyard for heat exposure through canopy-management practices or the installation of heat-reducing

shade netting (Parker et al. 2023). Providing information at meaningful scales has been identified as a key component to encouraging the adoption of adaptation practices (Johnson et al. 2023). In earlier work, Babin et al. (2022) showed that the presentation of climate change projections at the local scale to vineyard managers and technical service providers (TSPs) promotes the consideration of adaptation strategies in vineyard management planning. Here we address both AVA-scale climate projections and projected phenology, providing variety-specific information at a meaningful spatial scale that can empower growers to identify and adopt the most appropriate adaptation actions for their situation.

## Conclusions

Through quantifying these projected changes in phenology and agroclimatic metrics at the variety-specific and AVA scale, we offer information at a resolution that can support grower and industry decision-making. While the methodological approach employed here can be applied to other varieties and regions within and beyond California's borders, we recommend continued field trials to not only ensure accurate variety-specific bioclimate information, but also to attempt to elucidate the complex relationships between climate, variety, and other aspects of the vine (e.g., rootstock), the vineyard system (e.g., soils), and adaptive water and nutrient management practices. Ultimately, model outputs are only as good as their inputs and better understanding of these complex relationships will be needed to improve modeling for decision support and the long-term resilience of viticulture under climate change.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s00484-024-02684-8>.

**Acknowledgements** Funding for this publication was made possible by the United States Department of Agriculture's Agricultural Marketing Service through grant AM22SCBPCA1133. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the USDA. Additional funding was received from the United States Department of Agriculture, National Institute of Food Agriculture Award # 2021-68012-35914 and # 2021-69012-35916; the United States Department of Agriculture, Office of the Chief Economist IAA # 60-2032-2-001; and the United States Department of Agriculture, Agricultural Research Service CRIS Project 2032-21220-008-000-D. The authors acknowledge the role of the California Sustainable Winegrowing Alliance for their support of this work. The authors also wish to thank two anonymous reviewers whose feedback improved the quality of the manuscript.

**Funding** United States Department of Agriculture Agricultural Marketing Service grant AM22SCBPCA1133; United States Department of Agriculture, National Institute of Food Agriculture Award # 2021-68012-35914; United States Department of Agriculture, National Institute of Food and Agriculture Award # 2021-69012-35916; United

States Department of Agriculture, Office of the Chief Economist IAA # 60-2032-2-001; United States Department of Agriculture, Agricultural Research Service CRIS Project 2032-21220-008-000-D.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- [AgRisk Viewer] USDA Southwest Climate Hub AgRisk Viewer. <https://swclimatehub.info/rma/> Accessed 17 August 2023
- [CDFA] California Department of Food and Agriculture (2022) California Grape Acreage Report, 2021 Summary. [https://www.nass.usda.gov/Statistics\\_by\\_State/California/Publications/Specialty\\_and\\_Other\\_Releases/Grapes/Acreage/2022/grpacSUMMARY-2021Crop.pdf](https://www.nass.usda.gov/Statistics_by_State/California/Publications/Specialty_and_Other_Releases/Grapes/Acreage/2022/grpacSUMMARY-2021Crop.pdf) [Accessed 4 April 2023].
- [CDFA] California Department of Food and Agriculture (2023) California Grape Crush 2022. [https://www.nass.usda.gov/Statistics\\_by\\_State/California/Publications/Specialty\\_and\\_Other\\_Releases/Grapes/Crush/Final/2022/Grape\\_Crush\\_2022\\_Final.pdf](https://www.nass.usda.gov/Statistics_by_State/California/Publications/Specialty_and_Other_Releases/Grapes/Crush/Final/2022/Grape_Crush_2022_Final.pdf) [Accessed 4 April 2023]
- [CWI] California Wine Institute (2022) The economic impact of California wines. <https://wineinstitute.org/wp-content/uploads/2022/11/ca-wine-economic-impact-highlights-dec2022.pdf> [Accessed 4 April 2023]
- Abatzoglou JT (2013) Development of gridded surface meteorological data for ecological applications and modelling. *Int J Climatol* 33:121–131
- Abatzoglou JT, Brown TJ (2012) A comparison of statistical downscaling methods suited for wildfire applications. *Int J Climatol* 32:772–780
- Allen RG, Pereira LS, Raes D, Smith M (1998) Crop evapotranspiration-guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. *Fao Rome* 300(9):D05109
- Anderson K, Norman D, Wittwer G (2003) Globalisation of the world's wine markets. *World Econ* 26(5):659–687
- Auseil AGE, Law RM, Parker AK, Teixeira EI, Sood A (2021) Projected wine grape cultivar shifts due to climate change in New Zealand. *Frontiers in Plant Science*, 12, p.618039
- Babin N, Guerrero J, Rivera D, Singh A (2022) Vineyard-specific climate projections help growers manage risk and plan adaptation in the Paso Robles AVA. *Calif Agric* 75(3):142–150
- Bouby L, Figueiral I, Bouchette A, Rovira N, Ivorra S, Lacombe T, Pastor T, Picq S, Marinval P, Terral JF (2013) Bioarchaeological insights into the process of domestication of grapevine (*Vitis vinifera* L.) during roman times in Southern France. *PLoS ONE* 8(5):e63195
- Cayan DR, Maurer EP, Dettinger MD, Tyree M, Hayhoe K (2008) Climate change scenarios for the California region. *Clim Change* 87:21–42
- Cayan DR, DeHaan L, Tyree M, Nicholas KA (2023) A 4-week advance in the growing season in Napa Valley, California, USA. *Int J Climatol*

- Cohen SD, Tarara JM, Gambetta GA, Matthews MA, Kennedy JA (2012a) Impact of diurnal temperature variation on grape berry development, proanthocyanidin accumulation, and the expression of flavonoid pathway genes. *J Exp Bot* 63(7):2655–2665
- Cohen SD, Tarara JM, Kennedy JA (2012b) Diurnal temperature range compression hastens berry development and modifies flavonoid partitioning in grapes. *Am J Enol Viticult* 63(1):112–120
- Cook BI, Mankin JS, Marvel K, Williams AP, Smerdon JE, Anchukaitis KJ (2020) Twenty-first century drought projections in the CMIP6 forcing scenarios. *Earth's Future*, 8(6), e2019EF001461. <https://doi.org/10.1029/2019EF001461>
- Daly C, Taylor GH, Gibson WP (1997) The PRISM approach to mapping precipitation and temperature. In *Proc., 10th AMS Conf. on Applied Climatology* (pp. 20–23)
- Daly C, Halbleib M, Smith JI, Gibson WP, Doggett MK, Taylor GH, Curtis J, Pasteris PP (2008) Physiographically sensitive mapping of climatological temperature and precipitation across the conterminous United States. *Int J Climatol: J Royal Meteorological Soc* 28(15):2031–2064
- Diffenbaugh NS, White MA, Jones GV, Ashfaq M (2011) Climate adaptation wedges: a case study of premium wine in the western United States. *Environ Res Lett* 6(2):p024024
- Ferguson JC, Moyer MM, Mills LJ, Hoogenboom G, Keller M (2014) Modeling dormant bud cold hardiness and budbreak in twenty-three *Vitis* genotypes reveals variation by region of origin. *Am J Enol Viticult* 65(1):59–71
- Fraga H, García de Cortázar Atauri I, Malheiro AC, Santos JA (2016) Modelling climate change impacts on viticultural yield, phenology and stress conditions in Europe. *Glob Change Biol* 22(11):3774–3788
- Gershunov A, Guirguis K (2012) California heat waves in the present and future. *Geophys Res Lett*, 39(18)
- Gershunov A, Cayan DR, Iacobellis SF (2009) The great 2006 heat wave over California and Nevada: Signal of an increasing trend. *J Clim* 22(23):6181–6203
- Gladstones J (1992) *Viticulture and environment* Winetitles
- Gordo O, Sanz JJ (2010) Impact of climate change on plant phenology in Mediterranean ecosystems. *Glob Change Biol* 16(3):1082–1106
- Greer DH, Weston C (2010) Heat stress affects flowering, berry growth, sugar accumulation and photosynthesis of *Vitis vinifera* Cv. Semillon grapevines grown in a controlled environment. *Funct Plant Biol* 37(3):206–214
- Gross L (February 2021) Warmer California Winters May Fuel Grapevine-Killing Pierce's Disease, vol 17. Inside Climate News
- Hall A, Jones GV (2010) Spatial analysis of climate in winegrape-growing regions in Australia. *Aust J Grape Wine Res* 16(3):389–404
- Hannah L, Roehrdanz PR, Ikegami M, Shepard AV, Shaw MR, Tabor G, Zhi L, Marquet PA, Hijmans RJ (2013) Climate change, wine, and conservation. *Proc Natl Acad Sci* 110(17):6907–6912 <https://insideclimatenews.org/news/17022021/warmer-california-winters-may-fuel-grapevine-killing-pierces-disease/> [Accessed 30 August 2023]
- Huglin P (1978) Nouveau Mode d'Évaluation Des Possibilités Héliothermiques D'un Milieu Viticole. *C R Acad Agr France*, p. 1117–1126
- Johnson D, Parker LE, Pathak TB, Crothers L, Ostojica SM (2023) Technical assistance providers identify Climate Change Adaptation practices and barriers to Adoption among California Agricultural Producers. *Sustainability* 15(7):5973
- Jolly WM, Nemani R, Running SW (2005) A generalized, bioclimatic index to predict foliar phenology in response to climate. *Glob Change Biol* 11(4):619–632
- Jones GV (2003) Winegrape phenology. *Phenology: an integrative environmental science*, 523–539
- Jones G (2015) Climate, Grapes, and Wine. *Terroir and the importance of climate on grapevine production*
- Jones GV, Webb LB (2010) Climate change, viticulture, and wine: challenges and opportunities. *J Wine Res* 21(2–3):103–106
- Jones GV, Duff AA, Hall A, Myers JW (2010) Spatial analysis of climate in winegrape growing regions in the western United States. *Am J Enol Viticult* 61(3):313–326
- Jones GV, Reid R, Vilks A (2012) Climate, grapes, and wine: structure and suitability in a variable and changing climate. The geography of wine: regions, Terroir and techniques. Springer, Amsterdam, The Netherlands, pp 109–133
- Kharin VV, Zwiers FW, Zhang X, Wehner M (2013) Changes in temperature and precipitation extremes in the CMIP5 ensemble. *Clim Change* 119:345–357
- Krantz W, Pierce D, Goldenson N, Cayan D (2021) Memorandum on evaluating global climate models for studying regional climate change in California. [https://www.energy.ca.gov/sites/default/files/2022-09/20220907\\_CDAWG\\_MemoEvaluating\\_GCMs\\_EPC-20-006\\_Nov2021-ADA.pdf](https://www.energy.ca.gov/sites/default/files/2022-09/20220907_CDAWG_MemoEvaluating_GCMs_EPC-20-006_Nov2021-ADA.pdf)
- Li H, Sheffield J, Wood EF (2010) Bias correction of monthly precipitation and temperature fields from Intergovernmental Panel on Climate Change AR4 models using equidistant quantile matching. *J Geophys Res: Atmos* 115(D10)
- Lobell DB, Torney A, Field CB (2011) Climate extremes in California agriculture. *Clim Change* 109:355–363
- Luedeling E, Zhang M, Girvetz EH (2009) Climatic changes lead to declining winter chill for fruit and nut trees in California during 1950–2099. *PLoS ONE* 4(7):e6166
- Martínez-Lüscher J, Chen CCL, Brillante L, Kurtural SK (2020) Mitigating heat Wave and exposure damage to Cabernet Sauvignon Wine grape with partial shading under two irrigation amounts. *Front Plant Sci* 11:1760
- Monteverde C, De Sales F (2020) Impacts of global warming on southern California's winegrape climate suitability. *Adv Clim Change Res* 11(3):279–293
- Morales-Castilla I, García de Cortázar-Atauri I, Cook BI, Lacombe T, Parker A, Van Leeuwen C, Nicholas KA, Wolkovich EM (2020) Diversity buffers winegrowing regions from climate change losses. *Proc Natl Acad Sci* 117(6):2864–2869
- Mosedale JR, Wilson RJ, Maclean IM (2015) Climate change and crop exposure to adverse weather: changes to frost risk and grapevine flowering conditions. *PLoS ONE*, 10(10), e0141218
- Parker LE, Abatzoglou JT (2018) Shifts in the thermal niche of almond under climate change. *Clim Change* 147(1–2):211–224
- Parker LE, McElrone AJ, Ostojica SM, Forrester EJ (2020a) Extreme heat effects on perennial crops and strategies for sustaining future production. *Plant Science*, 295, p.110397
- Parker AK, de Cortázar-Atauri IG, Trought MC, Destrac A, Agnew R, Sturman A, Van Leeuwen C (2020b) Adaptation to climate change by determining grapevine cultivar differences using temperature-based phenology models. *Oeno One* 54(4):955–974
- Parker LE, Johnson D, Pathak TB, Wolff M, Jameson V, Ostojica SM (2023) Adaptation Resources Workbook for California Specialty Crops. USDA California Climate Hub Technical Report CACH-2023-1. Davis, CA: U.S. Department of Agriculture, Climate Hubs. 55 p
- Pathak TB, Stoddard CS (2018) Climate change effects on the processing tomato growing season in California using growing degree day model. *Model Earth Syst Environ* 4:765–775
- Pathak TB, Maskey ML, Dahlberg JA, Kearns F, Bali KM, Zaccaria D (2018) Climate change trends and impacts on California agriculture: A detailed review. *Agronomy*, 8(3), p.25
- Polgar CA, Primack RB (2011) Leaf-out phenology of temperate woody plants: from trees to ecosystems. *New Phytol* 191(4):926–941
- Pope KS, Dose V, Da Silva D, Brown PH, Leslie CA, DeJong TM (2013) Detecting nonlinear response of spring phenology to climate change by bayesian analysis. *Glob Change Biol* 19(5):1518–1525

- Reyes J.J., Elias E (2019) Spatio-temporal variation of crop loss in the United States from 2001 to 2016. *Environ Res Lett* 14(7):p074017
- Sadras VO, Moran MA (2013) Nonlinear effects of elevated temperature on grapevine phenology. *Agric for Meteorol* 173:107–115
- Sheridan SC, Lee C (2018) Temporal trends in Absolute and relative Extreme temperature events across North America. *J Geophys Res Atmos* 123:11–889
- Walter IA, Allen RG, Elliott R, Jensen ME, Itenfisu D, Mechem B, Howell TA, Snyder R, Brown P, Echings S, Spofford T (2000) ASCE's standardized reference evapotranspiration equation. In *Watershed management and operations management 2000* (pp. 1–11)
- Webb LB, Whetton PH, Barlow EWR (2007) Modelled impact of future climate change on the phenology of winegrapes in Australia. *Aust J Grape Wine Res* 13(3):165–175
- White MA, Diffenbaugh NS, Jones GV, Pal JS, Giorgi F (2006) Extreme heat reduces and shifts United States premium wine production in the 21st century. *Proceedings of the National Academy of Sciences*, 103(30), 11217–11222
- Wolkovich EM, Cook BI, Allen JM, Crimmins TM, Betancourt JL, Travers SE, Pau S, Regetz J, Davies TJ, Kraft NJ, Ault TR (2012) Warming experiments underpredict plant phenological responses to climate change. *Nature* 485(7399):494–497
- Wolkovich EM, Burge DO, Walker MA, Nicholas KA (2017) Phenological diversity provides opportunities for climate change adaptation in winegrapes. *J Ecol* 105(4):905–912
- Xia Y, Mitchell K, Ek M, Sheffield J, Cosgrove B, Wood E, Luo L, Alonge C, Wei H, Meng J, Livneh B (2012) Continental-scale water and energy flux analysis and validation for the North American Land Data Assimilation System project phase 2 (NLDAS-2): 1. Intercomparison and application of model products. *J Geophys Research: Atmos* 117:D3
- Zapata D, Salazar-Gutierrez M, Chaves B, Keller M, Hoogenboom G (2017) Predicting key phenological stages for 17 grapevine cultivars (*Vitis vinifera* L). *Am J Enol Viticult* 68(1):60–72
- Zhang J, Guan K, Peng B, Jiang C, Zhou W, Yang Y, Pan M, Franz TE, Heeren DM, Rudnick DR, Abimbola O (2021) Challenges and opportunities in precision irrigation decision-support systems for center pivots. *Environ Res Lett* 16(5):053003

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.