UC Berkeley

UC Berkeley Previously Published Works

Title

Accounting for sub-pixel variability of clouds and/or unresolved spectral variability, as needed, with generalized radiative transfer theory

Permalink

<https://escholarship.org/uc/item/3cr846t1>

ISBN

9781557528148

Authors

Davis, AB

Xu, F

Collins, WD

Publication Date

2015

DOI

10.1364/hise.2015.ht4b.3

Peer reviewed

Accounting for Sub-Pixel Variability of Clouds and/or Unresolved Spectral Variability, as Needed, with Generalized Radiative Transfer Theory

Anthony B. Davis,^{*} Feng Xu^{*} and William D. Collins[†]

^{*}Jet Propulsion Laboratory / California Institute of Technology, 4800 Oak Grove Drive (Mail Stop 233-200), Pasadena, CA 91109, USA

[†]Earth Sciences Division, Lawrence Berkeley National Laboratory & Dept. of Earth and Planetary Science, University of California - Berkeley

Anthony.B.Davis@jpl.nasa.gov; Feng.Xu@jpl.nasa.gov; wcollins@berkeley.edu

Abstract: Atmospheric hyperspectral VNIR sensing struggles with sub-pixel variability of clouds and/or limited spectral resolution mixing molecular lines. Our generalized radiative transfer model addresses both issues with new propagation kernels characterized by power-law decay in space.

OCIS codes: (290.1090) Aerosol and cloud effects; (290.4210) Multiple scattering; (280.4991) Passive remote sensing.

1. Motivation & Approach

A common tradeoff in hyperspectral sensor design is spatial-for-spectral resolution, to the point where it may no longer be an imager (cf. Fig. 1a). Conversely, in multispectral VNIR/SWIR imager design, SNR and/or sensitivity considerations argue for the inclusion of some or many molecular spectral absorption lines to be mixed in with the scattered and reflected signals and preserve the potentially fine pixel size and broad enough swath (cf. Fig. 1b). Either way, the forward and inverse radiative transfer (RT) modeling used in physics-based retrieval algorithms is challenged. Serendipitous developments in spectral [1] and spatial [2] aspects of RT theory have been unified to address either or both of these issues. This involves however a deep modification of the foundational RT equation (cf. Fig. 1c). In short, averaging Beer's transmission law $T(\ell) = \exp(-\sigma\ell)$ for a fixed geometrical path ℓ over the unresolved variability of the extinction σ , parameterized by a Gamma distribution [3], leads to $T_a(\ell) = (1 + \sigma_m \ell/a)^{-a}$, where σ_m is now the mean extinction and $a = \sigma_m^2/\text{Variance}[\sigma]$. The limit $a \rightarrow \infty$ reverts to the usual exponential law.

Fig. 1. (a) Possible sub-pixel clouds. (b) Spectral line mixing, e.g., O₂ A-band. (c) 2D Monte Carlo RT with old and new propagation kernels.

2. Standard & Generalized Radiative Transfer Equations (RTEs), and Their Numerical Solution

Let $I(\tau, \Omega)$ be mean radiance in direction Ω at mean optical depth τ in a plane-parallel medium of total optical depth τ^* (with $\sigma_m = 1$). The RTE in integral form is $I(\tau, \Omega) = \int_{4\pi} \int_{(0, \tau^*)} K(\tau, \Omega; \tau', \Omega') I(\tau', \Omega') d\tau' d\Omega' + I_0(\tau, \Omega)$, where $I_0(\tau, \Omega)$ is uncollided radiance and, for simplicity, the surface below is assumed black. Taking single scattering albedo ω and phase function $p(\Omega' \cdot \Omega)$ as constant, standard ($a = \infty$) and generalized ($0 < a < \infty$) RT models have kernels and sources

$$K_a(\tau, \Omega; \tau', \Omega') = \omega p(\Omega' \cdot \Omega) \Theta((\tau - \tau')/\mu) dT_a/d\tau((\tau - \tau')/\mu) / |\mu| \quad \text{and} \quad I_0(\tau, \Omega) = T_a(\tau/\mu_0) \delta(\Omega - \Omega_0) \quad (1)$$

where $\Theta(\cdot)$ and $\delta(\cdot)$ are respectively the Heaviside and Dirac functions, $\mu = \Omega_z$, and Ω_0 is incoming solar direction. We show that (1) is efficiently solved numerically with Markov chain methods [2,4], and we illustrate some important characteristics of generalized RT drawing on practical examples in spatial- and/or spectral-domains.

3. References

- [1] A.J. Conley & W.D. Collins, "Extension of the weak-line approximation and application to correlated- k methods," JQSRT **112**, 1525- (2011).
- [2] A.B. Davis & F. Xu, "A generalized linear transport model for spatially-correlated stochastic media," J. Comput. & Theor. Transp. (in press).
- [3] H.W. Barker, B.A. Wielicki, and L. Parker, "A parameterization for computing grid-averaged solar fluxes for inhomogeneous marine boundary layer clouds - Part 2, Validation using satellite data," JAS **53**, 2304- (1996).
- [4] F. Xu, A.B. Davis, R.A. West, and L.W. Esposito, "Markov chain formalism for polarized light transfer in plane-parallel atmospheres, with numerical comparison to the Monte Carlo method," Opt. Exp. **19**, 946- (2011).