

UC Irvine

UC Irvine Previously Published Works

Title

Enhanced pre-synaptic glutamate release in deep-dorsal horn contributes to calcium channel alpha-2-delta-1 protein-mediated spinal sensitization and behavioral hypersensitivity

Permalink

<https://escholarship.org/uc/item/3ct3b394>

Journal

Molecular Pain, 5(1)

ISSN

1744-8069

Authors

Nguyen, David
Deng, Ping
Matthews, Elizabeth A
et al.

Publication Date

2009

DOI

10.1186/1744-8069-5-6

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Research

Open Access

Enhanced pre-synaptic glutamate release in deep-dorsal horn contributes to calcium channel alpha-2-delta-1 protein-mediated spinal sensitization and behavioral hypersensitivity

David Nguyen¹, Ping Deng², Elizabeth A Matthews³, Doo-Sik Kim¹, Guoping Feng⁴, Anthony H Dickenson³, Zao C Xu² and Z David Luo^{*1,5}

Address: ¹Department of Anesthesiology & Perioperative Care, School of Medicine, University of California Irvine, Irvine, CA 92697, USA, ²Department of Anatomy & Cell Biology, School of Medicine, Indiana University, Indianapolis, IN 46202, USA, ³Department of Pharmacology, University College London, London, WC1E 6BT, UK, ⁴Department of Neurobiology, Duke University Medical Center, Durham, NC 27710, USA and ⁵Department of Pharmacology, School of Medicine, University of California Irvine, Irvine, CA 92697, USA

Email: David Nguyen - davnguyn@yahoo.com; Ping Deng - pideng@iupui.edu; Elizabeth A Matthews - elizabeth.matthews@ucl.ac.uk; Doo-Sik Kim - kds0728md@gmail.com; Guoping Feng - feng@neuro.duke.edu; Anthony H Dickenson - anthony.dickenson@ucl.ac.uk; Zao C Xu - zxu@anatomy.iupui.edu; Z David Luo* - zluo@uci.edu

* Corresponding author

Published: 12 February 2009

Received: 5 December 2008

Molecular Pain 2009, **5**:6 doi:10.1186/1744-8069-5-6

Accepted: 12 February 2009

This article is available from: <http://www.molecularpain.com/content/5/1/6>

© 2009 Nguyen et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Nerve injury-induced expression of the spinal calcium channel alpha-2-delta-1 subunit ($Ca_v\alpha_2\delta_1$) has been shown to mediate behavioral hypersensitivity through a yet identified mechanism. We examined if this neuroplasticity modulates behavioral hypersensitivity by regulating spinal glutamatergic neurotransmission in injury-free transgenic mice overexpressing the $Ca_v\alpha_2\delta_1$ proteins in neuronal tissues. The transgenic mice exhibited hypersensitivity to mechanical stimulation (allodynia) similar to the spinal nerve ligation injury model. Intrathecally delivered antagonists for *N*-methyl-D-aspartate (NMDA) and α -amino-3-hydroxyl-5-methylisoxazole-4-propionic acid (AMPA)/kainate receptors, but not for the metabotropic glutamate receptors, caused a dose-dependent allodynia reversal in the transgenic mice without changing the behavioral sensitivity in wild-type mice. This suggests that elevated spinal $Ca_v\alpha_2\delta_1$ mediates allodynia through a pathway involving activation of selective glutamate receptors. To determine if this is mediated by enhanced spinal neuronal excitability or pre-synaptic glutamate release in deep-dorsal horn, we examined wide-dynamic-range (WDR) neuron excitability with extracellular recording and glutamate-mediated excitatory postsynaptic currents with whole-cell patch recording in deep-dorsal horn of the $Ca_v\alpha_2\delta_1$ transgenic mice. Our data indicated that overexpression of $Ca_v\alpha_2\delta_1$ in neuronal tissues led to increased frequency, but not amplitude, of miniature excitatory post synaptic currents mediated mainly by AMPA/kainate receptors at physiological membrane potentials, and also by NMDA receptors upon depolarization, without changing the excitability of WDR neurons to high intensity stimulation. Together, these findings support a mechanism of $Ca_v\alpha_2\delta_1$ -mediated spinal sensitization in which elevated $Ca_v\alpha_2\delta_1$ causes increased pre-synaptic glutamate release that leads to reduced excitation thresholds of post-synaptic dorsal horn neurons to innocuous stimuli. This spinal sensitization mechanism may mediate at least partially the neuropathic pain states derived from increased pre-synaptic $Ca_v\alpha_2\delta_1$ expression.

Background

Chronic pain derived from nerve injuries, or neuropathic pain, is a common clinical problem affecting over 50 million people in the United States [1]. Neuropathic pain may manifest as spontaneous pain, pain resulting from stimuli that are normally innocuous (allodynia), and increased pain responses to suprathreshold stimuli (hyperalgesia), all of which affect adversely the quality of patients' daily-life. Unfortunately, neuropathic pain tends to be long-term, and difficult to manage due to the poor efficacies and severe side effects associated with current conventional analgesic treatments. It appears that neuropathic pain with various etiologies can lead to similar behavioral endpoints that respond differentially to pharmacological agents, suggesting that different mechanisms may underlie specific neuropathic pain states deriving from different pathological conditions. Thus, the development of target specific, safe and effective agents for neuropathic pain management relies on new insights into mechanisms of different neuropathic pain states.

Findings from previous studies have indicated that peripheral nerve injury induces a dramatic upregulation of the voltage-gated calcium channel (VGCC) $\alpha_2\delta_1$ subunit ($\text{Ca}_v\alpha_2\delta_1$) in sensory neurons and spinal dorsal horn [2-6], which correlates with neuropathic pain development [2,3,7]. Knocking down of injury-induced $\text{Ca}_v\alpha_2\delta_1$ upregulation in dorsal spinal cord with intrathecal antisense oligodeoxynucleotides results in allodynia reversal in spinal nerve injured rats [7]. In addition, transgenic mice overexpressing $\text{Ca}_v\alpha_2\delta_1$ in neuronal tissue display tactile allodynia similar to nerve injury models that can be blocked by intrathecal gabapentin [8], a drug that binds to the $\text{Ca}_v\alpha_2\delta_1$ proteins with high affinity [9] and has anti-neuropathic pain properties in patients and experimental animal models [2,3,10-16]. Furthermore, blocking axonal transport of injury-induced $\text{Ca}_v\alpha_2\delta_1$ from DRG neurons to their pre-synaptic terminals in spinal dorsal horn diminishes tactile allodynia in nerve-injured animals [7]. Together, these findings suggest that elevated $\text{Ca}_v\alpha_2\delta_1$ likely contributes to behavioral hypersensitivity through a pre-synaptic mechanism at the spinal cord level. This hypothesis, however, has not been tested directly.

Since elevated $\text{Ca}_v\alpha_2\delta_1$ in the transgenic mice has been shown to cause dorsal horn neuron hyperexcitability, behavioral hypersensitivity [8]; enhanced glutamate neurotransmission is an important mediator in neuropathic pain processing [17,18]; and elevated $\text{Ca}_v\alpha_2\delta_1$ may modulate various calcium channels that are known to regulate spinal pre-synaptic neurotransmission [19-25], we examined if elevated $\text{Ca}_v\alpha_2\delta_1$ mediated deep-dorsal horn neuron excitability and behavioral hypersensitivity through activation of glutamate receptors, and if so, whether it was mediated by a pre-synaptic or post-synaptic mechanism.

We tested these hypotheses in the injury-free $\text{Ca}_v\alpha_2\delta_1$ -overexpressing transgenic mouse model showing dorsal horn neuron hyperexcitability and behavioral hypersensitivity without the influence from other nerve injury factors [8].

Results

A similar tactile allodynia state is developed in spinal nerve injured mice and injury-free $\text{Ca}_v\alpha_2\delta_1$ transgenic mice

Our previous studies have indicated that unilateral spinal nerve ligation (SNL) induces upregulation of the $\text{Ca}_v\alpha_2\delta_1$ subunit in DRG and spinal dorsal horn that plays a critical role in tactile allodynia development [2,7]. However, nerve injury may also induce altered expression of other factors and activation of other pathways independent of the $\text{Ca}_v\alpha_2\delta_1$ induction, which may also contribute to tactile allodynia development. To determine if injury-induced $\text{Ca}_v\alpha_2\delta_1$ alone is sufficient to induce tactile allodynia, we examined the behavioral response of transgenic (TG) mice overexpressing the $\text{Ca}_v\alpha_2\delta_1$ proteins in neuronal tissues [8]. Similar to the SNL mice (Fig. 1 left), the injury-free $\text{Ca}_v\alpha_2\delta_1$ TG mice also developed a similar allodynic state compared with their wild-type (WT) littermates (Fig. 1, right). Since other factors in these two models,




Figure 1
Increased $\text{Ca}_v\alpha_2\delta_1$ expression in spinal cord and DRG correlates with tactile allodynia in spinal nerve ligated and injury-free $\text{Ca}_v\alpha_2\delta_1$ TG mice. Unilateral L5 spinal nerve ligation (SNL) surgeries were performed in mice, and paw withdrawal thresholds (PWT) to von Frey filament stimulation were measured two-weeks post injury in SNL mice or in injury-free, adult male $\text{Ca}_v\alpha_2\delta_1$ TG and WT mice as described in the Methods. Data presented are the means \pm SEM from $n = 6$ for the SNL group, and $n = 12$ each for the injury-free WT and TG groups. C – contralateral (non-injury) side. Ip – ipsilateral (injury) side. * $p < 0.05$ compared to contralateral side or WT control using unpaired Students' *t* test.

e.g. other injury-induced factors in the SNL model and compensatory factors in the $Ca_v\alpha_2\delta_1$ TG mice, are not likely the same, these data support that elevated $Ca_v\alpha_2\delta_1$ is highly likely a common factor mediating the abnormal sensations seen in these two models. This is supported by the findings that the tactile allodynia in both models can be blocked by treatment with intrathecal gabapentin [2,8,10], which binds to the $Ca_v\alpha_2\delta_1$ subunit [9].

A spinal glutamatergic pathway mediates behavioral hypersensitivity in the $Ca_v\alpha_2\delta_1$ -transgenic mice

The pathway underlying $Ca_v\alpha_2\delta_1$ -mediated behavioral hypersensitivity is not known. Since activation of glutamate receptors plays a role in central sensitization in pain processing [18,26-29] and activation of selective glutamate receptors is believed to play a critical role in SNL-induced tactile allodynia [17,18], we tested the hypothesis that glutamate receptor activation might underlie the $Ca_v\alpha_2\delta_1$ -mediated behavioral hypersensitivity in the TG mice by examining the effects of intrathecal glutamate receptor antagonists on allodynia reversal in the TG mice. Intrathecal injection of memantine, a selective *N*-methyl-D-aspartate (NMDA) receptor antagonist, caused a dose-dependent allodynia reversal in the $Ca_v\alpha_2\delta_1$ TG mice (Fig. 2A). Its anti-allodynic effects at the highest dose tested (2 mg/kg) lasted for two-three hrs (Fig. 2B), and diminished completely overnight (data not shown). Similar drug treatment in the WT littermates led to a slight increase of behavioral thresholds to the same type of mechanical stimulation, which was not statistically significant compared with the pretreatment level (Fig. 2). The allodynia reversal by memantine was similar to that reported in the SNL model [17].

Intrathecal injection of 6,7-dinitroquinoxaline-2,3-dione (DNQX), a non-NMDA receptor antagonist for the α -amino-3-hydroxyl-5-methylisoxazole-4-propionic acid (AMPA)/kainate receptors, at a dose of 50 μ g/kg provided little allodynia reversal in the TG mice. However, intrathecal DNQX at a higher dose (0.15 mg/kg) provided a rapid onset and transient allodynia reversal in the TG mice (Fig. 3), similar to that reported in the SNL model [17]. Similar DNQX treatment in the WT littermates led to a slight increase of behavioral thresholds to the same type of mechanical stimulation, which was not statistically significant compared with the pretreatment level (Fig. 3).

Intrathecal injection of 2-amino-3-phosphonopropionic acid (AP3), a non-competitive metabotropic glutamate receptor antagonist, at a maximum tolerable dose reported (50 μ g/kg) [17], did not significantly reverse tactile allodynia in the TG mice (Fig. 4), which was in agreement with findings from the SNL model [17].

$Ca_v\alpha_2\delta_1$ -mediated spinal hyperexcitability through enhanced pre-synaptic glutamate release, but not likely through enhanced intrinsic excitability of dorsal horn neurons

We reported previously that elevated $Ca_v\alpha_2\delta_1$ in the TG mice, without other injury induced factors, was sufficient to cause WDR neuron hyperexcitability and behavioral hypersensitivity [8]. Data from our previous studies have indicated that intrathecal injection of gabapentin, a drug that binds to $Ca_v\alpha_2\delta_1$ proteins and has anti-neuropathic pain efficacy in the SNL model [2,3,10-12] and patients [13-15], can reverse the behavioral hypersensitivity in the $Ca_v\alpha_2\delta_1$ TG mice [8], supporting that $Ca_v\alpha_2\delta_1$ -mediated WDR neuron activation contributes at least partially to the behavioral hypersensitivity. Since the wind-up response, a phenomenon of progressively exaggerated WDR neuron excitability to repetitive C-fiber stimulation, is similar between the TG and WT littermates [8], the hyperexcitability of the TG dorsal horn neurons is not likely mediated by a facilitated state mediated by the wind-up mechanism.

To test if $Ca_v\alpha_2\delta_1$ overexpression affects post-synaptic dorsal horn neuron excitability to acute stimuli, we compared C-fiber responses, post discharges, and non-potentiated responses (input) of WDR neurons from TG and WT littermates after brief transcutaneous electric stimulation with the C-fiber intensity. As shown in Fig. 5, similar levels of end point measures were observed between the TG and WT neurons. Even though our data do not allow the complete exclusion about possible influences from changes in strength of C-fiber synaptic inputs onto TG neurons, the occurrence of these changes, if any, would require opposite compensations in the excitability of TG neurons to maintain a level of excitability similar to that observed in WT mice, which are not supported by reported findings [8]. Thus, these findings suggest that the intrinsic excitability of WDR neurons, at least that to acute C-fiber inputs, is not changed in the $Ca_v\alpha_2\delta_1$ TG mice.

This is further supported by our findings that elevated AMPA/kainate and NMDA receptor-mediated miniature excitatory postsynaptic currents (mEPSCs) in TG deep-dorsal horn neurons are at least partially mediated by enhanced pre-synaptic glutamate release, but not by changes in post-synaptic components, including enhanced intrinsic excitability of post-synaptic deep dorsal horn neurons. As shown in Fig. 6A and 6B, at a holding potential of -70 mV, the frequency of AMPA/kainate receptor mediated mEPSCs was significantly increased from 0.22 ± 0.06 Hz for WT neurons ($n = 7$) to 0.79 ± 0.08 Hz for the TG neurons ($n = 9$). Application of AMPA/kainate receptor antagonist CNQX (10 μ M) could reduce mEPSCs by $91.3 \pm 5.3\%$ and $93.9 \pm 3.3\%$ in WT and TG neurons ($n = 5$ in each group, data not shown), respectively. In the presence of CNQX (10 μ M), the frequency of





Figure 2 (see legend on next page)

Figure 2 (see previous page)

Intrathecal NMDA receptor antagonists reverse tactile allodynia in the TG mice. Memantine, an NMDA receptor antagonist, was injected intrathecally between L5/6 lumbar regions into adult TG and WT mice, and behavioral sensitivities to mechanical stimulation before or after drug treatments were assessed blindly as paw withdrawal thresholds (PWT) to von Frey filament stimulation in the hindpaw. **A.** Dose-dependent effects of intrathecal memantine for PWT in the TG and WT mice measured 120 min post i.t. injection. Data shown represent the means \pm SEM from $n = 7$ for each group. ### $p < 0.01$, #### $p < 0.001$ compared with WT mice, *** $p < 0.001$ compared with the pretreatment level using two-way ANOVA followed by Bonferroni post-tests. **B.** Time-dependent effects of intrathecal memantine (2.0 mg/kg) on allodynia reversal in TG mice compared with saline treatment and similar treatments in WT mice. Data shown represent the means \pm SEM from $n = 7$ for each group. * $p < 0.05$, ** $p < 0.01$ compared with the pretreatment level, \$ $p < 0.05$, \$\$\$ $p < 0.001$ compared with saline treatment in TG mice using two-way ANOVA followed by Bonferroni post-tests.

NMDA receptor-mediated mEPSCs, recorded at a holding potential of +50 mV, was also increased in the TG neurons (WT: 0.03 ± 0.02 Hz, $n = 5$; TG: 0.12 ± 0.01 , $n = 3$). However, the amplitudes of AMPA/kainate and NMDA receptor mediated mEPSCs were about the same between the WT and TG neurons. In addition, the kinetics of the AMPA/kainate and NMDA receptor mediated mEPSCs showed no obvious difference between the WT and TG neurons (Fig. 6C and 6D). The rising time of AMPA/kainate receptor mediated mEPSCs was 1.19 ± 0.10 ms ($n = 7$)

and 1.26 ± 0.09 ms ($n = 9$) in WT and TG neurons, respectively, and the decay time constant was 5.05 ± 0.43 ms ($n = 7$) and 5.12 ± 0.39 ms ($n = 9$) in WT and TG neurons, respectively. The rising time of NMDA receptor mediated mEPSCs was 1.82 ± 0.11 ms ($n = 3$) in WT and 1.70 ± 0.05 ms ($n = 3$) in TG neurons, and the decay time constant was 15.13 ± 0.34 ms ($n = 3$) in WT and 14.70 ± 1.29 ms ($n = 3$) in TG neurons.

**Figure 3**

Intrathecal AMPA/kainate receptor antagonists reverse tactile allodynia in the TG mice. Different doses of DNQX, an antagonist of AMPA/kainate receptors, were injected intrathecally between L5/6 lumbar regions into adult TG and WT mice, and behavioral sensitivities to mechanical stimulation before or after drug treatments were assessed blindly as paw withdrawal thresholds (PWT) to von Frey filament stimulation in the hindpaw. Data shown are the dose- and time-dependent effects of intrathecal DNQX on allodynia reversal in TG mice compared with saline treatment or similar treatment in the WT mice. For the purpose of clarity, effects of the lower dose of DNQX (50 μ g/kg) on PWT of WT mice, which were similar to that in saline treated WT mice, were not shown. Means \pm SEM from $n = 7$ for each group. * $p < 0.01$, *** $p < 0.001$ compared with the pretreatment level, \$ $p < 0.05$, \$\$\$ $p < 0.001$ compared with saline treatment in TG mice using two-way ANOVA followed by Bonferroni post-tests.




Figure 4

Intrathecal antagonism of metabotropic glutamate receptors does not cause allodynia reversal in the TG

mice. AP3, a mGluR antagonist, was injected intrathecally between L5/6 lumbar regions into adult TG and WT mice, and behavioral sensitivities to mechanical stimulation before or after drug treatments were assessed blindly as paw withdrawal thresholds (PWT) to von Frey filament stimulation in the hindpaw. Data shown are time-dependent effects of intrathecal AP3, at the maximum tolerable dose (50 μ g/kg), on PWT of TG and WT mice. Means \pm SEM from $n = 7$ for each group.

Discussion

While previous evidences have indicated a clear correlation between nerve injury-induced $Ca_v\alpha_2\delta_1$ upregulation and neuropathic pain states [3,4,7], pathways underlying $Ca_v\alpha_2\delta_1$ -mediated behavioral hypersensitivity have not been identified. Since cellular changes in spinal cord and DRG after a prolonged peripheral nerve injury include altered expression of large numbers of genes [5,6,30], multiple factors induced by nerve injury may contribute to neuropathic hypersensitivity. In an attempt to study if upregulated $Ca_v\alpha_2\delta_1$ is sufficient to mediate behavioral hypersensitivity, and if so, what is the underlying mechanism, we have generated TG mice overexpressing $Ca_v\alpha_2\delta_1$ in neuronal tissues and compared their behavioral sensitivity to mechanical stimulation with the SNL model. Our data indicated that the TG mice have tactile allodynia similar to that observed in the SNL model. Since similar behavioral hypersensitivity and increased $Ca_v\alpha_2\delta_1$ expression occur in both models, and other factors such as compensatory changes in the TG mice and other injury-induced factors in the SNL model are not likely the same, our data, while do not allow the complete exclusion of other factors, strongly support that increased $Ca_v\alpha_2\delta_1$ pro-

teins are sufficient in mediating tactile allodynia observed in these models.

The pharmacology profiles in allodynia reversal by the spinally delivered, selective glutamate receptor antagonists are consistent with the pharmacology profiles of allodynia reversal by these agents in the SNL model as reported by others [17]. These data suggest that increased $Ca_v\alpha_2\delta_1$ mediates allodynia through a pathway related to a selective activation of AMPA/kainate and NMDA receptors. In contrast, metabotropic glutamate receptor antagonist, AP3, was without effect in allodynia reversal. This is consistent with the notion that metabotropic glutamate receptors are mainly involved in acute and persistent pain states associated with input from afferent C-fibers, but less so on tactile allodynia likely mediated by input from A β fibers [31].

Since the TG mice have elevated $Ca_v\alpha_2\delta_1$ in different neuronal tissues ranging from DRG in the peripheral to spinal cord and supraspinal cord locations in the central nervous system [7], it is possible that increased $Ca_v\alpha_2\delta_1$ expression in peripheral as well as central locations in the TG mice




Figure 5
Similar excitability to high intensity electrical stimulation between WT and TG wide-dynamic-range neurons. Extracellular recordings were performed blindly to record dorsal horn single WDR neuron activity in response to C-fiber strength electrical stimulation applied to the receptive fields over the hindpaw of the WT and TG mice. Numbers of spikes from each neuron were pooled and the data are presented as the means \pm SEM from $n = 10$ in each group. C – C-fiber responses. PD – post discharge.


contributes to the behavioral hypersensitivity through a glutamate receptor-dependent mechanism. Therefore, the enhanced dorsal horn neuronal excitability could derive from altered pre-synaptic neurotransmission, intrinsic post-synaptic neuron excitability or descending modulation from supra-spinal locations in the TG mice. While our data do not allow the exclusion of potential contributions to tactile allodynia by activated AMPA/kainate and NMDA receptors at supra-spinal locations in the TG mice, Chaplan et al. [17] reported that only spinal, but not i.c.v., injection of NMDA receptor antagonists reversed injury-induced allodynia in the SNL model. This suggests that activation of a spinal glutamatergic pathway is critical in the development of neuropathic pain states post peripheral nerve injury.

Data from this study further demonstrate that, in contrast to the enhanced excitability of TG WDR neurons to natural stimuli reported previously [8], acutely evoked excitability of WDR neurons to C-fiber-strength electric stimulation is not altered in the TG mice. While mechanisms underlying this discrepancy are not clear, the simplest explanation is that detected WDR neuron excitability to natural stimuli derives from neuronal activation by inputs from multiple sensory fibers including A-fibers,

and elevated $Ca_v\alpha_2\delta_1$ expression results in a higher excitability of the WDR neuron to A-fiber-, but not C-fiber-, strength stimulation (see discussion below). Thus, it is likely that increased $Ca_v\alpha_2\delta_1$ expression mediates tactile allodynia through a $Ca_v\alpha_2\delta_1$ -dependent pathway by enhancing synaptic strength of A-fibers. Indeed, this hypothesis is consistent with reported findings that: (a) The basal level of $Ca_v\alpha_2\delta_1$ expression is relatively high in small DRG neurons, low in large DRG neurons [6,8], and there is a dramatic increase in $Ca_v\alpha_2\delta_1$ expression in large DRG neurons after SNL injury [6] or in the $Ca_v\alpha_2\delta_1$ TG mice [8]; (b) increased $Ca_v\alpha_2\delta_1$ expression correlates with development of allodynia mainly mediated by large A β fibers, but not formalin- and carrageenan-induced acute pain states mainly mediated by nociceptive C-fiber inputs in the TG mice [8], and formalin- or carrageenan-induced pain models do not have increased $Ca_v\alpha_2\delta_1$ expression in DRG and spinal cord (data not shown); (c) increased $Ca_v\alpha_2\delta_1$ expression in the TG mice does not alter the receptive field sizes of WDR neurons, nor the excitability of these neurons to repetitive C-fiber stimulation (wind-up) [8]; (d) In the model of sciatic nerve chronic constriction injury with increased $Ca_v\alpha_2\delta_1$ expression [3], a subpopulation of dorsal horn neurons exhibits abnormal afterdischarges to mechanical and thermal stimuli, but the number of dorsal horn neurons showing wind-up phenomenon to repetitive C-fiber stimulation remains unchanged [32].

The increase in frequency of AMPA/kainate and NMDA receptor mediated mEPSCs in deep dorsal horn neurons from the TG mice suggests that elevated spinal $Ca_v\alpha_2\delta_1$ proteins lead to an increased rate of quanta release of excitatory neurotransmitter glutamate from pre-synaptic terminals at the resting state, which in turn, enhances excitability of post-synaptic dorsal horn neurons through mainly AMPA/kainate receptor activation. Since the deep dorsal horn neurons for in vitro and vivo recordings were selected from a similar dorsal horn lamina, and very few non-WDR neurons (which were excluded) were detected in this lamina during neuron sampling in vivo, it is highly likely that a similar population of WDR neurons was examined in our in vivo and in vitro recording. Our data indicate that elevated $Ca_v\alpha_2\delta_1$ proteins do not change the amplitudes of AMPA/kainate and NMDA receptor-mediated mEPSCs. This suggests that neither the quantum size of glutamate release nor the intrinsic excitability of post-synaptic dorsal horn neurons to glutamate receptor activation is changed by increased $Ca_v\alpha_2\delta_1$ expression. Direct measurements of glutamate release in deep dorsal horn and membrane excitability of WDR neurons from TG mice would further help to clarify these issues.

It remains elusive that AMPA/kainate receptor antagonists only normalize behavioral hypersensitivity in the TG mice

**Figure 6**

Increased frequency, but not amplitude, of AMPA/kainate- and NMDA-receptor mediated mEPSCs in deep-dorsal horn neurons of the TG mice. Spinal cord slices were prepared from the WT and TG mice, and spontaneous mEPSCs were recorded blindly from deep-dorsal horn neurons. **A.** Representative traces mediated by AMPA/kainate and NMDA receptors, respectively. **B.** Summarized mEPSC frequency (left) and amplitude (right) from WT and TG neurons mediated by AMPA/kainate and NMDA receptors, respectively. **C.** Representative traces of AMPA/kainate receptor mediated mEPSCs recorded from WT and TG neurons. **D.** Summarized mEPSC rising time (left) and decay time constant (right) from WT and TG neurons mediated by AMPA/kainate and NMDA receptors, respectively. Data presented are the means \pm SEM from the number of independent neurons indicated in the text. Statistic significance is indicated by $p < 0.01$ using unpaired Students' *t* test. n.s. – not significant

without affecting the behavioral sensitivity in the WT mice significantly. It is likely that binding of glutamate to AMPA receptors at a resting state causes a conformational change of the receptors that is linked to channel opening followed by desensitization of the receptors, a mechanism controlling glutamate-induced neuron activation under normal physiological conditions [33]. It is possible that DNQX has lower binding affinity to AMPA receptors at the conformation under physiological conditions, thus, is less effective in blocking AMPA receptors. In the transgenic mice or under a pathological condition in which pre-synaptic glutamate release is enhanced, the AMPA receptor channels could be in a different conformational state due to partial membrane depolarization and channel opening [33], a primed state as suggested in Fig. 7B. DNQX could have a higher affinity to this state, thus, be

more effective in blocking AMPA receptors in the TG mice or pathological conditions. Alternatively, glutamate activation of kainate receptors at the pre-synaptic sensory neuron terminals can reduce glutamate release [34], and that at spinal interneurons can induce inhibitory transmitter release [35], which together help to terminate transient activation of post-synaptic dorsal horn neurons under normal physiological conditions. This heterosynaptic regulation is transient due to kainate receptor desensitization [35]. It is possible that this heterosynaptic regulation is less sensitive to DNQX. However, in the TG mice or under a pathological condition in which pre-synaptic glutamate release is enhanced, prolonged activation of kainate receptors at inhibitory interneurons can enhance GABA release that may activate pre-synaptic GABA_B autoreceptors, which in turn, reduces potential-




Figure 7

Proposed mechanism of $Ca_v\alpha_2\delta_1$ -mediated spinal hypersensitivity. A model of enhanced spinal glutamate release illustrates the possible mechanism of $Ca_v\alpha_2\delta_1$ -mediated spinal hypersensitivity. **A.** Under normal conditions, the $Ca_v\alpha_2\delta_1$ expression level and quantum release of excitatory neurotransmitter glutamate from pre-synaptic A-fiber terminals is low so that post-synaptic deep dorsal horn neurons maintain a normal excitation threshold. As a consequence, glutamate release in response to stimuli results in a physiological activation of deep dorsal horn neurons mainly by activating AMPA/kainate receptors. The $Ca_v\alpha_2\delta_1$ expression level at C-fiber terminals is relatively high (causing a ceiling effect for C-fiber activation), but there is little C-fiber input onto deep dorsal horn neurons at a resting state. **B.** Under a pathological condition with an elevated $Ca_v\alpha_2\delta_1$ -level at the A-fiber pre-synaptic terminals, such as post peripheral nerve injuries, the quantum release of glutamate from these terminals at a resting membrane potential is dramatically enhanced. This glutamate "leaking" causes a partial depolarization of post-synaptic deep dorsal horn neurons by activating AMPA/kainate receptors. As a consequence, these primed deep dorsal horn neurons have lowered excitation thresholds to subsequently evoked glutamate release, and also hyperexcitability due to lowered thresholds to NMDA receptor activation. Neither the partial depolarization of deep dorsal horn neurons nor the increased $Ca_v\alpha_2\delta_1$ expression in C-fiber terminals has any detectable effect to high intensity C-fiber stimulations. **C.** Proposed potential mechanisms mediating the $Ca_v\alpha_2\delta_1$ -enhancement of spinal glutamate release.

dependent inhibitory transmitter release and causes disinhibition [35]. It is possible that DNQX is more effective in blocking this prolonged activation of kainate receptors. Finally, it is possible that the WT mice are relatively insensitive to ($\text{A}\beta$ -mediated) mechanical stimulation, possibly due to the low level $\text{Ca}_v\alpha_2\delta_1$ expression in large neurons as discussed previously, so that it is difficult to detect any change in mechanical sensitivity. These possibilities may be not mutually exclusive.

Nevertheless, findings from this study allow us to propose the following mechanism of $\text{Ca}_v\alpha_2\delta_1$ -mediated tactile allodynia (Fig. 7): At the resting state under normal synaptic functions, the expression levels of $\text{Ca}_v\alpha_2\delta_1$ in large DRG neurons and A-fiber terminals are relatively low, so as the rate of quanta glutamate release from A-fiber terminals that maintains a normal neuronal excitation threshold to innocuous stimulation. The $\text{Ca}_v\alpha_2\delta_1$ expression levels in small DRG neurons and C-fiber terminals are relatively high, but may have little effect on glutamate release from C-fiber terminals at a resting state due to the lack of C-fiber inputs. This high level of basal $\text{Ca}_v\alpha_2\delta_1$ expression in small DRG neurons may set up a ceiling effect for the influence of $\text{Ca}_v\alpha_2\delta_1$ on WDR neuron excitability to C-fiber or windup stimulations (discussed below).

Mild pre-synaptic membrane depolarization in response to an action potential leads to the opening of VGCC, triggering calcium influx and a quantum release of excitatory neurotransmitter glutamate. Binding of glutamate to post-synaptic AMPA receptors at a resting state causes a conformational change of the receptors that is linked to channel opening, which is terminated by desensitization of the receptors [33]. In addition, glutamate activation of kainate receptors at the pre-synaptic sensory neuron terminals can reduce glutamate release [34], and that at spinal interneurons can induce inhibitory transmitter release [35]. This heterosynaptic regulation, which is transient due to kainate receptor desensitization, also plays a role in terminating transient activation of post-synaptic dorsal horn neurons under normal physiological conditions [35]. Under normal physiological conditions, released glutamate is insufficient in activating post-synaptic NMDA receptors (Fig. 7A).

After peripheral nerve injuries, injury-induced $\text{Ca}_v\alpha_2\delta_1$ upregulation in DRG neurons, more so in large DRG neurons, leads to increased axonal transport of $\text{Ca}_v\alpha_2\delta_1$ proteins to central pre-synaptic terminals, which in turn causes an increased rate of basal release of glutamate through a yet identified mechanism. This "glutamate leaking" causes partial sensitization of deep dorsal horn neurons by activating mainly AMPA/kainate receptors at the resting state. These "sensitized" dorsal horn neurons

therefore have lowered excitation thresholds to a subsequent release of glutamate in response to innocuous or mild nociceptive stimuli that could evoke further conformational change of the AMPA/kainate receptors and prolonged channel opening, leading to spinal hyperexcitability. In addition, signals of mild noxious stimulation could be further amplified in these sensitized dorsal horn neurons due to their low thresholds to NMDA receptor activation, leading to exaggerated painful sensations (Fig. 7B). These hyperexcitability states are sensitive to antagonists of AMPA/kainate and NMDA receptors. However, neither the partial neuron sensitization by glutamate leaking from A-fiber terminals nor the increased $\text{Ca}_v\alpha_2\delta_1$ expression in C-fiber terminals has any detectable effect on WDR neuron excitability to high intensity C-fiber stimulations (the ceiling effect discussed above).

The detail mechanisms regarding how elevated $\text{Ca}_v\alpha_2\delta_1$ mediates enhanced pre-synaptic glutamate release remain elusive (Fig. 7C). Based on *in vitro* data indicating that $\text{Ca}_v\alpha_2\delta_1$ expression is required for normal membrane expression of functional VGCC [36-40], which is critical in mediating pre-synaptic neurotransmission, it is possible that elevated $\text{Ca}_v\alpha_2\delta_1$ leads to an enhanced VGCC expression at the pre-synaptic terminals. However, there is no direct evidence so far to support this hypothesis. Alternatively, recent findings in *Drosophila* point to the direction that $\text{Ca}_v\alpha_2\delta$ plays a critical role in formation of pre-synaptic structures and active zones related to pre-synaptic neurotransmission [41]. Indeed, it has been proposed that the ability of descending excitatory 5HT3 mediated projections to induce a state-dependency in the actions of gabapentin may also occur with the monoamine receptor interacting with pre-synaptic mechanisms at the spinal level [16]. Thus, it is possible that increased $\text{Ca}_v\alpha_2\delta_1$ causes enhanced synaptic strength and/or synapse numbers by promoting formation of more active zones through a yet identified mechanism. These two possibilities may not be mutually exclusive, and deserve further investigation.

Conclusion

In conclusion, findings from this study support that elevated $\text{Ca}_v\alpha_2\delta_1$ causes spinal hypersensitivity and tactile allodynia through at least partially enhanced pre-synaptic glutamate release, which may lead to sensitization of post-synaptic dorsal horn neurons by lowering their thresholds to innocuous stimuli. This spinal sensitization may contribute to the neuropathic pain states derived from increased pre-synaptic $\text{Ca}_v\alpha_2\delta_1$ expression.

Methods

Animals

Transgenic mice over-expressing the $Ca_v\alpha_2\delta_1$ gene were generated with a transgenic vector containing the mouse brain $Ca_v\alpha_2\delta_1$ cDNA (Genbank accession number [U73484](#)) down-stream of a 6.5 kb murine *thy-1.2* gene extending from the promoter region to the intron after exon 4 without exon 3 and its flanking introns as described previously [42]. Thy-1 is a member of the immunoglobulin superfamily that is expressed in both neuronal and non-neuronal tissues [43], and deletion of exon 3 and its flanking introns has been shown to eliminate expression in non-neuronal cells [44]. Only adult male TG mice and their WT littermates were used for the experiments. All the mice were fertile, appeared normal with respect to grooming, social interactions and feeding, and showed no signs of abnormality or any obvious motor defects, tremor, seizure, or ataxia [8].

All animal care and experimental procedures were performed based on protocols approved by the Institutional Animal Care Committees of the University of California, Irvine, University College London and Indiana University.

Neuropathic lesions

The surgical procedure of SNL was performed as described [45]. Briefly, under isoflurane anesthesia, the left L5 spinal nerve of mice were exposed and ligated with a silk suture distal to DRG and proximal to their conjunction to form the sciatic nerve. Sham operations were performed in the same way except that spinal nerves were not ligated.

Behavioral assays

Behavioral response to mechanical stimulation was tested blindly as described before and post drug treatments [8]. Briefly, mice were allowed to acclimatize for 1 hr in a clear plastic cage with a wire mesh bottom. The 50% paw withdrawal thresholds (PWT) to calibrated von Frey filament (Stoelting, Wood Dale, IL) stimulation were determined using a modified up-down method of Dixon [46]. A series of filaments, starting with one that has a buckling weight of 0.41 g, were applied in consecutive sequence to the plantar surface of the left hindpaw with a pressure causing the filament to buckle. Paw lifting within 5 s was considered a positive response and led to the use of the next weaker filament. Absence of a paw lifting after 5 s led to the use of the next filament with increasing weight. The 50% paw withdrawal threshold was calculated as described before [2] from the resulting scores of six measurements starting from the one prior to the first positive response; or until four consecutive positive (assigned a score of 0.01 g) or five consecutive negative (assigned a score of 3 g) responses had occurred. Paw withdrawal thresholds from each hindpaw of a SNL mouse were recorded separately, or averaged paw withdrawal thresh-

olds from both hindpaws of the WT and injury-free $Ca_v\alpha_2\delta_1$ TG mice were used for comparing sensitivities between injury and non-injury side in SNL mice or between injury-free WT and TG mice, respectively, with or without drug treatment.

Drug administration

Drug working solutions were prepared in sterile saline before intrathecal injections between lumbar regions 5–6 (5 μ L/mouse) [47]. In some cases that multiple drugs were used in the same group of animals, a three-day drug-free period was introduced between each drug to ensure that the previous drug was completely out, even though all the drug effects in this study were terminated within 24 hr post injection.

Electrophysiological recordings in dorsal horn neurons in vivo

Recordings were performed by examiners blinded to the mouse genotype. Animals were placed in a stereotaxic frame under anesthesia and the L3–L6 spinal segments were exposed with a laminectomy. Extracellular recordings were made from individual wide dynamic range neurons using parylene coated tungsten electrodes (A-M Systems, USA). Neurons were visualized, isolated, and discriminated based on their spike amplitudes and waveforms. All neurons used for recording had receptive fields over the hindpaw. Depth of the neuron was measured from the surface of the spinal cord. Spontaneous activity was quantified over a period of at least 100 s in the absence of any evoked stimulus on isolation of a neuron.

Electrical stimulation was given via two needles inserted into the receptive field and the evoked activity was characterized and quantified as follows. A train of 16 stimuli was given (2 ms pulse duration, 0.5 Hz at three-times the C-fiber threshold), and the evoked neuronal responses were quantified using online analysis (Spike 2 software, CED Cambridge UK). The evoked neuronal responses were superimposed as a post-stimulus histogram and responses identified – based on fiber conduction velocity and the resulting latency – as C-fiber-evoked (50–250 msec) action potentials. Neuronal responses occurring after the C-fiber latency band of the neuron (250–800 ms) were classed as post discharge. The 'input' (non-potentiated response) was determined as: Input = [action potentials (50–800 ms) evoked by first pulse at 3 times C-fiber threshold] \times total number of pulses.

Electrophysiological recordings in dorsal horn of spinal cord slices

Recordings were performed by examiners blinded to the mouse genotype. The animals were anesthetized with ketamine-HCl (1 mg/kg, i.p.) and the spinal cord was extracted by hydraulic extrusion. Transverse slices (330

μm) were cut from the L5 lumbar enlargement using a vibratome (VT1000S; Leica, Nussloch, Germany) in an ice-cold (4°C) sucrose solution containing (in mM): 230 sucrose, 26 NaHCO_3 , 2.5 KCl, 1.25 NaH_2PO_4 , 0.5 CaCl_2 , 10 MgSO_4 , 10 glucose, pH 7.4, 290–305 mOsm/L, equilibrated with 95% O_2 and 5% CO_2 . The slices were maintained in an artificial cerebrospinal fluid (ACSF) containing (in mM): 130 NaCl, 3 KCl, 2 CaCl_2 , 2 MgCl_2 , 1.25 NaH_2PO_4 , 26 NaHCO_3 , and 10 glucose, pH 7.4, 295–305 mOsm/L. The ACSF was continuously equilibrated with 95% O_2 and 5% CO_2 , and slices were incubated for > 1 h before recording. Recording electrodes were prepared from borosilicate glass (World Precision Instruments, Sarasota, FL) using a horizontal electrode puller (P-97; Sutter Instruments, Novato, CA). Electrodes had resistances of 2–4 M Ω when filled with an intracellular solution containing (in mM): 92 CsMeSO₄, 43 CsCl, 1 MgCl_2 , 2 EGTA, 5 TEA, 10 HEPES and 2 Mg-ATP, pH 7.4, 295–300 mOsm/L. Oxygenated ACSF was used as bath solution, and the flow rate was adjusted to 2–3 mL/min. Depth of the neuron was measured from the surface of the spinal cord and neurons in the deep dorsal horn were visualized with an infrared-differential interference contrast (DIC) microscope (BX50WI; Olympus Optical, Tokyo, Japan) and a CCD camera. All recordings were performed at $25 \pm 1^{\circ}\text{C}$ with an Axopatch 200B amplifier (Molecular Devices, Foster City, CA). During whole-cell recordings, series resistance (8–15 M Ω) was monitored periodically, and cells with $> 15\%$ changes were excluded from the analysis. Signals were filtered at 1 kHz and digitized at a sampling rate of 5 kHz using a data-acquisition program (Axograph 4.6; Molecular Devices).

To record mEPSCs, tetrodotoxin (1 μM) and bicuculline (30 μM) were continuously applied in the bath solution. At a holding potential of -70 mV, the recorded mEPSCs were mainly mediated by AMPA/kainate receptors because application of AMPA/kainate receptor antagonist CNQX (10 μM) could reduce mEPSCs by $91.3 \pm 5.3\%$ and $93.9 \pm 3.3\%$ in WT and TG neurons ($n = 5$ in each group), respectively. In the presence of CNQX (10 μM), NMDA receptor-mediated mEPSCs were recorded at a holding potential of +50 mV.

Statistical Analyses

Indicated significant changes were defined by the two-tailed p values as determined with unpaired Student's t test, or two-way ANOVA followed by Bonferroni post-tests.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

DN participated in the design, carried out the behavioral pharmacology studies, and participated in drafting the manuscript. PD participated in the design and carried out the electrophysiology studies in spinal cord slices. EAM participated in the design and carried out the electrophysiology studies in dorsal horn neurons in vivo. DSK participated in the experimental design and animal model studies. GF participated in the design and generation of the transgenic mice. AHD participated in the experimental design, coordination and data analysis of the in vivo electrophysiology studies, and in drafting the manuscript. ZCX participated in the experimental design, coordination and data analysis of the electrophysiology studies in spinal cord slices, and in drafting the manuscript. ZDL conceived of the study and participated in its design, coordination and drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Supported in part by NIH grants: NS40135, DE14545 (Z. D. Luo). We would like to thank Mr. Timothy Yolo for his assistant in maintaining the transgenic mice. David Nguyen was supported by a Medical Student Anesthesia Research Fellowship from the Foundation for Anesthesia Education and Research.

References

1. Mitka M: "Virtual textbook" on pain developed: effort seeks to remedy gap in medical education. *Jama* 2003, **290**:2395.
2. Luo ZD, Chaplan SR, Higuera ES, Sorkin LS, Stauderman KA, Williams ME, Yaksh TL: **Upregulation of dorsal root ganglion (alpha)2(delta) calcium channel subunit and its correlation with allodynia in spinal nerve-injured rats.** *J Neurosci* 2001, **21**:1868-1875.
3. Luo ZD, Calcutt NA, Higuera ES, Valder CR, Song YH, Svensson CI, Myers RR: **Injury type-specific calcium channel alpha 2 delta-1 subunit up-regulation in rat neuropathic pain models correlates with allodynic effects of gabapentin.** *J Pharmacol Exp Ther* 2002, **303**:1199-1205.
4. Newton RA, Bingham S, Case PC, Sanger GJ, Lawson SN: **Dorsal root ganglion neurons show increased expression of the calcium channel alpha2delta-1 subunit following partial sciatic nerve injury.** *Brain Res Mol Brain Res* 2001, **95**:1-8.
5. Wang H, Sun H, Della Penna K, Benz RJ, Xu J, Gerhold DL, Holder DJ, Koblan KS: **Chronic neuropathic pain is accompanied by global changes in gene expression and shares pathobiology with neurodegenerative diseases.** *Neuroscience* 2002, **114**:529-546.
6. Valder CR, Liu JJ, Song YH, Luo ZD: **Coupling gene chip analyses and rat genetic variances in identifying potential target genes that may contribute to neuropathic allodynia development.** *J Neurochem* 2003, **87**:560-573.
7. Li CY, Song YH, Higuera ES, Luo ZD: **Spinal dorsal horn calcium channel alpha2delta-1 subunit upregulation contributes to peripheral nerve injury-induced tactile allodynia.** *J Neurosci* 2004, **24**:8494-8499.
8. Li CY, Zhang XL, Matthews EA, Li KW, Kurwa A, Boroujerdi A, Gross J, Gold MS, Dickenson AH, Feng G, Luo ZD: **Calcium channel alpha2delta1 subunit mediates spinal hyperexcitability in pain modulation.** *Pain* 2006, **125**:20-34.
9. Gee NS, Brown JP, Dissanayake VU, Offord J, Thurlow R, Woodruff GN: **The novel anticonvulsant drug, gabapentin (Neurontin), binds to the alpha2delta subunit of a calcium channel.** *J Biol Chem* 1996, **271**:5768-5776.
10. Hwang JH, Yaksh TL: **Effect of subarachnoid gabapentin on tactile-evoked allodynia in a surgically induced neuropathic pain model in the rat.** *Regional Anesthesia* 1997, **22**:249-256.

11. Abdi S, Lee DH, Chung JM: **The anti-allodynic effects of amitriptyline, gabapentin, and lidocaine in a rat model of neuropathic pain.** *Anesthesia and Analgesia* 1998, **87**:1360-1366.
12. Field MJ, McCleary S, Hughes J, Singh L: **Gabapentin and pregabalin, but not morphine and amitriptyline, block both static and dynamic components of mechanical allodynia induced by streptozocin in the rat.** *Pain* 1999, **80**:391-398.
13. Backonja M, Beydoun A, Edwards KR, Schwartz SL, Fonseca V, Hes M, LaMoreaux L, Garofalo E: **Gabapentin for the symptomatic treatment of painful neuropathy in patients with diabetes mellitus: a randomized controlled trial [see comments].** *Jama* 1998, **280**:1831-1836.
14. Rosner H, Rubin L, Kestenbaum A: **Gabapentin adjunctive therapy in neuropathic pain states.** *Clinical Journal of Pain* 1996, **12**:56-58.
15. Rosenberg JM, Harrell C, Ristic H, Werner RA, de Rosayro AM: **The effect of gabapentin on neuropathic pain.** *Clinical Journal of Pain* 1997, **13**:251-255.
16. Suzuki R, Rahman W, Rygh LJ, Webber M, Hunt SP, Dickenson AH: **Spinal-supraspinal serotonergic circuits regulating neuropathic pain and its treatment with gabapentin.** *Pain* 2005, **117**:292-303.
17. Chaplain SR, Malmberg AB, Yaksh TL: **Efficacy of spinal NMDA receptor antagonism in formalin hyperalgesia and nerve injury evoked allodynia in the rat.** *J Pharmacol Exp Ther* 1997, **280**:829-838.
18. Gao X, Kim HK, Chung JM, Chung K: **Enhancement of NMDA receptor phosphorylation of the spinal dorsal horn and nucleus gracilis neurons in neuropathic rats.** *Pain* 2005, **116**:62-72.
19. Nudler S, Piriz J, Urbano FJ, Rosato-Siri MD, Renteria ES, Uchitel OD: **Ca²⁺ channels and synaptic transmission at the adult, neonatal, and P/Q-type deficient neuromuscular junction.** *Ann N Y Acad Sci* 2003, **998**:11-17.
20. Perrier JF, Alaburda A, Hounsgaard J: **Spinal plasticity mediated by postsynaptic L-type Ca²⁺ channels.** *Brain Res Brain Res Rev* 2002, **40**:223-229.
21. Westenbroek RE, Hoskins L, Catterall WA: **Localization of Ca²⁺ channel subtypes on rat spinal motor neurons, interneurons, and nerve terminals.** *J Neurosci* 1998, **18**:6319-6330.
22. Kato A, Ohkubo T, Kitamura K: **Allosteric pain processing in mouse spinal cord: differential involvement of voltage-dependent Ca(2+) channels in synaptic transmission.** *Br J Pharmacol* 2002, **135**:1336-1342.
23. Matthews EA, Dickenson AH: **Effects of spinally delivered N- and P-type voltage-dependent calcium channel antagonists on dorsal horn neuronal responses in a rat model of neuropathy.** *Pain* 2001, **92**:235-246.
24. Reid CA, Bekkers JM, Clements JD: **Presynaptic Ca²⁺ channels: a functional patchwork.** *Trends Neurosci* 2003, **26**:683-687.
25. Talley EM, Cribbs LL, Lee JH, Daud A, Perez-Reyes E, Bayliss DA: **Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels.** *J Neurosci* 1999, **19**:1895-1911.
26. Ji RR, Kohno T, Moore KA, Woolf CJ: **Central sensitization and LTP: do pain and memory share similar mechanisms?** *Trends Neurosci* 2003, **26**:696-705.
27. Zou X, Lin Q, Willis WD: **Role of protein kinase A in phosphorylation of NMDA receptor I subunits in dorsal horn and spinothalamic tract neurons after intradermal injection of capsaicin in rats.** *Neuroscience* 2002, **115**:775-786.
28. Zhang X, Wu J, Lei Y, Fang L, Willis WD: **Protein phosphatase modulates the phosphorylation of spinal cord NMDA receptors in rats following intradermal injection of capsaicin.** *Brain Res Mol Brain Res* 2005, **138**:264-272.
29. Li P, Zhuo M: **Silent glutamatergic synapses and nociception in mammalian spinal cord.** *Nature* 1998, **393**:695-698.
30. Costigan M, Befort K, Karchewski L, Griffin RS, D'Urso D, Allchorne A, Sitariski J, Mannion JW, Pratt RE, Woolf CJ: **Replicate high-density rat genome oligonucleotide microarrays reveal hundreds of regulated genes in the dorsal root ganglion after peripheral nerve injury.** *BMC Neurosci* 2002, **3**:16. Print 2002 Oct 2025.
31. Varney MA, Gereau RW: **Metabotropic glutamate receptor involvement in models of acute and persistent pain: prospects for the development of novel analgesics.** *Curr Drug Targets CNS Neurol Disord* 2002, **1**:283-296.
32. Laird JM, Bennett GJ: **An electrophysiological study of dorsal horn neurons in the spinal cord of rats with an experimental peripheral neuropathy.** *J Neurophysiol* 1993, **69**:2072-2085.
33. Sun Y, Olson R, Horning M, Armstrong N, Mayer M, Gouaux E: **Mechanism of glutamate receptor desensitization.** *Nature* 2002, **417**:245-253.
34. Kerchner GA, Wilding TJ, Li P, Zhuo M, Huettner JE: **Presynaptic kainate receptors regulate spinal sensory transmission.** *J Neurosci* 2001, **21**:59-66.
35. Kerchner GA, Wang G-D, Qiu C-S, Huettner JE, Zhuo M: **Direct presynaptic regulation of GABA/glycine release by kainate receptors in the dorsal horn: An ionotropic mechanism.** *Neuron* 2001, **32**:477-488.
36. Gurnett CA, Felix R, Campbell KP: **Extracellular interaction of the voltage-dependent Ca²⁺ channel alpha2delta and alpha1 subunits.** *J Biol Chem* 1997, **272**:18508-18512.
37. Gurnett CA, De Waard M, Campbell KP: **Dual function of the voltage-dependent Ca²⁺ channel alpha 2 delta subunit in current stimulation and subunit interaction.** *Neuron* 1996, **16**:431-440.
38. Felix R, Gurnett CA, De Waard M, Campbell KP: **Dissection of functional domains of the voltage-dependent Ca²⁺ channel alpha2delta subunit.** *J Neurosci* 1997, **17**:6884-6891.
39. Wiser O, Trus M, Tobi D, Halevi S, Giladi E, Atlas D: **The alpha 2/delta subunit of voltage sensitive Ca²⁺ channels is a single transmembrane extracellular protein which is involved in regulated secretion.** *Febs Letters* 1996, **379**:15-20.
40. Williams ME, Feldman DH, McCue AF, Brenner R, Velicelebi G, Ellis SB, Harpold MM: **Structure and functional expression of alpha 1, alpha 2, and beta subunits of a novel human neuronal calcium channel subtype.** *Neuron* 1992, **8**:71-84.
41. Dickman DK, Kurshan PT, Schwarz TL: **Mutations in a Drosophila alpha2delta voltage-gated calcium channel subunit reveal a crucial synaptic function.** *J Neurosci* 2008, **28**:31-38.
42. Feng G, Mellor RH, Bernstein M, Keller-Peck C, Nguyen QT, Wallace M, Nerbonne JM, Lichtman JW, Sanes JR: **Imaging Neuronal Subsets in Transgenic Mice Expressing Multiple Spectral Variants of GFP.** *Neuron* 2000, **28**:41-51.
43. Gordon JW, Chesa PG, Nishimura H, Rettig WJ, Maccari JE, Endo T, Seravalli E, Seki T, Silver J: **Regulation of Thy-1 gene expression in transgenic mice.** *Cell* 1987, **50**:445-452.
44. Vidal M, Morris R, Grosveld F, Spanopoulou E: **Tissue-specific control elements of the Thy-1 gene.** *Embo Journal* 1990, **9**:833-840.
45. Kim SH, Chung JM: **An experimental model for peripheral neuropathy produced by segmental spinal nerve ligation in the rat.** *Pain* 1992, **50**:355-363.
46. Dixon WJ: **Efficient analysis of experimental observations.** *Annual Review of Pharmacology and Toxicology* 1980, **20**:441-462.
47. Inoue M, Rashid MH, Fujita R, Contos JJ, Chun J, Ueda H: **Initiation of neuropathic pain requires lysophosphatidic acid receptor signaling.** *Nat Med* 2004, **10**:712-718.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

