

Lawrence Berkeley National Laboratory

Recent Work

Title

THE 2p FINAL-STATE INTERACTION IN THE $3\text{He}(3\text{He},o\<)2p$ REACTION

Permalink

<https://escholarship.org/uc/item/3dv1q62r>

Authors

Slobodrian, R.J.

McKee, J.S.C.

Tivol, W.P.

et al.

Publication Date

1967-03-02

University of California
Ernest O. Lawrence
Radiation Laboratory

THE 2p FINAL-STATE INTERACTION IN THE ${}^3\text{He}({}^3\text{He}, \alpha)2p$ REACTION

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Physical Review Letters

UCRL-17421
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

THE $2p$ FINAL-STATE INTERACTION IN THE ${}^3\text{He}({}^3\text{He},\alpha)2p$ REACTION

R. J. Slobodrian, J. S. C. McKee, W. F. Tivol,
D. J. Clark and T. A. Tombrello

March 2, 1967

THE 2p FINAL-STATE INTERACTION IN THE ${}^3\text{He}({}^3\text{He},\alpha)2\text{p}$ REACTION

R. J. Slobodrian, J. S. C. McKee[†], W. F. Tivol,[‡]
D. J. Clark and T. A. Tombrello

Lawrence Radiation Laboratory
University of California
Berkeley, California

March 2, 1967

ABSTRACT

The angular variation of the α -particle spectra from the reaction ${}^3\text{He}({}^3\text{He},\alpha)2\text{p}$ has been measured at 43.7 MeV and 53.0 MeV laboratory energy. The energy spectra show a prominent peak at the high energy end and the angular distributions of this peak exhibit a very pronounced diffraction pattern. Good fits to the energy spectra are obtained with the Watson-Migdal final-state formalism using the known p-p scattering parameters.

Recent Letters^{1,2} have reported spectra of the ${}^3\text{He}({}^3\text{He},\alpha)2\text{p}$ reaction at 26 MeV¹ and at 11.96 MeV² laboratory energy, bearing qualitative evidence for the p-p interaction. The spectra at 26 MeV have not been analyzed theoretically, but at 6° in the laboratory system the 2p interaction seems to dominate the spectrum. A fit using the Watson-Migdal formalism³ was produced for the 11.96 MeV spectra,² but the agreement is partially obscured by the contribution of the p- α interaction (Li^5), prominent at the angles investigated.

There is continued interest in the study of reactions leading to the ${}^1\text{S}_0$ state of two nucleons, particularly with regard to the determination of the scattering parameters of the n-n interaction.⁴ The usefulness of the deuteron break-up reaction $n + d \rightarrow n + n + p$ for such purpose has been recently

[†]On leave from the University of Birmingham, England.

[‡]Summer visitor from California Institute of Technology.

questioned, and the Watson theory seems to be misleading in this reaction.⁵ The existing data on the mirror reaction $p + d \rightarrow p + p + n$ indicate that the p-n interaction obscures the effects of the p-p interaction.^{6,7} On the other hand there are indications that the peaks observed in angular correlation measurements are sharper than can be predicted by the Watson-Migdal formalism.⁸ This would indicate an effect in the right direction to explain the discrepancy between the value obtained for the n-n scattering length a_n from the $d(\pi^-, \gamma)2n$ ⁹ the ${}^3\text{H}(d, {}^3\text{He})2n$ reactions,⁴ and the values obtained from the n+d experiments.^{10,11} Recent work on the reactions ${}^2\text{H}({}^3\text{He}, t)2p$ and $p({}^3\text{He}, d)2p$ has been reported,¹² showing the spectra of the third particle to be sharper than predicted by the known p-p scattering parameters in the context of the Watson-Migdal formalism. In each reaction qualitative reasons for such an observation can be found, like the charge exchange effect that may dominate the ${}^2\text{H}({}^3\text{He}, t)2p$ reaction,¹³ and the known diffuseness of the outgoing deuteron in the $p({}^3\text{He}, d)2p$ reaction. It should be noted that the scattering lengths required to fit the spectra from the $p({}^3\text{He}, d)2p$ ¹² and ${}^3\text{H}(n, d)2n$ ¹⁴ reactions seem larger than the currently accepted values of these quantities. The same is true for the ${}^2\text{H}({}^3\text{He}, t)2p$ reaction¹² mentioned above, indicating that spectra from the reaction $d(t, {}^3\text{He})2n$ may also require a large value of a_n . No such data are presently available. Evidence of a similar correlation might be obtained from a study of the reaction ${}^2\text{H}(t, {}^3\text{He})2n$, as it could also require a high value for a_n . The physical fact is that all such reactions lead asymptotically to three particles in the final state, and this is true for Breit-Wigner resonances between pairs of particles, as well as for virtual or antibound "states". Thus, neither the Breit-Wigner resonances nor the virtual states are true quantum mechanical states, and they are described by wave functions $\psi \notin L_2(0, \infty)$.¹⁵ The Eisenbud¹⁶

interpretation of a rapidly increasing phase shift as a time delay, leads to the definition of a physical lifetime for both resonant and virtual states, and both manifest themselves by peaks in the energy spectra of the third particle. They can be elegantly interpreted in terms of poles in the scattering amplitude of two particles, and the question is how well can one extract the pole parameters from the spectrum shape (or "line" shape) of the third particle. Hence reactions leading to the 2p system and a third particle are useful to test a given theoretical formalism, if there is dominance of the p-p pole. Such comparison was made for the Watson-Migdal formalism using the ${}^3\text{He}(d,t)2p$ reaction over a wide range of energies.^{2,17} Since there was good agreement between the calculated spectra and the experimental results, the mirror reaction ${}^3\text{H}(d,{}^3\text{He})2n$ was used to determine a value and sign for the 1S_0 n-n scattering length a_n .⁴

We have studied the ${}^3\text{He}({}^3\text{He},\alpha)2p$ reaction at 43.7 and 53.0 MeV laboratory energy (at target center), using the ${}^3\text{He}$ beam of the Berkeley 88-inch variable energy cyclotron. The detection was accomplished using solid state detector telescopes in conjunction with electronic particle identifier circuits.¹⁸ Calibration spectra were obtained using the reaction ${}^{14}\text{N}({}^3\text{He},{}^4\text{He}){}^{13}\text{N}^*$. The energy resolution was about 370 keV, in the relevant region of the spectra. The beam energy was determined by measuring its range in aluminum, using a Faraday cup and an electrometer.¹⁹ The experimental techniques were basically the same as described elsewhere.¹⁷ The large positive Q is quite helpful for the detection of the identified ${}^4\text{He}$ spectra over a sizable angular range, between 5° and 42° lab (10° - 90° cm). A prominent peak near the high energy end of the alpha particle spectra was observed (fig. 1(a) and 1(c)). Its angular distribution follows a pronounced diffraction pattern with minima (fig. 1(b)) deeper than usual even for particle transfers leading to particle bound states or long lived

resonances. The angular distributions at both energies are shown in fig. 2(a) and 2(b) together with fits based on a diffraction picture of the reaction mechanism.²⁰ The P.W.B.A.²¹ is inadequate in fitting the relative values of successive maxima as well as in the positions of the minima. The rather large "interaction radii" r_i that fit the angular distributions, in excess of 4 F, lend support to the direct or peripheral picture of the reaction mechanism. Therefore it should be reasonable to expect that the transition amplitude can be factored into the product of a term containing a pole in momentum transfer and a term containing a pole in the nucleon-nucleon momentum. In the present case the rearrangement collision consists of an $l=0$ nucleon transfer. More complicated processes, like charge exchange, are excluded. The energy spectra are compared with the Watson-Migdal formalism predictions, using the formulae contained in ref. 4 and the known p-p scattering length, excluding the vacuum-polarization correction, $a_p \cong -7.7 F$.²² Figure 3 contains several sample spectra showing a good agreement between the theory and the experimental results, over a considerable angular range. Our results indicate that the study of the reaction ${}^3\text{H}(t, \alpha)2n$ should prove relevant in obtaining additional information on the n-n interaction. In particular it should be interesting to ascertain whether the ${}^3\text{H}(t, \alpha)2n$ reaction yields a scattering length in agreement with the ${}^3\text{H}(d, {}^3\text{He})2n$ reaction,⁴ and thus also with the $\pi^- + d$ experiment.⁹ If there is agreement then, hopefully, properly chosen reactions, with an outgoing third particle having a sharply defined boundary, may allow in the future a more precise and complete determination of the 1S_0 n-n scattering parameters.

We gratefully acknowledge the support of D. Landis with the electronic equipment, R. Lothrop who made the solid state detectors and J. Meneghetti for his part with the mechanical set up of the experiment.

REFERENCES

1. K. P. Artjomov, V. I. Chuev, V. Z. Goldberg, A. A. Ogloblin V. P. Rudakov, and J. N. Serikov, Phys. Letters 12, 53 (1964).
2. T. A. Tombrello and A. D. Bacher, Phys. Letters 17, 37 (1965).
3. K. M. Watson, Phys. Rev. 88, 1163 (1952); A. B. Migdal, Zh. Eksperim. i. Teor. Fiz. 28, 3 (1955)[translation: Soviet Phys. JETP 1, 2 (1955)].
4. E. Baumgartner, H. E. Conzett, E. Shield and R. J. Slobodrian, Phys. Rev. Letters 14, 318 (1965) and references therein.
5. R. Aaron and R. D. Amado, Phys. Rev. 150, 857 (1966) and references therein.
6. B. V. Rybakov, V. A. Sidorov and V. A. Vlasov, Nucl. Phys. 23, 491 (1961).
7. J. L. Friedes and M. K. Brussel, Phys. Rev. 131, 1194 (1963).
8. W. D. Simpson, J. D. Bronson, W. J. Jackson, G. C. Phillips, Rev. Mod. Phys. 37, 523 (1965); G. C. Phillips, paper 9-A International Conference on Nuclear Physics, Gatlinburg, Tennessee, September. 1966.
9. R. P. Haddock, R. M. Salter, Jr., M. Zeller, J. B. Czirr and D. R. Nygren, Phys. Rev. Letters 14, 318 (1965).
10. M. Cerineo, K. Ilakovac, J. Šlaus, P. Tomaš, and V. Valković, Phys. Rev. 133, B948 (1964).
11. V. K. Voitovetskii, I. L. Korsunskii, and V. F. Pazhin, Phys. Letters 10, 109 (1964) and Nucl. Phys. 64, 513 (1965).
12. B. J. Morton, M. P. Fricke, R. O. Ginaven, E. E. Gross, J. J. Malanify and A. Zucker, Bull. Am. Phys. Soc. Ser. II 12, 16 (1967).
13. E. M. Henley, F. C. Richards, D. V. L. Yu., Phys. Letters 16, 333 (1965).
14. V. Ajdačić, M. Cerineo, B. Lalović, G. Paić, I. Šlaus, and P. Thomaš. Phys. Rev. Letters 14, 442 (1965).

15. V. De Alfaro and T. Regge, Potential Scattering (North-Holland Publishing Company, Amsterdam 1965).
16. L. Eisenbud Thesis (1948) Princeton University. M. L. Goldberger and K. M. Watson, Collision Theory, (John Wiley and Sons, Inc., New York, 1964).
17. H. E. Conzett, E. Shield, R. J. Slobodrian, and S. Yamabe, Phys. Rev. Letters 13, 625 (1964).
18. F. S. Goulding, D. A. Landis, J. Cerny III, and R. Pehl, I.E.E.E. Trans. Nucl. Sc. 11, 388 (1964).
19. C. Williamson and P. Boujot, Rapport CEA 2189 (1962) (unpublished).
20. A. Dar, Nucl. Phys. 55, 305 (1964) and reference therein.
21. M. H. MacFarlane and J. B. French, Rev. Mod. Phys. 32, 567 (1960) and references therein.
22. R. J. Slobodrian, Nuovo Cimento 40, 443 (1965).

FIGURE CAPTIONS

Fig. 1. ${}^4\text{He}$ spectra from the reaction ${}^3\text{He} + {}^3\text{He} \rightarrow {}^4\text{He} + 2p$. A pulser group is seen at the far right.

a) At 17° Lab with 43.7 MeV bombarding energy. The arrow indicates the interval expanded in Fig. 3b).

b) At 10° Lab (near the first minimum of the 2p enhancement), with 43.7 MeV bombarding energy.

c) At 16° Lab with 53.0 MeV bombarding energy.

Fig. 2. Angular distribution of the ${}^3\text{He} + {}^3\text{He} \rightarrow {}^4\text{He} + 2p$ peaks integrated to a value where they drop to $1/e$ of the value at the maximum. The solid line corresponds to a fit with $J_0^2(q, r_1)$, the dashed line is the PWBA fit.

a) Data at 43.7 MeV. The $J_0^2(q, r_1)$ fit was calculated with $r_1 = 4.6$ F. The PWBA fit was obtained with $r_1 = 5.5$ F.


b) Data at 53.0 MeV. The interaction radii are respectively $r_1 = 4.2$ F and $r_1 = 5.0$ F for the $J_0^2(q, r_1)$ and PWBA fits.

Fig. 3. The solid lines are fits to the ${}^4\text{He}$ energy spectra obtained with the Watson-Migdal formalism using $a_p = -7.7$ F, after folding the energy resolution and the effect of the finite angular resolution of the detector (approximately $\pm 0.3^\circ$). The solid dots are the experimental points.

a) Spectrum at $\theta_{\text{Lab}} = 5^\circ$ and 43.7 MeV bombarding energy.


b) Spectrum at $\theta_{\text{Lab}} = 17^\circ$ and 43.7 MeV bombarding energy.

c) Spectrum at $\theta_{\text{Lab}} = 5^\circ$ and 53.0 MeV bombarding energy.



XBL673-984

Fig. 1



XBL673-985

Fig. 2




Fig. 3

RL673-996

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

