

UC Davis

UC Davis Previously Published Works

Title

Rational Dyck Paths in the Non Relatively Prime Case

Permalink

<https://escholarship.org/uc/item/3f29k218>

Journal

The Electronic Journal of Combinatorics, 24(3)

ISSN

1097-1440

Authors

Gorsky, Eugene

Mazin, Mikhail

Vazirani, Monica

Publication Date

2017

DOI

10.37236/6901

Peer reviewed

RATIONAL DYCK PATHS IN THE NON RELATIVELY PRIME CASE

EUGENE GORSKY, MIKHAIL MAZIN, AND MONICA VAZIRANI

ABSTRACT. We study the relationship between rational slope Dyck paths and invariant subsets of \mathbb{Z} , extending the work of the first two authors in the relatively prime case. We also find a bijection between (dn, dm) -Dyck paths and d -tuples of (n, m) -Dyck paths endowed with certain gluing data. These are the first steps towards understanding the relationship between rational slope Catalan combinatorics and the geometry of affine Springer fibers and knot invariants in the non relatively prime case.

1. INTRODUCTION

Catalan numbers, in one of their incarnations, count the number of Dyck paths, that is, the lattice paths in a square which never cross the diagonal. In recent years, a number of interesting results and conjectures [3, 4, 7, 9, 12, 13, 14, 16, 24, 28] about “rational Catalan combinatorics” have been formulated. An (n, m) -Dyck path is a lattice path in an $n \times m$ rectangle, going from the bottom-right corner $(m, 0)$ to the top-left corner $(0, n)$ and never going above the diagonal, which is the line that connects them. We will denote the set of all (n, m) -Dyck paths by $Y_{n,m}$. For coprime m and n there are a number of interesting maps involving $Y_{n,m}$, see Figure 1:

- (a) J. Anderson constructed a bijection \mathcal{A} between $Y_{n,m}$ and the set $\text{Core}_{n,m}$ of simultaneous (n, m) -core partitions.
- (b) Armstrong, Loehr, and Warrington defined a “sweep” map $\zeta : Y_{n,m} \rightarrow Y_{n,m}$ and conjectured that it is bijective. This conjecture was proved by Thomas and Williams in [26].
- (c) The first two authors defined two maps \mathcal{D} and \mathcal{G} between $Y_{n,m}$ and the set $\mathbf{M}_{n,m}$ of (n, m) -invariant subsets of $\mathbb{Z}_{\geq 0}$ containing 0. If combined with a natural bijection between $\text{Core}_{n,m}$ and $\mathbf{M}_{n,m}$, the map \mathcal{D} coincides with \mathcal{A} . Furthermore, one can prove that $\zeta = \mathcal{G} \circ \mathcal{D}^{-1}$. As a consequence, the map \mathcal{G} is also bijective.




FIGURE 1. Rational Catalan maps in the coprime case

Date: September 24, 2018.

Key words and phrases. rational Dyck paths, rational Catalan combinatorics, simultaneous core partitions, invariant integer subsets, semigroups.

The goal of the present paper is a partial generalization of the diagram in Figure 1 to the non-coprime case. Let (n, m) be relatively prime, and d be a positive integer. Let $N = dn$ and $M = dm$. The set $Y_{N,M}$ is well defined for all n, m, d , and the definition of ζ can be carried over with minimal changes. However, while the sets $\text{Core}_{N,M}$ and $\mathbf{M}_{N,M}$ are still in bijection with each other, the sets become infinite. Indeed, an (N, M) -invariant subset of $\mathbb{Z}_{\geq 0}$ can be identified with a collection of d (n, m) -invariant subsets, one for each remainder mod d . These subsets won't necessarily have minimum element 0, and so we will want to shift or translate each a fixed amount. We will want to shift or translate each a fixed amount. More abstractly, this defines a map $\epsilon : \mathbf{M}_{N,M} \rightarrow (\mathbf{M}_{n,m})^d$ and different shifts correspond to different preimages under ϵ .

To resolve this problem, we introduce a certain equivalence relation \sim on $\mathbf{M}_{N,M}$. It satisfies that $\Delta_1 \sim \Delta_2$ implies $\bar{\epsilon}(\Delta_1) = \bar{\epsilon}(\Delta_2)$, where $\bar{\epsilon} : \mathbf{M}_{N,M} \rightarrow (\mathbf{M}_{n,m})^d \rightarrow (\mathbf{M}_{n,m})^d / \mathcal{S}_d$, so $\bar{\epsilon}$ is well defined on $\mathbf{M}_{N,M} / \sim$. The following theorem is the main result of the paper.

Theorem 1.1. *For all positive N, M one can define maps*

$$\mathcal{D}, \mathcal{G} : \mathbf{M}_{N,M} / \sim \longrightarrow Y_{N,M}$$

such that the following results hold:

- (a) *The maps \mathcal{D} and \mathcal{G} are bijective.*
- (b) *The ‘‘sweep’’ map factorizes similarly to the coprime case: $\zeta = \mathcal{G} \circ \mathcal{D}^{-1}$.*
- (c) *Let $d = \gcd(N, M)$, $n = N/d$, and $m = M/d$. The composition*

$$\text{col}_d := \mathcal{D}^d \circ \epsilon \circ \mathcal{D}^{-1} : Y_{N,M} \rightarrow (Y_{n,m})^d / \mathcal{S}_d,$$

can be described as follows: color the $N + M$ steps in an (N, M) -Dyck path with d colors, i.e., by $\mathbb{Z}/d\mathbb{Z}$, so that there are $n + m$ steps of the same color i , and these steps will form an (n, m) -Dyck path after possibly translating connected components by integer multiples of $(m, -n)$ to make the i -colored steps connected.

As we do not have a canonical way of assigning colors, we must pass to \mathcal{S}_d orbits above. We shall see in Section 3.2 that the coloring is finer than \mathcal{S}_d orbits and in fact corresponds to an isomorphism class of a labeled directed graph with d nodes.




FIGURE 2. Rational Catalan maps in the non-coprime case.

We illustrate all these maps in Figure 2. We also give an explicit description of the ‘‘coloring map’’ col_d , as well as its inverse given proper gluing data. In the ‘‘classical’’ case $M = N$ we get $d = N$ and $m = n = 1$, therefore col_d colors a Dyck path in n colors such that the pairs of steps of the same color form a $(1, 1)$ -Dyck path. In this case the coloring is equivalent to presenting a Dyck path as a regular sequence of parentheses, with every opening and its corresponding closing parenthesis corresponding to the pair of steps of the same color.

We conjecture a relation between the constructions of this paper, combinatorial identities and link invariants. Recall that the ‘‘compositional rational shuffle conjecture’’ of [7] (proved in [24]) relates a certain sum over (N, M) -Dyck paths to certain matrix elements of operators acting on symmetric functions. Here we propose a different sum over (N, M) -invariant subsets, and plan to clarify the relation between the two in the future work. We define the generating series:

$$(1) \quad C_{N,M}(q, t) = \sum_{\Delta \in \mathcal{M}_{N,M}} q^{\text{gap}(\Delta)} t^{\text{dinv}(\Delta)},$$

where

$$\text{gap}(\Delta) = |\mathbb{Z}_{\geq 0} \setminus \Delta|.$$

For $d = 1$ it agrees with the rational q, t -Catalan polynomial [12, 3]

$$c_{N,M}(q, t) = \sum_{D \in \mathcal{Y}_{N,M}} q^{\text{area}(D)} t^{\text{dinv}(D)},$$

and it follows from the results of [24] that:

$$(2) \quad C_{n,m}(q, t) = c_{n,m}(q, t) = \sum_{D \in \mathcal{Y}_{n,m}} q^{\text{area}(D)} t^{\text{dinv}(D)} = (P_{n,m}(1), h_n).$$

Here $P_{n,m}$ is a certain operator defined in [16, 7] and acting on the space of symmetric functions. In particular, the left hand side of (2) is symmetric in q and t . It was also proved in [16] that the right hand side of (2) equals the ‘‘refined Chern-Simons invariant’’ (in the sense of [1]) of the (n, m) torus knot, and conjectured that it equals the Poincaré polynomial of the $(a = 0)$ part of the Khovanov-Rozansky homology [19] of this knot.

For $d > 1$, the formula for $C_{N,M}(q, t)$ generalizing (2) was conjectured in [7] and proved in [24]. However, $C_{N,M}$ is now an infinite power series while $c_{n,m}$ is a finite polynomial.

Conjecture 1.2. *For general $d \geq 1$, the following statements hold:*

- (a) *One has $C_{N,M}(q, t) = \frac{1}{(1-q)^{d-1}} (P_{n,m}^d(1), h_N)$, where $P_{n,m}$ is the same operator as in (2).*
- (b) *The series $C_{N,M}(q, t)/(1-q)$ agrees with the Poincaré series of the $(a = 0)$ part of the Khovanov-Rozansky homology of the (N, M) torus link.*

The part (a) immediately implies that $C_{N,M}(q, t)(1-q)^{d-1}$ is symmetric in q and t . To support the conjecture, we use a recent result of Elias and Hogancamp [10] to prove the following:

Theorem 1.3. *Conjecture 1.2(b) holds for $M = N$.*

In the case $M = N$, part (a) of the conjecture is equivalent to [10, Conjecture 1.15] (see also [27]), but, to our knowledge, it is still open. For general M and N , it fits into the framework of conjectures of [1, 16, 17], and we refer the reader to these references for more details.

ACKNOWLEDGEMENTS

We would like to thank François Bergeron and Nathan Williams for the useful discussions. A preliminary version of the paper was reported at the FPSAC 2016 conference [15]. The work of E. G. was partially supported by the NSF grant DMS-1559338, Hellman fellowship, grant RSF 16-11-10160 and Russian Academic Excellence Project 5-100. NSF grant DMS-1559338 partially supported collaborative visits by M. M. M.M. participation in FPSAC 2016 was supported by a KSU start-up grant.

2. RELATIVELY PRIME CASE

Let (n, m) be a pair of relatively prime positive integers. Consider an $n \times m$ rectangle $R_{n,m}$. Let $Y_{n,m}$ be the set of Young diagrams that fit under the diagonal in $R_{n,m}$. We will often abuse notation by identifying a diagram $D \in Y_{n,m}$ with its boundary path (sometimes also called a *rational Dyck path*), and with the corresponding partition. We will also think about the rectangle $R_{n,m}$ as a set of boxes, identified with a subset in $\mathbb{Z}_{\geq 0}$ with the bottom-left corner box identified with $(0, 0)$. In our convention, n is the height of $R_{n,m}$ and m is its width; and the boundary path of $D \subset R_{n,m}$ follows the boundary from the bottom-right corner to the top-left corner. See Example 2.14 below. In Section 3.3.1, it will also be convenient to identify the path D with a function (or its plot) $[0, n + m] \rightarrow \mathbb{R}^2$.

There are two important combinatorial statistics on the set $Y_{n,m}$: area and dinv .

Definition 2.1. Let $D \in Y_{n,m}$. Then $\text{area}(D)$ is equal to the number of whole boxes that fit between the diagonal of $R_{n,m}$ and the boundary path of D .

Note that $\text{area}(D)$ ranges from 0 for the full diagram to

$$\delta = \frac{(m-1)(n-1)}{2}$$

for the empty diagram. The $\text{co-area}(D) = \delta - \text{area}(D)$ is then just the number of boxes in the Young diagram D . One natural approach to the dinv statistic is to define the map $\zeta : Y_{n,m} \rightarrow Y_{n,m}$ and then set $\text{dinv}(D) := \text{area}(\zeta(D))$. In the case $m = n + 1$ the map ζ was first defined by Haglund ([18]), then it was generalized by Loehr to the case $m = kn + 1$ for any $k \in \mathbb{Z}_{\geq 0}$ ([20]), and to the general case of any relatively prime (n, m) by Gorsky and Mazin in [12]. In [4] it was put into even larger framework of so called sweep maps. Below is one of the equivalent possible definitions.

Definition 2.2. The *rank* of a box $(x, y) \in \mathbb{Z}^2$ is given by the linear function

$$\text{rank}(x, y) = mn - m - n - nx - my.$$

Note that the boxes of non-negative ranks are exactly those that fit under the bottom-right to top-left diagonal of $R_{n,m}$. Let $D \in Y_{n,m}$. One ranks the steps of the boundary path of D as follows.

Definition 2.3. The rank of a vertical step of D is equal to the rank of the box immediately to the left of it. The rank of a horizontal step is equal to the rank of the box immediately above it.

In other words, the ranks of steps can be defined inductively as follows. We follow the boundary path of D starting from the bottom-right corner. The first step is ranked $-m$. Otherwise, the rank of each step equal to the rank of the previous step plus n , if the previous step is horizontal, and it equals to the rank of the previous step minus m , if the previous step is vertical. Note the last step is ranked 0 (and is vertical).

Note that for relatively prime (n, m) all the ranks of the steps of a diagram $D \in Y_{n,m}$ are distinct.

Definition 2.4. The boundary path of the diagram $\zeta(D)$ is obtained from the boundary path of D by rearranging the steps in the increasing order of ranks.

The definition of the map ζ is illustrated in Example 2.14. One can verify that the diagram $\zeta(D)$ fits under the diagonal of $R_{n,m}$ (see [12] and [4]). The following result is considerably harder, see also [13, 18, 20, 28] for partial results in this direction.

Theorem 2.5 ([26]). *The map ζ is bijective.*

The following approach to studying the map ζ was suggested in [12].

Definition 2.6. We say that a subset $\Delta \subset \mathbb{Z}_{\geq 0}$ is (n, m) -invariant and 0-normalized if $\Delta + m \subset \Delta$, $\Delta + n \subset \Delta$, and $\min(\Delta) = 0$. Let $\mathbf{M}_{n,m}$ be the set of all such subsets Δ .

In [12] two maps \mathcal{D} and \mathcal{G} from the set $\mathbf{M}_{n,m}$ to $Y_{n,m}$ were constructed.

Definition 2.7. Let $\Delta \in \mathbf{M}_{n,m}$. The diagram $\mathcal{D}(\Delta)$ consists of all boxes in $R_{n,m}$ whose ranks belong to Δ .

Clearly, $\mathcal{D}(\Delta)$ fits under the diagonal. In particular, one gets that $\mathcal{D}(\Gamma_{n,m}) = \emptyset$, where $\Gamma_{n,m} := \{an + bm \mid a, b \in \mathbb{Z}_{\geq 0}\}$ is the semigroup generated by n and m , and $\mathcal{D}(\mathbb{Z}_{\geq 0})$ is the full diagram containing all the boxes below the diagonal. Note that the (n, m) -invariance of Δ implies that $\mathcal{D}(\Delta)$ is indeed a Young diagram. Note also that \mathcal{D} is a bijection. Indeed, it is not hard to see that rank provides a bijection between the boxes below the diagonal in $R_{n,m}$ and the integers in $\mathbb{Z}_{\geq 0} \setminus \Gamma_{n,m}$.

It is also important to sometimes consider the *periodic extension* $P(\Delta)$ of the boundary path of $\mathcal{D}(\Delta)$. Equivalently, it can be defined as the infinite lattice path separating the boxes in \mathbb{Z}^2 which ranks belong to Δ from the boxes which ranks belong to the complement $\mathbb{Z} \setminus \Delta$. We will call such paths (n, m) -periodic. See Figure 3 for an example.

Remark 2.8. J. Anderson in [2] defined a bijection between $Y_{n,m}$ and the set $\text{Core}_{n,m}$ of (n, m) -cores, that is, Young diagrams with no hooks of length n or m . The standard bijection between $\text{Core}_{n,m}$ and $\mathbf{M}_{n,m}$ identifies Anderson's bijection with the map \mathcal{D} , see e.g [13] for details.

Definition 2.9. The numbers $0 = a_0 < a_1 < \dots < a_{n-1}$, such that

$$\{a_0, \dots, a_{n-1}\} = \Delta \setminus (\Delta + n)$$

are called *the n -generators of Δ* . The numbers $\{b_0 < b_1 < \dots < b_{m-1}\}$ such that

$$\{b_0, \dots, b_{m-1}\} = (\Delta - m) \setminus \Delta$$

are called *the m -cogenerators of Δ* .

Remark 2.10. Let $D = \mathcal{D}(\Delta)$. The ranks of the vertical steps of D are exactly the n -generators of Δ , and the ranks of the horizontal steps of D are exactly the m -cogenerators of Δ . We will often mark n -generators by \times and m -cogenerators by \square .

Definition 2.11. The diagram $\mathcal{G}(\Delta)$ has row lengths g_0, \dots, g_{n-1} given by the following formula:

$$g_k = \#\{b_i \mid b_i > a_k\}.$$

Equivalently, the boundary path of $\mathcal{G}(\Delta)$ can be obtained by rearranging the set

$$S = \{a_0, \dots, a_{n-1}, b_0, \dots, b_{m-1}\}$$

in increasing order and replacing n -generators by vertical steps and m -cogenerators by horizontal steps, from bottom right to top left.

The next result follows from the above definitions.

Proposition 2.12. [12, 13] *The following identity holds:*

$$\zeta(D) = \mathcal{G} \circ \mathcal{D}^{-1}(D).$$

Corollary 2.13. *Since ζ and \mathcal{D} are bijective, the map \mathcal{G} is a bijection too.*

Example 2.14. For example, if $n = 5$, $m = 3$, and $\Delta = \{0, 3, 5, 6, 7, 8, \dots\}$ then the 5-generators of Δ are 0, 3, 6, 7, 9 and 3-cogenerators are $-3, 2, 4$. The diagram $\mathcal{D}(\Delta)$ consists of one box, which has rank 7. The ranked boundary path of D is

$$\begin{array}{cccccccc} \text{h} & \text{h} & \text{v} & \text{h} & \text{v} & \text{v} & \text{v} & \text{v} \\ -3 & 2 & 7 & 4 & 9 & 6 & 3 & 0 \end{array}$$




FIGURE 3. Here $n = 5$ and $m = 3$. On the left is (a fragment of) the periodic path $P = P(\{0, 3, 5, 6, 7, 8, \dots\})$, center is the diagram $D = \mathcal{D}(\{0, 3, 5, 6, 7, 8, \dots\})$, and on the right is the diagram $\zeta(D)$.

read bottom to top, which we sort to the boundary path of $\zeta(D)$

$$\begin{array}{cccccccc} \mathbf{h} & \mathbf{v} & \mathbf{h} & \mathbf{v} & \mathbf{h} & \mathbf{v} & \mathbf{v} & \mathbf{v} \\ -3 & 0 & 2 & 3 & 4 & 6 & 7 & 9 \end{array}$$

See Figure 3 for the diagrams D and $\zeta(D)$. Note, that if one takes the union of the 5-generators and 3-cogenerators and reads them in the increasing order, then one gets $-3, 0, 2, 3, 4, 6, 7, 9$. Replacing generators by “v” and cogenerators by “h”, one gets hvhvhvvhv, which is the boundary path of $\zeta(D)$.

The approach with invariant subsets allows one to relate the dinv statistic to geometry. Let $V = \mathbb{C}[t]/t^{2\delta}\mathbb{C}[t]$ be the ring of polynomials of degree less than $2\delta = (m - 1)(n - 1)$. Let $Gr(\delta, V)$ be the Grassmannian of half-dimensional subspaces in V . Consider the subvariety $J_{n,m} \subset Gr(\delta, V)$ consisting of subspaces in V invariant under multiplication by t^m and t^n :

$$J_{n,m} = \{U \in Gr(\delta, V) : t^m U \subset U, t^n U \subset U\}.$$

These varieties appear in algebraic geometry as local versions of the compactified Jacobians (see Beauville [5] and Piontkowski [25]), and in representation theory as homogeneous affine Springer fibers, where they were first considered by Lusztig and Smelt in [22] and then by Piontkowski [25]. Both Lusztig and Smelt, and Piontkowski showed that $J_{n,m}$ has a natural decomposition into complex affine cells enumerated by elements of $\mathbf{M}_{n,m}$. Moreover, the dimension of the cell C_Δ corresponding to an invariant subset $\Delta \in \mathbf{M}_{n,m}$ is given by

$$\dim_{\mathbb{C}} C_\Delta = |\mathcal{G}(\Delta)| = \delta - \text{dinv}(\Delta).$$

Therefore, one gets the following theorem.

Theorem 2.15 ([12]). *The Poincaré polynomial $P_{n,m}(t)$ of the variety $J_{n,m}$ is given by*

$$P_{n,m}(t) = \sum_{D \in Y_{n,m}} t^{2(\delta - \text{dinv}(D))}.$$

Moreover, bijectivity of the map ζ (or, equivalently, the map \mathcal{G}) implies a simpler formula:

$$P_{n,m}(t) = \sum_{D \in Y_{n,m}} t^{2|D|},$$

where $|D| = \delta - \text{area}(D)$ is simply the number of boxes in D .




FIGURE 4. Here $n = 9$ and $m = 6$. On the left is the diagram D with the boundary path $hvhvvhhhvhvwww$ marked with ranks; and on the right is the diagram $\zeta(D)$.

3. NON-RELATIVELY PRIME CASE.

3.1. **Sweep map.** The notion of a rational Dyck path naturally generalizes to the non relatively prime case. Let (n, m) be relatively prime, and d be a positive integer. Let $N = dn$ and $M = dm$. Consider an $N \times M$ rectangle $R_{N,M}$ and the set $Y_{N,M}$ of Young diagrams that fit under the diagonal in $R_{N,M}$. The area statistic can be generalized directly. The dinv statistic and the map ζ are a bit more tricky. It is convenient to adjust the rank function on the boxes in the following way:

$$\text{rank}(x, y) = dmn - m - n - nx - my.$$

The steps of the boundary path of a diagram $D \in Y_{N,M}$ are ranked as before with respect to the new rank function. The first step is still ranked $-m$ and the inductive description of the ranks still holds with respect to $+n, -m$; it still holds that the boxes with non-negative rank are those below the diagonal. However, for $d > 1$ some distinct steps might have the same rank, therefore rearranging the steps of the path according to their rank is problematic. The following idea for overcoming this difficulty was suggested by François Bergeron. It can also be found in [4].

Definition 3.1. Let $D \in Y_{N,M}$. The boundary path of the diagram $\zeta(D)$ is obtained from the boundary path of D by rearranging the steps so their ranks are weakly increasing. If two steps have the same rank, then they are ordered in the reversed order of appearance in the boundary path of D .

Example 3.2. Consider the diagram $D \in Y_{9,6}$ with the boundary path $hvhvvhhhvhvwww$ (see Figure 4). The ranked boundary path of D is

$$\begin{matrix} h & v & h & v & v & h & h & h & v & h & v & v & v & v & v \\ -2 & 1 & -1 & 2 & 0 & -2 & 1 & 4 & 7 & 5 & 8 & 6 & 4 & 2 & 0 \end{matrix}$$

which we sort to the boundary path of $\zeta(D)$

$$\begin{matrix} h & h & h & v & v & h & v & v & v & v & h & h & v & v & v \\ -2 & -2 & -1 & 0 & 0 & 1 & 1 & 2 & 2 & 4 & 4 & 5 & 6 & 7 & 8. \end{matrix}$$

Note there are two steps of rank 4 in the boundary path of D : when read bottom to top on the path but left to right above, first there is a horizontal step, and then there is a vertical step. In the boundary path of $\zeta(D)$ the order of these two steps is reversed. Similarly for the two steps of rank 1.

Now the statistic dinv can be defined as

$$\text{dinv}(D) := \text{area}(\zeta(D)).$$

Note that in [7] a different definition of dinv for the non relatively prime case is used:

$$\text{dinv}'(D) := \# \left\{ \square \in D : \frac{\text{leg}(\square)}{\text{arm}(\square) + 1} < \frac{n}{m} \leq \frac{\text{leg}(\square) + 1}{\text{arm}(\square)} \right\}.$$

Lemma 3.3. *One has*

$$\text{dinv}(D) = \text{dinv}'(D)$$

for any $D \in Y_{N,M}$.

Proof. This result essentially follows from Corollary 1 on page 8 in [23]. For every box $\square \in R_{N,M}$ there is exactly one horizontal step h_\square of the Dyck path D in the same column, and exactly one vertical step v_\square of D in the same row. This provides a bijection between the boxes in $R_{N,M}$ and the couples: one vertical step of D and one horizontal step of D . The reordering of the steps according to ζ gives rise to a bijective map $\phi : R_{N,M} \rightarrow R_{N,M}$, where the box $\phi(\square)$ corresponds to the pair of steps of $\zeta(D)$ obtained from h_\square and v_\square by reordering according to ζ .

With the terminology above, $\text{arm}(\square)$ is the number of boxes strictly between the box $\square \in R_{N,M}$ and the horizontal step h_\square of the boundary of D , whereas its $\text{leg}(\square)$ is the number of boxes strictly between the box \square and v_\square . Observe, v_\square appears in the path before h_\square if and only if $\square \in D$. Consider two cases:

(1) Suppose $\square \in D$, then one has

$$\text{rank}(h_\square) = \text{rank}(\square) - m(\text{leg}(\square) + 1),$$

and

$$\text{rank}(v_\square) = \text{rank}(\square) - n\text{arm}(\square).$$

One gets $\phi(\square) \in \zeta(D)$ if and only if after the reordering the step in $\zeta(D)$ corresponding to v_\square comes before the step corresponding to h_\square . According to the definition of ζ , in this case it is equivalent to $\text{rank}(v_\square) < \text{rank}(h_\square)$, which is in turn equivalent to

$$\frac{\text{leg}(\square) + 1}{\text{arm}(\square)} < \frac{n}{m}.$$

(2) Suppose $\square \in R_{N,M} \setminus D$. Similarly, one gets

$$\text{rank}(h_\square) = \text{rank}(\square) + m\text{leg}(\square),$$

and

$$\text{rank}(v_\square) = \text{rank}(\square) + n(\text{arm}(\square) + 1).$$

In this case,

$$\frac{\text{leg}(\square)}{\text{arm}(\square) + 1} \geq \frac{n}{m}$$

if and only if $\text{rank}(v_\square) \leq \text{rank}(h_\square)$, if and only if $\phi(\square) \in \zeta(D)$.

Since by definition $\text{dinv}(D) = \#R_{N,M}^+ - \#\zeta(D)$, where $R_{N,M}^+$ is the set of boxes in $R_{N,M}$ that fit under the diagonal, one gets

$$\text{dinv}(D) = \#R_{N,M}^+ - \# \left\{ \square \in D : \frac{\text{leg}(\square) + 1}{\text{arm}(\square)} < \frac{n}{m} \right\} - \# \left\{ \square \in R_{N,M} \setminus D : \frac{\text{leg}(\square)}{\text{arm}(\square) + 1} \geq \frac{n}{m} \right\}$$

by the above considerations. Corollary 1 on page 8 in [23] proves

$$\# \left\{ \square \in R_{N,M} \setminus D : \frac{\text{leg}(\square)}{\text{arm}(\square) + 1} \geq \frac{n}{m} \right\} = \# \left\{ \square \in D : \frac{\text{leg}(\square)}{\text{arm}(\square) + 1} \geq \frac{n}{m} \right\} + \#(R_{N,M}^+ \setminus D).$$

Therefore, we conclude that

$$\text{dinv}(D) = \#D - \# \left\{ \square \in D : \frac{\text{leg}(\square) + 1}{\text{arm}(\square)} < \frac{n}{m} \right\} - \# \left\{ \square \in D : \frac{\text{leg}(\square)}{\text{arm}(\square) + 1} \geq \frac{n}{m} \right\}$$

$$= \# \left\{ \square \in D : \frac{\text{leg}(\square)}{\text{arm}(\square) + 1} < \frac{n}{m} \leq \frac{\text{leg}(\square) + 1}{\text{arm}(\square)} \right\} = \text{dinv}'(D).$$

□

The cardinality of the sets $Y_{N,M}$ of Dyck paths get more complicated in the non-relatively prime. In [8] Bizley shows that

$$\exp\left(\sum_{d \geq 1} \frac{1}{d(m+n)} \binom{d(m+n)}{dm} x^d\right)$$

is the generating function whose coefficients give the cardinalities of $Y_{N,M}$, where $(N, M) = (dn, dm)$ for $\gcd(n, m) = 1$.

On the other hand, the set $\mathbf{M}_{N,M}$ of subsets $0 \in \Delta \subset \mathbb{Z}_{\geq 0}$ invariant under addition of M and N is infinite when $\gcd(N, M) = d > 1$. Therefore, there is no hope to construct a bijection between the set of such subsets and $Y_{N,M}$. However, the map $\mathcal{G} : \mathbf{M}_{N,M} \rightarrow Y_{N,M}$ is still well defined. We define an equivalence relation \sim on the set $\mathbf{M}_{N,M}$, so that the relative order of N -generators and M -cogenerators, and hence the value of \mathcal{G} , is the same within each equivalence class. We then construct a bijection \mathcal{D} between the equivalent classes $\mathbf{M}_{N,M}/\sim$ and $Y_{N,M}$, so that one gets $\zeta = \mathcal{G} \circ \mathcal{D}^{-1}$ as in the $d = 1$ case.

Comparing the definition of \mathcal{G} to Lemma 3.3, one can see that dinv is constant on the fibres of \mathcal{G} . Further, each fiber of \mathcal{G} is a union of \sim equivalence classes. In fact, each equivalence class is precisely one fiber by the result of Thomas and Williams[26] showing that ζ is always bijective.

Given an (N, M) -invariant subset $\Delta \in \mathbf{M}_{N,M}$ one can extract d many (n, m) -invariant subsets from it by the following procedure: for each $r \in \{0, 1, \dots, d-1\}$ consider the subset in Δ consisting of all integers congruent to r modulo d , subtract r from all these elements and then divide by d . In other words one has

$$(3) \quad \Delta_r = [(\Delta \cap (d\mathbb{Z} + r)) - r] / d.$$

Note that the subsets Δ_r for $r > 0$ might not be 0-normalized. Note also that Δ can be uniquely reconstructed from $\Delta_0, \dots, \Delta_{d-1}$, so we have a bijection between the set of 0-normalized (N, M) -invariant subsets and (ordered or $\mathbb{Z}/d\mathbb{Z}$ -colored) collections of d many (n, m) -invariant subsets, such that Δ_0 is zero normalized and $\Delta_i \subset \mathbb{Z}_{\geq 0}$ for all i .

Remark 3.4. There is a natural bijection, extending Anderson's construction, between the set of (N, M) -invariant subsets and the set of (N, M) -cores. If λ is an (N, M) -core corresponding to Δ then one can check that the d -quotient of λ ([21]) consists of d diagrams each of which are (n, m) -cores. They naturally correspond to $\Delta_0, \dots, \Delta_{d-1}$.

3.2. Equivalence relation. The idea of the equivalence relation is that one should fix the collection $\Delta_0, \dots, \Delta_{d-1}$ up to shifts, but allow them to slide with respect to each other as long as the N -generators and M -cogenerators of Δ do not "jump" over each other. It is motivated by making the invariant sets in the same fiber of \mathcal{G} equivalent. Recall the map \mathcal{G} only cares about the relative order of the N -generators and M -cogenerators. We will analyze an equivalence class by understanding all the positions the generators and cogenerators can fill while retaining this relative order. This analysis will allow us to construct a representative in the equivalence class of $\Delta \in \mathbf{M}_{N,M}$ which has the minimal number of gaps, and it is on that representative that we can define \mathcal{D} . Later, we will describe the equivalence class of Δ in terms of rank data from the Δ_r along with appropriate gluing data.

Let us first explain the equivalence relation with an example.

Example 3.5. Let $(N, M) = (6, 4)$. The following two elements of $Y_{N,M}$ are equivalent. Let Δ^1 be given by:

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 ...
 × ○ □ ○ × □ ● ○ × × ● □ ● × ● ...

and Δ^2 :

0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 ...
 × ○ □ ○ × ○ ● □ × ○ ● × ● □ ● ...

Here \times 's are 6-generators, \square 's are the 4-cogenerators, \bullet 's are other elements of the subset, and \circ 's are the other elements of the complement. Note that not all 6 generators and 4 cogenerators fit in the pictures. It is more illustrative to split Δ^1 into its even and odd parts:

$r = 0$ 0 2 4 6 8 10 12 14 ...
 × □ × ● × ● ● ...
 $r = 1$ 1 3 5 7 9 11 13 ...
 ○ ○ □ ○ × □ ...

It is more compact to then stack them as

$r = 0$ -4 -2 0 2 4 6 8 10 12 14 16 18 ...
 □ ○ × □ × ● × ● ● ● ● ...
 $r = 1$ -3 -1 1 3 5 7 9 11 13 15 17 19 ...
 ○ ○ ○ ○ □ ○ × □ × ● × ...

reminiscent of a d -abacus.

Finally we just record Δ_0^1 and Δ_1^1 :

-2 -1 0 1 2 3 4 5 6 7 8 9 10 ...
 Δ_0^1 □ ○ × □ × ● × ● ● ● ● ...
 Δ_1^1 ○ ○ ○ ○ □ ○ × □ × ● × ...

To restore Δ^1 one should multiply both Δ_0^1 and Δ_1^1 by two, add one to Δ_1^1 , and merge them together. In other words, $\Delta^1 = 2\Delta_0^1 \cup (1 + 2\Delta_1^1)$. Similarly, for Δ^2 one gets

-2 -1 0 1 2 3 4 5 6 7 8 9 10 ...
 Δ_0^2 □ ○ × □ × ● × ● ● ● ● ● ...
 Δ_1^2 ○ ○ ○ ○ ○ □ ○ × □ × ● × ...

Note that the sequences of N -generators and M -cogenerators are the same for Δ^1 and Δ^2 , even if we take into account the remainder modulo 2. In both cases one gets

(4) □ × □ × □ × × □ × ×

where red is for even generators and cogenerators ($r = 0$), and blue is for odd ($r = 1$). This is the reason $\Delta^1 \sim \Delta^2$. If we only knew the even and odd parts, then, in this example, the odd part can be shifted by 1 with respect to the even part without changing the sequence or parity of generators and cogenerators. Note that one cannot shift further: in Δ^1 one cannot shift the odd part to the left, and in Δ^2 one cannot shift the odd part to the right and still yield an invariant set equivalent to Δ^1 . Also note that while $\Delta_0^1 = \Delta_0^2$, $\Delta_1^1 = -1 + \Delta_1^2$.

Let us give a formal definition of the equivalence classes.

Definition 3.6. The *skeleton* of an (N, M) -invariant subset Δ is the set consisting of its N -generators and M -cogenerators.

Example 3.7. The skeleton of Δ^1 from Example 3.5 above is $\{-4, 0, 2, 4, 5, 8, 9, 11, 13, 17\}$. Note it has $10 = 6 + 4$ elements.

Note that one can uniquely reconstruct an invariant subset Δ from its skeleton. Indeed, the skeleton contains all the N -generators of Δ , and to distinguish the N -generators from the M -cogenerators one should simply choose the biggest elements in each congruence class mod N .

An attentive reader may have noticed that the above definition of the skeleton are not obviously symmetric in N and M . In fact, it is (almost) symmetric by the following lemma.

Lemma 3.8. *Let Δ be some (N, M) -invariant subset. An integer x is either an N -generator or an M -cogenerator of Δ if and only if $x + M$ is an $(N + M)$ -generator of Δ .*

Proof. Indeed, $x + M$ is an $(N + M)$ -generator of Δ if and only if $x + M \in \Delta$ and $x - N \notin \Delta$.

If x is an N -generator then $x \in \Delta$, so $x + M \in \Delta$, but $x - N \notin \Delta$. Hence by the above it is an $(N + M)$ -generator. Assume that x is an M -cogenerator. Then $x \notin \Delta$ but $x + M \in \Delta$. If $x - N \in \Delta$ then $x \in \Delta$, contradiction, therefore $x - N \notin \Delta$, and again $x + M$ is an $(N + M)$ -generator.

Conversely, assume that $x - N = x + M - (N + M) \notin \Delta$ and $x + M \in \Delta$. If $x \in \Delta$ then x is an N -generator, and if $x \notin \Delta$ then x is a M -cogenerator. \square

Remark 3.9. One can also prove Lemma 3.8 using generating functions. Let $f(t) = \sum_{s \in \Delta} t^s$ be the generating function for Δ , then the generating function for the set of N -generators equals $(1 - t^N)f(t)$ while the generating function for the set of M -generators equals $(t^{-M} - 1)f(t)$. Therefore the generating function for the skeleton equals:

$$(1 - t^N)f(t) + (t^{-M} - 1)f(t) = (t^{-M} - t^N)f(t) = t^{-M}(1 - t^{M+N})f(t).$$

Corollary 3.10. *Let $\Delta \in \mathbf{M}_{N,M}$. Then x is in the (N, M) -skeleton of Δ if and only if $x - N + M$ is in the (M, N) -skeleton of Δ .*

Remark 3.11. Indeed, the distribution of generators and cogenerators in the (N, M) and (M, N) skeletons is different (say, there are N generators in the former and M in the latter). One can be obtained from the other by reading the distribution backwards and swapping \times s and \square s. In Example 3.5 the $(6, 4)$ -skeleton of Δ^1 is $\{-4, 2, 5, 11\} \cup \{0, 4, 8, 9, 13, 17\}$ whereas its $(4, 6)$ -skeleton is $\{-6, -2, 2, 3, 7, 11\} \cup \{0, 6, 9, 15\}$, pictured as

$$\square \times \times \square \square \times \square \times \square \times ,$$

which one should compare to (4).

In other words, the (N, M) -skeleton and the (M, N) -skeleton of Δ differ by a shift by $(M - N)$ which does not depend on Δ . In particular, all the constructions are symmetric (up to an overall shift) in M and N . From now on we will continue to use notation as in Definitions 3.6 above and 3.13 below.

Let Δ be an (N, M) -invariant set, let S be its skeleton, and S_0, \dots, S_{d-1} be the parts of the skeleton in different remainders modulo $d = \gcd(M, N)$ (i.e., $S_i = S \cap (d\mathbb{Z} + i)$).

Definition 3.12. A shift $S_0, S_1 + a_1, \dots, S_{d-1} + a_{d-1}$ is called *acceptable* (relative to S) if there exists a continuous path $\bar{\phi} = (\phi_1, \dots, \phi_{d-1}) : [0, 1] \rightarrow \mathbb{R}^{d-1}$ with $\bar{\phi}(0) = (0, \dots, 0)$ and $\bar{\phi}(1) = (a_1, \dots, a_{d-1})$, such that for any $0 \leq t \leq 1$ the sets $S_0, S_1 + \phi_1(t), \dots, S_{d-1} + \phi_{d-1}(t)$ are pairwise disjoint. In other words, we allow S_1, \dots, S_{d-1} to shift by translations as long as the elements of different S_i 's do not intersect. In this case we will call the tuple (a_1, \dots, a_{d-1}) an acceptable shifting of S and an integral shifting when all $a_i \in \mathbb{Z}$.

When S is understood we will lighten notation by not specifying the shifts are relative to S or the shiftings are of S . In fact, in the rest of this section we will assume everything is relative to some fixed skeleton S unless stated otherwise.

Definition 3.13. Let $\Delta, \Delta' \in \mathbf{M}_{N,M}$ with skeletons $S = \bigsqcup S_i, S' = \bigsqcup S'_i$, respectively. We say that Δ is equivalent to Δ' or $\Delta \sim \Delta'$ if there exist a permutation $\sigma \in \mathcal{S}_{d-1}$ such that $S'_0, S'_{\sigma(1)}, \dots, S'_{\sigma(d-1)}$ is an acceptable shift of S_0, S_1, \dots, S_{d-1} .

Equivalence class will always mean \sim equivalence class.

Dealing with equivalence classes is complicated. Instead, we want to choose one representative from each class—a minimal one, as defined below. Our goal is to shift S_1, \dots, S_{d-1} down as much as possible, so that the parts of the skeleton are “stuck” on one another. In fact, this will minimize the size of $\mathbb{Z}_{\geq 0} \setminus \Delta$. Note that not every integral acceptable shift of a skeleton is again a skeleton of some (N, M) -invariant subset, because different parts of the skeleton might end up in the same congruence class modulo d . However, it is convenient to consider the set of all acceptable integral shifts. We will show that there always exists a minimal integral acceptable shift and use it as an intermediate step in the construction of the bijection $\mathcal{D} : \mathbf{M}_{N,M}/\sim \rightarrow Y_{N,M}$.

Lemma 3.14. *The acceptability condition on a shifting a_1, \dots, a_{d-1} is equivalent to satisfying a system of linear inequalities of the form*

$$a_i - a_j < \tilde{b}_{ij},$$

where $\tilde{b}_{ij} \in \mathbb{Z}_{>0} \cup \infty$ for $0 \leq i, j < d$, are fixed and the condition $a_0 = 0$. In particular, the set of acceptable shiftings is convex.

Proof. Fix a skeleton S . Set

$$(5) \quad \tilde{b}_{ij} := \min_{x \in S_i, y \in S_j, y > x} y - x,$$

if $\{x, y : x \in S_i, y \in S_j, y > x\} \neq \emptyset$, and $\tilde{b}_{ij} = \infty$ otherwise. Suppose that (a_1, \dots, a_{d-1}) is an acceptable shifting of S . It follows that for any i, j one has $a_i - a_j < \tilde{b}_{ij}$. Indeed, assume otherwise, i.e., $a_i - a_j \geq \tilde{b}_{ij}$. By definition, there exist $x \in S_i$ and $y \in S_j$ such that $y - x = \tilde{b}_{ij} > 0$. However, after shifting one has

$$(y + a_j) - (x + a_i) = \tilde{b}_{ij} - (a_i - a_j) \leq 0.$$

Therefore, for any continuous path $\bar{\phi} : [0, 1] \rightarrow \mathbb{R}^{d-1}$ such that $\bar{\phi}(0) = (0, \dots, 0)$ and $\bar{\phi}(1) = (a_1, \dots, a_{d-1})$, there exists $t \in (0, 1]$ such that $y + \phi_i(t) = x + \phi_j(t)$. Contradiction.

Conversely, suppose that (a_1, \dots, a_{d-1}) is such that all the inequalities $a_i - a_j < \tilde{b}_{ij}$ are satisfied. Take the path $\bar{\phi}$ to be the line segment connecting $(0, \dots, 0)$ and (a_1, \dots, a_{d-1}) , i.e., take

$$\bar{\phi}(t) := (ta_1, \dots, ta_{d-1}).$$

Then for any $t \in [0, 1]$, any $1 \leq i, j \leq d-1$, $i \neq j$, and any $x \in S_i$ and $y \in S_j$ with $y > x$ one has $x + ta_i \neq y + ta_j$. Indeed,

$$(y + ta_j) - (x + ta_i) = (y - x) - t(a_i - a_j) > \tilde{b}_{ij} - t\tilde{b}_{ij} \geq 0.$$

(The case $y < x$ is covered similarly by the inequality $a_j - a_i < \tilde{b}_{ji}$.) □

We are interested in integral acceptable shiftings, so we can set $b_{ij} = \tilde{b}_{ij} - 1$, and then all integral acceptable shiftings satisfy

$$(6) \quad a_i - a_j \leq b_{ij}.$$

Let $A = A_S \subset \mathbb{R}^{d-1}$ be the set defined by the inequalities (6).

Example 3.15. Let $(n, m) = (3, 2)$ and $d = 4$. Consider the $(12, 8)$ -invariant subset

$$\Delta = \{0, 1, 5, 8, 9, 12, 13, 16, 17, 20, 21, 24, 25, 27, 28, 29, 30\} \cup (\mathbb{Z}_{\geq 32}).$$

As in Example 3.5, it is convenient to reserve a separate row for each remainder modulo $d = 4$:

	-8	-4	0	4	8	12	16	20	24	28	32	36	40	44	...
0	□	○	×	□	×	●	×	●	●	●	●	●	●	●	...
1	□	□	×	×	×	●	●	●	●	●	●	●	●	●	...
2	○	○	○	○	○	○	○	□	□	×	×	×	●	●	...
3	○	○	○	○	○	○	○	□	○	×	□	×	●	×	●

Here each box in the table correspond to the sum of the numbers at the top of the column and at the left and of the row, so to recover the subset Δ one should read the table column by column, top to bottom, then left to right. As usual, \times denotes a 12-generator, and \square an 8-cogenerator, while \bullet are the other elements of Δ and \circ the other elements in the complement. The parts S_0 , S_1 , S_2 , and S_3 of the skeleton of Δ are given by

$$S_0 = \{-8, 0, 4, 8, 16\}, S_1 = \{-7, -3, 1, 5, 9\}, S_2 = \{22, 26, 30, 34, 38\},$$

$$\text{and } S_3 = \{19, 27, 31, 35, 43\}.$$

We can compute the numbers $b_{ij} = \tilde{b}_{ij} - 1$, $i \neq j$ in this example:

$$(b_{ij})_{i,j=0}^3 = \begin{pmatrix} & 0 & 5 & 2 \\ 2 & & 12 & 9 \\ \infty & \infty & & 0 \\ \infty & \infty & 2 & \end{pmatrix}.$$

Therefore, the set $A_S \subset \mathbb{R}^3$ is given by the shiftings (a_1, a_2, a_3) satisfying

$$-2 \leq a_0 - a_1 \leq 0, \quad -\infty \leq a_0 - a_2 \leq 5, \quad -\infty \leq a_0 - a_3 \leq 2,$$

$$-\infty \leq a_1 - a_2 \leq 12, \quad -\infty \leq a_1 - a_3 \leq 9, \quad -2 \leq a_2 - a_3 \leq 0,$$

where $a_0 = 0$. This simplifies to

$$0 \leq a_1 \leq 2, \quad a_3 \geq -2, \quad 0 \leq a_3 - a_2 \leq 2.$$

Lemma 3.16. *Define*

$$(7) \quad m_i = \max_{i=i_1, i_2, \dots, i_k=0} \sum_{\ell=1}^{k-1} (-b_{i_{\ell+1}i_\ell})$$

where the maximum is taken over all sequences $\{i_1, i_2, \dots, i_k\}$ of integers between 0 and $d-1$, such that $i_1 = i$ and $i_k = 0$. Then $(m_1, \dots, m_{d-1}) \in A$ and for any i and any integral acceptable shifting $(a_1, \dots, a_{d-1}) \in A \cap \mathbb{Z}^{d-1}$ we have $a_i \geq m_i$.

Proof. By definition, there exists a sequence of integers $i = i_1, i_2, \dots, i_{k-1}, i_k = 0$ such that $m_i = \sum_{\ell=1}^{k-1} (-b_{i_{\ell+1}i_\ell})$. Then one has

$$a_i = (a_{i_1} - a_{i_2}) + (a_{i_2} - a_{i_3}) + \dots + (a_{i_{k-1}} - a_0) \geq -b_{i_2i_1} - \dots - b_{i_ki_{k-1}} = m_i.$$

Suppose $(m_1, \dots, m_{d-1}) \notin A$. Then $m_i - m_j > b_{ij}$ for some $0 \leq i, j < d$. By definition, there exists a sequence $i = i_1, i_2, \dots, i_k = 0$ such that $m_i = \sum_{\ell=1}^{k-1} (-b_{i_{\ell+1}i_\ell})$. But then

$$m_j < m_i - b_{ij} = -b_{ij} + \sum_{\ell=1}^{k-1} (-b_{i_{\ell+1}i_\ell}),$$

which contradicts the maximality of m_j (consider the sequence $j, i = i_1, i_2, \dots, i_k = 0$). In other words, (m_1, \dots, m_{d-1}) is the minimal integral acceptable shifting. \square

Note all $m_i \leq 0$ as $(0, \dots, 0) \in A$. Set $M_0 = S_0, M_1 = S_1 + m_1, \dots, M_{d-1} = S_{d-1} + m_{d-1}$ to be the shifted parts of the skeleton corresponding to the minimal integral acceptable shift relative to S .

Definition 3.17. Let $f(i)$ be the remainder of any element of M_i modulo d (recall that all elements of M_i have the same remainder). This defines a function $f : \{0, \dots, d-1\} \rightarrow \{0, \dots, d-1\}$.

For every $0 \leq i < d$ set $s_i := \lfloor \frac{M_i}{d} \rfloor = \{\lfloor \frac{x}{d} \rfloor \mid x \in M_i\}$ and let Δ_i be the (n, m) -invariant subset such that s_i is the skeleton of Δ_i . Note Δ_i might not be 0-normalized.

Definition 3.18. Let the directed graph (digraph) $G = G_S$ on the vertex set $\{0, \dots, d-1\}$ be defined in the following way: vertices i and j are connected by an edge $i \rightarrow j$ if $f(i) < f(j)$ and the intersection $s_i \cap s_j$ is not empty.

Lemma 3.19. *The value $f(i)$ equals the length of the longest oriented path from 0 to i in the digraph G .*

Proof. By definition, we have $f(i) < f(j)$ for any edge $i \rightarrow j$. Therefore, it suffices to prove the following two conditions:

- (1) If $f(j) > 0$ then there exists i such that $f(i) = f(j) - 1$ and G contains the edge $i \rightarrow j$.
- (2) $f(i) = 0$ implies $i = 0$.

Both conditions follow immediately from the minimality of the shift. Indeed, if the first property is not satisfied for a vertex i of G then $(m_1, \dots, m_i - 1, \dots, m_{d-1})$ is an acceptable shifting, which contradicts Lemma 3.16.

Suppose now that $f(i) = 0, i \neq 0$. This and the first property imply that $f(j) \neq d-1$ for any $j \in \{0, \dots, d-1\}$. Therefore, again, $(m_1, \dots, m_i - 1, \dots, m_{d-1})$ is an acceptable shifting. Contradiction. \square

Corollary 3.20. *The function f can be recovered from the orientation of the graph G .*

Example 3.21. Continuing Example 3.15, we compute

$$m_1 = -b_{01} = 0, m_2 = -b_{32} - b_{03} = -4, m_3 = -b_{03} = -2,$$

so the minimal integral acceptable shifting of S is $(0, -4, -3)$. The minimal integral acceptable shift is then given by

$$M_0 = S_0 = \{-8, 0, 4, 8, 16\}, M_1 = S_1 + 0 = \{-7, -3, 1, 5, 9\}, \\ M_2 = S_2 - 4 = \{18, 22, 26, 30, 34\}, M_3 = S_3 - 2 = \{17, 25, 29, 33, 41\}.$$

Note that elements of both M_1 and M_3 have remainder 1 modulo $d = 4$. Therefore, this shift does not correspond to any $(12, 8)$ -invariant subset. The skeletons $s_i = \lfloor \frac{M_i}{4} \rfloor$ are given by

$$s_0 = \{-2, 0, 1, 2, 4\}, s_1 = \{-2, -1, 0, 1, 2\}, \\ s_2 = \{4, 5, 6, 7, 8\}, s_3 = \{4, 6, 7, 8, 10\},$$

and we get $f(0) = 0, f(1) = f(3) = 1$, and $f(2) = 2$. Note $M_i = ds_i + f(i)$. See Figure 5 for the graph G . We will also consider the (n, m) -periodic lattice paths corresponding to s_0, s_1, s_2, s_3 (see Figure 6).

Definition 3.22. Let $T_{n,m}^d$ denote the set of acyclically oriented graphs G on d vertices with a unique source v_0 , and vertices labeled by skeletons of (n, m) -invariant subsets, such that

- (1) All labels are non-negatively normalized, and the label of v_0 is zero normalized,
- (2) Two skeletons intersect if and only if the corresponding vertices are connected by an edge.

Elements of $T_{n,m}^d$ are considered up to label preserving isomorphisms.




FIGURE 5. The digraph G with vertices labelled by the skeletons of the $(3, 2)$ -invariant subsets. The function f corresponds to the levels: 0 at the blue vertex, 1 at the green and orange vertices, and 2 at the red vertex.




FIGURE 6. The four $(3, 2)$ -periodic paths corresponding to the skeletons from Figure 5. Note that the elements of the skeletons are exactly the ranks of the boxes above the horizontal steps and to the left of the vertical steps. Equivalently, they are the ranks of the steps of the paths.

Note that the underlying (not oriented) graph is determined by the d -tuple of (n, m) -invariant subsets. We will refer to the orientation of the digraph G as the *gluing data* on the d -tuple of invariant subsets.

The construction above provides a map

$$A : \mathbf{M}_{N,M}/\sim \rightarrow T_{n,m}^d.$$

Moreover, the map is injective by construction, because given a labeled graph $G \in T_{n,m}^d$ one can use Lemma 3.19 to reconstruct the function $f(i)$ and then recover the sets M_0, \dots, M_{d-1} by setting $M_i = ds_i + f(i)$. We need to show that this map is also surjective, i.e., show that for any labeled digraph $G \in T_{n,m}^d$ the corresponding sets M_0, \dots, M_{d-1} form a minimal integral acceptable shift of the skeleton of an (N, M) -invariant subset.

Let $G \in T_{n,m}^d$ be a labeled graph. Let f be the function on the vertex set of G constructed as in Lemma 3.19, i.e., for a vertex v , $f(v)$ equals to the length of the longest oriented path from the source v_0 to v . Since elements of $T_{n,m}^d$ are considered up to label preserving isomorphisms,

one can assume that the vertex set of G is $V_G = \{0, 1, \dots, d-1\}$ and the function f is weakly monotone:

$$i < j \Rightarrow f(i) \leq f(j).$$

In particular, one gets $v_0 = 0$. Note, that two different labeled digraphs on the vertex set $\{0, 1, \dots, d-1\}$ might be related by a label preserving isomorphism, in which case they correspond to the same element of $T_{n,m}^d$ (see Figure 7 for an example).

Let the vertices $0, \dots, d-1$ of G have labels s_0, \dots, s_{d-1} respectively.

Lemma 3.23. *Given $G \in T_{n,m}^d$, there exists an (N, M) -invariant subset $\Delta = \Delta^G$ with skeleton $S = \bigsqcup S_i$, such that*

$$M_0 := ds_0, M_1 := ds_1 + f(1), \dots, M_{d-1} := ds_{d-1} + f(d-1)$$

is the minimal integral acceptable shift of S_0, \dots, S_{d-1} . More over, we can choose Δ by setting

$$S_0 := ds_0, S_1 := ds_1 + 1, \dots, S_{d-1} := ds_{d-1} + (d-1).$$

Proof. By construction, for every i every element of S_i has remainder i modulo d . It follows that $S := \bigsqcup S_i$ is the skeleton of an (N, M) -invariant set Δ . The non-negative normalization of the s_i imply $\Delta \in \mathbf{M}_{N,M}$. It remains to show that M_0, M_1, \dots, M_{d-1} is the minimal integral acceptable shift of S_0, S_1, \dots, S_{d-1} . By construction, for every i we have $M_i = S_i + a_i$ where $a_i := f(i) - i$. Recall that the minimal integral acceptable shifting is given by (7), and by (5) the integers b_{ij} are given by

$$b_{ij} := \left(\min_{x \in S_i, y \in S_j, y > x} y - x \right) - 1.$$

Suppose $i \rightarrow j$ is an edge of G . Then as f is monotone we must have $i < j$. Since $s_i \cap s_j \neq \emptyset$ it follows that there are $x \in S_i$ and $y \in S_j$ such that $\lfloor \frac{x}{d} \rfloor = \lfloor \frac{y}{d} \rfloor$, so we get $b_{ij} = (j - i) - 1$. On the other hand, suppose there is no directed edge $i \rightarrow j$. If $i < j$ we immediately get that $b_{ij} \geq (d + j - i) - 1 \geq j$. Similarly, if $i > j$ (even if $j \rightarrow i$) we get $b_{ij} \geq (j + d - i) - 1 \geq j$.

It follows that to maximize $\sum_{\ell=1}^{k-1} (-b_{i_{\ell+1}i_\ell})$ in (7) one has to consider the longest directed path in G from 0 to i . Such a path has $f(i)$ steps, so we get

$$m_i = f(i) - i = a_i.$$

Therefore, M_0, M_1, \dots, M_{d-1} is the minimal integral acceptable shift of S_0, S_1, \dots, S_{d-1} . \square

Note that the representative constructed in Lemma 3.23 above has the following property: $\lfloor \frac{S_i}{d} \rfloor = \lfloor \frac{M_i}{d} \rfloor$ where, as before, S_0, \dots, S_{d-1} are the parts of the skeleton of the representative Δ of the equivalence class in the corresponding remainders modulo d , and M_0, \dots, M_{d-1} is the minimal integral acceptable shift of S_0, \dots, S_{d-1} . We call such representatives Δ *the minimal representatives*. Note that an equivalence class might contain more than one minimal representative (see Example 3.24). Recall the sets M_0, \dots, M_{d-1} might not correspond to an element of $\mathbf{M}_{N,M}$, but the S_0, \dots, S_{d-1} will.

Example 3.24. Continuing Example 3.21 and using the graph on the left of the Figure 7, one gets

$$\begin{aligned} S_0 &= 4\{-2, 0, 1, 2, 4\} = \{-8, 0, 4, 8, 16\}, \\ S_1 &= 4\{4, 6, 7, 8, 10\} + 1 = \{17, 25, 29, 33, 41\}, \\ S_2 &= 4\{-2, -1, 0, 1, 2\} + 2 = \{-6, -2, 0, 6, 10\}, \\ S_3 &= 4\{4, 5, 6, 7, 8\} + 3 = \{19, 23, 27, 31, 35\}. \end{aligned}$$

Therefore, the $(12, 8)$ -invariant subset Δ^G we constructed is depicted below.




FIGURE 7. Two graphs corresponding to the same point of $T_{3,2}^4$: the isomorphism switching vertices 1 and 2 preserves the labels.

	-8	-4	0	4	8	12	16	20	24	28	32	36	40	44	...
0	□	○	×	□	×	●	×	●	●	●	●	●	●	●	...
1	○	○	○	○	○	○	□	○	×	□	×	●	×	●	...
2	□	□	×	×	×	●	●	●	●	●	●	●	●	●	...
3	○	○	○	○	○	○	□	□	×	×	×	●	●	●	...

Note that if we had used the graph on the right of Figure 7 instead, we would get a different invariant subset in the same equivalence class. Both these subsets are minimal, as they are constructed according to the algorithm in Lemma 3.23. Note that they are both different from the invariant subset we started from in Example 3.15, which was not minimal.

Also note this set has 14 gaps, which is the area of the rational Dyck path constructed in Figure 8.

3.3. From equivalence classes to Dyck paths. The last step is to construct a bijection $B : T_{n,m}^d \rightarrow Y_{N,M}$ so that we can set $\mathcal{D} = B \circ A$. Let $G \in T_{n,m}^d$ be a labeled graph, and let P_0, \dots, P_{d-1} be the (m, n) -periodic lattice paths corresponding to the labels s_0, \dots, s_{d-1} of G .

Lemma 3.25. *Let $0 \leq i < j < d$. Then the paths P_i and P_j intersect if and only if the skeletons s_i and s_j intersect or, equivalently, if and only if the graph G has an edge between the corresponding vertices.*

Proof. Suppose that $x \in s_i \cap s_j$, and let \square be a box in \mathbb{Z}^2 with $\text{rank}(\square) = x$. Then P_i contains either the step v_\square that is to the right (if x is an n -generator) or the step h_\square at its bottom (if x is an m -cogenerator). (Here we extend the notation from Lemma 3.3 to periodic paths.) In both cases, P_i passes through the right-bottom corner of \square . The same is true for P_j . Hence P_i intersects P_j at that corner.

Conversely, if P_i contains a lattice point p let \square be the box with p at its bottom right corner. Then P_i must contain either the step v_\square going up from p , or the the step h_\square going left from p . In both cases, it implies $x = \text{rank}(\square)$ has to be in the skeleton s_i . The same holds for P_j and s_j , so $x \in s_i \cap s_j$. Finally, the graph G has an edge between two vertices if and only if the corresponding skeletons intersect. \square

3.3.1. *Gluing algorithm.* We will glue together paths P_0, \dots, P_{d-1} (more precisely, a union of possibly disconnected intervals of total length $(n+m)$ of these paths) to get an (N, M) -Dyck path D in the following way, which we call our gluing algorithm. We start by taking the interval of P_0 that is an (n, m) -Dyck path (there is a unique way to choose such an interval, up to a periodic shift). At each step we glue in an interval of length $(n+m)$ of one of the periodic paths P_1, \dots, P_{d-1} into our path. This is done using the following procedure.

Let \hat{D} be a (kn, km) -Dyck path and let \hat{P} be an (n, m) -periodic path, such that the intersection $\hat{D} \cap \hat{P}$ is not empty. Let p be the first point of intersection of \hat{D} and \hat{P} relative to \hat{D} (recall that we orient all lattice paths from bottom-right to top-left, i.e., p is the point of intersection, closest to the bottom-right end of \hat{D}). The new $((k+1)n, (k+1)m)$ -Dyck path $\hat{D} \vee \hat{P}$ is the union of three lattice paths:

- (1) First we follow the path \hat{D} from its start up to p ;
- (2) Then we follow \hat{P} for $(n+m)$ steps starting at p ;
- (3) Finally, we follow the remaining part of \hat{D} translated by n up and m to the left, i.e., by $(-m, n)$.

More precisely, let us now also identify a (kn, km) -Dyck path \hat{D} with the function $\hat{D} : [0, k(n+m)] \rightarrow \mathbb{R}^2$, so that \hat{D} is its plot and the function is an isometry to the image. Similarly, a periodic path can be regarded as a function $P : \mathbb{R} \rightarrow \mathbb{R}^2$ satisfying $P(z+m+n) = P(z) + (-m, n)$. Given $r \in \mathbb{Z}$ and a function $I : [r, r+n+m] \rightarrow \mathbb{R}^2$ satisfying $I(r+n+m) = I(r) + (-m, n)$, we extend I periodically to $P(I) : \mathbb{R} \rightarrow \mathbb{R}^2$ by $P(I)(z+k(n+m)) = I(z) + k(-m, n)$, for $r \leq z \leq r+n+m$. Note if I was an interval of a (kn, km) -Dyck path then $P(I)$ is normalized so $0 \leq z \leq n+m$ implies $P(z)$ is between the lines $y=0$ and $y=n$. However, it is convenient to treat periodic paths so their parameterization might not be normalized in this way (i.e., so that we need not have $a=b$ below, or so that we can consider an interval of it as a function with domain $[0, n+m]$). Using this function notation, we may describe

$$\hat{D} \vee \hat{P}(z) = \begin{cases} \hat{D}(z) & 0 \leq z \leq a \\ \hat{P}(z+b-a) & a \leq z \leq a+n+m \\ \hat{D}(z-(m+n)) + (-m, n) & a+n+m \leq z \leq (k+1)(n+m) \end{cases}$$

where $a, b \in \mathbb{R}$ are the parameters such that $\hat{D}(a) = \hat{P}(b) = p$ and $p \in \mathbb{R}^2$ is the first point of \hat{D} that is also in \hat{P} .

We apply the above procedure $d-1$ times in the following order. Let $k_j = \#\{i \mid f(i) \leq j\}$. We start by setting D_0 to be the interval of P_0 such that D_0 is an (n, m) -Dyck path. Take all paths P_i , such that $f(i) = 1$. Note that all such paths intersect D_0 and do not intersect each other. Therefore, we can glue them in using the above procedure, and the order in which we do it does not matter, i.e. the path created is independent of gluing order for these i . Let D_1 be the resulting rational Dyck path. Note it is a (k_1n, k_1m) -Dyck path.

At the $(j+1)$ th step we start with the (k_jn, k_jm) -Dyck path D_j obtained from D_0 by gluing in intervals of all paths P_i such that $f(i) \leq j$, one level of G at a time, and we glue in intervals of all P_i 's, such that $f(i) = j+1$. Again, all such paths intersect at least one of the intervals we glued in on the previous step, and they do not intersect each other. We proceed in the same manner until we glued in intervals of all periodic paths P_1, \dots, P_{d-1} . (See Figure 8 for an example.)

We need to show that this process is invertible. Consider an (N, M) -Dyck path D .

First we will define *removal* of intervals. Let D be a (kn, km) -Dyck path. We call I a *balanced* interval of D if it consists of $n+m$ consecutive steps of D of which n are vertical and m are horizontal. Using our function notation, this means I is the restriction of D to




FIGURE 8. We apply the gluing algorithm to the periodic paths from Figure 6, using the graph from Figure 5. On each step we indicate the gluing point: the first intersection of the Dyck path built so far with the next periodic path.

$[r, r + n + m]$, with $r \in \mathbb{Z}$ and $D(r) = D(r + n + m) + (m, -n)$. We will say D' is obtained from D by removing a balanced interval I if it corresponds to the function given by

$$D'(z) = \begin{cases} D(z) & 0 \leq z \leq r \\ D(z + (m + n)) + (m, -n) & r \leq z \leq (k - 1)(n + m) \end{cases}.$$

Definition 3.26. An interval I of D is called *good* if it is of length $n + m$, balanced, and its (n, m) -periodic extension does not intersect the part of D before I .

Our definition of good is motivated by the need to invert the gluing process. Thus good intervals must have the following properties. Suppose we remove a good interval I from a (kn, km) -Dyck path D which yields D' . If we now glue the periodic extension of I into D' by taking the first (lowest rightmost) point of D' that intersects the periodic path, this should yield the original D , i.e., $D = D' \vee P(I)$. That good removal inverts the gluing algorithm is based on the following Lemmas.

Lemma 3.27. *There always exists at least one good interval in a (kn, km) -Dyck path D .*

Proof. The proof goes in two steps. First, one can show that there always exists at least one balanced interval of D of length $(n + m)$. This is equivalent to showing that the intersection of D with D translated by n down and m to the right, i.e. D intersect $D + (m, -n)$, is non-empty. Since D is a Dyck path, it stays weakly below the diagonal connecting its start with its end. It follows that $D + (m, -n)$ intersects the vertical line through the start of D (weakly) below the start of D , and ends (weakly) above D . Therefore, it has to intersect D .

Second, the balanced interval closest to the bottom-right end of D is always good. \square

Lemma 3.28. *Periodic extensions of the good intervals of D do not intersect each other.*

Proof. Indeed, otherwise the one further away from the bottom-right end of D is not good. \square

Lemma 3.29. *If I and J are good intervals of D , and D' is obtained from D by removing I , then the image of J in D' is still a good interval of D' .*

Proof. Indeed, if the periodic extension of J intersects the part of D' before it, then it also intersects the part of D before it. \square

Lemma 3.30. *Let $G \in T_{n,m}^d$ and suppose that the (N, M) -Dyck path D was obtained from the periodic paths P_1, \dots, P_{d-1} according to the gluing algorithm given by G . The periodic extension $P(I)$ of good intervals I of D agree with the P_i for i the sinks of G .*

Proof. In this proof we will use “before” and “after” according to the steps of the gluing algorithm (temporally), and switch to “higher” or “lower” to refer to locations of steps or lattice points of paths.

Let I be a good interval of D . Then, according to Lemma 3.29 either it was glued in on the last step of our algorithm, or it was already a good interval before the last step. In the latter case, the same holds for the second to the last step and so on. Therefore, I was glued in at some point.

Suppose that I corresponds to the vertex i of G , and suppose that there is an edge $i \rightarrow j$ in G . Then the interval corresponding to j was glued in after I was. Therefore, either it was glued in in the middle of I , or lower than I , in which case after that gluing I is not a good interval any more, because periodic paths P_i and P_j intersect. Contradiction. \square

Theorem 3.31. *The map $B : T_{n,m}^d \rightarrow Y_{N,M}$ is a bijection.*

Proof. We will induct on d . The case $d = 1$ corresponds to the relatively prime case.

There is some flexibility in the gluing algorithm: if two periodic paths do not intersect each other, then as noted previously it does not matter in which order we glue in intervals of these paths. In particular, we can change the order so that the paths corresponding to the sink vertices i of the graph G are glued in at the last step of the gluing algorithm. Suppose that G has k sink vertices $\{i_1, \dots, i_k\}$ and let $G' \in T_{n,m}^{d-k}$ be the labeled graph obtained from G by removing the sink vertices. Let also $(s_{i_1}, \dots, s_{i_k})$ be the skeletons corresponding to the sink vertices of G . Note that the following two properties are satisfied:

- (1) The skeletons s_{i_1}, \dots, s_{i_k} are pairwise disjoint,
- (2) Every skeleton corresponding to a sink vertex of the graph G' intersects at least one of the skeletons $s_i, i \notin \{i_1, \dots, i_k\}$.

Indeed, if the first property is not satisfied then the corresponding two vertices of G are connected by an edge and cannot both be sinks. If the second property is not satisfied, then the corresponding vertex is also a sink of G , which is a contradiction. Conversely, for any $0 < k < d$, any labeled graph $G' \in T_{n,m}^{d-k}$ and a collection of skeletons s_{i_1}, \dots, s_{i_k} satisfying the above two conditions there is a unique graph $G \in T_{n,m}^d$, such that s_{i_1}, \dots, s_{i_k} are the labels of the sink vertices of G , and G' is obtained from G by removing the sink vertices.

Exactly the same situation happens on the Dyck path side. Let $D \in Y_{N,M}$ be an (N, M) -Dyck path. Suppose it has k good intervals, and let $D' \in Y_{(N-kn, M-km)}$ be the Dyck path obtained from D by removing the good intervals. Let P_1, \dots, P_k be the periodic extensions of the good intervals of D . The following two properties are satisfied:

- (1) The periodic paths P_1, \dots, P_k are pairwise disjoint,
- (2) The periodic extension of any good interval of the Dyck path D' intersects at least one of the paths P_1, \dots, P_k .

Indeed, if the first property is not satisfied then the corresponding two intervals of D cannot both be good. If the second property is not satisfied, then the corresponding good interval of D' is also a good interval of D , which is a contradiction. Conversely, for any $0 < k < d$, any Dyck path $D' \in Y_{(N-kn, M-km)}$ and a collection of (n, m) -periodic paths P_1, \dots, P_k satisfying the above two conditions there is a unique Dyck path $D \in Y_{N,M}$, such that P_1, \dots, P_k are the periodic extensions of the good intervals of D , and D' is obtained from D by removing all good intervals.

Using Lemma 3.25 and induction on d we now can build the inverse map $B^{-1} : Y_{N,M} \rightarrow T_{n,m}^d$. See Figure 9 for an example. \square

We conclude that since the maps $A : M_{N,M}/\sim \rightarrow T_{n,m}^d$ and $B : T_{n,m}^d \rightarrow Y_{N,M}$ are bijections, the map $D = B \circ A : M_{N,M}/\sim \rightarrow Y_{N,M}$ is a bijection as well.




FIGURE 9. On the first step we remove two good intervals and record the corresponding skeletons: $\{-2, -1, 0, 1, 2\}$ and $\{4, 5, 6, 7, 8\}$. On the second step there is only one good interval, with the corresponding skeleton $\{4, 6, 7, 8, 10\}$. Finally, on the last step we are left with a $(3, 2)$ -Dyck path, which is its own good interval. The corresponding skeleton is $\{-2, 0, 1, 2, 4\}$. On the right we have the reconstructed labeled graph. Note the sinks are the first intervals removed. Note, that it is isomorphic to the graph in Figure 5.

Theorem 3.32. *The sweep map $\zeta : Y_{N,M} \rightarrow Y_{N,M}$ factorizes according to Figure 2:*

$$\zeta = \mathcal{G} \circ \mathcal{D}^{-1}$$

for all positive N, M .

Proof. Let $D \in Y_{N,M}$ be a Dyck path, and let $\Delta \in \mathbf{M}_{N,M}$ be a minimal representative of the equivalence class $\mathcal{D}^{-1}(D) \subset \mathbf{M}_{N,M}$. Similar to the $d = 1$ case, the steps of the path D correspond to the elements of the skeleton S of Δ . However, the correspondence is a bit trickier. Since the Δ is a minimal representative, the rank of a step of D equals $\lfloor \frac{x}{d} \rfloor$, where x is the corresponding element of the skeleton of Δ . However, according to the gluing algorithm, if $x, y \in S$ are two elements of the skeleton of Δ , such that $x < y$ and $\lfloor \frac{x}{d} \rfloor = \lfloor \frac{y}{d} \rfloor$, then the step corresponding to x is glued in lower than the step corresponding to y . In turn, that implies that the step in D corresponding to x appears higher than the step corresponding to y , which matches with the ‘‘tie breaking’’ adjustment in the construction of the sweep map in the non relatively prime case (see Example 3.2). \square

The following proposition gives a simple interpretation of the area statistic for rational Dyck paths in terms of (N, M) -invariant subsets.

Proposition 3.33. *Let $\Delta \in \mathbf{M}_{N,M}$. Then*

$$\text{area}(\mathcal{D}(\Delta)) = \min_{\Delta' \sim \Delta} \text{gap}(\Delta'),$$

where, as above, $\text{gap}(\Delta') = |\mathbb{Z}_{\geq 0} \setminus \Delta'|$.

Proof. As before, let $\Delta_0, \dots, \Delta_{d-1}$ be the d -tuple of (n, m) -invariant subsets defined by

$$\Delta_r = [(\Delta \cap (d\mathbb{Z} + r)) - r] / d.$$

Let also $m_r = \min \Delta_r$. Then

$$\text{gap}(\Delta) = \sum_r \text{gap}(\Delta_r) = \sum_r [m_r + \text{gap}(\Delta_r - m_r)].$$

Note that $(\Delta_r - m_r) \in \mathbf{M}_{n,m}$ and, in particular, $\text{gap}(\Delta_r - m_r) = \text{area}[\mathcal{D}(\Delta_r - m_r)]$, because in the relatively prime case the area statistic counts the boxes whose ranks are exactly the gaps, and each gap is counted exactly once. It follows that to obtain the Δ' with minimal gap over the equivalence class of Δ one should consider an invariant subset with the minimal $\sum_r m_r$, which is equivalent to considering one of the minimal representatives. Therefore, it is sufficient to prove that if Δ is a minimal representative, then $\text{area}(\mathcal{D}(\Delta)) = \text{gap}(\Delta)$. Let us compute the area between $\mathcal{D}(\Delta)$ and the diagonal in the (N, M) rectangle $R_{N,M}$. It consists of the areas between the Dyck paths (possibly shifted and disconnected) for the (n, m) -invariant subsets Δ_r and their local diagonals, and the parallelograms between these small diagonals and the big diagonal. Since Δ is a minimal representative, the smallest rank of a box that fits under the local diagonal corresponding to Δ_r is m_r . Therefore, such a parallelogram contains the boxes with all possible ranks between 0 and m_r , each rank appearing exactly once. Therefore $\text{area}(\mathcal{D}(\Delta)) = \text{gap}(\Delta)$. \square

3.4. Example: k -Catalan arrangement. Let us describe the equivalence relation in the case $M = kN$, $k \in \mathbb{Z}_{>0}$. In this case, $d = N$, $n = 1$ and $m = k$. A module is (N, M) -invariant if and only if it is N -invariant, and therefore has the form

$$\Delta(k_0, \dots, k_{N-1}) = \{k_i + Nj : i = 0, \dots, N-1, j \geq 0\},$$

where k_i is an arbitrary integer with remainder i modulo N . To be 0-normalized we further require $k_0 = 0$ and $k_i \geq 0$. Now $\Delta_i = \{k_i + Nj : j \geq 0\}$, so the skeleton S_i has a unique N -generator k_i and has k M -cogenerators $k_i - N, \dots, k_i - kN$. Therefore the i -th skeleton of Δ has the form

$$S_i = \{k_i, k_i - N, \dots, k_i - kN\}.$$

Recall that the k -Catalan arrangement in \mathbb{R}^N is defined by the equations $x_i - x_j = s$ where $i < j$ and s runs through $\{-k, \dots, k\}$, and the k -Shi arrangement is defined by the same equations with $s \in \{-(k-1), \dots, k\}$. We will call the connected components of their complements k -Catalan and k -Shi regions, respectively. Clearly, in the dominant cone where $x_1 < \dots < x_N$ the arrangements agree and it is known that the number of dominant k -Shi regions is equal to the n th Fuss-Catalan number

$$c_N(k) := \frac{((k+1)N)!}{(kN+1)!N!},$$

which is also equal to the number of Dyck paths in the $N \times kN$ rectangle. Since the k -Catalan arrangement is S_N -invariant, the total number of k -Catalan regions equals $N!c_N(k)$.

If we pass to $V = \mathbb{R}^N / \text{span}(1, 1, \dots, 1)$, the connected components of the complement of the hyperplane arrangement $\{x_i - x_j = s | s \in \mathbb{Z}\}$ are called alcoves. Observe that while these regions are unbounded in \mathbb{R}^N , in V they are bounded and each alcove has centroid of the form $(\frac{a_1}{N}, \dots, \frac{a_N}{N})$ with $a_i \in \mathbb{Z}$ and $\{a_i \bmod N\}$ distinct. We will always take our representative of $(\frac{a_1}{N}, \dots, \frac{a_N}{N}) + \text{span}(1, 1, \dots, 1)$ to be such that $\min\{a_i\} = 0$. This is compatible with taking Δ to be 0-normalized. (Note that in the literature one often normalizes to be “balanced,” so that $\sum a_i = 0$ and $\sum k_i = \binom{N+1}{2}$.)

Note further that to each $\Delta(k_0, \dots, k_{N-1})$ we can associate the alcove that has centroid $p_\Delta = (\frac{k_0}{N}, \dots, \frac{k_{N-1}}{N})$. Since Δ is independent of the order of the k_i , we could just as easily associate to it the alcove in the dominant cone $x_1 < \dots < x_N$ that has centroid $p_\Delta^+ = (\frac{k_{\sigma(0)}}{N}, \dots, \frac{k_{\sigma(N-1)}}{N})$, where $\sigma \in \text{Perm}\{0, 1, \dots, N-1\}$ is chosen so that $k_{\sigma(i)} < k_{\sigma(i+1)}$.

Proposition 3.34. *The set of integral acceptable shifts for Δ for which the shifting is by distinct integers mod N is in bijection with the set of alcoves that are in the same k -Catalan region as p_Δ .*

Proof. Indeed, the shifting (a_0, \dots, a_{N-1}) is acceptable if and only if for all i and j the order of (colored) points in the sets $S_i \cup S_j$ and $S_i + a_i \cup S_j + a_j$ is the same. As the order of x and y (i.e., whether $x < y$) is determined by whether $x - y < 0$ and since our shifting is by integers distinct mod N , it suffices to consider the signs of such differences. More algebraically, for all pairs $(x = k_i - tN, y = k_j - t'N) \in S_i \times S_j$ the sign of $x - y$ and the sign of $(x + a_i) - (y + a_j)$ is the same. The sign the $x - y$ is determined by the sign of $k_i/N - k_j/N - (t - t')$, so we require that the points $(k_0/N, \dots, k_{N-1}/N)$ and $((k_0 + a_0)/N, \dots, (k_{N-1} - a_{N-1})/N)$ are on the same side of the hyperplane $x_i - x_j = t - t'$. (This is still true if we look at points whose coordinates are sorted to lie in the dominant cone, i.e., the alcoves in the same region as p_Δ^+ .) It remains to notice that possible values of $t - t'$ run between $-k$ and k . \square

We conclude that the set of equivalence classes of (N, kN) -invariant subsets is in bijection with the set of dominant k -Catalan (or, equivalently, dominant k -Shi) regions, and both sets have $c_N(k)$ elements. Therefore our main construction provides yet another bijection [FV10] between dominant k -Shi regions and Dyck paths in $N \times kN$ rectangle.

4. RELATION TO KNOT INVARIANTS

In this section we prove Theorem 1.3. We will use the following result:

Theorem 4.1. ([10, Theorem 1.9]) *The Poincaré series of the $(a = 0)$ part of the Khovanov-Rozansky homology of the (n, n) torus link equals*

$$F_n(q, t) = \sum_{a=(a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n} q^{\sum a_i} t^{d(a)},$$

where $d(a) = |\{i < j : a_i = a_j \text{ or } a_j = a_i + 1\}|$.

Lemma 4.2. *One has*

$$(1 - q)F_n(q, t) = \sum_{a=(a_1, \dots, a_{n-1}, 0) \in \mathbb{Z}_{\geq 0}^n} q^{\sum a_i} t^{d(a)}.$$

Proof. Let us define the cyclic shift operator $\pi : (a_1, \dots, a_n) \mapsto (a_n - 1, a_1, \dots, a_{n-1})$, which is well defined if $a_n > 0$. By applying π repeatedly, we can transform a given tuple a to a tuple with $a_n = 0$. Clearly, $\sum \pi(a) = \sum a_i - 1$ and one can check that $d(\pi(a)) = d(a)$. Therefore:

$$F_n(q, t) = \sum_{k \geq 0} \sum_{a=(a_1, \dots, a_{n-1}, 0) \in \mathbb{Z}_{\geq 0}^n} q^{k + \sum a_i} t^{d(a)} = \frac{1}{1 - q} \sum_{a=(a_1, \dots, a_{n-1}, 0) \in \mathbb{Z}_{\geq 0}^n} q^{\sum a_i} t^{d(a)}.$$

\square

We are ready to prove Theorem 1.3.

Proof of Theorem 1.3. By Lemma 4.2 and Theorem 4.1 we need to prove the identity

$$C_{n,n}(q, t) = \sum_{a=(a_1, \dots, a_{n-1}, 0) \in \mathbb{Z}_{\geq 0}^n} q^{\sum a_i} t^{d(a)}.$$

A subset $\Delta \subset \mathbb{Z}_{\geq 0}$ is (n, n) -invariant if and only if it is n -invariant. In remainder i it has an n -generator $x_i = i + na_i$ and an n -cogenerator $y_i = i + na_i - n$, for some $a_i \geq 0$. It is 0-normalized if $a_n = 0$. It is easy to check that $\text{gap}(\Delta) = \sum a_i$. Now

$$\text{dinv}(\Delta) = \binom{n}{2} - \#\{i, j : y_j > x_i\} = \binom{n}{2} - \#\{i < j : a_j > a_i + 1\} - \#\{i > j : a_j > a_i\} = d(a).$$

\square

Example 4.3. Let us compute $C_{2,2}(q, t)$. All 2-invariant 0-normalized subsets have the form

$$\Delta_k = \{0, 2, \dots, 2k, 2k + 1, 2k + 2, \dots\}.$$

Clearly, $\text{gap}(\Delta_k) = k$ and

$$\text{dinv}(\Delta_k) = \begin{cases} 1, & \text{if } k = 0, \\ 0, & \text{if } k > 0. \end{cases}$$

Therefore

$$C_{2,2}(q, t) = \sum_{k=0}^{\infty} q^{\text{gap}(\Delta_k)} t^{\text{dinv}(\Delta_k)} = t + \frac{q}{1-q} = \frac{q+t-qt}{1-q}.$$

Note that $c_{2,2}(q, t) = q + t$.

REFERENCES

- [1] M. Aganagic, S. Shukirov. Refined Chern-Simons theory and knot homology. String-Math 2011, 3–31, Proc. Sympos. Pure Math., 85, Amer. Math. Soc., Providence, RI, 2012.
- [2] J. Anderson. Partitions which are simultaneously t_1 - and t_2 -core. Discrete Math. **248** (1-3) (2002). 237–243.
- [3] D. Armstrong. Hyperplane arrangements and diagonal harmonics. 23rd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2011), 39–50, Discrete Math. Theor. Comput. Sci. Proc., AO, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2011.
- [4] D. Armstrong, N. Loehr, G. Warrington, Sweep maps: A continuous family of sorting algorithms. Adv. Math. **284** (2015), 159–185.
- [5] A. Beauville, Counting rational curves on K3-surfaces. Duke Math. J. **97** (1999), 99–108.
- [6] F. Bergeron. Open Questions for operators related to Rectangular Catalan Combinatorics, arXiv:1603.04476. (To appear: Journal of Combinatorics.)
- [7] F. Bergeron, A. Garsia, E. Leven, G. Xin, Compositional (km, kn) -Shuffle Conjectures. Int. Math. Res. Not. IMRN 2016, no. 14, 4229–4270.
- [8] M. T. L. Bizley, Derivation of a new formula for the number of minimal lattice paths from $(0, 0)$ to (km, kn) having just t contacts with the line $my = nx$ and having no points above this line; and a proof of Grossman’s formula for the number of paths which may touch but do not rise above this line, J. Inst. Actuar., **80** (1954), 55–62.
- [9] C. Ceballos, T. Denton, C. R. H. Hanusa, Combinatorics of the zeta map on rational Dyck paths. J. Combin. Theory Ser. A **141** (2016), 33–77.
- [10] B. Elias, M. Hogancamp. On the computation of torus link homology. arXiv:1603.00407
- [11] S. Fishel, M. Vazirani. A bijection between dominant Shi regions and core partitions. European J. Combin. **31** (2010), no. 8, 2087–2101.
- [12] E. Gorsky, M. Mazin. Compactified Jacobians and q, t -Catalan Numbers, I. Journal of Combinatorial Theory, Series A **120** (2013), pp. 49–63.
- [13] E. Gorsky, M. Mazin. Compactified Jacobians and q, t -Catalan numbers, II. J. Algebraic Combin. **39** (2014), no. 1, pp. 153–186.
- [14] E. Gorsky, M. Mazin, M. Vazirani, Affine Permutations And Rational Parking Functions. Trans. Amer. Math. Soc. **368** (2016), no. 12, 8403–8445.
- [15] E. Gorsky, M. Mazin, M. Vazirani, Rational Dyck Paths in the Non Relatively Prime Case. 28th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2016), 587–598, Discrete Math. Theor. Comput. Sci. Proc., AO, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2016.
- [16] E. Gorsky, A. Negut, Refined knot invariants and Hilbert schemes, Journal de mathématiques pures et appliquées 104 (2015), pp. 403–435.
- [17] E. Gorsky, A. Negut, J. Rasmussen. Flag Hilbert schemes, colored projectors and Khovanov-Rozansky homology. arXiv:1608.07308
- [18] J. Haglund, The q, t -Catalan Numbers and the Space of Diagonal Harmonics: With an Appendix on the Combinatorics of Macdonald Polynomials. AMS University lecture series, 2008.
- [19] M. Khovanov, L. Rozansky. Matrix factorizations and link homology. II. Geom. Topol. **12** (2008), no. 3, 1387–1425.
- [20] N. Loehr. Conjectured Statistics for the Higher q, t -Catalan Sequences. The Electronic Journal of Combinatorics **12** (2005), Research Paper 9, 54 pp.
- [21] D. E. Littlewood. Modular representations of symmetric groups. Proc. Roy. Soc. London. Ser. A., **209** (1951), 333–353.

- [22] G. Lusztig, J. M. Smelt. Fixed point varieties on the space of lattices. Bull. London Math. Soc. **23** (1991), no. 3, 213–218.
- [23] M. Mazin, A bijective proof of Loehr-Warrington’s formulas for statistics $\text{ctot}_{\frac{q}{p}}$ and $\text{mid}_{\frac{q}{p}}$, Ann. Comb. 18 (2014), no. 4, pp. 709-722.
- [24] A. Mellit. Toric braids and (m, n) -parking functions. arXiv:1604.07456
- [25] J. Piontowski, Topology of the compactified Jacobians of singular curves. Math. Z. **255** (2007), no. 1, 195-226.
- [26] H. Thomas, N. Williams, Sweeping up zeta, arXiv:1512.01483.
- [27] A. T. Wilson. Torus link homology and the nabla operator. arXiv:1606.00764
- [28] G. Xin, An efficient search algorithm for inverting the sweep map on rational Dyck paths, arXiv:1505.00823.

UNIVERSITY OF CALIFORNIA AT DAVIS, DAVIS, CALIFORNIA, US

INTERNATIONAL LABORATORY OF REPRESENTATION THEORY AND MATHEMATICAL PHYSICS, NRU-HSE, MOSCOW, RUSSIA

E-mail address: egorskiy@math.ucdavis.edu

KANSAS STATE UNIVERSITY, MANHATTAN, KANSAS, US

E-mail address: mmazin@math.ksu.edu

UNIVERSITY OF CALIFORNIA AT DAVIS, DAVIS, CALIFORNIA, US

E-mail address: vazirani@math.ucdavis.edu