

UC Riverside

UC Riverside Previously Published Works

Title

The Strongest Isolable Acid

Permalink

<https://escholarship.org/uc/item/3g85n9gp>

Journal

Angewandte Chemie International Edition, 43(40)

ISSN

1433-7851

Authors

Juhasz, Mark
Hoffmann, Stephan
Stoyanov, Evgenii
[et al.](#)

Publication Date

2004-10-11

DOI

10.1002/anie.200460005

Peer reviewed

Brønsted Acids**The Strongest Isolable Acid****

*Mark Juhasz, Stephan Hoffmann, Evgenii Stoyanov,
Kee-Chan Kim, and Christopher A. Reed**

Acids based on carborane anions as conjugate bases (Figure 1) are a new class of Brønsted (protic) acids, notable for their “strong yet gentle” qualities.^[1] For example, whereas conventional strong acids (e.g. H₂SO₄) and superacids (e.g. HFSO₃/SbF₅) decompose fullerenes even at low temperatures, the carborane acid H(CHB₁₁H₅Cl₆) cleanly protonates C₆₀ at room temperature to give the isolable salt [HC₆₀][CHB₁₁H₅Cl₆]. We now show that carborane acids are the

[*] M. Juhasz, S. Hoffmann, Dr. K.-C. Kim, Prof. Dr. C. A. Reed
Department of Chemistry
University of California
Riverside, CA 92521-0403 (USA)
Fax: (+1) 909-787-2027
E-mail: chris.reed@ucr.edu

Dr. E. Stoyanov
Boreskov Institute of Catalysis
Prospekt Lavrentieva 5, Novosibirsk 630090 (Russia)

[**] This work was supported by NSF grant CHE-0095206 and NIH grant GM 23851.



Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.





Figure 1. The $[\text{CHB}_{11}\text{R}_5\text{X}_6]^-$ carborane ions.

strongest isolable (Lewis-free) Brønsted acids presently known.


Two reactivity considerations have led us to believe that carborane acids must have uncommon strength. First is their stabilization of the benzenium ion in salts such as $[\text{C}_6\text{H}_7]$ $[\text{CHB}_{11}\text{Me}_5\text{Br}_6]$. The strongest known neat oxyacid, HFSO_3 , (whose Hammett acidity function $H_0 = -15.1$) is not strong enough to fully protonate benzene,^[2] yet carborane acids do so readily at the one-equivalent level in dilute solution.^[3] Second, we have recently shown that carborane anions stabilize *tert*-butyl cation in dichloromethane or liquid SO_2 solution, as well as in isolable crystalline salts.^[4] This means that carborane acids are strong enough to shift the equilibrium of Equation (1) completely to the right hand side. If this were not the case, deprotonation of *tert*-butyl cation to give *iso*-butene and subsequent decomposition by cationic oligomerization would occur rapidly.



Since mixed Lewis/Brønsted superacid media such as “Magic Acid” ($\text{HFSO}_3/\text{SbF}_5$) having H_0 somewhere in the range -17 to -27 are used to stabilize *tert*-butyl cation,^[5] it is apparent that carborane acids have intrinsic Brønsted acidities comparable to those only found previously in Lewis/Brønsted acid mixtures.

The acidities of carborane acids cannot be measured in the conventional manner of an H_0 Hammett acidity function because carborane acids are solids not liquids. They sublime under vacuum, or melt at atmospheric pressure, at temperatures well above 150°C . Gas-phase acidities have been calculated for carborane acids, which rank them the strongest of any known isolable acid,^[6] but it is presently not possible to translate gas-phase ΔG data into measures of condensed-phase acidity.

The need to compare the strengths of solid acids with liquid acids therefore led us to obtain a measure of acidity in dilute solution, conditions where most acid catalysis is carried out. The chosen solvent is liquid SO_2 because of its low basicity, relatively high dielectric constant for solvation of ions, and ease of drying. The chosen method is adapted from that developed by Fărcașiu and Ghenciu.^[7] It is based on the ^{13}C NMR spectroscopy chemical shift difference ($\Delta\delta$) between the C_α and C_β carbon atoms of mesityl oxide which increases with increasing protonation as Equation (2) is shifted to the right hand side.



^{13}C NMR spectroscopy data for 0.15 M solutions of various acids and 0.10 M mesityl oxide were obtained at room temperature and are given in Table 1. The choice of oxyacids

Table 1: Acid-strength ranking from protonation of mesityl oxide in liquid SO_2 .

Acid	$\Delta\delta$ (^{13}C) [ppm]	H_0
H($\text{CHB}_{11}\text{Cl}_{11}$)	84.0 ± 0.1	[a]
H($\text{CHB}_{11}\text{H}_5\text{Cl}_6$)	83.8 ± 0.1	[a]
H($\text{CHB}_{11}\text{H}_5\text{Br}_6$)	83.8 ± 0.1	[a]
H($\text{CHB}_{11}\text{H}_5\text{I}_6$)	83.3 ± 0.1	[a]
FSO_3H	73.8 ± 0.5	-15.1
$\text{CF}_3\text{SO}_3\text{H}$	72.9 ± 0.4	-14.1
$\text{HN}(\text{SO}_2\text{CF}_3)_2$	72.0 ± 0.4	[a]
H_2SO_4	64.3 ± 3.1 ^[b]	-12.1
Unprotonated mesityl oxide	32.4 ± 0.1	

[a] H_0 values unavailable because acids are solids, not liquids. [b] Incomplete miscibility of H_2SO_4 in liquid SO_2 leads to higher error limits and possible underestimate of $\Delta\delta$.

for comparison to carborane acids is limited by their availability in pure form and the need for miscibility or dissolution in liquid SO_2 . Nevertheless, it is immediately evident from their high chemical shift values that, as a class, carborane acids are stronger than conventional oxyacids. They outrank fluorosulfuric acid, the strongest known pure Brønsted acid on the H_0 acidity scale (-15.1), as well as triflic acid ($H_0 = -14.1$). A strict correlation between $\Delta\delta$ values in liquid SO_2 and the H_0 values of neat acids is not expected because acidity is a solvation-dependent phenomenon that changes as the medium changes. Nevertheless, for the limited data set available, the ranking of acid strength by $\Delta\delta$ values follows the H_0 order (Table 1).

It is also evident from the data of Table 1 that whereas oxyacids only partially protonate mesityl oxide, carborane acids are strong enough to move Equation (2) completely to the right hand side, that is, their acidities are leveled. The $\Delta\delta$ value maximizes at $\delta = 84.0$ ppm. Though close to the limits of discrimination, the hexa-iodo carborane acid, $[\text{H}(\text{CHB}_{11}\text{H}_5\text{I}_6)]$ with a value of $\delta = 83.3$ ppm, appears to be perceptibly weaker than its chloro and bromo counterparts.

To discriminate between the acidities of the different carborane acids, we have developed a new, qualitative ranking of acidity based on the $\nu\text{N}-\text{H}$ frequencies of the ammonium salts of their conjugate base anions. The ranking is based on the influence of the anion A^- on the $\text{N}-\text{H}$ bond in a contact ion pair (Figure 2).



Figure 2. Ammonium salt contact ion pair.

The weaker the interaction of the base A^- , the stronger the N–H bonding, and the higher the $\nu N-H$ frequency.^[8] Since $\nu N-H$ frequencies ($> 3000\text{ cm}^{-1}$) are energetically far removed from $\nu H-A$ frequencies ($< 300\text{ cm}^{-1}$), vibrational coupling or mass differences in A^- will have an insignificant effect on $\nu N-H$, that is, $\nu N-H$ frequencies should be a good measure of N–H bond strength. Since hydrogen bonding is a predominantly electrostatic phenomenon,^[9] N–H bond strength should correlate strongly with A^- basicity, that is, with HA acidity. $\nu N-H$ frequencies will decrease with increasing A^- basicity, correlating with HA acidity. In a constant solvation environment within a series of isostructural anions, the correlation should be quite direct. Although the direct correlation of acidity with IR stretching frequency has been questioned for solid surfaces,^[10–12] acidity rankings based on IR data have been shown to have excellent validity within a number of classes of discrete compounds, particularly when structurally related.^[13–17]

Tri-*n*-octylammonium salts of carborane anions were prepared by silver-salt metathesis reactions and characterized as detailed in the Supporting Information. Octyl groups in the cation were chosen to provide good solubility of [*n*-Oct₃NH]⁺ [carborane][−] ion pairs in CCl₄ solution. To avoid possible effects from aggregation, $\nu N-H$ frequencies were measured in dilute solution where there was no concentration dependence (Table 2). The strong dependence of $\nu N-H$ on the

Table 2: $\nu N-H$ frequencies of tri-*n*-octylammonium salts in CCl₄.

Anion	$\nu N-H$ [$\pm 1\text{ cm}^{-1}$]	$\Delta\nu$ [cm^{-1}]
[CHB ₁₁ Cl ₁₁] ^{−[a]}	3163	0
[CHB ₁₁ Me ₁₁] [−]	3156	7
[CHB ₁₁ H ₅ Cl ₆] ^{−[a]}	3148	15
[Co ^{III} (C ₂ B ₉ H ₈ Cl ₃) ₂] [−]	3145	18
[CHB ₁₁ Me ₅ Cl ₆] [−]	3143	20
[CHB ₁₁ H ₁₁] [−]	3129	34
[CHB ₁₁ H ₅ Br ₆] ^{−[a]}	3125	38
[CHB ₁₁ Me ₅ Br ₆] [−]	3120	43
[CHB ₁₁ H ₅ I ₆] ^{−[a]}	3097	66
[CHB ₁₁ Me ₅ I ₆] [−]	3091	72

[a] Denotes presently known isolable acids.

nature of the anion indicates that contact ion pairs are formed. In the more polar solvent 1,2-dichloroethane, the rank order is the same but some salts showed a second band of invariable frequency ($3179 \pm 1\text{ cm}^{-1}$) which is assigned to solvent-separated ion pairs. For the corresponding deuterated [Oct₃N–D]⁺ salts the $\nu N-D/\nu N-H$ ratio was 1.339 ± 0.004 , indicating the absence of anharmonicity effects or Fermi resonance.

The $\nu N-H$ ranking brings out a number of features: a) For each of the pairs of hexahalo carboranes [CHB₁₁H₅X₆][−] (X = Cl, Br, I) and their pentamethylated counterparts [CHB₁₁Me₅X₆][−],^[18] the difference in $\nu N-H$ values is 5 or 6 cm^{-1} . This consistency gives confidence in the scale, suggesting that the CCl₄ solvation environment remains constant over the series and that mass differences and steric effects have negligible influence. It is evident that small differences in anion basicity can be detected quite accurately

and the scale should be useful for a much wider range of weakly coordinating anions. b) The ranking of the hexahalo carborane acids, $H(CHB_{11}H_5Cl_6) > H(CHB_{11}H_5Br_6) > H(CHB_{11}H_5I_6)$, makes sense on the basis of electronegativity and polarizability considerations for the halogen substituents. It is consistent with conceptually related deductions drawn recently from $\nu C-H$ frequencies in benzenium ion salts having the same series of carborane counterions.^[3] c) The top ranking (i.e. least basic) carborane anion measured to date is the undecachlorinated ion, [CHB₁₁Cl₁₁][−],^[19] which leads to the prediction that its conjugate acid will be the strongest. As detailed in the Supporting Information, H(CHB₁₁Cl₁₁) can be prepared in a similar manner to the hexahalo carborane acids as a sublimable, analytically-pure, colorless solid by the reaction of Et₃Si(CHB₁₁Cl₁₁) with anhydrous HCl. Thus, H(CHB₁₁Cl₁₁) can lay claim to be the strongest isolable acid presently known.

Carborane acids are strong acids because their anions are exceptionally weak bases. This arises from the large size of the anions, the delocalized nature of the negative charge in the [CB₁₁][−] cluster, and the shield of halide substituents. However, there is an important additional necessity for preparing such a strong acid, namely, the anion must be exceptionally inert from a chemical point of view. The inertness of carborane anions arises from σ aromaticity within the [CB₁₁][−] icosahedral cage, a notable feature of boron cluster chemistry. The [CB₁₁][−] framework resists chemical disruption to a truly exceptional degree. There are anions that probably have lower basicity than carboranes, for example, organofluoro anions, such as [B(C₆F₅)₄][−],^[20] but their conjugate acids cannot be prepared because the anion decomposes at high acidity.^[21] This situation is mirrored in that traditional fluoroacids, commonly written as HBF₄, HPF₆, HSbF₆ etc., are also nonexistent compounds. They are all unstable with respect to formation of HF and the corresponding Lewis acid. The superacid commonly written “HSbF₆” is actually a mixture of [H(HF)_{*x*}][SbF₆·*n*SbF₅] ions in HF/SbF₅ solvent and cannot be isolated as a pure Brønsted acid. Thus, present limits to acid strength are seen to be more dependent on considerations of anion stability than basicity, although low basicity is clearly required.

An important and potentially very useful feature of carborane acids is that their high protic acidity is expressed in the absence of an added Lewis acid. The presence of Lewis acids in traditional superacids (e.g. HFSO₃/SbF₅) can interfere with protonation chemistry and limit their usefulness in two identifiable ways. As alluded to earlier for C₆₀, Lewis acids (particularly SbF₅) can supply oxidizing equivalents and halide nucleophiles that together, conspire to decompose many protonated unsaturated molecules. A more subtle effect arises from the likely formation of Lewis acid/base complexes when substrates are investigated in mixed Brønsted/Lewis superacids. If a substrate forms an adduct with a Lewis acid, its basicity will be lowered and it will become much more difficult to protonate. This phenomenon can be called “basicity suppression”. It suggests that the basicities of weakly basic molecules may have been significantly underestimated, and makes the pursuit of even stronger pure Brønsted acids an important endeavor.

Carborane acids may lead not only to the isolation and stabilization of protonated species hitherto unattainable, but also to a reassessment of the intrinsic basicity of weakly basic molecules. Experiments to test this hypothesis are underway.

Received: March 16, 2004

Revised: June 30, 2004

Keywords: acids · basicity suppression · Brønsted acids · carboranes · IR spectroscopy

-
- [1] C. A. Reed, K.-C. Kim, R. D. Bolskar, L. Mueller, *Science* **2000**, *289*, 101.
- [2] D. M. Brouwer, E. L. Mackor, C. MacLean in *Carbonium Ions, Vol. 1* (Eds.: G. A. Olah, P. v. R. Schleyer), Wiley-Interscience, New York, **1970**, p. 837.
- [3] C. A. Reed, K.-C. Kim, E. S. Stoyanov, D. Stasko, F. S. Tham, L. J. Mueller, P. D. W. Boyd, *J. Am. Chem. Soc.* **2003**, *125*, 1796.
- [4] T. Kato, C. A. Reed, *Angew. Chem.* **2004**, *116*, 2967–2971; *Angew. Chem. Int. Ed.* **2004**, *43*, 2907–2911.
- [5] G. A. Olah, G. K. S. Prakash, J. Sommer, *Superacids*, Wiley, New York, **1985**.
- [6] I. A. Koppel, P. Burk, I. Koppel, I. Leito, T. Sonoda, M. Mishima, *J. Am. Chem. Soc.* **2000**, *122*, 5114.
- [7] D. Fărcașiu, A. Ghenciu, *J. Am. Chem. Soc.* **1993**, *115*, 10901.
- [8] C. Fischer, H. Wagner, V. V. Bagreev, E. S. Stoyanov, *J. Inorg. Nucl. Chem.* **1977**, *39*, 513.
- [9] M. M. Davis, *Acid–Base Behavior in Aprotic Organic Solvents*, Department of Commerce, National Bureau of Standards, Washington, DC, **1968**, 113.
- [10] W. E. Farneth, R. J. Gorte, *Chem. Rev.* **1995**, *95*, 615.
- [11] R. A. v. Santen, G. J. Kramer, *Chem. Rev.* **1995**, *95*, 637.
- [12] D. Fărcașiu, D. Hâncu, *Catal. Lett.* **1998**, *53*, 3.
- [13] A. V. Iogansen in *The Hydrogen Bond* (Ed.: N. D. Sokolov), Nauka, Moscow, **1981**, pp. 112–155.
- [14] M. D. Joesten, R. S. Drago, *J. Am. Chem. Soc.* **1962**, *84*, 3817.
- [15] E. M. Arnett, L. Joris, E. Mitchell, T. S. S. R. Murty, T. M. Gorrie, P. v. R. Schleyer, *J. Am. Chem. Soc.* **1970**, *92*, 2365.
- [16] R. S. Drago, N. O'Bryan, G. C. Vogel, *J. Am. Chem. Soc.* **1970**, *92*, 3924.
- [17] A. D. Sherry, K. F. Purcell, *J. Am. Chem. Soc.* **1972**, *94*, 1853.
- [18] D. Stasko, C. A. Reed, *J. Am. Chem. Soc.* **2002**, *124*, 1148.
- [19] Z. Xie, C.-W. Tsang, E. T.-P. Sze, Q. Yang, D. T. W. Chan, T. C. W. Mak, *Inorg. Chem.* **1998**, *37*, 6444.
- [20] The N–H frequency of the trioctylammonium salt of the $[\text{B}(\text{C}_6\text{F}_5)_4]^-$ ion (3233 cm^{-1}) is even higher than that of the carborane salts.
- [21] C. A. Reed, N. L. P. Fackler, K.-C. Kim, D. Stasko, D. R. Evans, P. D. W. Boyd, C. E. F. Rickard, *J. Am. Chem. Soc.* **1999**, *121*, 6314.
-