UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

Analysis of microtubule dynamics using growth curve models

Permalink

<https://escholarship.org/uc/item/3g86q60c>

Journal

Journal of Applied Statistics, 36(6)

ISSN

0266-4763

Authors

Jammalamadaka, SR
Siddiqi, MA
Ghosh, K

Publication Date

2009-06-01

DOI

10.1080/02664760802479131

Peer reviewed

This article was downloaded by: [Jammalamadaka, S. Rao]

On: 31 May 2009

Access details: Access Details: [subscription number 911180831]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK

Journal of Applied Statistics

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title-content=t713428038>

Analysis of microtubule dynamics using growth curve models

S. Rao Jammalamadaka ^a; Md. Aleemuddin Siddiqi ^b; Kaushik Ghosh ^c

^a Department of Statistics and Applied Probability, University of California Santa Barbara, Santa Barbara, CA, USA ^b Symbiance, Inc., Princeton Junction, NJ, USA ^c Department of Mathematical Sciences, University of Nevada Las Vegas, Las Vegas, NV, USA

First Published: June 2009

To cite this Article Jammalamadaka, S. Rao, Siddiqi, Md. Aleemuddin and Ghosh, Kaushik (2009) 'Analysis of microtubule dynamics using growth curve models', *Journal of Applied Statistics*, 36:6, 621 — 631

To link to this Article: DOI: 10.1080/02664760802479131

URL: <http://dx.doi.org/10.1080/02664760802479131>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Analysis of microtubule dynamics using growth curve models

S. Rao Jammalamadaka^{a*}, Md. Aleemuddin Siddiqi^b and Kaushik Ghosh^c

^aDepartment of Statistics and Applied Probability, University of California Santa Barbara, Santa Barbara, CA, USA; ^bSymbiance, Inc., Princeton Junction, NJ, USA; ^cDepartment of Mathematical Sciences, University of Nevada Las Vegas, Las Vegas, NV, USA

Microtubules are part of the structural network within a cell's cytoplasm, providing structural support as well as taking part in many of the cellular processes. A large body of data provide evidence that dynamics of microtubules in a cell is responsible for the performance of many critical cellular functions such as cell division. In this article, we study the effect of four different isoforms of a protein tau on microtubule dynamics using growth curve models. The results show that a linear growth curve model is sufficient to explain the data. Moreover, we find that a mutated version of a 3-repeat tau protein has a similar effect as a 4-repeat tau protein on microtubule dynamics. The latter findings conform with the biological understanding of the effect of the protein tau on microtubule dynamics.

Keywords: growth curves; microtubule dynamics; polynomial regression; splines; MANOVA; Wilks' Lambda

1. Introduction

Microtubules are sub-cellular structures present in most plant and animal cells and form what is known as the cytoskeleton (see [1,2]). They are cytoplasmic fibers with a tubular form, composed of several strands linked by lateral interactions. A combination of biochemical, structural and morphological analyses during the past three decades have shown that the cytoplasm of a cell is not a disorganized mass of jelly but a highly structured cell compartment formed of a cytoskeleton, one of whose principal components are microtubules.

Studies have revealed that microtubules are not only well organized, but also highly dynamic (see [6,9]). By dynamic, we mean that they undergo changes in length over time. A microtubule which grows and/or shortens very rapidly is said to be highly dynamic. The dynamics of the microtubules are induced by their polymerization and depolymerization. This, in turn, affects all the microtubule-induced cellular functions such as cell shape, intracellular transport, chromosome segregation and flagella-mediated cell motility (see [3]).

The dynamics of microtubules in living cells has been of great importance in studying certain diseases and their treatments. For instance, promising anti-cancer drugs suppress cell division by

*Corresponding author. Email: kaushik.ghosh@unlv.edu

stabilizing microtubule activity. It is thus of interest to study the dynamics of these microtubules under different treatments and conditions.

In this article, we study the effect of various isoforms of a protein called tau on microtubule dynamics using statistical growth curve analysis. Using rigorous statistical analyses and tests, we are able to confirm or refute various biological hypotheses about the microtubules under differing treatment conditions. We observe in our analysis that a mutated 4-repeat (4R) tau, i.e. where G272 has been mutated to V, has a similar effect on microtubule dynamics as the 3-repeat (3R) tau. This supports the structure function model of the biochemical interaction of the tau protein on the microtubules.

2. Microtubule dynamics

Microtubule images in living cells are acquired by fluorescence microscopy. The images are in the form of a video which is essentially a stack of still-images taken over successive time intervals. Figure 1 gives one such still-image of various microtubules. In each stack, a certain number of microtubules are selected and tracked, i.e. their individual coordinate locations are noted for each image in the stack by the experimenter.

By observing a stack/video of the microtubules, the experimenter chooses some microtubules whose tips are easy to track. These microtubule tips are then tracked in each frame. For each microtubule, one fixes the initial position $\mathbf{p}_0 = (a_0, b_0)$ (i.e. where the microtubule is attached and seems to grow out of) as the initial observation or the origin. Then for each succeeding frame i , the tip's pixel location $\mathbf{p}_i = (a_i, b_i)$ is noted. The Euclidean distance between the points $\mathbf{p}_0$ and $\mathbf{p}_i$, i.e., $\|\mathbf{p}_i - \mathbf{p}_0\|$, represents the approximate length of the microtubule in frame number i .

Figure 1. Microtubule tips are tracked manually or automatically. Picture courtesy Professors Stuart Feinstein, Leslie Wilson and Dr Janis Bunker of the Department of Molecular Cellular and Developmental Biology, UCSB.

Figure 2. The path traversed by a typical microtubule tip. The first point (origin) is the location of one end of the microtubule. Data courtesy Professors Stuart Feinstein, Leslie Wilson and Dr Janis Bunker of the Department of Molecular Cellular and Developmental Biology, UCSB.

Figure 2 gives an example of the path taken by the sequence $\mathbf{p}_0, \mathbf{p}_1, \mathbf{p}_2, \dots$ in a typical microtubule. In practice, not all microtubules can be tracked for the same number of frames, as some microtubules are lost or become denatured before others. Hence, for each tip, the tracking sequence can be of different length. As a preliminary analysis, for our illustrative example, we consider only those microtubules that could be tracked for the full set of frames (the number of which is denoted by n). For each microtubule, we compute the length sequence $\{l_i\}$ given by

$$l_i = \{(a_i - a_0)^2 + (b_i - b_0)^2\}^{1/2}; \quad i = 1, 2, \dots, n. \quad (1)$$

For most in-vivo microtubules, the tip locations $\mathbf{p}_1, \mathbf{p}_2, \dots$ are reasonably along a straight line, as can be seen in Figure 2. In view of this, we define our ‘growth variable’ $x(i)$, $i = 1, 2, \dots, n$ as the cumulative change in length up till frame i . Hence,

$$x(i) = \sum_{j=1}^i |l_j - l_{j-1}|, \quad (2)$$

where we take $l_0 = l_1$, making $x(1) = 0$. This growth variable is our measure of the dynamics of each microtubule and will be used for our growth curve analyses in this paper. Our growth curve for a particular microtubule is thus defined by $x(i)$, $i = 1, 2, 3, \dots, n$.

Note that an alternative measure of dynamics can be given by

$$y(i) = \sum_{j=1}^i \|\mathbf{p}_j - \mathbf{p}_{j-1}\|.$$

Such a measure would perhaps be more appropriate when the path is not as linear as in Figure 2 and is hence not used in this research.

3. Growth curve modeling

Suppose that there are r different treatments (or groups) and x is the real-valued growth variable measured at p different time points: $t_1, t_2, \dots, t_p$ for n_j individuals chosen at random from the j th treatment. Such correlated data over time, labeled ‘growth curves’, can be analysed using the methods introduced in Potthoff and Roy [12]. It has been discussed and analysed later by several authors, significant among them being Khatri [7], Grizzle and Allen [4], Rao [13], Pan and Fang [11] and Sengupta and Jammalamadaka [14]. Using specialized packages, one can also do exact tests for growth curve models, details of which can be found in Weerahandi [15].

We consider the following polynomial regression model of degree $(q - 1)$ for the mean growth of x for the j th treatment on the time variable t ,

$$E(x_j(t)) = \psi_{j0}t^0 + \psi_{j1}t^1 + \dots + \psi_{j,q-1}t^{q-1} \quad (3)$$

for $t = t_1, \dots, t_p$; $j = 1, \dots, r$ with $p > q - 1$. Let

$$\boldsymbol{\psi}'_j = (\psi_{j0} \psi_{j1} \dots \psi_{j,q-1}) \quad (4)$$

denote the vector of the growth curve coefficients for the j th treatment. The observations $x_j(t_1), \dots, x_j(t_p)$, being on the same specimen, are correlated, and we denote their variance-covariance matrix by $\boldsymbol{\Sigma}$. We assume $\boldsymbol{\Sigma}$ to be the same for all the r groups. Assuming n_j observations in the j th group, let $\mathbf{X}_j$ denote the $p \times n_j$ matrix of the observations for the j th group. Since our growth variable sequence obtained for each of the microtubules starts with $x_1 = 0$, this first value is redundant. It will make the design matrix $\mathbf{X}$ [Equation (5)] singular and is thus omitted. Let

$$\mathbf{X} = [\mathbf{X}_1 \mathbf{X}_2 \dots \mathbf{X}_r]. \quad (5)$$

$\mathbf{X}$ is a $p \times N$ matrix of all the observations, where

$$N = n_1 + n_2 + \dots + n_r. \quad (6)$$

Therefore from Equation (3) we get

$$\begin{aligned} E(\mathbf{X}_j) &= [\mathbf{B} \boldsymbol{\psi}_j \mathbf{B} \boldsymbol{\psi}_j \dots \mathbf{B} \boldsymbol{\psi}_j] \\ &= \mathbf{B} \boldsymbol{\psi}_j \mathbf{E}_{1n_j} \quad (j = 1, 2, \dots, r), \end{aligned} \quad (7)$$

where

$$\mathbf{B} = \begin{bmatrix} t_1^0 & t_1^1 & \dots & t_1^{q-1} \\ t_2^0 & t_2^1 & \dots & t_2^{q-1} \\ \dots & \dots & \dots & \dots \\ t_p^0 & t_p^1 & \dots & t_p^{q-1} \end{bmatrix} \quad (8)$$

and $\mathbf{E}_{ab}$ denotes a matrix of order $a \times b$ with all elements equal to 1. $\mathbf{B}_{p \times q}$ is called the ‘‘design matrix’’. From Equation (7), we get

$$\begin{aligned} E(\mathbf{X}) &= [\mathbf{B} \boldsymbol{\psi}_1 \mathbf{E}_{1n_1} | \mathbf{B} \boldsymbol{\psi}_2 \mathbf{E}_{1n_2} | \dots | \mathbf{B} \boldsymbol{\psi}_r \mathbf{E}_{1n_r}], \\ &= \mathbf{B} \boldsymbol{\Psi} \mathbf{A}, \end{aligned} \quad (9)$$

where

$$\boldsymbol{\Psi} = [\boldsymbol{\psi}_1 \dots \boldsymbol{\psi}_r] \quad (10)$$

is the $q \times r$ matrix of the growth curve coefficients, and

$$\mathbf{A} = \text{diag}[\mathbf{E}_{1n_1}, \mathbf{E}_{1n_2}, \dots, \mathbf{E}_{1n_r}] \quad (11)$$

is a block diagonal matrix with $\mathbf{E}_{1n_j}$ ($j = 1, 2, \dots, r$) along the diagonal blocks and zeros elsewhere. $\mathbf{A}$ is of order $r \times N$.

Let $\text{Vec}\mathbf{X}$ be defined as the column vector obtained by stacking the columns of $\mathbf{X}$ one below the other. Denoting $\text{Var}(\text{Vec}\mathbf{X})$ by $\text{Var}(\mathbf{X})$, we find that

$$\text{Var}(\mathbf{X}) = \mathbf{I}_N \otimes \Sigma, \quad (12)$$

where $\otimes$ denotes the Kronecker product of two matrices. Equations (9) and (12) define what is called the growth curve model and considerable detail on this topic can be found in Kshirsagar and Smith [8] and Pan and Fang [11].

4. Data description

Tau is a protein which plays a major role in microtubule dynamics and stability. Biologists have a biochemical structural model for the mechanisms by which the various isoforms of tau attach to the microtubules and affect their dynamics. Under these models, the 4R units of tau are the main units by which it attaches to the microtubule (Figure 3). Similarly 3R units of tau play the main role in attachment to the microtubule. 3R tau is the dominant tau isoform found in human fetuses whereas 4R tau appears mostly in adults. 4RGV is a mutant of 4R in the second repeat unit, i.e. G272, has been mutated to V (denoted by G272V). This mutation is known to cause fronto-temporal dementia with Parkinsonism-like symptoms (FTDP-17) in humans and to affect the ability of 4R-tau to regulate microtubule dynamics. Under the biochemical models, 4RGV is expected to behave like 3R tau in binding to the microtubules. If true, the microtubules treated with 3R should have similar dynamics as those treated with 4RGV.

Data on microtubules were collected by Professors Stuart Feinstein, Leslie Wilson and Dr Janis Bunker of the Department of Molecular Cellular and Developmental Biology, UCSB. In this paper we analyse the microtubule dynamics for four different classes; the control (or un-injected untreated microtubules) and three treatments, which we describe briefly. In our modest study, we have four treatment groups (including the control) as follows:

- (1) Control: These are for the normal cells with no treatments. There are 27 observations in this class.
- (2) 3R: These are for cells injected with 3R tau. There are 22 observations in this class.
- (3) 4R: These are for cells injected with 4R tau. There are 20 observations in this class.
- (4) 4RGV: These are for cells injected with the G272V mutant of 4R tau. There are 16 observations in this class.

Hence, there are a total of $N = 85$ observations in all the categories combined. Figure 4 shows the plots for the growth variable x for these different treatment groups.

Figure 3. 4R and 3R tau schematic.

Figure 4. Growth variables for the different treatment groups. Blue: control, red: 3R, green: 4R, black: 4RGV. Data courtesy Professors Stuart Feinstein, Leslie Wilson and Dr Janis Bunker of the Department of Molecular Cellular and Developmental Biology, UCSB. Available in color online.

5. Growth curve analysis

To fit the growth curve model and to test various hypotheses, we perform the following computations, which are described more fully in [8]. For our analysis, we chose Σ to have the Rao's simple covariance structure:

$$\Sigma = B\Gamma B' + Q\Theta Q',$$

where Γ is a $q \times q$ positive definite matrix, Θ a $(p - q) \times (p - q)$ positive definite matrix and Q a $p \times (p - q)$ matrix of rank $p - q$ with $B'Q = \mathbf{0}$. This ensures that the maximum-likelihood estimates and the generalized least-squares estimates coincide. Other general structures of the covariance matrix Σ are possible, but were not used, since they require iteratively reweighted least squares procedures, which need large amounts of data. Under this assumption, the maximum likelihood estimator of the coefficient matrix ψ is given by (see [11])

$$\hat{\psi} = (B'S^{-1}B)^{-1}(B'S^{-1}X)A'(AA')^{-1}, \quad (13)$$

where

$$S = X[I - A'(AA')^{-1}A]X'.$$

5.1 Model selection

The first step would be to determine the degree $q - 1$ of the polynomial in the growth curve model that adequately fits the data. To do this, we test the hypothesis

$$H_0^*: \text{the degree } (q - 1) \text{ of the growth curves is adequate.} \quad (14)$$

To do this, we first obtain a matrix B_2 of order $p \times (p - q)$ such that

$$B_2'B = 0, \quad (15)$$

where B is as in Equation (8). This can be done by choosing $(p - q)$ linearly independent columns of $[I_p - B(B'B)^{-1}B']$.

Next we compute

$$\mathbf{S} = \mathbf{X}(\mathbf{I} - \mathbf{A}'(\mathbf{A}\mathbf{A}')^{-1}\mathbf{A})\mathbf{X}', \quad (16)$$

where $\mathbf{A}$ is defined in Equation (11).

For each hypothesis testing problem, we need the following degrees of freedom (d.f.),

- d_m order of the error or hypothesis matrix,
- d_E d.f. associated with the error matrix, and
- d_H d.f. associated with the hypothesis matrix.

For this purpose, we construct the MANOVA table given in Table 1.

We construct the matrix of observations $\mathbf{X}_{24 \times 85}$ and obtain the design matrix $\mathbf{B}_{24 \times q}$ from Equation (8). To perform the test we construct the Wilks' Λ statistic defined by

$$\Lambda_0 = \frac{|\mathbf{E}_0|}{|\mathbf{E}_0 + \mathbf{H}_0|}. \quad (17)$$

To test the null hypothesis we use the fact (see, e.g. [13, Section 8c.5]) that the null distribution of

$$F = \frac{1 - \Lambda_0^{1/s}}{\Lambda_0^{1/s}} \times \frac{ms - 2\lambda}{d_H d_m} \quad (18)$$

is approximately an F_{v_1, v_2} where

$$\begin{aligned} v_1 &= d_m d_H, \\ v_2 &= ms - 2\lambda \end{aligned}$$

and

$$\begin{aligned} m &= N - \frac{d_m + d_H + 1}{2}, \\ s &= \left(\frac{(d_m d_H)^2 - 4}{d_m^2 + d_H^2 - 5} \right)^{1/2}, \\ \lambda &= \frac{(d_m d_H - 2)}{4}. \end{aligned}$$

In our case, $p = 24$ and $N = n_1 + n_2 + n_3 + n_4 = 27 + 22 + 20 + 16 = 85$.

The results of the tests for $q = 1$ and $q = 2$ are summarized in Table 2.

Based on the results, we see that a constant model (no change in time) is rejected but a linear fit for the growth curve provides an adequate fit, and the model is well specified. Under the model that the mean growth function is linear, we show the plots for the mean curves in Figure 5. Figure 5 also shows how these linear fitted means for each class compare with the (non-parametric) observed

Table 1. MANOVA for testing model adequacy.

Source	Degrees of freedom	Dispersion, order ($p - q = 22$)
H_0^*	$r = 4$	$\mathbf{H}_0 = \mathbf{B}_2' \mathbf{X} [\mathbf{A}' (\mathbf{A} \mathbf{A}')^{-1} \mathbf{A}] \mathbf{X}' \mathbf{B}_2$
Error	$N - r = 81$	$\mathbf{E}_0 = \mathbf{B}_2' \mathbf{S} \mathbf{B}_2$
Total	$N = 85$	$\mathbf{H}_0 + \mathbf{E}_0 = \mathbf{B}_2' \mathbf{X} \mathbf{X}' \mathbf{B}_2$

Table 2. Summary of model adequacy tests for various growth curve models.

q	Λ_0	F	p -value
1	0.0291	3.7062	4.4409E-16
2	0.3228	0.9018	0.7099

$q - 1$ is the degree of polynomial, Λ_0 is the Wilks' Lambda statistic and F is the corresponding F approximation.

Figure 5. The observed means (solid lines) and the fitted means (dotted lines) using a linear (first-order polynomial) growth curve.

means, computed as the mean at each time point for all the functions in that class. Figure 6 gives the error band (formed from the maximum and minimum observed value at each timepoint) and the fitted curve for the control group. Based on the picture, the model fits the observed data quite well.

Remark Most curve-fitting techniques, such as the polynomials, do not provide for local control of shape. Consequently, local change (for example change in the observed function at one time point) in a polynomial fit affects the entire fitted curve. The B -spline curve avoids this problem by using a special set of blending functions that has only local influence and depends on only a few neighboring control points.

The *spline regression* model for x on the time variable t is given by

$$E(x_j(t)) = \psi_{j0}s_0(t) + \psi_{j1}s_1(t) + \cdots + \psi_{jq-1}s_{q-1}(t) \tag{19}$$

Figure 6. The error band from the observed data (o) and linear fit (*) for the control group.

for $t = t_1, t_2, \dots, t_p$; $p > q - 1$; $j = 1, 2, \dots, r$. Here $s_0(\cdot), \dots, s_{q-1}(\cdot)$ are the basis functions. Then, the growth curve coefficients for the j th treatment becomes

$$\psi'_j = (\psi_{j0}\psi_{j1}\cdots\psi_{jq-1}). \quad (20)$$

See, for example, James and Hastie [5], Mortenson [10] for further details. Such B-splines can also be used to fit the data. Most of the analysis, as well as the testing is very similar to what has been described so far, except that the design matrix $\mathbf{B}$ given in Equation (8) is to be replaced by

$$\mathbf{B} = \begin{bmatrix} s_0(t_1) & s_1(t_1) & \cdots & s_{q-1}(t_1) \\ s_0(t_2) & s_1(t_2) & \cdots & s_{q-1}(t_2) \\ \vdots & \vdots & \ddots & \vdots \\ s_0(t_p) & s_1(t_p) & \cdots & s_{q-1}(t_p) \end{bmatrix}. \quad (21)$$

Such spline bases are likely to be useful when dealing with non-steady-state data. For the steady-state data that we have dealt with in this analysis, linear fits seem to do perfectly fine as expected.

5.2 Comparison of treatments

Next we test if the four groups, namely the control and the three treatments are significantly different. Hence we test the hypothesis

$$\begin{aligned} H_1^*: \psi_1 &= \psi_2 = \cdots = \psi_4 \\ &: \mathbf{L}\psi\mathbf{M} = 0, \end{aligned} \quad (22)$$

where,

$$\mathbf{L} = \mathbf{I}_2 \quad \text{and} \quad \mathbf{M} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & -1 & -1 \end{bmatrix}.$$

Table 3. MANOVA table for Testing H_1^* .

Source	Degrees of freedom	Dispersion, order ($p - q = 22$)
H_1^*	$m = 3$	$\mathbf{H}_1 = (\mathbf{L}\hat{\boldsymbol{\psi}}\mathbf{M})(\mathbf{M}'\mathbf{R}_{11}\mathbf{M})^{-1}(\mathbf{L}\hat{\boldsymbol{\psi}}\mathbf{M})'$
Error	$N - r - (p - q) = 59$	$\mathbf{E}_1 = \mathbf{L}(\mathbf{B}'\mathbf{S}^{-1}\mathbf{B})^{-1}\mathbf{L}'$
Total	$N - r - (p - q) + m = 62$	$\mathbf{H}_1 + \mathbf{E}_1$

Table 4. p -values of various pairwise comparisons.

Pair	Wilks' Λ	p -value
*3R vs control	0.6839	0.0185
*4R vs control	0.4112	1.3836E-4
*4RGV vs control	0.4842	0.0043
3R vs 4R	0.8655	0.3150
3R vs 4RGV	0.9004	0.5616
4R vs 4RGV	0.8399	0.4179

* denotes a significant pair.

To test this hypothesis we construct the MANOVA table given in Table 3. $\mathbf{R}_{11}$ in Table 3 is defined by,

$$\mathbf{R}_{11} = (\mathbf{A}\mathbf{A}')^{-1}[\mathbf{I} + \mathbf{A}\mathbf{X}' \times \{\mathbf{S}^{-1}\mathbf{B}(\mathbf{B}\mathbf{S}^{-1}\mathbf{B})^{-1}\}\mathbf{X}\mathbf{A}'(\mathbf{A}\mathbf{A}')^{-1}]. \tag{23}$$

Based on Table 2, we calculate the following Wilks' Λ statistic,

$$\Lambda_1 = \frac{|\mathbf{E}_1|}{|\mathbf{E}_1 + \mathbf{H}_1|} = 0.5564. \tag{24}$$

and the corresponding F -statistic (with $d_m = 2$) to be

$$F_1 = \frac{1 - \sqrt{\Lambda_1}}{\sqrt{\Lambda_1}} \cdot \frac{(d_E - 1)}{d_H} = 6.5851. \tag{25}$$

Under the hypothesis H_1^* , F_1 has an F distribution with $d.f. = (2d_H, 2d_E - 2) = (6, 116)$. The p -value of this test thus turns out to be approximately = 0.0. Hence H_1^* is rejected, indicating at least one of the growth curves significantly differ from the rest.

Next, we perform pairwise comparisons to see how the classes compare with one another. These were carried out by fitting the model to two classes at a time and testing the appropriate linear hypotheses for the equality of the polynomial coefficients (taken to be linear here). Table 4 summarizes these results.

These p -values show that there is significant difference between the following pairs: 3R vs control, 4R vs control and 4RGV vs control. It is worth noting that the p -value for the pair (3R, 4RGV) is the highest, indicating that they are quite similar in their dynamics—confirming the widely believed bio-chemical structure function model of the biologists.

6. Conclusion

We have provided a novel statistical analysis of the dynamics of microtubules. Our analysis supports the widely held beliefs of biochemical functioning of microtubules and their responses to various treatments.

Acknowledgements

The authors would like to thank Professors Stuart Feinstein and Leslie Wilson as well as Dr Janis Bunker of the Department of Molecular Cellular and Developmental Biology, UCSB for helping us with the biological interpretations and for providing the original data sets. They also wish to express thanks to Professor B.S. Manjunath of the Electrical and Computer Engineering department at UCSB for the opportunity to collaborate on this bioinformatics project.

The authors would also like to thank two anonymous referees for their detailed comments which led to several improvements in the presentation.

The work of Jammalamadaka and Siddiqi was supported by National Science Foundation Grant # ITR-0331697.

References

- [1] B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts, and P. Walters, *Molecular Biology of the Cell*, Garland Science, London, 2002.
- [2] J. Avila, *Microtubule dynamics*, FASEB J. 4 (1990), pp. 3284–3290.
- [3] S.C. Feinstein and L. Wilson, *Inability of tau to properly regulate neuronal microtubule dynamics: a loss-of-function mechanism by which tau might mediate neuronal cell death*, Biochim. Biophys. Acta 1739 (2005), pp. 268–279.
- [4] J.E. Grizzle and D.M. Allen, *Analysis of growth and dose response curves*, Biometrics 25 (1969), pp. 357–381.
- [5] G. James and T. Hastie, *Functional linear discriminant analysis for irregularly sampled curves*, J. R. Statist. Soc. Ser. B 63 (2001), pp. 533–550.
- [6] M.A. Jordan and L. Wilson, *Microtubules as target for anticancer drugs*, Nat. Rev. Cancer 4 (2004), pp. 253–265.
- [7] C.G. Khatri, *A note on MANOVA model applied to problems in growth curves*, Ann. Inst. Statist. Math. 18 (1966), pp. 75–86.
- [8] A.M. Kshirsagar and W.B. Smith, *Growth Curves*, Marcel Dekker, New York, 1995.
- [9] T. Mitchison and M. Kirschner, *Dynamic instability of microtubule growth*, Nature 312 (1984), pp. 237–242.
- [10] M.E. Mortenson, *Geometric Modeling*, John Wiley and Sons, New York, 1985.
- [11] J.-X. Pan and K.-T. Fang, *Growth Curve Models and Statistical Diagnostics*, Springer-Verlag, New York, 2001.
- [12] R. Potthoff and S.N. Roy, *A generalized multivariate analysis of variance model useful especially for growth curve problems*, Biometrika 51 (1964), pp. 313–326.
- [13] C.R. Rao, *Linear Statistical Inference*, 2nd ed., John Wiley and Sons, New York, 1973.
- [14] D. Sengupta and S.R. Jammalamadaka, *Linear Models – An Integrated Approach*, World Scientific, Singapore, 2003.
- [15] S. Weerahandi, *Generalized Inference in Repeated Measures*, John Wiley and Sons, New York, 2004.