

# Lawrence Berkeley National Laboratory

## LBL Publications

### Title

Neutron Spectra as a Function of Depth in Uranium and Iron-Faced Concrete Shielding for Medium-Energy Cyclotron Beams on Tantalum

### Permalink

<https://escholarship.org/uc/item/3gd7n0qv>

### Authors

Wadman, William W

Miller, A Jerry

### Publication Date

1968

# University of California

## Ernest O. Lawrence Radiation Laboratory

NEUTRON SPECTRA AS A FUNCTION OF DEPTH  
IN URANIUM AND IRON-FACED CONCRETE SHIELDING  
FOR MEDIUM-ENERGY CYCLOTRON BEAMS ON TANTALUM

William W. Wadman III and A. Jerry Miller

January 25, 1968

**TWO-WEEK LOAN COPY**  
*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

RECEIVED  
LAWRENCE  
RADIATION LABORATORY  
LIBRARY AND  
DOCUMENTS SECTION

Berkeley, California

UCRL-18050-Summary  
ey J

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted for Am. Nucl. Soc.  
Annual Mtg., Toronto, Canada,  
June 9-13, 1968

UCRL-18050  
Summary  
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory  
Berkeley, California

AEC Contract No. W-7405-eng-48

NEUTRON SPECTRA AS A FUNCTION OF DEPTH  
IN URANIUM AND IRON-FACED CONCRETE SHIELDING  
FOR MEDIUM-ENERGY CYCLOTRON BEAMS ON TANTALUM

William W. Wadman III and A. Jerry Miller

January 25, 1968

NEUTRON SPECTRA AS A FUNCTION OF DEPTH  
IN URANIUM AND IRON-FACED CONCRETE SHIELDING  
FOR MEDIUM-ENERGY CYCLOTRON BEAMS ON TANTALUM

William W. Wadman III and A. Jerry Miller

Lawrence Radiation Laboratory  
University of California  
Berkeley, California

January 25, 1968

ABSTRACT

Shielding against neutrons produced by medium-energy cyclotron ion beams bombarding elemental tantalum has been studied at the Berkeley 88-inch cyclotron. Beams of 110-MeV alphas, 55-MeV deuterons, and 50-MeV protons were used to bombard a thick tantalum target. The resultant neutrons were attenuated by a stack of precision-cast ordinary concrete blocks into which threshold reaction detectors were placed at 3-in. intervals. The first 12 in. of concrete shielding were replaced with iron and depleted uranium to measure the effect of each of the face materials upon the neutron spectra and upon attenuation lengths of neutrons from the three different ion beams.

The detector materials used were nickel, aluminum, and thallium for the threshold reactions, and gold for thermal and epithermal neutron capture reactions. Properties of the threshold detectors are given in Table I.

The activities produced in each detector were counted with a  $\gamma$ -ray spectrometer. The computer program SAMPO was used to locate the

Table I. Threshold detector properties.

Reaction	Calculated threshold (MeV) <sup>a</sup>	Peak cross section (barns)	Product half-life	Target isotope (%)	γ-Ray energy of product (MeV)
$^{58}\text{Ni}(n, p)^{58}\text{Co}$	4.1	0.556	71 days	67.8	0.81
$^{27}\text{Al}(n, \alpha)^{24}\text{Na}$	6.7	0.243	15 hr	100	1.37, 2.74
$^{203}\text{Tl}(n, 2n)^{202}\text{Tl}$	8.5	2.78	12 days	29.5	0.44
$^{58}\text{Ni}(n, 2n)^{57}\text{Ni}$	12.4	0.25	37 hr	67.8	1.36
$^{27}\text{Al}(n, \alpha 2n)^{22}\text{Na}$	23.4	0.030	2.6 yr	100	0.51, 1.27
$^{203}\text{Tl}(n, 4n)^{200}\text{Tl}$	24.7	≈1.3	27 hr	29.5	1.207

a. The energy at which the cross section value is 2% of the maximum cross section value.

$\gamma$ -ray peaks, fit them to a Gaussian shape, determine their energy, strip each peak and calculate the area under it. The counts under the peak were corrected for counting geometry and detection efficiencies to obtain absolute counting rates. The activity data from SAMPO were then put into the computer program FLUXPOS along with information regarding irradiation, decay and counting times, half-life, foil and atomic weights, fraction of isotope, atoms per molecule, and the fraction of activity produced by neutrons below 30 MeV (the program's energy cutoff). With this information, the reaction cross section as a function of neutron energy was used to compute the neutron spectra. The relationships between detector activity, cross section, and neutron flux are expressed as

$$A_i' \int_0^{\infty} \phi(E) \sigma_i(E) d(E),$$

where  $A_i'$  = detector saturation activity or disintegration rate per target nucleus in  $\text{sec}^{-1}$ ,

$\phi(E)$  = neutron flux in  $\text{cm}^{-2} \text{sec}^{-1} \text{MeV}^{-1}$ ,

$\sigma(E)$  = reaction cross section in  $\text{cm}^2$ ,

$E$  = neutron energy in MeV,

and  $i$  = identification of the reaction and the residual nucleus.

One equation for each threshold reaction is used to form a set of integral equations that describe the unknown neutron flux. By iteration, the computer produces a set of calculated activities that best fit the set of measured activities. It then calculates the neutron spectrum that would be required to achieve the calculated ratios between the activities and the neutron flux necessary to produce the measured activities. Both programs are run on the CDC-6600 computer. The nine different experimental configurations produced 156 neutron spectra.

The results produce neutron spectra and attenuation data as a function of the measurement depth in the shielding and the material used for the first 12 in. Table II summarizes the attenuation for the nine configurations.

This study is the only known work that experimentally compares three beam ions, their forward neutron spectra, and the attenuation of forward neutrons, for the same geometry.

This work done under the auspices of the U. S. Atomic Energy Commission.



Table II. Attenuation length ( $1/e$  in  $\text{g/cm}^2$ ).

Energy particle	Shielding configuration		
	All concrete	12-in. iron faced	12-in. uranium (depleted)
110-MeV alpha	35	29.0	26.8
55-MeV deuteron	36.8	24.4	23.8
50-MeV proton	30.5	24.4	25.0

This paper is sponsored by Dr. Hans Mark and Dr. Lawrence Ruby.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

