UC San Diego

UC San Diego Previously Published Works

Title

Response to Cohen and Germain: Defining a Good Death.

Permalink

<https://escholarship.org/uc/item/3gp395nq>

Journal

The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry, 24(10)

ISSN

1064-7481

Authors

Meier, Emily A
Gallegos, Jarred V
Montross-Thomas, Lori P
et al.

Publication Date

2016-10-01

DOI

10.1016/j.jagp.2016.04.014

Peer reviewed

Response to Cohen and Germain: Defining a Good Death

TO THE EDITOR:

We appreciate the opportunity to respond to the points that Drs. Cohen and Germain have raised. First, we commend these investigators on their important work in end-stage renal disease and palliative medicine, highlighting the value of discontinuing dialysis earlier in a patient's treatment trajectory as a means of reducing suffering and thus allowing for a good death.^{1,2} We applaud their prospective study design to determine whether dialysis discontinuation increased quality of life and decreased unnecessary suffering.² Additionally, we compliment these authors on their poignant case study of a patient who chose to discontinue his dialysis, seeking a good death, and preferring *quality* of life over *quantity* of life.¹

The reason for our not including these articles in our review³ was that they did not meet our selection criteria. As we stated in our article, "... there are far fewer studies that have specifically defined, rather than conceptualized, what a good death is according to patients, family members, and health care providers (HCPs). The goal of this paper is to review the literature that examined the definitions of a good death from the perspectives of such patients, their family members, and HCPs" (p.262). Of the three articles by these investigators that were cited in their letter, one is a description of the renal palliative care initiative⁴ and one is a case study of a 73-year-old man who voluntarily decided to ter-

minate his dialysis and died 11 days later.¹ As can be seen in Figure 1 of our article, we had chosen to exclude "10 clinical case reports" because these reports conceptualized components of a good death from the author's perspective but did not specifically define good death according to the individual. Similarly, the study of 11 patients who discontinued dialysis and died subsequently² was excluded because the definition of good death used therein was researcher-derived rather than provided specifically by patients, their family members, and HCPs. Nonetheless, we are pleased to note that several factors associated with good death in our review were common to those reported by Drs. Cohen and Germain, that is, being pain-free, process of death (brevity and place of death), existential factors, and presence of loved ones.

Drs. Cohen and Germain have conducted valuable research in end-stage renal disease and in developing tools such as the Dialysis Quality of Dying Apgar.⁴ We share with them the goal of ensuring that patients have a good death by reducing unnecessary aggressive treatments that often prolong life but diminish quality of life. We too believe that further defining what constitutes a good death is an area of increasing importance for the medical community as well as for the general population.

Emily A. Meier, Ph.D.

Department of Psychiatry, Sam and Rose Stein Institute for Research on Aging, Moores Cancer Center, Psychiatry & Psychosocial Services; Patient & Family Support Services, University of California, San Diego, CA

Jarred V. Gallegos, M.A.

Department of Psychiatry, Sam and Rose Stein Institute for Research on Aging, Moores Cancer Center, Psychiatry & Psychosocial Services; Patient & Family Support Services, University of California, San Diego, CA

Lori P. Montross-Thomas, Ph.D.

Departments of Psychiatry and Family Medicine and Public Health, Sam and Rose Stein Institute for Research on Aging, Moores Cancer Center, Psychiatry & Psychosocial Services; Patient & Family Support Services, University of California, San Diego, CA

Colin A. Depp, Ph.D.

Department of Psychiatry, Sam and Rose Stein Institute for Research on Aging, University of California, San Diego, CA

Scott A. Irwin, M.D., Ph.D.

Department of Psychiatry, Moores Cancer Center, Psychiatry & Psychosocial Services; Patient & Family Support Services, University of California, San Diego, CA

Dilip V. Jeste, M.D.

Department of Psychiatry, Sam and Rose Stein Institute for Research on Aging, University of California, San Diego, CA

References

1. Cohen LM, Poppel DM, Germain GM, et al: A very good death: measuring quality of dying in end-stage renal disease. *J Palliat Med* 2001; 4:167-172
2. Cohen LM, McCue JD, Germain M, et al: Dialysis discontinuation: a good death? *Arch Intern Med* 1995; 155:42-47
3. Meier E, Gallegos JV, Montross-Thomas LP, et al: Defining a good death (successful dying): literature review and a call for research and public dialogue. *Am J Geriatr Psychiatry* 2016; 24:261-271
4. Poppel D, Cohen L, Germain M: The renal palliative care initiative. *J Palliat Med* 2003; 6:321-326