Lawrence Berkeley National Laboratory

Recent Work

Title

STUDY OF ANISOTROPIC DIFFUSION OF ORIENTED MOLECULES BY MULTIPLE QUANTUM SPIN ECHOES

Permalink

<https://escholarship.org/uc/item/3jq7b6w6>

Authors

Zax, D.

Pines, A.

Publication Date

1982-09-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

RECEIVED
LAWRENCE
BERKELEY LABORATORY

Materials & Molecular Research Division

JAN 11 1983

LIBRARY AND
DOCUMENTS SECTION

Submitted to the Journal of the American Chemical
Society

STUDY OF ANISOTROPIC DIFFUSION OF ORIENTED MOLECULES
BY MULTIPLE QUANTUM SPIN ECHOES

D. Zax and A. Pines

September 1982

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782.*

LBL-14997

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

STUDY OF ANISOTROPIC DIFFUSION OF ORIENTED MOLECULES
BY MULTIPLE QUANTUM SPIN ECHOES

D. Zax and A. Pines

Department of Chemistry and Materials

and

Molecular Research Division, Lawrence Berkeley Laboratory

University of California, Berkeley, California

94720

This work was supported by the Director, Office of Energy Research,
Office of Basic Energy Sciences, Materials Science Division of the
U.S. Department of Energy under Contract Number DE-AC03-76SF00098.

Abstract

Experiments have been completed which demonstrate the greatly enhanced sensitivity of the multiple quantum version of the NMR spin echo diffusion experiment. These experiments are particularly well-suited to studies of anisotropic diffusion where the presence of dipolar couplings may make the standard experiment technically infeasible.

A variety of methods have been used to analyze the diffusive motion of molecules. In this communication we report on the application of multiple quantum NMR to the problem of high sensitivity measurements of diffusion in anisotropic systems. The multiple quantum diffusion experiment is related to the standard NMR spin echo diffusion experiment¹ and is a generalization of the 2-quantum experiments of Morton et al.² to arbitrary dipolar coupled proton systems. It shares those qualities which have made the NMR diffusion experiment a favorite method of measuring diffusion constants. It measures diffusion in a system at equilibrium without the need to introduce any external probes or labels. Individual molecules, instead, are labeled by their position in a magnetic field gradient after an initial radio frequency (r.f.) pulse, and the motion is followed by monitoring the amplitude of the spin echo following a second r.f. pulse some time later, as depicted in Figure 1(a). Those nuclear spins which have moved little contribute more to the bulk echo signal than those which have strayed far from the field in which they were labeled.

While the NMR spin echo diffusion experiment is routinely performed on isotropic systems, the interesting case of anisotropic diffusion, as is found in liquid crystals, bilayers, membranes, or solids, is studied less frequently. Though the same conceptually simple method is applicable, it is complicated by two factors: firstly, the diffusion tends to be slower (which necessitates larger field gradients) and secondly, the dipolar interactions, which are absent in isotropic systems, interfere with the formation of a spin echo at times long compared to the inverse of the dipolar spectral width.

The multiple quantum spin echo diffusion experiment substantially

alleviates both these problems. Normal Fourier transform NMR techniques involve the excitation and detection of coherences where the difference in Zeeman quantum numbers, Δm , is ± 1 . Multiple quantum NMR utilizes a sequence of r.f. pulses to prepare and detect coherences where $n = \Delta m$ is arbitrary³. We exploit the fact that an n quantum coherence dephases in a field gradient n times faster than does a single quantum coherence⁴. This effect is obviously of considerable advantage in diffusion studies and has previously been demonstrated by Morton et al.² for the 2 quantum coherence of CH_2Cl_2 dissolved in various liquid crystals. For N coupled protons, the N -quantum transition is both free of dipolar couplings and most sensitive to dephasing by the field gradient. This approach eliminates the need for those line narrowing techniques typically employed for diffusion measurements in dipolar coupled systems.⁵ Our generalization of the NMR spin echo diffusion experiment to multiple quantum echoes is shown in Figure 1(b).

Gradients of up to 15 gauss/cm (calibrated against H_2O) were applied through a quadrupole coil and homemade current pulser capable of switching 2 amp in 10 μsec . The gradient coil was rotated to produce gradients either parallel or perpendicular to the main magnetic field. Multiple quantum diffusion experiments have been performed on samples of benzene and 2-butyne dissolved in Eastman liquid crystal #15320, forming a nematic phase, which demonstrate the increasing sensitivity of n quantum echoes as n increases. Figure 2 shows the order dependence of the decay of the echo amplitude for all orders of benzene. The straight lines represent best fits to the Stejskal-Tanner equation modified for n quantum echoes:

$$\ln \frac{A(t_1)}{A_0(t_1)} = -\gamma^2 n^2 g^2 D (\delta^2 (\Delta - \frac{\delta}{3})) \quad (1)$$

where $A(t_1)$ is the observed echo amplitude, $A_0(t_1)$ is the amplitude in the absence of any gradient, γ is the magnetogyric ratio, n is the multiple quantum order, D is the diffusion constant and δ and Δ are as defined in Figure 1.

Results are shown in Table 1, and represent an average over all n quantum orders. None of the orders gave results outside of the error bars listed. The primary uncertainty remains the gradient calibration. Experiments performed in a time-independent gradient and corrected for T_2 relaxation gave essentially identical results.

TABLE 1

	$D_{\perp}$ (cm ² /sec)	$D_{\parallel}$ (cm ² /sec)	$D_{\parallel}/D_{\perp}$
Benzene	$.9 \times 10^{-6} \pm .1$	$1.1 \times 10^{-6} \pm .1$	$1.25 \pm .05$
2-Butyne	$3.1 \times 10^{-6} \pm .2$	$3.5 \times 10^{-6} \pm .2$	$1.1 \pm .05$

It is clear that the high n -quantum echoes form much more sensitive measures of diffusion. The technique is likely to find its greatest applicability in the measurement of slow self diffusion of oriented molecules where current methods fail either due to short T_2 's or the need for unrealistically large field gradients. The combination of heightened sensitivity and narrowed lines achievable in the multiple quantum spin echo diffusion experiment seems the ideal solution to current experimental limitations.

Acknowledgements

We thank R. Tycko for work on the current pulsing apparatus.

D. Zax gratefully acknowledges receipt of a National Science Foundation Graduate Fellowship. This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Materials Science Division of the U.S. Department of Energy under Contract Number DE-AC03-76SF00098.

Figure 1. NMR pulsed gradient diffusion experiment. Cross hatched areas indicate gradient pulses. The experiment is typically performed with Δ fixed and δ regularly incremented. Only the peak height of the single quantum echo is digitized. The results are fitted to Equation 1 and the diffusion constant, D , extracted. (a) The single quantum version. (b) The multiple quantum version. This sequence prepares multiple quantum orders nonselectively and detects selectively only magnetization which evolved during t_1 as n quantum.⁶ Preparation and mixing times (τ and T) were optimized for the multiple quantum order of interest using parameter proportional phase incrementation (PPPI).⁷ The fixed multiple quantum evolution time, t_1 , was similarly chosen to maximize the amplitude of the observed coherence transfer echo.

Figure 2. Results of the NMR diffusion experiment for all n quantum orders of benzene (25 mol %) dissolved in Eastman liquid crystal #15320. The applied gradient is parallel to the magnetic field axis. The plots show the normalized echo amplitude vs. the gradient pulse timing parameter of equation 1. Straight lines are linear least-squares fits to the accumulated data. The slopes of these lines vary as n^2 , as is predicted by theory. This illustrates the vastly increased sensitivity of the multiple quantum diffusion experiment.

References

- (1) (a) Hahn, E. L. Phys. Rev. 1950, 80, 580
(b) Carr, H. Y.; Purcell, E. M. Phys. Rev. 1954, 94, 630
(c) Stejskal, E. O.; Tanner, J. E. J. Chem. Phys. 1965, 42, 288
- (2) Morton, J. F.; Selwyn, L. S.; Vold, R. L.; Vold, R. R.
J. Chem. Phys. 1982, 76, 2632
- (3) Drobny, G.; Pines, A.; Sinton, S.; Weitekamp, D.; Wemmer, D.
Faraday Div. Chem. Soc. Symp. 1979, 13, 49
- (4) (a) Vega, S.; Pines, A. J. Chem. Phys. 1977, 66, 5624
(b) Bax, A.; De Jong, P. G.; Mehlkopf, A. F.; Smidt, J.
Chem. Phys. Lett. 1980, 69, 567
(c) Wokaun, A.; Ernst, R. R. Mol. Phys. 1978, 36, 317
(d) Bodenhausen, G.; Vold, R. L.; Vold, R. R. J. Magn. Reson.
1980, 37, 93
- (5) (a) Blinc, R.; Pirs, J.; Zupancic, I. Phys. Rev. Lett. 1973,
30, 546
(b) Zupancic, I.; Pirs, J.; Luzar, M.; Blinc, R.; Doane, J. W.
Sol. State. Comm. 1974, 15, 227
(c) Karlicek, R. F.; Lowe, I. J. J. Magn. Reson. 1980, 37, 75.
(d) Kuo, A.; Wade, C. G. Biochemistry 1979, 18, 2301
- (6) (a) Hatanaka, H.; Terao, T.; Hashi, T. J. Phys. Soc. Japan 1975,
39, 835
(b) Maudsley, A. A.; Wokaun, A.; Ernst, R. R.; Chem. Phys. Lett.
1978, 55, 9
- (7) Weitekamp, D. P.; Garbow, J. R.; Pines, A. J. Magn. Reson. 1982,
46, 529

(a)

(b)

XBL 828-10789

FIGURE 1

XBL 828-10830

FIGURE 2

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720