

UCSF

UC San Francisco Previously Published Works

Title

Newborn screening for SCID identifies patients with ataxia telangiectasia.

Permalink

<https://escholarship.org/uc/item/3k46b1xw>

Journal

Journal of Clinical Immunology, 33(3)

Authors

Mallott, Jacob

Kwan, Antonia

Church, Joseph

[et al.](#)

Publication Date

2013-04-01

DOI

10.1007/s10875-012-9846-1

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Newborn Screening for SCID Identifies Patients with Ataxia Telangiectasia

Jacob Mallott · Antonia Kwan · Joseph Church ·
Diana Gonzalez-Espinosa · Fred Lorey ·
Ling Fung Tang · Uma Sunderam · Sadhna Rana ·
Rajgopal Srinivasan · Steven E. Brenner · Jennifer Puck

Received: 22 October 2012 / Accepted: 27 November 2012 / Published online: 20 December 2012
© Springer Science+Business Media New York 2012

Abstract

Purpose Severe combined immunodeficiency (SCID) is characterized by failure of T lymphocyte development and absent or very low T cell receptor excision circles (TRECs), DNA byproducts of T cell maturation. Newborn screening for TRECs to identify SCID is now performed in several states using PCR of DNA from universally collected dried blood spots (DBS). In addition to infants with typical SCID, TREC screening identifies infants with T lymphocytopenia who appear healthy and in whom a SCID diagnosis cannot be confirmed. Deep sequencing was employed to find causes of T lymphocytopenia in such infants.

Jacob Mallott and Antonia Kwan contributed equally to the work.

J. Mallott · A. Kwan · D. Gonzalez-Espinosa · J. Puck (✉)
Department of Pediatrics, University of California San Francisco,
513 Parnassus Avenue, HSE 301A, Box 0519, San Francisco,
CA 94143-0519, USA
e-mail: puckj@peds.ucsf.edu

J. Church
Department of Pediatrics, Keck School of Medicine, University
of Southern California and Children's Hospital Los Angeles,
Los Angeles, CA, USA

F. Lorey
Genetic Disease Laboratory, California Department of Public
Health, Richmond, CA, USA

L. F. Tang · J. Puck
Institute for Human Genetics, University of California San
Francisco, San Francisco, CA, USA

U. Sunderam · S. Rana · R. Srinivasan
Innovations Labs, Tata Consulting Services, Hyderabad, AP, India

S. E. Brenner
Department of Plant and Microbial Biology,
University of California, Berkeley, CA, USA

Methods Whole exome sequencing and analysis were performed in infants and their parents. Upon finding deleterious mutations in the ataxia telangiectasia mutated (*ATM*) gene, we confirmed the diagnosis of ataxia telangiectasia (AT) in two infants and then tested archival newborn DBS of additional AT patients for TREC copy number.

Results Exome sequencing and analysis led to 2 unsuspected gene diagnoses of AT. Of 13 older AT patients for whom newborn DBS had been stored, 7 samples tested positive for SCID under the criteria of California's newborn screening program. AT children with low neonatal TRECs had low CD4 T cell counts subsequently detected ($R=0.64$).

Conclusions T lymphocytopenia in newborns can be a feature of AT, as revealed by TREC screening and exome sequencing. Although there is no current cure for the progressive neurological impairment of AT, early detection permits avoidance of infectious complications, while providing information for families regarding reproductive recurrence risks and increased cancer risks in patients and carriers.

Keywords Ataxia telangiectasia · SCID · newborn screening · TREC · whole exome sequencing

Introduction

Severe combined immunodeficiency (SCID), characterized by extremely low or absent T cell production, defective T cell function and absent antibody responses, can be caused by defects in any of several genes and if untreated leads to early death due to infections [1, 2]. Population based newborn screening for SCID has been recommended to identify affected infants before the onset of devastating infections so that effective treatment can be provided [3–6]. A newborn screening test for SCID, now implemented in several states,

ascertains T-cell receptor excision circles (TRECs), DNA byproducts of T cell antigen receptor gene rearrangement, as a biomarker of normal T cell development [5, 7–10]. TRECs are measured by quantitative PCR (qPCR) of DNA isolated from infant dried blood spot (DBS) samples universally collected in nurseries. For infants with undetectable or low TRECs, or with unsatisfactory DNA amplification, a differential white blood count and analysis of lymphocyte subsets by flow cytometry are obtained to establish the absolute number of naïve T cells, after which further clinical and laboratory evaluations are performed to arrive at a definitive diagnosis [9].

Beyond typical SCID cases, TREC screening has detected a spectrum of infants with inadequate numbers of diverse, autologous T cells. As predicted before the start of screening, “leaky” SCID and Omenn syndrome, both due to hypomorphic mutations in SCID genes, have been found in infants with low TRECs, as have cases of DiGeorge syndrome/chromosome 22q11 deletion in which a substantial degree of thymic insufficiency exists. In addition, secondary causes of T lymphocytopenia have included abnormal loss of T cells from the peripheral circulation, such as with chylothorax or hydrops. A challenging and less anticipated category of cases with abnormal TREC results has been the infants with persistent T lymphocytopenia of 300–1,500 T cells/ μ L, no maternal T cell engraftment, and absence of identified deleterious mutations in common SCID genes. These infants have been designated as combined immunodeficiency (CID) or SCID variants by TREC newborn screening programs [9]. CID or SCID variant cases have been of particular interest, providing an opportunity to discover previously unappreciated causes of newborn T lymphocytopenia. In the absence of clues to narrow the number of potential candidate genes to account for variant SCID in asymptomatic infants who appear healthy, high throughput deep sequencing may be useful; this approach has led to gene identification in other primary immunodeficiencies [11–13].

Using whole exome sequencing (WES), we found two infants with variant SCID who had deleterious mutations in the Ataxia Telangiectasia Mutated (*ATM*) gene. Prompted by the prospective discovery of these patients’ diagnoses and a recent report of low TRECs in archived DBS from cases of ataxia telangiectasia (AT) [14], we reviewed 13 cases of AT in our clinical cohorts. By retrieving their residual DBS samples taken in the newborn nursery and measuring their TREC numbers, we showed that over half of AT patients could be identified as abnormal, making AT a secondary target of SCID screening.

Methods

Subjects Infants V003 and V004 were identified as positive by routine California SCID screening by TREC test and

confirmed to have T lymphocytopenia. Informed consent for research, including cellular immune studies and WES, was obtained for the infants and their parents under approved protocols at Children’s Hospital Los Angeles (CHLA) and the University of California San Francisco (UCSF). Additional patients from the pediatric immunology services at CHLA and UCSF were enrolled with institutional review board approval.

DNA Samples Genomic DNA from EDTA anticoagulated whole blood was prepared using a Gentra Puregene Blood kit (Qiagen USA: Germantown, MD).

Exome Sequencing Libraries were prepared by ligating a pair of TruSeq adaptors (Illumina: San Diego, CA) to genomic DNA sheared to a mean fragment size of 200–300 bp (S2 sonicator, Covaris: Woburn, MA). Specific sequence tags were added to different samples to differentiate each individual of origin. Libraries with these adaptors and barcode sequences were enriched with 10 cycles of PCR. For infant V003 and parents, exon capture was performed by pooling 500 ng of each of 6 libraries incubated with Illumina TruSeq version 2 biotinylated exon-encoded DNA oligonucleotides for 20 h. For V004 and parents, exon capture was performed by incubation with a Roche Nimblegen version 3 capture array. Exon-enriched DNA was captured with streptavidin-labeled magnetic beads, washed and eluted. Capture reactions were repeated to enhance specificity. After 10 cycles of DNA amplification, the exome libraries were sequenced (HiSeq2000, Illumina). Paired 100 bp end reads were generated (>50 M reads/subject), to yield an average of >65 reads covering the targeted regions with >90 % covered by at least 10 reads.

Whole Exome Sequence Analysis Raw reads were aligned against reference genome hg19 using BWA (0.5.9) software [15]. The resulting files were converted to compressed binary format (BAM), sorted by coordinate, indexed, and marked for PCR duplicate reads using the Picard toolkit (<http://picard.sourceforge.net>). BAM files were processed to reduce artifacts and improve call accuracy using GATK software (v 1.4.15) [16, 17]. Specifically, local realignment was performed around known insertion or deletion (indel) locations, and base quality scores were re-calibrated using co-variates such as position in read and sequencing chemistry effect.

Variants were called using the GATK UnifiedGenotyper. The called single nucleotide polymorphisms (SNPs) had their scores re-calibrated by variant quality score recalibration (VQSR) using the exomes in this report plus 24 others sequenced at our site. HapMap v3.3 and the Omni chip array sets from the 1,000 genomes project (October, 2011 release) were training data, and HapMap 3.3 provided truth sites [18,

19]. A truth sensitivity cutoff of 99 % was used. For indel recalibration and quality selection, we used QD <2.0, ReadPosRankSum <-20.0, Fischer Strand >200.0. Additional annotations including region, effect, dbSNP 135 and 1,000 genomes membership and OMIM phenotype were added using custom scripts. Filtering of exome data was performed using a combination of custom scripts and vcftools [20]. Aligned sequences were viewed using the Savant Genome Browser [21].

PCR and Genomic Sequencing *ATM* exons 7, 10, 39, and 46 were amplified from genomic DNA using published primers (labeled as exons 9, 12, 41, and 48 in Thorstenson et al. [22]) with an M13 extension 5' to each reverse primer as follows:

ATMe7_Forward 5' - GTA AAA CGA CGG CCA GTC AGC ATA CCA CTT CAT AAC TG
ATMe7_Reverse 5' - TCA TAT CCT CCT AAA GAA CAC
ATMe10_Forward 5' -TGT GAT GGA ATA GTT TTC AA
ATMe10_Reverse 5' -GTA AAA CGA CGG CCA GTT GTG ATG GAA TAG TTT TCA A
ATMe39_Forward 5' -TGT GGT TTT TGG GAA TTT GTA
ATMe39_Reverse 5' -GTA AAA CGACGG CCA GTT GTG GTT TTT GGG AAT TTG TA
ATMe46_Forward 5' - GTA AAA CGA CGG CCA GTT CTT GTC ACT ACA AAA GTT CCT TT
ATMe46_Reverse 5' - TCT TTT TCC CTC AGG CTT TC.

Sequencing was performed with the M13 forward primer, and results were compared with reference *ATM*NG_009830.1, using Sequencher 4.10.1 software (Gene Codes Cooperation: Ann Arbor, MI).

Neonatal Dried Blood Samples Residual DBS originally collected for routine NBS and stored at -20° by the Genetic Disease Laboratory (GDL) of the California Department of Public Health (CDPH) were retrieved, and TREC and β -actin gene copy number determined by the GDL newborn screening laboratory, using the protocol of Chan and Puck [5] modified for high throughput and implemented by PerkinElmer, Inc (lab within a lab at California Department of Public Health, Richmond, CA; parent company based in Hershey, PA) with cutoff values as reported [9].

Results

Infant Clinical and Immunologic Findings Newborns in California are screened for SCID and classified as positive

if TREC copy number is ≤ 5 with β -actin >5,000 copies, or TREC copy number is between 6 and 25 with β -actin >10,000 copies. The tests are classified as incomplete and are repeated if there are low TRECs, but also low copies of the β -actin gene segment amplified as a control. T cells are measured by flow cytometry in cases that are positive or that have two incomplete DBS samples.

Infants V003 and V004 were unrelated, healthy females born at term following normal pregnancies. Family history for both was negative for immune disease or consanguinity. Lymphocyte flow cytometry was ordered for infant V003 after two DBS yielding incomplete results, while V004 had an initial positive result with 21 TRECs and 13,300 β -actin copies (Table I).

At age 3 months Patient V003 had a normal total white blood cell count, but only 4 TRECs/ μ L (with normal β -actin copies) and only 1,600 lymphocytes/ μ L (Table I). There were 996 T and 52 B cells/ μ L (normal >2,000 and >300, respectively), and the number of CD45RA naïve CD4 T cells was low. NK cell number was normal. Low T and B cell numbers persisted, and low IgG levels with failure to produce antibodies after vaccination led to institution of immunoglobulin replacement and trimethoprim-sulfamethoxazole antibiotic prophylaxis. Lymphocyte proliferation to phytohemagglutinin and TCR V β diversity assessed by spectratyping [23] were normal, but an Epstein Barr virus transduced B cell line from the patient had only half normal phosphorylation of STAT5 in response to IL-2 [24], suggesting an intrinsic lymphocyte impairment. Later, at age 14–16 months, V003 was reported by her mother to have “unsteady” gait; physical examination first showed mild truncal ataxia at 20 months.

Infant V004 had flow cytometry at 21 days of age, showing only 1,060 T cells, with low CD45RA naïve helper CD4 T cells (Table I). B and NK cell numbers and lymphocyte proliferation were normal, but V β spectratyping showed decreased T cell diversity (not shown). As with infant V003, in vitro phosphorylation of STAT5 after IL-2 activation was diminished, but not absent, as would be the case in SCID due to defects in the IL-2 receptor common γ chain or Janus kinase 3 [9, 24]. Physical examination of infant V004 has been normal to date, but she did not mount robust antibody to T-cell dependent protein-conjugated *H. influenzae* vaccination.

Exome Analysis and Gene Confirmation To investigate the genetic etiology underlying their observed immunodeficient status, DNA samples from V003, V004 and their parents were subjected to WES to generate a list of small nucleotide polymorphism (SNP) and small insertion or deletion (indel) variants. These lists were filtered to retain successively fewer candidate variants as shown for SNPs and indels separately for each infant in Fig. 1b. After initial quality

Table 1 Immunologic Phenotype of Infants Identified by SCID Newborn Screening

Infant	V003		V004	
Age	3 m ^a	5.5 m	21 days ^b	7 m
TREC (normal >25)	4^c		21	
Beta-actin (>10,000)	10,000		13,300	
WBC (5,000–19,500 cells/ μ L)	7,100	3,530	4,500	5,320
ALC (2,500–16,500)	1,600	2,189	2,000	3,032
CD3 T cells (2,550–5,500)	996	615	1,060	1,397
CD4 T-helper cells (1,600–4,000)	749	502	600	777
CD8 T-cytotoxic cells (560–1,700)	185	172	340	400
CD3/CD4/CD45RA (1,200–3,700)	456		360	
CD3/CD4/CD45RO (60–900)	299		180	
CD19 B cells (300–2,000)	52	84	440	1,232
CD16/56 NK cell (170–1,100)	347	1,605	300	381
IgG (165–781 mg/dL)		156		414
IgA (25–154)		< 7		17
IgM (31–103)		37		66
IgE (<17 IU/mL)		< 2		
Anti-tetanus (>0.1 IU/mL) ^d		0.07		3.73
Anti- <i>H. influenzae</i> type b (>1 mcg/ml) ^d		< 0.15		0.68
Proliferation to PHA		normal		normal
Diversity of TCR V β repertoire ^e		normal		decreased
IL-2 induced STAT5 phosphorylation ^f		55% of control		75% of control

^a Initial and follow-up dried blood spot samples from infant V003 were incomplete, with low TRECs, but also low β -actin; despite multiple notifications from the newborn screening program, there were delays between the initial screening result and a follow-up blood sample being submitted for both flow cytometry and a third TREC screen, which was positive

^b Initial screening result from infant V004 was positive at age 16 days; follow-up flow cytometry was performed at age 21 days; this timeline is typical for the CA SCID screening program

^c Abnormal values displayed in **bold type**

^d Serum antibody concentrations following 3 immunizations

^e Spectratyping according to Sarzotti et al. [23]

^f EBV transformed cells exposed to IL-2, normal control=100 % [24]

filtering and removal of variants common enough to be found in dbSNP 135 and the 1,000 Genomes database, further filters were applied to keep only the non-synonymous variants lying in captured exonic and splice-site regions of genes, and only those variants with a high (>30) genotype quality score (Fig. 1a, step 4).

We then further limited the disease gene candidates either by function of gene products (Fig. 1a, yellow) or by genetic segregation (red). For functional selection, we used a list of 49 candidate genes involved in T cell development or reported to be defective in human primary T cell deficiencies by the International Union of Immunological Sciences Committee on Primary Immunodeficiency [25]. In infant V003, this filtering left three heterozygous variants, of which two were frameshift deletions within the *ATM* gene. Aligned sequence reads supporting one of these, variant c1787delAA (K468fs), a two base deletion in *ATM* exon 10, are illustrated in Fig. 2a. Evidence for the second *ATM*

variant of V003, c6238delA (F1952fs), a single base deletion in exon 39, was equally robust (not shown).

In infant V004, inclusion in the T cell gene list yielded only two heterozygous SNPs, both in *ATM*, c.1260C>T (P292L), and c.7064C>T (R2227C). These results suggested disease-causing compound heterozygosity in both V003 and V004.

By the segregation filtering method, we retained gene altering variants fitting a homozygous or compound heterozygous model of recessive inheritance - that is, those genes for which the patient inherited one rare allele from each parent - using exome data from each infant/parent trio. By this method, 5 candidate genes remained in the family of V003 and 9 in the family of V004. By filtering out variants also found in the unrelated local exomes used for VQSR, these numbers were reduced to 3 and 7 genes, respectively. Infant V003 shared the *ATM* mutation K468fs with her mother and F1952fs with her father, while V004 shared




Fig. 1 WES variants filtering paths. **a** Trapezoids represent filters with resulting numbers of variants retained after each step indicated by a circled digit. Starting with initial total variant lists (1), filters were applied for quality (2) and then to keep rare alleles (3) that alter splice sites or produce non-synonymous codon changes and are absent in local exomes (4). Subsequent strategies were: focusing on variants from a list of genes associated with T cell phenotypes (yellow shading, 5); or demanding a recessive inheritance pattern (red shading, 6). **b** Numbers of variants retained for the exome of each proband, V003 and

V004, after each filtering step in A, showing individual numbers of SNPs/indels, left, and genes harboring variants, right. For steps (7) and (8), number of genes containing candidate variants in each proband are shown. The final lists of genes at step (8) are as follows: for V003 – *ATM*, *PCDH15*, *PHF2*; for V004 – *ATM*, *EYS*, *PCDP1*, *PRUNE2*, *SH3D21*, *TSHZ3*, *TTN*. *, 2 variants, both in the *ATM* gene. **, genes with rare variants conforming to a recessive disease model in the family trio

P292L with her mother and R2227C with her father. The other genes harboring parentally shared variants were not associated with any recognized immunologic phenotype.

Sanger sequencing confirmed the *ATM* mutations seen by exome sequencing for both V003 (Fig. 2b) and V004, as well as in both sets of parents (results not shown).

Subsequent to the sequence findings, both infants underwent measurement of serum alpha fetoprotein (AFP) levels; elevated AFP compared to age-adjusted normal ranges is a reliable marker for AT in children. V003 at age 16 months had AFP 307 $\mu\text{g/L}$, while V004 at 7 months had 112 $\mu\text{g/L}$ (normal range for these ages, 8–80 $\mu\text{g/L}$ [26]). Western blot showed absent *ATM* protein in both patients (data not shown).

TREC Analysis in Additional AT Cases To test whether T lymphocytopenia detected by low TRECs is common in infants with *ATM* mutations, medical records of California-born patients with AT followed at CHLA and UCSF over the past 25 years were reviewed, and their residual neonatal DBS were retrieved by the CDPH for TREC testing. As summarized in Table II, 13 patients with AT, 9 females and 4 males, none of whom had been suspected to have AT at birth, had their newborn DBS samples retrieved. Upon testing, 7 samples had TRECs ≤ 25 ; thus newborn screening would have flagged these AT patients in infancy to receive follow-up lymphocyte immunophenotyping. The AT patients' ethnic distribution was not different from the overall distribution of California births.

All 13 AT patients initially presented with symptoms of ataxia and abnormal gait between ages 12 months and 8 years (median 17 months). Patients experienced a delay from 2 months to 10 years between onset of symptoms and AT diagnosis, which occurred between ages 1.5 and 12 years (median age 3 years 5 months). At diagnosis all 13 AT patients had high serum AFP concentrations, from 16.8 $\mu\text{g/L}$ at 1 year 7 months (Patient 6) to 310 $\mu\text{g/L}$ at 12 years 3 months (Patient 3). AFP levels in utero and at birth are high, but fall to < 8 $\mu\text{g/L}$ for children over age 2 [26]. We confirmed the correlation between AFP and age ($R=0.63$), as reported [26, 27]. Furthermore, all 13 patients had T lymphocytopenia with $< 1,500$ T cells/ μL at the time of their diagnosis of AT; subsequently, 8 had recurrent immunological manifestations including soft tissue infections (1 patient), cutaneous and visceral granulomas (3 patients), and chronic respiratory infections (4 patients). Out of 13 patients, 4 also had hematological malignancies, leading to fatality in Patients 1 and 10.

Comparing the 7 patients whose SCID newborn screens were positive (≤ 25 TRECs with normal β -actin) vs. the 6 whose were negative (TREC > 25 copies), there were no significant differences in age at presentation with neurological symptoms, AFP levels, total CD3 T cell counts, or time between symptom onset and diagnosis, though our T cell information from retrospective chart review did not provide lymphocyte subset data in infancy; the median age for the first recorded immune panel was 4 years. Interestingly, there was a correlation between TREC copy numbers in the




Fig. 2 Sequence evidence for a heterozygous exon 10 2-bp deletion of *ATM* in infant V003. **a** Aligned paired-end reads from whole exome sequence, chr11:108121571–108121612, viewed with Savant Genome Browser.¹⁸ Center black bars indicate deletion of c1787delAA, K468fs, found in 27 of the 54 reads that include this sequence. Dark blue, forward reads; light blue, reverse reads. Colored rectangles, mismatch base calls compared to reference genome (judged to be artifacts because of singular occurrence). **b** Sanger genomic reverse sequence confirming the heterozygous deletion. **c** Reference and mutated amino acid codons, showing the frameshift, which led to 17 missense codons followed by a termination

patients' archived newborn DBS and their subsequently measured CD4 T cell counts ($R=0.64$). Further analysis with a larger sample of AT patients might reveal additional relationships between newborn TREC numbers and phenotypic clinical and laboratory features of AT.

Discussion

Newborn screening by TRECs was developed with SCID as its primary target, but a spectrum of conditions are also identified that feature clinically significant T lymphocytopenia, defined by the California newborn screening program as $<1,500$ T cells/ μL or a lack of CD45RA naïve T cells. Through screening with the TREC assay, two apparently healthy California newborns with unexplained lymphocytopenia were identified to have AT, with deleterious mutations in the *ATM* gene detected by WES and confirmed by Sanger

sequencing, AFP elevation and undetectable ATM protein expression. The abnormal TREC screening results in infants V003 and V004 allowed physicians to avoid exposure to live attenuated rotavirus vaccine, contraindicated in infants with T cell immunodeficiency. Although T and B cell immunodeficiency is a well recognized, but variable feature of AT patients of older ages, the degree of immune compromise in early infancy has not been documented. Our early diagnosis of AT has provided an opportunity to observe prospectively the evolution of immunological, neurological and malignant features of AT.

Clinical features of AT include the loss of motor milestones between 1 and 2 years of age, with falls, slurred speech, oculomotor apraxia, and truncal instability. Ocular telangiectasias appear between 2 and 8 years. Family history, if positive, may assist diagnosis. Laboratory features often include absent IgA and always include elevated serum AFP, though the interpretation of AFP requires age-adjustment because this fetal serum protein remains abundant in infancy [26, 27]. Magnetic resonance imaging of the brain is unhelpful in early disease, as cerebellar abnormalities become apparent only after 2 years [28]. Increased cellular radiosensitivity is found in AT, though assays may not differentiate AT from other defects with defective DNA repair [29]. Thus, low TRECs and T lymphocytopenia detected by newborn screening can be the earliest and simplest warning that AT may be present, and validates the hypothesis of Borte et al., who suggested that AT might be detectable by prospective TREC screening, based on their survey measuring TREC and kappa chain B cell excision circles in archival DBS from patients with known primary immunodeficiencies [14]. The large size of the *ATM* gene, with 63 exons encoding an mRNA of 13,147 nucleotides, makes genomic sequencing costly and laborious; thus, WES may become a cost effective first line approach to examine this gene.

Deep Sequencing in the Context of Newborn Screening with TRECs

Our study demonstrates the value of a “short-list” of gene candidates of interest when searching deep sequencing data for rare, disease-causing variants. By focusing on a list of genes known to be associated with T-cell deficiencies, we were able to single out clinically relevant mutations with a minimum of other filters. The variants in our families occurred in regions with excellent coverage, as illustrated by the abundant bidirectional reads in Fig. 2a. However, high-quality coverage of all AT exons by WES or other deep sequencing is not guaranteed. In the exome datasets of our infants and their parents, between one and four *ATM* exons had $<15\times$ coverage, arguably inadequate to make accurate

Table II Characteristics of AT patients whose newborn dried blood spots were tested for TRECS

Patient	Ethnicity	Sex	Newborn DBS	Age at first symptoms	Age at diagnosis of AT	First immune panel, showing absolute lymphocyte number/ μ L at indicated age				Earliest AFP, μ g/L (age)	Immune and hematologic manifestations	Current age or age at death			
						T	CD4	B	NK						
1	White	F	3	7,690	Positive	2y	2y 9m	1,162	995	n/a	n/a	n/a	131 (2y)	Lymphoma (6 y), granulomatous skin lesions (3 y)	d. 6y 3m
2	Multiple	F	7	65,500	Positive	1y	2y 8m	681	373	n/a	n/a	n/a	60 (2y 8m)	bronchiectasis	16y
3	White	M	14	16,400	Positive	5y	5y 8m	406	70	42	175	310 (12y 3m)	Splenic granulomas (24 y), osteomyelitis (19 y), abdominal Burkitt lymphoma (14 y)	25y	
4	Hispanic	F	19	33,500	Positive	8y	11y 6m	740	475	142	n/a	62.8 (11y 7m)	None	14y 11m	
5	Hispanic	M	22	20,500	Positive	1y	3y 4m	443	227	173	421	108.1 (3y 4m)	Granulomatous skin lesions	6y 2m	
6	Hispanic	M	24	36,800	Positive	1y 5m	1y 7m	295	125	100	210	16.8 (1y 7m)	None	8y 0m	
7	Asian	F	25	31,900	Positive	1y	n/a	n/a	n/a	n/a	n/a	164 (11y 11m)	Large B-cell lymphoma (13 y); fungal pneumonia (14 y)	17y	
8	Hispanic	F	28	14,900	Normal	1y 4m	3y 5m	338	180	128	255	86.8 (3y 7m)	None	6y 2m	
9	Hispanic	F	30	61,300	Normal	3y	11y 10m	463	277	47	n/a	174 (11y 10m)	Bronchiectasis	12y 2m	
10	Hispanic	M	39	34,700	Normal	2y	7y 3m	1,069	534	196	n/a	127 (7y 6m)	Recurrent HSV; relapsed AML (10 y)	d. 10y 3m	
11	Asian	F	41	26,100	Normal	10m	1y 6m	176	132	522	n/a	48.7 (1y 9m)	None	2y 6m	
12	Hispanic	F	101	88,100	Normal	1y 2m	2y 11m	861	369	284	n/a	210 (6y 8m)	None	8y 4m	
13	Hispanic	F	216	37,300	Normal	2y 6m	4y 6m	1,571	428	238	n/a	37.6 (4y 6m)	Chronic sinusitis and otitis	7y 6m	

variant calls. As quality, coverage and affordability of deep sequencing improve, the chance of missing a variant will decline, but should be considered when using WES clinically.

Another consideration regarding deep sequencing analysis is the utility of filtering against local exome data other than dbSNP and 1,000 Genomes. Many indels are not annotated in existing databases despite being relatively common. Thus we observed large numbers of indels compared to SNPs persisting through the first 3 steps of our analysis (Fig. 1b). Most indels were removed by genotype quality filtering against local exomes. Also, indel calling is less reliable than SNP calling; filtering indel variants against unrelated local exomes reduces the rate of local errors and artifacts.

Infant V003's gene mutations have not been previously reported, but, like most *ATM* mutations, produce early stop codons, predicted to lead to nonsense mediated decay of mRNA [30]. One of infant V004's mutations has been reported [31]; the other is predicted to be damaging by SNAP and Polyphen-2 algorithms [32, 33].

Epidemiology and Public Health Considerations.

We have demonstrated the ability to diagnose AT within the context of a newborn screening program and have provided documentation for the first time of sufficiently low T cell counts in two infants with AT in the first months of life that live vaccines should be avoided. The incidence of AT has been reported to be between 1:40,000 and 1:100,000 births [34] with high rates in some populations due to founder mutations [35]. A more recent estimate was lower, 1:300,000 births [36]. The two infants reported here are the only newborns found to have AT in the first 18 months of California's SCID newborn screening program, in which over 740,000 infants were screened. With 7 of 13 AT cases (54 %) from archived samples having abnormal TREC screening, our data in the highly diverse California population suggest an incidence figure around 1:200,000. California has the largest number of annual births of any state in the U.S., and population-based newborn screening with TRECs will provide a prospective, unbiased method to establish the incidence of AT.

The detection of AT by newborn screening illuminates challenges surrounding the inclusion of new tests into a public health screening program. Currently there is no cure for AT; the neurological deterioration is progressive and irreversible such that patients usually become wheelchair dependent by their teenage years, and lifespan is decreased. In addition, patients with AT have increased risk for malignancies, attributed to compromise of DNA repair mechanisms. Consistent with the reported 30–40 % lifetime risk

and 10–15 % childhood or early adulthood risk of lymphoid malignancy, 3 of our 13 retrospective patients had lymphomas and one had leukemia [37, 38]. *ATM* heterozygous mutation carriers also have an increased risk of breast cancer as well as other epithelial malignancies [39, 40]. Thus, although screening programs are designed to identify newborns with diseases for which there are treatments, the early diagnosis of AT as a secondary target of TREC screening for SCID can provide important information for family and genetic counseling. Patients should avoid undue irradiation and should be monitored for malignancy as well as protected from infection, while carriers of *ATM* mutations should be made aware of their own increased risk of cancer, and any anti-cancer therapy they require should be tailored in light of their increased sensitivity to radiation [41]. A diagnosis of AT by newborn screening can also help parents plan for the care of the affected child and obtain genetic counseling when considering future pregnancies.

Furthermore, while there is currently no effective treatment for AT, multiple lines of research are aiming towards that goal. Antioxidant therapies [42–44], and more recently HDAC4 inhibition [45], have shown promise in mouse models; and in vitro experiments with human cells suggest that for some patients *ATM* function could be restored using mutation-targeted therapy [46]. For any of the potential therapies under development, early intervention made possible by identification through newborn screening would be most likely to show benefit by allowing therapy to take effect before extensive degeneration of relevant tissues has occurred.

Conclusion

This study demonstrates that T lymphocytopenia revealed by newborns TREC screening can be an identifying feature of AT. We also show the utility of exome sequencing to arrive at a gene diagnosis for infants with variant SCID or CID. Although there is no current cure for the progressive neurological impairment of AT, early detection provides information to improve patient management and offer family genetic counseling. Whole exome sequencing coupled with newborn screening in an ethnically diverse, large population will reveal unbiased data about rare diseases associated with T lymphocytopenia.

Acknowledgments The authors thank Ajit Bhandal, Constantino Aznar, Robert Currier, Leslie Gaffney and the PerkinElmer SCID Screening team at the California Genetic Disease Laboratory and Stanley Naides at Quest Nichols Laboratory; we also thank Catherine Chu and Richard Lau for technical assistance with whole exome sequencing and Stephen Yee at UC Berkeley and Kunal Kundu and Sushma Motarmari at TCS for helping with scripts for annotating and filtering of variant files.

Dr. Kwan received funding from Hospital Corporation of America (HCA) International Foundation Travelling Scholarship. Dr. Brenner received support from Tata Consultancy Services (TCS) and unrestricted research funds from Celera. Dr. Puck received support from National Institutes of Health (NIH) National Institute of Allergy and Infectious Diseases (NIAID) RO1 078248 and National Center of Research Resources (NCRR) UCSF-Clinical & Translational Science Institute (CTSI) UL1 RR024131 and from the UCSF Jeffrey Modell Diagnostic Center for Primary Immunodeficiencies. The California SCID Newborn Screening program acknowledges support from the Jeffrey Modell Foundation; the NIH Eunice Kennedy Shriver National Institute of Child Health and Human Development contract HHSN267200603430C to New York State; and PerkinElmer, Inc.

References

- Buckley RH. The multiple causes of human SCID. *J Clin Inv.* 2004;114(10):1409–11. Epub 2004/11/17.
- Puck JM. Population-based newborn screening for severe combined immunodeficiency: steps toward implementation. *J Allergy Clin Immunol.* 2007;120(4):760–8. Epub 2007/10/13.
- Myers LA, Patel DD, Puck JM, Buckley RH. Hematopoietic stem cell transplantation for severe combined immunodeficiency in the neonatal period leads to superior thymic output and improved survival. *Blood.* 2002;99(3):872–8. Epub 2002/01/25.
- Lindgren ML, Kobrynski L, Rasmussen SA, Moore CA, Grosse SD, Vanderford ML, et al. Applying public health strategies to primary immunodeficiency diseases: a potential approach to genetic disorders. *MMWR Recomm Rep: Morb Mortal Wkly Rep Recomm Rep/Centers Dis Control.* 2004;53(RR-1):1–29. Epub 2004/01/16.
- Chan K, Puck JM. Development of population-based newborn screening for severe combined immunodeficiency. *J Allergy Clin Immunol.* 2005;115(2):391–8. Epub 2005/02/08.
- Douek DC, Vescio RA, Betts MR, Brenchley JM, Hill BJ, Zhang L, et al. Assessment of thymic output in adults after haematopoietic stem-cell transplantation and prediction of T-cell reconstitution. *Lancet.* 2000;355(9218):1875–81. Epub 2000/06/24.
- Routes JM, Grossman WJ, Verbsky J, Laessig RH, Hoffman GL, Brokopp CD, et al. Statewide newborn screening for severe T-cell lymphopenia. *JAMA: J Am Med Assoc.* 2009;302(22):2465–70. Epub 2009/12/10.
- Buckley RH. The long quest for neonatal screening for severe combined immunodeficiency. *J Allergy Clin Immunol.* 2012;129(3):597–604. quiz 5–6. Epub 2012/01/27.
- Puck JM. Laboratory technology for population-based screening for severe combined immunodeficiency in neonates: the winner is T-cell receptor excision circles. *J Allergy Clin Immunol.* 2012;129(3):607–16. Epub 2012/01/31.
- Verbsky J, Thakar M, Routes J. The Wisconsin approach to newborn screening for severe combined immunodeficiency. *J Allergy Clin Immunol.* 2012;129(3):622–7. Epub 2012/01/17.
- Bolze A, Byun M, McDonald D, Morgan NV, Abhyankar A, Premkumar L, et al. Whole-exome-sequencing-based discovery of human FADD deficiency. *Am J Hum Genet.* 2010;87(6):873–81. Epub 2010/11/27.
- Byun M, Abhyankar A, Lelarge V, Plancoulaine S, Palanduz A, Telhan L, et al. Whole-exome sequencing-based discovery of STIM1 deficiency in a child with fatal classic Kaposi sarcoma. *J Exp Med.* 2010;207(11):2307–12. Epub 2010/09/30.
- Conley ME, Dobbs AK, Quintana AM, Bosompem A, Wang YD, Coustan-Smith E, et al. Agammaglobulinemia and absent B lineage cells in a patient lacking the p85alpha subunit of PI3K. *J Exp Med.* 2012;209(3):463–70. Epub 2012/02/22.
- Borte S, von Döbeln U, Fasth A, Wang N, Janzi M, Winiarski J, et al. Neonatal screening for severe primary immunodeficiency diseases using high-throughput triplex real-time PCR. *Blood.* 2012;119(11):2552–5. Epub 2011/12/02.
- Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics.* 2009;25(14):1754–60. Epub 2009/05/20.
- McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The genome analysis toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* 2010;20(9):1297–303. Epub 2010/07/21.
- DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet.* 2011;43(5):491–8. Epub 2011/04/12.
- A map of human genome variation from population-scale sequencing. *Nature.* 2010;467(7319):1061–73. Epub 2010/10/29.
- Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL, Gibbs RA, et al. A second generation human haplotype map of over 3.1 million SNPs. *Nature.* 2007;449(7164):851–61. Epub 2007/10/19.
- Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call format and VCFtools. *Bioinformatics.* 2011;27(15):2156–8. Epub 2011/06/10.
- Fiume M, Williams V, Brook A, Brudno M. Savant: genome browser for high-throughput sequencing data. *Bioinformatics.* 2010;26(16):1938–44. Epub 2010/06/22.
- Thorstenson YR, Roxas A, Kroiss R, Jenkins MA, Yu KM, Bachrich T, et al. Contributions of ATM mutations to familial breast and ovarian cancer. *Cancer Res.* 2003;63(12):3325–33. Epub 2003/06/18.
- Sarzotti M, Patel DD, Li X, Ozaki DA, Cao S, Langdon S, et al. T cell repertoire development in humans with SCID after nonablative allogeneic marrow transplantation. *J Immunol.* 2003;170(5):2711–8. Epub 2003/02/21.
- Kellermayer R, Hsu AP, Stankovics J, Balogh P, Hadzsiev K, Vojcek A, et al. A novel IL2RG mutation associated with maternal T lymphocyte engraftment in a patient with severe combined immunodeficiency. *J Hum Genet.* 2006;51(5):495–7. Epub 2006/04/08.
- Al-Herz W, Bousfiha A, Casanova JL, Chapel H, Conley ME, Cunningham-Rundles C, et al. Primary immunodeficiency diseases: an update on the classification from the international union of immunological societies expert committee for primary immunodeficiency. *Front Immunol.* 2011;2:54. Epub 2011/01/01.
- Blohm ME, Vesterling-Horner D, Calaminus G, Gobel U. Alpha 1-fetoprotein (AFP) reference values in infants up to 2 years of age. *Pediatr Hematol Oncol.* 1998;15(2):135–42. Epub 1998/05/21.
- Stray-Pedersen A, Borresen-Dale AL, Paus E, Lindman CR, Burgers T, Abrahamsen TG. Alpha fetoprotein is increasing with age in ataxia-telangiectasia. *Eur J Paediatr Neurol: EJPN: Off J Eur Paediatr Neurol Soc.* 2007;11(6):375–80. Epub 2007/06/02.
- Tavani F, Zimmerman RA, Berry GT, Sullivan K, Gatti R, Bingham P. Ataxia-telangiectasia: the pattern of cerebellar atrophy on MRI. *Neuroradiology.* 2003;45(5):315–9. Epub 2003/05/13.
- Dvorak CC, Cowan MJ. Radiosensitive severe combined immunodeficiency disease. *Immunol Allergy Clin North Am.* 2010;30(1):125–42. Epub 2010/02/02.
- Boder E, Sedgwick RP. Ataxia-telangiectasia. (Clinical and immunological aspects). *Psychiatr Neurol Med Psychol Beih.* 1970;13–14:8–16. Epub 1970/01/01.
- Sandoval N, Platzer M, Rosenthal A, Dork T, Bendix R, Skawran B, et al. Characterization of ATM gene mutations in 66 ataxia telangiectasia families. *Hum Mol Genet.* 1999;8(1):69–79. Epub 1999/01/15.
- Bromberg Y, Rost B. SNAP: predict effect of non-synonymous polymorphisms on function. *Nucleic Acids Res.* 2007;35(11):3823–35. Epub 2007/05/29.

33. Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al. A method and server for predicting damaging missense mutations. *Nat Methods*. 2010;7(4):248–9. Epub 2010/04/01.
34. Swift M, Morrell D, Cromartie E, Chamberlin AR, Skolnick MH, Bishop DT. The incidence and gene frequency of ataxia-telangiectasia in the United States. *Am J Hum Genet*. 1986;39(5):573–83. Epub 1986/11/01.
35. Woods CG, Bunday SE, Taylor AM. Unusual features in the inheritance of ataxia telangiectasia. *Hum Genet*. 1990;84(6):555–62. Epub 1990/05/01.
36. Verhagen MM, Last JI, Hogervorst FB, Smeets DF, Roeleveld N, Verheijen F, et al. Presence of ATM protein and residual kinase activity correlates with the phenotype in ataxia-telangiectasia: a genotype-phenotype study. *Hum Mutat*. 2012;33(3):561–71. Epub 2012/01/04.
37. Lavin MF, Shiloh Y. Ataxia Telangiectasia. In: Ochs HD, Smith CIE, Puck JM, editors. *Primary immunodeficiency diseases: A molecular and genetic approach*. 2nd ed. New York: Oxford University Press; 2007. p. 402–26.
38. Gummy-Pause F, Wacker P, Sappino AP. ATM gene and lymphoid malignancies. *Leuk: Off J Leuk Soc Am Leuk Res Fund UK*. 2004;18(2):238–42. Epub 2003/11/25.
39. Swift M, Morrell D, Massey RB, Chase CL. Incidence of cancer in 161 families affected by ataxia-telangiectasia. *N Engl J Med*. 1991;325(26):1831–6. Epub 1991/12/26.
40. Renwick A, Thompson D, Seal S, Kelly P, Chagtai T, Ahmed M, et al. ATM mutations that cause ataxia-telangiectasia are breast cancer susceptibility alleles. *Nat Genet*. 2006;38(8):873–5. Epub 2006/07/13.
41. Byrd PJ, Srinivasan V, Last JI, Smith A, Biggs P, Carney EF, et al. Severe reaction to radiotherapy for breast cancer as the presenting feature of ataxia telangiectasia. *Br J Cancer*. 2012;106(2):262–8. Epub 2011/12/08.
42. Lavin MF, Gueven N, Bottle S, Gatti RA. Current and potential therapeutic strategies for the treatment of ataxia-telangiectasia. *Br Med Bull*. 2007;81–82:129–47. Epub 2007/06/26.
43. Reliene R, Schiestl RH. Experimental antioxidant therapy in ataxia telangiectasia. *Clin Medi Oncol*. 2008;2:431–6. Epub 2008/01/01.
44. Vos M, Esposito G, Edirisinghe JN, Vilain S, Haddad DM, Slabbaert JR, et al. Vitamin K2 is a mitochondrial electron carrier that rescues pink1 deficiency. *Science*. 2012;336(6086):1306–10. Epub 2012/05/15.
45. Li J, Chen J, Ricupero CL, Hart RP, Schwartz MS, Kusnecov A, et al. Nuclear accumulation of HDAC4 in ATM deficiency promotes neurodegeneration in ataxia telangiectasia. *Nat Med*. 2012. Epub 2012/04/03.
46. Nakamura K, Du L, Tunuguntla R, Fike F, Cavalieri S, Morio T, et al. Functional characterization and targeted correction of ATM mutations identified in Japanese patients with ataxia-telangiectasia. *Hum Mutat*. 2012;33(1):198–208. Epub 2011/10/19.